
Appendix E

Errata

Page 13, Exercise 1.5.6 (6):

Replace “vector space of functions \mathcal{V} .” with “vector space of functions \mathcal{V} which is closed under taking absolute values.”

Page 13, Exercise 1.5.6 (8) should read:

Prove that f is a regulated function on $I = [a, b]$ if and only if both of the limits

$$\lim_{x \rightarrow c+} f(x) \quad \text{and} \quad \lim_{x \rightarrow d-} f(x)$$

exist for every $c \in [a, b)$ and every $d \in (a, b]$.

Page 76, Exercise 4.3.9 (2) should read:

Let X be an uncountable set and let \mathcal{A} be the σ -algebra of all subsets of X . Prove there is no finite measure on X which assigns a positive measure to every finite non-empty set.

Page 88, Exercise 5.1.10 (1) should read: Give an example of a function $f : [a, b] \rightarrow \mathbb{R}$ such that f is integrable but f^2 is not.

Page 91, line 5 should read:

$$\dots \leq 4n^2\mu(A) + \delta^2\mu(A^c) \leq 4n^2\delta + (b-a)\delta^2.$$

Page 91, line 7 should read:

$$\|f_n - g\| \leq \sqrt{4n^2\delta + (b-a)\delta^2} < \frac{\varepsilon}{2}.$$

Page 102, line 7 should read:

$s_n = \sum_{i=0}^n \langle w, u_i \rangle u_i$. Then since the family is orthogonal, we know

Page 108, line 5 should read:

$$= \|w\|^2 - \sum_{n=0}^N |a_n|^2 + \sum_{n=0}^N (a_n - c_n) \overline{(a_n - c_n)}$$

Page 109, line -4 should read:

$$\sum_{i=0}^{\infty} |\langle w, u_i \rangle|^2 = \|w\|^2$$

Page 116, lines 13–14 should read:

with $|g(x) - p(x)| < \varepsilon/4$ for all x . So

$$\|g - p\|^2 = \frac{1}{\pi} \int (g - p)^2 d\mu \leq \frac{1}{\pi} \int \frac{\varepsilon^2}{16} d\mu = \frac{\varepsilon^2}{8}.$$

Hence, $\|f - p\| \leq \|f - g\| + \|g - p\| < \varepsilon/2 + \varepsilon/\sqrt{8} < \varepsilon$.

Page 118, the last line of exercise 6.1.10 (1) should read:

$$\sin \theta_1 \sin \theta_2 = \frac{1}{2} (\cos(\theta_1 - \theta_2) - \cos(\theta_1 + \theta_2)).$$

Page 135, the second paragraph of the proof of 7.2.2 should read:

Conversely, if f is a T -invariant measurable function then its real part, $u(x)$, and imaginary part, $v(x)$, are T -invariant. If f is not ν -almost everywhere constant, then at least one of $u(x)$ and $v(x)$ is not ν -almost everywhere constant. Suppose, without loss of generality, that it is $u(x)$. Then there is a $c \in \mathbb{R}$ such that the set $A = u^{-1}([-\infty, c])$ satisfies $\nu(A) > 0$ and $\nu(A^c) > 0$. The set A is T -invariant.

Page 138, line 4 of Corollary 7.3.2 should read:

$A \cap \{T^k(x)\}_{k=0}^{m-1}$, i.e., the number of points in A from the orbit segment

Page 169, line 2 should read:

$$|\langle v, w \rangle|^2 = \|\langle v, w \rangle w\|^2 \leq \|\langle v, w \rangle w\|^2 + \|v - \langle v, w \rangle w\|^2 = 1$$

Page 187, Lemma B.3.9 and its proof should read:

Lemma B.3.9. *Suppose A_0 and B_0 are disjoint sets in \mathcal{M}_0 and X_0 is an arbitrary subset of \mathbb{R} . If $A = A_0 \cap X_0$ and $B = B_0 \cap X_0$ then*

$$\mu^*(A \cup B) = \mu^*(A) + \mu^*(B).$$

The analogous result for a finite union of disjoint sets is also valid.

Proof. Since A_0 and B_0 are disjoint, so are A_0 and B , and therefore $A_0 \cap B = \emptyset$ and $A_0^c \cap B = B$. Hence

$$A_0 \cap (A \cup B) = (A_0 \cap X_0) \cup (A_0 \cap B \cap X_0) = A_0 \cap X_0 = A.$$

Likewise,

$$A_0^c \cap (A \cup B) = (A_0^c \cap A_0 \cap X_0) \cup (A_0^c \cap B) = B.$$

Hence, the fact that A_0 is in \mathcal{M}_0 (using $A \cup B$ for X in the definition of \mathcal{M}_0) tells us

$$\begin{aligned} \mu^*(A \cup B) &= \mu^*(A_0 \cap (A \cup B)) + \mu^*(A_0^c \cap (A \cup B)) \\ &= \mu^*(A) + \mu^*(B). \end{aligned}$$

The result for a finite collection $A_0 \cap X_0, A_1 \cap X_0, \dots, A_n \cap X_0$, where $A_i \in \mathcal{M}_0$ follows immediately by induction on n . \square

Page 188, the first displayed equation should read:

$$B_{n+1} = A_{n+1} \setminus \bigcup_{i=1}^n A_i = A_{n+1} \cap \left(\bigcup_{i=1}^n A_i \right)^c.$$

Page 197, item [C]:

Replace “sumas” with “sums”

Page 197, item [M]:

Replace “Erdodic” with “Ergodic”

April 30, 2017.