

ERRATA / ADDENDA TO ERGODIC THEORY VIA JOININGS

PART I:

Page	Line	Replace by ...
26	5	Show that the system (X, Γ) is minimal, admits no non-trivial equicontinuous factor, but the relation Q is not an equivalence relation ...
57	3	$\hat{d}(\phi, \psi) = \sum_{n=1}^{\infty} 2^{-n} \frac{1}{2} (\mu(\phi A_n \Delta \psi A_n) + \mu(\phi^{-1} A_n \Delta \psi^{-1} A_n))$.
58	3	$\alpha(n, y) = \begin{cases} \alpha(T^{n-1}y) \cdots \alpha(Ty) \alpha(y) & \text{for } n \geq 1 \\ \text{id} & \text{for } n = 0 \\ \alpha(T^n y)^{-1} \cdots \alpha(T^{-1}y)^{-1} & \text{for } n < 0. \end{cases}$
65	-13	Conversely, assume that π admits no non-zero invariant vectors and suppose that $\mathbf{m}(f) \neq 0$ for some function $0 \neq f \in B_\pi$. By Theorem 1.51.2(c) we can assume that f is in $AP(\Gamma)$ and, therefore, that it arises as $f(\gamma) = \langle \pi(\gamma)x, x \rangle$ for a finite dimensional irreducible sub-representation of π . This however contradicts the previous lemma and we conclude that $\mathbf{m}(f) = 0$ for every function $f \in B_\pi$.
68	-5	Denote weak-closure $\pi(X) = Y \subset L^2(\mu)$, and $\nu = \pi_*(\mu)$, then, since the action of Γ on the compact space Y is WAP and topologically transitive, we can apply Lemma 1.50 to deduce that (Y, ν, Γ) is a nontrivial isometric factor.
81	11	This is wrong; condition 2 does not imply mixing.
149	-6	... we must have $\hat{\pi}^{k+1}(\hat{\lambda}) = \nu^{k+1}$. The assumption that Γ is abelian implies that $\hat{\lambda}$ is Γ -invariant. Set $\lambda = \sigma^{k+1}(\hat{\lambda})$; ...
218	-10	Here and in the sequel replace $C(X)$ by $\text{Aut}(\mathbf{X})$.
223	6	Here and in the sequel replace n by r .
223	17	Here and in the sequel replace 3 by r .
292	1	A subset $A \subset X$ is called <i>uniform</i> if $\lim_{N \rightarrow \infty} \text{ess sup}_x \left \frac{1}{N} \sum_0^{N-1} \mathbf{1}_A(T^i x) - \mu(A) \right = 0.$ A partition α is uniform if every set in $\cup_{n=0}^{\infty} \alpha_{-n}^n$ is uniform.

PART II

Replacement for Chapter 15, Section 2.

2. KAKUTANI, ROHLIN AND K-R TOWERS

Let \mathbf{X} be a dynamical system and $B \subset X$; an array $\mathbf{c} = \{B, TB, \dots, T^{N-1}B\}$ with $T^j B$, $0 \leq j < N$ pairwise disjoint is called a *Rohlin tower* or a *column over B of height N* . The set B is called the *base* of the tower and $T^{N-1}B$ is its *roof*. Let $|\mathbf{c}| = \cup_{j=0}^{N-1} T^j B$, the *carrier* of the tower \mathbf{t} . A collection (finite or countable) \mathbf{t} of disjoint columns \mathbf{c}_k (with bases B_k and heights N_k) is called a *tower* and we let $|\mathbf{t}| = \cup_{k=0}^{\infty} |\mathbf{c}_k|$. The union of the bases $B = \cup_k B_k$ is the *base* of \mathbf{t} , and the union of the roofs is the *roof* of \mathbf{t} . We sometimes write (a bit imprecisely) $\mathbf{t} = \{\mathbf{c}_k : k = 1, 2, \dots\}$. The sets $\{T^i x : 0 \leq i < N_k(x)\}$ for $x \in B$ are called the *fibers* of \mathbf{t} .

Given a tower \mathbf{t} with columns $\{\mathbf{c}_k : k = 1, 2, \dots\}$, base $B = \cup_k B_k$, and a finite (or countable) partition $\alpha = \{A_1, \dots, A_t\}$, we define an equivalence relation on B as follows: $x \sim y$ iff x and y are in the same B_k and for every $0 \leq j < N_k$, $T^j x$ and $T^j y$ are in the same element of α ; i.e. x and y have the same (α, N_k) -name. We now consider each equivalence class $B_{k,\mathbf{a}}$, with \mathbf{a} a name in $\alpha_0^{N_k-1}$, as a basis of a column $\mathbf{c}_{k,\mathbf{a}} = \{B_{k,\mathbf{a}}, TB_{k,\mathbf{a}}, \dots, T^{N_k-1}B_{k,\mathbf{a}}\}$ and say that the resulting tower $\mathbf{t}_\alpha = \{\mathbf{c}_{k,\mathbf{a}} : \mathbf{a} \in \alpha_0^{N_k-1}, k = 1, \dots\}$ is the tower \mathbf{t} *refined* according to α . When $\alpha = \{C, X \setminus C\}$, C a subset of X , we write $\mathbf{t}_\alpha = \mathbf{t}_C$ and say that \mathbf{t} is refined according to C .

A subset B of X is called a *sweeping set* if $\cup_{n \geq 0} T^n B = X$. For an ergodic system every set of positive measure is sweeping. Given any set B of positive measure, define the *return time function* $r_B : B \rightarrow \mathbb{N} \cup \{\infty\}$ by

$$r_B(x) = \min\{n \geq 1 : T^n x \in B\},$$

when this minimum is finite and $r_B(x) = \infty$ otherwise. Let $B_k = \{x \in B : r_B(x) = k\}$ and note that by Poincaré's recurrence theorem B_∞ is a null set. We let \mathbf{c}_k be the column $\{B_k, TB_k, \dots, T^{k-1}B_k\}$ and we call the tower $\mathbf{t} = \mathbf{t}(B) = \{\mathbf{c}_k : k = 1, 2, \dots\}$, the *Kakutani tower over B* . Thus the Kakutani tower \mathbf{t} is a partition of the set $|\mathbf{t}|$. For a sweeping set B , $|\mathbf{t}| = X$ and the Kakutani tower over B is then a partition of the whole space. If the Kakutani tower over B has finitely many columns (i.e. the function r_B is bounded) we say that B has a *finite height* and we call the Kakutani tower over B a *K-R tower*. The number $\max r_B$ is called the *height* of B or the *height* of the K-R tower.

Note that for an ergodic \mathbb{Z} -system \mathbf{X} , either the space X consists of a finite set of points on which μ is equidistributed, or the measure μ is atom-less. In the first case the system is called *periodic*, and it is called *non-periodic* in the latter. The next we present the famous "Rohlin lemma". We shall prove it in the ergodic case but in fact it holds for every aperiodic \mathbb{Z} -system (see Halmos' book [113]).

Theorem 2.1 (Rohlin's lemma). *Let \mathbf{X} be a non-periodic ergodic system, N a positive integer and $\epsilon > 0$, then there exists a subset B such that the sets $B, TB, \dots, T^{N-1}B$ are pairwise disjoint and $\mu(\cup_{j=0}^{N-1} T^j B) > 1 - \epsilon$.*

Proof. Let $C \subset X$ be a set with measure $0 < \mu(C) < \epsilon/N$. Consider the Kakutani tower $\mathbf{t}(C)$ over C . For every $k \geq N$ divide the column $\mathbf{c}_k = \{T^i C_k : 0 \leq i < k\}$, starting from its base $C_k = \{x \in C : r_C(x) = k\}$, into blocks of size N . Mark

the first level of each of these blocks as belonging to B . Taking the union of these marked levels over the columns $\mathbf{c}_k, k = N, N + 1, \dots$, gives us a set B with $r_B \geq N$. Clearly $\mathbf{q} = \{B, TB, \dots, T^{N-1}B\}$ is a Rohlin tower; i.e. the sets $T^j B$ for $0 \leq j < N$ are disjoint. Now $|\mathbf{t}| \setminus |\mathbf{q}| = X \setminus |\mathbf{q}|$ is composed of the first N columns \mathbf{c}_k of height $k < N$ and some top levels from the other columns, with a contribution of at most $N - 1$ levels from each. A simple calculation shows therefore that

$$\mu(X \setminus |\mathbf{q}|) < N\mu(C) < \epsilon.$$

□

Theorem 2.2. *Let \mathbf{X} be a non-periodic ergodic system.*

- (1) *For any positive integer N there exists a set C of finite height such that the K-R tower $\mathbf{t}(C)$ satisfies $\text{range } r_C \subset \{N, N + 1\}$.*
- (2) *Given a K-R tower with base C and height N , for any sufficiently large n there is a bounded K-R tower with base D contained in C whose column heights are all at least n and at most $n + 4N$.*

Proof. 1. Let $n > 10N^2$ and use Rohlin's lemma to construct a Rohlin tower $\mathbf{q} = \{B, TB, \dots, T^{n-1}B\}$. Thus the return time function $r_B(x)$ is greater than $10 \cdot N^2$ on B . Let

$$B_k = \{x \in B : r_B(x) = k\}.$$

When B_k is non-empty one can therefore write k as a positive combination of N and $N + 1$, say

$$k = Nu_k + (N + 1)v_k.$$

Now consider the Kakutani tower over B and divide the column $\mathbf{c}_k = \{T^i B_k : 0 \leq i < k\}$ into u_k blocks of size N and v_k blocks of size $N + 1$. The set C is now defined as the union over the various columns, of the first levels of these blocks. Clearly the function r_C takes only two values, either N or $N + 1$ as required.

2. Start with a Rohlin tower $\mathbf{q} = \{B, TB, \dots, T^{M-1}B\}$ with $M = 10(n + 2N)^2$, and look at the unbounded (in general) Kakutani tower \mathbf{t} over B . As can be seen by the proof of Rohlin's lemma we can choose $B \subset C$. Refine this tower according to C and call the refined tower \mathbf{t}_C . For each $m \geq 10(n + 2N)^2$, the column \mathbf{q}_m over $B_m = \{x \in B : r_B = m\}$ in \mathbf{t} , is split in \mathbf{t}_C into a finite number of columns so that each level is either a subset of C or of C^c .

As in the first part of the proof we partition each column of the tower \mathbf{t}_C into blocks of sizes $n + 2N$ and $n + 2N + 1$. Since we want the base D of the K-R tower we construct to belong to C we move the base level of each block (up or down) to the nearest level that belongs to C . Since the height of C is N we do not move these levels more than $N - 1$ steps. The new blocks, with bases in C , are of size between n and $n + 4N$, and we let D be the union of these bases. □