

PIECEWISE DISTANCE PRESERVING MAPS

A. PETRUNIN AND A. YASHINSKI

ABSTRACT. This is part of a geometry course held during the Fall 2011 MASS Program at Penn State (www.math.psu.edu/mass/). The online version of these lectures also contains video illustrations, hints, and solutions for most of the exercises, and a minimalistic section covering preliminaries.

In the lectures, piecewise distance preserving maps are discussed that act from a 2-dimensional polyhedral space into the plane. Roughly speaking, a polyhedral space is a space that is glued together out of triangles, for example the surface of a polyhedron. If one imagines such a polyhedral space as a paper model, then a piecewise distance preserving map into the plane is essentially a way to fold the model so that it lays flat on a table.

Only the 2-dimensional case is considered, to keep things easy to visualize. However, most of the results admit generalizations to higher dimensions. These results are discussed in the Final Remarks, where proper credit and references are given.

INTRODUCTION

All the necessary background material is discussed in the first three chapters of the book “Metric Geometry” by Burago–Burago–Ivanov [10].

We list the less standard conventions that we use in the lectures.

- ◊ The distance between two points x and y in a metric space X will be denoted by

$$|x - y| \quad \text{or} \quad |x - y|_X;$$

the latter notation is used if we need to emphasize that the distance is measured in the metric space X .

- ◊ A metric space X is called a *length space* if for any two points $x, y \in X$ and any $\varepsilon > 0$, there is a curve α from x to y such that

$$\text{length } \alpha < |x - y|_X + \varepsilon.$$

- ◊ Let $f: X \rightarrow Y$ be a continuous map between two metric spaces. Then:
 - f is said to be *distance preserving* if

$$|f(x) - f(y)|_Y = |x - y|_X$$

for any pair of points $x, y \in X$;

- f is *distance nonexpanding* if

$$|f(x) - f(y)|_Y \leq |x - y|_X$$

for any pair of points $x, y \in X$;

- f is *length-preserving* if for any curve α in X , we have

$$\text{length } \alpha = \text{length}(f \circ \alpha).$$

- ◊ A length space P is called a *polyhedral space* if it admits a finite triangulation such that each simplex in P is isometric to a simplex in Euclidean space.

2010 *Mathematics Subject Classification*. Primary 52B70.

Key words and phrases. Polyhedral space, piecewise distance preserving maps, piecewise linear maps.

§1. ZALGALLER'S FOLDING THEOREM

Let P be a polyhedral space. A map $f: P \rightarrow \mathbb{R}^n$ is said to be *piecewise distance preserving* if there is a triangulation of P such that for any simplex Δ in the triangulation, the restriction $f|_{\Delta}$ is distance preserving.

1.1. Exercise. *Show that any piecewise distance preserving map is continuous, length-preserving, and distance nonexpanding.*

The following statement might look obvious, but try to prove it rigorously.

1.2. Exercise. *Suppose that an m -dimensional polyhedral space admits a piecewise distance preserving map into \mathbb{R}^n . Show that $n \geq m$.*

In fact, the converse of the statement in the exercise is true. In other words, dimension is the only obstruction to the existence of a piecewise distance preserving map into Euclidean space. The following theorem asserts this for the case where $m = 2$.

1.3. Zalgaller's theorem. *Any 2-dimensional polyhedral space admits a piecewise distance preserving map into the Euclidean plane.*

Imagine that you have a paper model of a 2-dimensional polyhedral space P in your hands, and you fold this model so that it lays flat on a table.¹ This is an intuitive way to think of a piecewise distance preserving map $f: P \rightarrow \mathbb{R}^2$. To make it closer to the actual definition, one has to imagine that the layers of paper can go through each other. Zalgaller's theorem says that such a "folding" is always possible; see also Exercise 5.3.

The following exercise shows that in this process, new folds may need to be introduced across the triangles of the given triangulation of P .

1.4. Exercise. *Let Δ be a nondegenerate 3-dimensional simplex in \mathbb{R}^3 , and let $\partial\Delta$ be its boundary equipped with the induced length metric. It is a polyhedral space glued from 4 triangles — the faces of Δ .*

Show that $\partial\Delta$ does not admit a map to \mathbb{R}^2 which is distance preserving on each of the 4 faces of Δ .

Describe explicitly a piecewise distance preserving map

$$f: \partial\Delta \rightarrow \mathbb{R}^2$$

that is distance preserving on 2 out of the 4 faces of Δ . (You will need to subdivide the other 2 faces into smaller triangles.)

Below we give two similar proofs of Zalgaller's theorem: the first with cheating and the second without. In the first proof, we use the following claim without proof.

❶ *Any 2-dimensional polyhedral space admits an acute triangulation, that is, a triangulation such that all of its triangles are acute.*

1.5. Exercise. *Show that any triangle admits an acute triangulation.*

Notice that there is more to proving ❶ than simply subdividing each triangle into acute triangles. One must be careful that the subdivisions of two triangles that share an edge are compatible on that common edge. Claim ❶ was essentially proved in [23], see also [7].

Proof using ❶. Fix an acute triangulation \mathcal{T}_0 of P provided by ❶. Mark all its vertices in white and denote them by $\{w_1, \dots, w_k\}$.

¹We recommend that you create such a paper model, say the surface of a cube, and then try to fold it on the table.

For each w_i , consider its *Voronoi domain* V_i , which is the subset

$$V_i = \{ x \in P \mid |x - w_i| \leq |x - w_j| \text{ for all } j \}.$$


Denote by $S(w_i)$ the *star* of w_i , which is the union of all simplexes of the triangulation \mathcal{T}_0 that contain w_i .

Since an acute triangle contains its own circumcenter, the Voronoi domain of a vertex of a triangle cannot cross the opposite edge, which implies that

$$V_i \subset S(w_i)$$

for all i . In particular, for any point $x \in V_i$ there is a unique geodesic $[w_i, x]$, which is a line segment in a single triangle or an edge of \mathcal{T}_0 .

Note that in each triangle of \mathcal{T}_0 , we have one point where three Voronoi domains meet and three points on the sides of the triangle where pairs of Voronoi domains meet. Let us bisect each edge of \mathcal{T}_0 and subdivide each triangle into 6 triangles as it is done on the picture above (solid lines only).



The Voronoi domains within one triangle.

In this way we obtain a new triangulation \mathcal{T}_1 . We mark the new vertices of \mathcal{T}_1 in black.

Note that

- 1) each V_i is a union of all triangles and edges of \mathcal{T}_1 that have w_i as a vertex;
- 2) each triangle in \mathcal{T}_1 has one white and two black vertices;
- 3) the triangles in \mathcal{T}_1 come in pairs of congruent triangles, they share two black vertices and have different white vertices.

Given a point $x \in P$, set

$$\varrho(x) = \min_i \{|w_i - x|\}.$$

Notice that if $x \in V_i$, then $\varrho(x) = |w_i - x|$. Given $x \in V_i$, we denote by $\vartheta_i(x)$ the smallest angle between $[w_i, x]$ and any edge of \mathcal{T}_1 coming from w_i .

By Property 1, if $x \in V_i \cap V_j$, then $\vartheta_i(x) = \vartheta_j(x)$. In other words, the function ϑ given by

$$\vartheta(x) = \vartheta_i(x), \quad x \in V_i,$$

is well defined on the set $P \setminus \{w_1, \dots, w_n\}$. Moreover, ϑ is a continuous function.

We now describe the map $f: P \rightarrow \mathbb{R}^2$ using polar coordinates on \mathbb{R}^2 . We define $f(w_i) = 0$ and $f(x) = (\varrho(x), \vartheta(x))$ if $x \in P \setminus \{w_1, \dots, w_n\}$.

Subdividing each triangle by the angle bisector at the white vertex (see the dashed lines on the picture above) gives a new triangulation \mathcal{T}_2 , which satisfies the conditions of the theorem for the constructed map f . \square

Now we modify the above proof so as to avoid using claim **1**. The only property we really need for the triangulation is that $V_i \subset S(w_i)$. On the one hand, this inclusion does not hold for a general triangulation. A simple example of this is obtained by gluing an equilateral triangle to the longer side of an obtuse triangle. On the other hand, by increasing the number of Voronoi domains, we can arrange that this inclusion be true without making the triangulation acute.

Proof without using 1. Fix a triangulation \mathcal{T}_0 of P that is not necessarily acute. We shall construct new triangulations \mathcal{T}_1 and \mathcal{T}_2 of P by subdividing the triangles of \mathcal{T}_0 , and we shall define a map $f: P \rightarrow \mathbb{R}^2$ which is distance preserving on each triangle of \mathcal{T}_2 . The vertices of \mathcal{T}_1 will be colored either white or black in such a way that each triangle of \mathcal{T}_1 will have two black vertices and one white vertex.

First, we describe the set of white vertices.

Fix a small number $\varepsilon > 0$. We mark in white all of the vertices of \mathcal{T}_0 , as well as the points on the edges of triangles of \mathcal{T}_0 with the property that the distance to the closest vertex of the edge is an integral multiple of ε . In this way we mark a finite number of points white. Label the white points by w_1, \dots, w_k .

As in the previous proof, let V_i be the Voronoi domain of w_i , so that

$$V_i = \{ x \in P \mid |x - w_i| \leq |x - w_j| \text{ for any } j \}.$$

We also let $S(w_i)$ be the star of w_i in \mathcal{T}_0 , which is the union of all simplexes of \mathcal{T}_0 that contain w_i . By the following exercise, we may assume that $V_i \subset S(w_i)$ for each i , by taking a suitably small ε .

1.6. Exercise. *Let ℓ be the minimal length of the edges in the triangulation, and let α be the smallest angle of all triangles in \mathcal{T}_0 . Show that if $\varepsilon < \frac{\ell \cdot \alpha}{100}$, then $V_i \subset S(w_i)$ for each i .*

Fix a triangle Δ of \mathcal{T}_0 . Note that for any $w_i \in \Delta$, the intersection $V_i \cap \Delta$ is a convex polygon. This follows because for any two points $w_i, w_j \in \Delta$, the inequality

$$|x - w_i| \leq |x - w_j|$$

describes the set of all points $x \in \Delta$ that lie on one side of the bisecting perpendicular to w_i and w_j . Let us color the vertices of all of the polygons $V_i \cap \Delta$ in black if they are not already white.

If \mathcal{T}_0 contains an edge E that is not a side of a triangle, then we color the midpoint of E in black.

Now we describe the triangulation \mathcal{T}_1 . The vertices of \mathcal{T}_1 are the black and white vertices. A white point w_i is connected by an edge to each black point in V_i . A pair of black vertices b and b' are connected by an edge if it forms a side of some $V_i \cap \Delta$ (for some Δ and $w_i \in \Delta$). In this case w_i, b , and b' also form a triangle of \mathcal{T}_1 . Notice that each black-black edge is a side of two congruent triangles of \mathcal{T}_1 with different white vertices.

The remaining part of the proof is the same as before. We define

$$\varrho(x) = \min_i \{ |w_i - x|_P \}$$

and let $\vartheta(x)$ for $x \in V_i$ be the smallest angle between $[w_i, x]$ and any edges in \mathcal{T}_1 coming from w_i . Then we define the map $f: P \rightarrow \mathbb{R}^2$ so that $f(w_i) = 0$ for each i and $f(x) = (\varrho(x), \vartheta(x))$ in the polar coordinates.


Further subdividing each triangle of \mathcal{T}_1 in two along the angle bisector from the white vertex produces a new triangulation \mathcal{T}_2 . It is straightforward to see that the constructed map f is distance preserving on each triangle of \mathcal{T}_2 . □

Use Zalgaller's theorem to show the following.

1.7. Advanced exercise. *Any 2-dimensional polyhedral space is isometric to the underlying set of a simplicial complex in \mathbb{R}^n , equipped with its induced length metric.*

We end this section with an entertaining exercise.

1.8. Exercise. *Let \mathcal{T} be a triangulation of a convex polygon Q in \mathbb{R}^2 such that each triangle is colored either black or white. Show that the following two conditions are equivalent.*



A triangle Δ of \mathcal{T}_0 with marked white points and the intersections of their Voronoi domains with Δ .

- a) *There is a piecewise distance preserving map $Q \rightarrow \mathbb{R}^2$ for this triangulation that preserves the orientation of each white triangle and reverses the orientation of each black triangle.*
- b) *The sum of black angles around any vertex of \mathcal{T} that lies in the interior of Q is either 0 , or π , or $2 \cdot \pi$.*

§2. BREHM'S EXTENSION THEOREM

2.1. Brehm's extension theorem. *Let a_1, \dots, a_n and b_1, \dots, b_n be two collections of points in \mathbb{R}^2 such that*

$$|a_i - a_j| \geq |b_i - b_j|$$

for all i and j , and let A be a convex polygon that contains a_1, \dots, a_n . Then there is a piecewise distance preserving map $f: A \rightarrow \mathbb{R}^2$ such that $f(a_i) = b_i$ for all i .

In other words, if $F = \{a_1, \dots, a_n\}$ is a finite subset of a convex polygon A , then any distance nonexpanding map $\varphi: F \rightarrow \mathbb{R}^2$ extends to a piecewise distance preserving map $f: A \rightarrow \mathbb{R}^2$.

Proof. The proof is by induction on n .

The base case $n = 1$ is trivial: we can take

$$f(x) = x + (b_1 - a_1),$$

which is distance preserving on all of A .

Applying the inductive hypothesis to the last $n - 1$ pairs of points, we get a piecewise distance preserving map $h: A \rightarrow \mathbb{R}^2$ such that $h(a_i) = b_i$ for all $i > 1$. We shall use h to construct the desired map $f: A \rightarrow \mathbb{R}^2$.

Consider the set

$$\Omega = \{x \in A \mid |a_1 - x| < |b_1 - h(x)|\}.$$

We may assume that $a_1 \in \Omega$; otherwise $h(a_1) = b_1$ and we can take $f = h$.

❶ *The set Ω is star-shaped with respect to a_1 . That is, if $x \in \Omega$, then the line segment $[a_1, x]$ lies in Ω .*

Indeed, if $y \in [a_1, x]$, then

$$|a_1 - y| + |y - x| = |a_1 - x|.$$

Since $x \in \Omega$, we have

$$|a_1 - x| < |b_1 - h(x)|.$$


Since h is distance nonexpanding (see Exercise 1.1), we have

$$|h(x) - h(y)| \leq |x - y|.$$

Combining the above with the triangle inequality, we see that

$$\begin{aligned} |a_1 - y| &= |a_1 - x| - |x - y| \\ &< |b_1 - h(x)| - |h(x) - h(y)| \\ &\leq |b_1 - h(y)|. \end{aligned}$$

This proves that $y \in \Omega$, which establishes Claim ❶.



Recall that $\partial_A \Omega$ denotes the boundary of Ω viewed as a subset of the space A . This may differ from $\partial_{\mathbb{R}^2} \Omega$. Note that

$$\textcircled{2} \quad |a_1 - x| = |b_1 - h(x)|$$

for any $x \in \partial_A \Omega$. To see this, consider a sequence of points in Ω that converges to x and another sequence of points in $A \setminus \Omega$ that converges to x , and then use the fact that h is continuous (Exercise 1.1).

$\textcircled{3}$ *The boundary $\partial_A \Omega$ is the union of a finite collection of line segments E_1, \dots, E_k that intersect each other only at the common endpoints. Moreover, h is distance preserving on each of these segments.*

Indeed, fix a triangulation of A so that h is distance preserving on each triangle. Note that for any point $x \in \partial_A \Omega$, this triangulation has a triangle $\Delta \ni x$ such that $\Delta \cap \Omega \neq \emptyset$. Fix such a triangle Δ . Since h is distance preserving on Δ , the restriction $h|_{\Delta}$ can be extended uniquely to an isometry $\iota: \mathbb{R}^2 \rightarrow \mathbb{R}^2$.

Set $b'_1 = \iota^{-1}(b_1)$. Note that

$$|b'_1 - x| = |b_1 - h(x)|$$

for any $x \in \Delta$, because ι is an isometry and $\iota|_{\Delta} = h|_{\Delta}$.

Observe that $a_1 \neq b'_1$. Indeed, otherwise we see that

$$|a_1 - x| = |b'_1 - x| = |b_1 - h(x)|$$

for any $x \in \Delta$, which contradicts the fact that $\Delta \cap \Omega = \emptyset$.

Denote by ℓ_{Δ} the perpendicular bisector to $[a_1, b'_1]$, which coincides with the set of all points equidistant from a_1 and b'_1 . By the definition of Ω , for any $x \in \Delta$ we see that $x \in \Omega$ if and only if x and a_1 lie on the same side from ℓ_{Δ} . Therefore, $\partial_A \Omega$ is the union of the intersections $\Delta \cap \ell_{\Delta}$ for all Δ as above. Hence $\textcircled{3}$ follows, because there are only finitely many such Δ .

For each edge E_i in $\partial_A \Omega$, consider the triangle T_i with vertex a_1 and base E_i . Condition $\textcircled{2}$ implies that there is an isometry ι_i of \mathbb{R}^2 such that $\iota_i(a_1) = b_1$ and $\iota_i(x) = h(x)$ for any $x \in E_i$.

We define $f(x) = h(x)$ for any $x \notin \Omega$ and $f(x) = \iota_i(x)$ for any $x \in T_i$. This defines f on $A \setminus \Omega$ and on all line segments from a_1 to $\partial_A \Omega$.

This completely determines f on A in the case where $\partial_A \Omega = \partial_{\mathbb{R}^2} \Omega$. If $Z = \partial_{\mathbb{R}^2} \Omega \setminus \partial_A \Omega$ is nonempty, then the points that lie on the lines between a_1 and the points in Z form a “blind zone” — this is the subset of A where f has yet to be defined.

Note that the closure of the blind zone is a union of a finite number of convex polygons Q_1, \dots, Q_m that intersect only at the common vertex a_1 . Each Q_i is bounded by a broken line in the closure of Z and two line segments from a_1 to the ends of this broken line.

So far, the distance nonexpanding map f is defined only on the two sides of each Q_i coming from a_1 , and by construction it is distance preserving on each of these two sides. From the exercise below it follows that one can extend f to each of Q_i keeping it piecewise distance preserving. \square

2.2. Exercise. Let $Q = [a_1 x_1 \dots x_k]$ be a convex polygon, and let b_1, y_1, y_k be points in the plane. Assume that

$$|b_1 - y_1| = |a_1 - x_1|, \quad |b_1 - y_k| = |a_1 - x_k|, \quad |y_1 - y_k| \leq |x_1 - x_k|.$$

Then there is a piecewise distance preserving map $f: Q \rightarrow \mathbb{R}^2$ such that $f(x_1) = y_1$, $f(x_k) = y_k$, and $f(a_1) = b_1$.

We finish this lecture with some additional exercises.

2.3. Exercise. Let a_1, \dots, a_n and b_1, \dots, b_n be two collections of points in \mathbb{R}^2 such that

$$|a_i - a_j| \geq |b_i - b_j|$$

for all i and j . Let $A = \text{Conv}\{a_1, \dots, a_n\}$ and $B = \text{Conv}\{b_1, \dots, b_n\}$ be their convex hulls. Show that

$$\text{perim } A \geq \text{perim } B,$$

where $\text{perim } A$ denotes the perimeter of A .

Is it true that

$$\text{area } A \geq \text{area } B?$$

The following exercise is a 2-dimensional case of Alexander’s theorem [4]. It has quite a simple solution, but it plays an important role in discrete geometry, check for example the paper [6] by Bezdek and Connelly.

2.4. Advanced exercise. Let a_1, \dots, a_n and b_1, \dots, b_n be two collections of points in \mathbb{R}^2 . We view \mathbb{R}^2 as a coordinate plane $\mathbb{R}^2 \times \{0\}$ in $\mathbb{R}^4 = \mathbb{R}^2 \times \mathbb{R}^2$.

Construct a collection of curves $\alpha_i: [0, 1] \rightarrow \mathbb{R}^4$ such that $\alpha_i(0) = a_i = (a_i, 0)$, $\alpha_i(1) = b_i = (b_i, 0)$ and the function $\ell_{i,j}(t) = |\alpha_i(t) - \alpha_j(t)|$ is monotone (i.e., increasing, decreasing or constant) for each i and j .

2.5. Exercise. Use Brehm’s extension theorem to prove Kirszbraun’s theorem, stated below, in the special case where Q is a finite set.

2.6. Kirszbraun’s theorem. Let $Q \subset \mathbb{R}^2$ be arbitrary subset and $f: Q \rightarrow \mathbb{R}^2$ a distance nonexpanding map. Then f admits a distance nonexpanding extension to all of \mathbb{R}^2 . In other words, there is a distance nonexpanding map $F: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that the restriction $F|_Q$ coincides with f .

§3. AKOPYAN’S APPROXIMATION THEOREM

Let P be a polyhedral space. A map $h: P \rightarrow \mathbb{R}^n$ is said to be *piecewise linear* if there is a triangulation of P such that the restriction of h to any simplex Δ is a linear map. This means that if v_0, \dots, v_k are the vertices of Δ , then for any $x \in \Delta$ we have

$$h(x) = \lambda_0 \cdot h(v_0) + \dots + \lambda_k \cdot h(v_k),$$

where $(\lambda_0, \dots, \lambda_k)$ are the barycentric coordinates of x .

3.1. Exercise. *Show that if P is a 2-dimensional polyhedral space, then any piecewise distance preserving map $f: P \rightarrow \mathbb{R}^2$ is piecewise linear.*

We shall be interested in approximating piecewise linear maps by piecewise distance preserving maps. Since all piecewise distance preserving maps are distance nonexpanding, it only makes sense to try this approximation for distance nonexpanding maps.

3.2. Akopyan’s theorem. *Assume P is a 2-dimensional polyhedral space. Then any distance nonexpanding piecewise linear map $h: P \rightarrow \mathbb{R}^2$ can be approximated by piecewise distance preserving maps.*

More precisely, for any $\varepsilon > 0$ there is a piecewise distance preserving map $f: P \rightarrow \mathbb{R}^2$ such that

$$|f(x) - h(x)| < \varepsilon$$

for all $x \in P$.

Akopyan’s theorem implies the existence of many piecewise distance preserving maps from P into \mathbb{R}^2 . In particular, it implies Zalgaller’s theorem (1.3). To see this, consider the constant map $h: P \rightarrow \mathbb{R}^2$; i.e., the map that sends the whole space P to a single point. Since h is piecewise linear, we can apply Akopyan’s theorem to produce a piecewise distancing preserving map f arbitrarily close to h .

As you will see below, the proof of Akopyan’s theorem will not involve Zalgaller’s theorem. We still consider the proof of Zalgaller’s theorem to be important because it gives a very clear geometric description of the piecewise distance preserving map. In contrast, the maps produced by Akopyan’s theorem will emerge from the recursive construction of Brehm’s theorem, which is harder to understand.


3.3. Exercise. *Show that if P is a convex polygon in \mathbb{R}^2 , then the above theorem follows from Brehm’s extension theorem (Theorem (2.1)).*

The main idea in the proof of Akopyan’s theorem is to triangulate P and use Brehm’s extension theorem on each triangle, as in the previous exercise.

Unfortunately, it is not that simple. The big technical issue that arises is that if two triangles share a common edge, we need to ensure that the maps produced by using Brehm’s theorem agree on that common edge.

To address this issue, we shall use the following *zigzag construction*. It produces a piecewise distance preserving map that is close to a given distance nonexpanding linear map defined on a line segment. For the construction, we fix a unit vector e in \mathbb{R}^2 . The choice of e does not matter, but one and the same e must be used uniformly in all zigzag constructions that follow.

Zigzag construction. Let E be a line segment and $h: E \rightarrow \mathbb{R}^2$ a distance nonexpanding linear map. Put $\ell = \text{length } E$ and $\ell' = \text{length } h(E)$. Since h is distance nonexpanding, we have $\ell' \leq \ell$.



Fix a positive integer n , and subdivide E into n equal intervals. Denote by z_0, \dots, z_n the endpoints of these intervals.

Note that the image $h(E)$ is either a line segment or a point. In the first case, let u be a unit normal vector to $h(E)$; otherwise, let $u = e$.

Given $x \in E$, set

$$s_n(x) = \min_i \{|z_i - x|\},$$

$$w_n(x) = k \cdot s_n(x) \cdot u + h(x),$$

where $k = \sqrt{1 - (\ell'/\ell)^2}$. If we subdivide E further by adding the midpoints between any two consecutive endpoints, then w_n will be distance preserving on each of the resulting subintervals. This shows that w_n is piecewise distance preserving. Moreover,

$$|w_n(x) - h(x)| \leq \frac{\ell}{2 \cdot n}$$

for any $x \in E$, because $k \leq 1$ and $s_n(x) \leq \frac{\ell}{2 \cdot n}$.

The piecewise distance preserving map w_n is the result of the n -step zigzag construction applied to h .

Given a triangulation \mathcal{T} of a polyhedral space P , let \mathcal{T}^1 denote the 1-skeleton of \mathcal{T} . Notice that \mathcal{T}^1 is a 1-dimensional polyhedral space when equipped with its induced length metric, which is different from the subspace metric it inherits from P .

The following proposition is the main technical step in the proof of Akopyan's theorem.

3.4. Proposition. *Let \mathcal{T} a triangulation of a 2-dimensional polyhedral space P , let \mathcal{T}^1 be its 1-skeleton, and let $h: \mathcal{T}^1 \rightarrow \mathbb{R}^2$ be a piecewise linear map such that*

$$|h(x) - h(y)|_{\mathbb{R}^2} \leq |x - y|_P$$

for any $x, y \in \mathcal{T}^1$. Then for any $\varepsilon > 0$, there is a piecewise distance preserving map $w: \mathcal{T}^1 \rightarrow \mathbb{R}^2$ with

$$|w(x) - w(y)|_{\mathbb{R}^2} \leq |x - y|_P$$

for any $x, y \in \mathcal{T}^1$, and

$$|w(x) - h(x)| < \varepsilon$$

for all $x \in \mathcal{T}^1$.

Proof. First we prove this proposition under the following additional assumption on h .

❶ For some fixed $\delta > 0$, we have

$$|h(x) - h(y)|_{\mathbb{R}^2} \leq (1 - \delta) \cdot |x - y|_P$$

for any $x, y \in \mathcal{T}^1$, and

$$h(v) = h(x)$$

for any vertex v of \mathcal{T}^1 and any point $x \in \mathcal{T}^1$ such that $|v - x|_P \leq \delta$.

Let \mathcal{S} denote the subdivision of \mathcal{T}^1 such that h is linear on each edge of \mathcal{S} . Subdividing \mathcal{S} further if necessary, we may assume without loss of generality that each edge of \mathcal{S} that comes from a vertex of \mathcal{T}^1 has length δ . (To perform this subdivision, we need to assume that δ in ❶ is sufficiently small.)

Denoting by ℓ the maximal length of the edges in \mathcal{T}^1 , we apply the n -step zigzag construction to each edge of \mathcal{S} . Since the maps occurring in the zigzag construction agree at the common vertices of different edges, we obtain a piecewise distance preserving map $w_n: \mathcal{T}^1 \rightarrow \mathbb{R}^2$ such that

❷
$$|w_n(x) - h(x)| \leq \frac{\ell}{2 \cdot n}$$

for all $x \in \mathcal{T}^1$.

For small $\delta > 0$ (say less than half the smallest edge length), consider the map

$$q_\delta: \mathcal{T}^1 \rightarrow \mathcal{T}^1$$

that smashes the δ -neighborhood of each vertex of \mathcal{T}^1 to that vertex and linearly stretches the remaining part of the edge, as in the figure.

Let L_δ be the optimal Lipschitz constant of q_δ , i.e., the smallest number such that

$$|q_\delta(x) - q_\delta(y)|_P \leq L_\delta \cdot |x - y|_P$$

for all $x, y \in \mathcal{T}^1$. Notice that $L_\delta \rightarrow 1$ as $\delta \rightarrow 0^+$. Then the map

$$h_\delta \stackrel{\text{def}}{=} \frac{1-\delta}{L_\delta} \cdot (h \circ q_\delta)$$

is piecewise linear and satisfies the condition stated in **1**. Moreover, we can choose δ so small that

$$|h_\delta(x) - h(x)| < \frac{\varepsilon}{2}$$

for all $x \in \mathcal{T}^1$.

By the previous part of the proof, there is a piecewise distance preserving map $w: \mathcal{T}^1 \rightarrow \mathbb{R}^2$ such that

$$|w(x) - h_\delta(x)| < \frac{\varepsilon}{2}, \quad |w(x) - w(y)|_{\mathbb{R}^2} \leq |x - y|_P$$

for all $x, y \in \mathcal{T}^1$. By the triangle inequality, we have

$$|w(x) - h(x)| < \varepsilon$$

for all $x \in \mathcal{T}^1$. □

Proof of Theorem 3.2. Fix a fine triangulation \mathcal{T} of P , one for which the diameter of each triangle is smaller than $\frac{\varepsilon}{3}$. Let \mathcal{T}^1 denotes the 1-skeleton of \mathcal{T} . By Proposition 3.4, there is a piecewise distance preserving map $w: \mathcal{T}^1 \rightarrow \mathbb{R}^2$ such that

$$|w(x) - h(x)|_{\mathbb{R}^2} < \frac{\varepsilon}{3}$$

for any $x \in \mathcal{T}^1$ and

$$|w(x) - w(y)|_{\mathbb{R}^2} \leq |x - y|_P$$

for any x and $y \in \mathcal{T}^1$.

We shall use Brehm's extension theorem (Theorem 2.1) to extend w up to a piecewise distance preserving map on P . To do this, let \mathcal{S} be a subdivision of \mathcal{T}^1 so that w is distance preserving on each edge of \mathcal{S} . Fix a triangle Δ of \mathcal{T} . Let a_1, \dots, a_n be the vertices of \mathcal{S} on the boundary of Δ , and let $b_i = w(a_i)$ for each i . By applying Brehm's theorem, we obtain a piecewise distance preserving map $f_\Delta: \Delta \rightarrow \mathbb{R}^2$.

Since w is distance preserving on each edge of \mathcal{S} , the maps f_Δ and w coincide on the boundary of Δ . In particular, if Δ and Δ' share a common edge, then f_Δ and $f_{\Delta'}$ agree on that common edge. Therefore, the collection of maps $\{f_\Delta\}$ determines a single piecewise distance preserving map $f: P \rightarrow \mathbb{R}^2$.

It remains to show that f satisfies the conclusion of the theorem. Let $x \in P$ be arbitrary and let y be a point on the edge of a triangle in \mathcal{T} that contains x . Then $|x - y| < \frac{\varepsilon}{3}$ by our choice of \mathcal{T} . Consequently,

$$\begin{aligned} |f(x) - h(x)| &\leq |f(x) - w(y)| + |w(y) - h(y)| + |h(y) - h(x)| \\ &= |f(x) - f(y)| + |w(y) - h(y)| + |h(y) - h(x)| \\ &\leq 2 \cdot |x - y| + |w(y) - h(y)| \\ &< \varepsilon, \end{aligned}$$

because $w(y) = f(y)$ and the maps f, h are distance nonexpanding. □

We conclude this section with a counterexample explaining one way in which we cannot improve Akopyan’s theorem. One might expect that a stronger statement holds, namely, the map f in Akopyan’s theorem can be constructed so that it coincides with h on a given finite set of points. The following exercise shows that this cannot be done in general.


3.5. Exercise. Consider the following 5 points in \mathbb{R}^3 :

$$o = (0, 0, 0), p = (0, 0, 1), a = (2, 0, 0), b = (-1, 2, 0), c = (-1, -2, 0).$$

Let P be the “tripod” that is the polyhedral space consisting of the three triangles $\triangle opa$, $\triangle opb$, and $\triangle opc$ in \mathbb{R}^3 , equipped with the induced length metric.

Note that the restriction of the coordinate projection $\pi(x, y, z) = (x, y, 0)$ to P is distance nonexpanding and piecewise linear. We have

$$\pi(o) = \pi(p) = o, \quad \pi(a) = a, \quad \pi(b) = b, \quad \pi(c) = c.$$



Show that there is no piecewise distance preserving map $f: P \rightarrow \mathbb{R}^2 = \mathbb{R}^2 \times \{0\}$ such that $f(a) = a$, $f(b) = b$, and $f(c) = c$.

§4. GROMOV’S RUMPLING THEOREM

Recall that \mathbb{S}^2 denotes the unit sphere in \mathbb{R}^3 , which we equip with its induced length metric. Here is our main theorem.

4.1. Theorem. There is a length-preserving map $f: \mathbb{S}^2 \rightarrow \mathbb{R}^2$.

Such a map f has to cease on an everywhere dense set in \mathbb{S}^2 . More precisely, the restriction of f to any open subset of \mathbb{S}^2 cannot be injective.³

In the proof of the theorem we will use the following statement.

4.2. Exercise. Let K be a convex polyhedron in \mathbb{R}^3 . Given a point x in \mathbb{R}^3 , show that there is a unique point $\bar{x} \in K$ that minimizes the distance $|x - \bar{x}|$. Moreover, show that the projection map

$$\varphi: \mathbb{R}^3 \rightarrow K, \quad \varphi(x) = \bar{x}$$

is distance nonexpanding.

Proof of Theorem 4.1. Consider a nested sequence $K_0 \subset K_1 \subset \dots$ of convex polyhedra in \mathbb{R}^3 whose union is the open unit ball. Let $P_n = \partial K_n$ denote the surface of K_n equipped with the induced length metric. Note that P_n is a 2-dimensional polyhedral space for each n .

Let φ_n denote the projection onto K_n , as in Exercise 4.2, which is a distance nonexpanding map. Since $K_n \subset K_{n+1}$, it follows that $\varphi_n(P_{n+1}) = P_n$. Note that P_n and P_{n+1} can be triangulated in such a way that the restriction of φ_n to any simplex of P_{n+1} is an orthogonal projection onto some simplex of P_n . In particular, the restriction of φ_n to P_{n+1} is piecewise linear⁴ and distance nonexpanding with respect to the length metrics on P_{n+1} and P_n .

We claim that for any point $x \in \mathbb{S}^2$, there is unique sequence of points $x_n \in P_n$ such that $x_n \rightarrow x$ as $n \rightarrow \infty$ and $\varphi_n(x_{n+1}) = x_n$ for all n . Uniqueness follows because the maps φ_n are distance nonexpanding. To show existence, fix any sequence $z_n \in P_n$ such

³To prove this, one can show that if f is injective and length-preserving on an open set $U \subset \mathbb{S}^2$, then f maps (sufficiently short) geodesics to straight lines (this requires the Domain Invariance Theorem, see for example Section 2.9 in [3].) It follows that the restriction of f to U is locally distance preserving, which is impossible.

⁴See the definition at the beginning of §3.

that $z_n \rightarrow x$. Consider the double sequence $y_{n,m} \in P_n$, defined for $n \leq m$, such that $y_{n,n} = z_n$ and $y_{n,m} = \varphi_n(y_{n+1,m})$ if $0 \leq n < m$. Then we set

$$x_n = \lim_{m \rightarrow \infty} y_{n,m}.$$

4.3. Exercise. Show that the limit above exists and $\varphi_n(x_{n+1}) = x_n$ for any n . Then show that $x_n \rightarrow x$ as $n \rightarrow \infty$.

Let $x_n \rightarrow x \in \mathbb{S}^2$ be the sequence as above. Define $\psi_n: \mathbb{S}^2 \rightarrow P_n$ by $\psi_n(x) = x_n$. Note that ψ_n is distance nonexpanding, $\psi_n = \varphi_n \circ \psi_{n+1}$ for all n , and that for any $p, q \in \mathbb{S}^2$ we have

①
$$|p_n - q_n|_{P_n} \rightarrow |p - q|_{\mathbb{S}^2} \text{ as } n \rightarrow \infty,$$

where $p_n = \psi_n(p)$ and $q_n = \psi_n(q)$.

The desired length-preserving map $f: \mathbb{S}^2 \rightarrow \mathbb{R}^2$ will be a “limit” in some sense of a sequence of piecewise distance preserving maps $f_n: P_n \rightarrow \mathbb{R}^2$. The maps will be constructed recursively to satisfy

$$|f_{n+1}(x) - f_n(\varphi_n(x))| < \varepsilon_n$$

for a carefully chosen sequence (ε_n) of positive numbers that decays rapidly to 0.

Recursive construction of $f_n: P_n \rightarrow \mathbb{R}^2$ and ε_n . Assume we have a piecewise distance preserving map $f_n: P_n \rightarrow \mathbb{R}^2$ and a given ε_n . The composition

$$f_n \circ \varphi_n: P_{n+1} \rightarrow \mathbb{R}^2$$

is piecewise linear and distance nonexpanding. So, we can apply Akopyan’s Theorem 3.2 to construct a piecewise distance preserving map $f_{n+1}: P_{n+1} \rightarrow \mathbb{R}^2$ that is ε_n -close to $f_n \circ \varphi_n$.

Let $M(n+1)$ denote the number of triangles in a triangulation of P_{n+1} such that f_{n+1} is distance preserving on each triangle. Set

②
$$\varepsilon_{n+1} = \frac{\varepsilon_n}{2 \cdot M(n+1)}.$$

Next, we recursively define the construction of f_n and ε_n . It goes as follows.

- (1) Choose an arbitrary $\varepsilon_0 > 0$ and take a piecewise distance preserving map $f_0: P_0 \rightarrow \mathbb{R}^2$, say, the one provided by Zalgaller’s folding theorem (1.3).
- (2) Use φ_0, f_0 , and ε_0 to construct f_1 .
- (3) Use f_1 to construct ε_1 .
- (4) Use φ_1, f_1 , and ε_1 to construct f_2 .
- (5) Use f_2 to construct ε_2 .
- (6) and so on.⁵

It remains to prove the following claim.

③ *The sequence of maps $f_n \circ \psi_n: \mathbb{S}^2 \rightarrow \mathbb{R}^2$ converges to a length-preserving map*

$$f: \mathbb{S}^2 \rightarrow \mathbb{R}^2.$$

Since ε_n decays faster than $\frac{\varepsilon_0}{2^n}$, the sequence $(f_n \circ \psi_n)(x) \in \mathbb{R}^2$ is Cauchy, hence convergent, for any fixed x . We define $f: \mathbb{S}^2 \rightarrow \mathbb{R}^2$ by

$$f(x) = \lim_{n \rightarrow \infty} (f_n \circ \psi_n)(x).$$

⁵The procedure is similar to walking on stairs: you take a right step, which makes it possible to take a left step, which in turn makes it possible to take a right step, and so on.

By the recursive construction of f_n , we have

$$|(f_n \circ \psi_n)(x) - f(x)| < \varepsilon_n$$

for any $x \in \mathbb{S}^2$ and any n . Since each $f_n \circ \psi_n$ is distance nonexpanding, f is distance nonexpanding as well.

It only remains to show that the constructed map $f: \mathbb{S}^2 \rightarrow \mathbb{R}^2$ is length-preserving. Note that it suffices to show that

$$\textcircled{4} \quad \text{length}(f \circ \alpha) \geq |p - q|_{\mathbb{S}^2}$$

for any curve α between two points $p, q \in \mathbb{S}^2$. For the remainder of the proof, we will need the following notion; it should be viewed as an analog of the length of a curve.

4.4. Definition. *Let X be a metric space and $\alpha: [a, b] \rightarrow X$ a curve. Set*

$$\ell_k(\alpha) \stackrel{\text{def}}{=} \sup \left\{ \sum_{i=1}^k |\alpha(t_i) - \alpha(t_{i-1})|_X \mid a = t_0 < t_1 < \dots < t_k = b \right\}.$$

Note that, given a curve $\alpha: [a, b] \rightarrow X$, we have

$$\begin{aligned} \ell_1(\alpha) &\leq \ell_2(\alpha) \leq \ell_3(\alpha) \leq \dots, \\ \ell_k(\alpha) &\rightarrow \text{length } \alpha \text{ as } k \rightarrow \infty, \\ \ell_k(\alpha) &\leq \text{length } \alpha \text{ for any } k. \end{aligned}$$

Moreover, if

$$\ell_k(\alpha) = \text{length } \alpha,$$

then α is a chain made from at most k geodesic segments.

The following exercise states that if two curves α and β are sufficiently close, then $\ell_k(\alpha) \approx \ell_k(\beta)$. Note that the value $|\text{length } \alpha - \text{length } \beta|$ might be large in this case.

4.5. Exercise. *Suppose that $\alpha, \beta: \mathbb{I} \rightarrow X$ are two curves that are close in the sense that*

$$|\alpha(t) - \beta(t)|_X < \varepsilon$$

for all $t \in \mathbb{I}$. Show that

$$|\ell_k(\alpha) - \ell_k(\beta)| \leq 2 \cdot k \cdot \varepsilon.$$

Now we come back to the proof of $\textcircled{4}$. Set $p_n = \psi_n(p)$ and $q_n = \psi_n(q)$. Let β be an arbitrary curve from p_n to q_n in P_n . Note that one can find a shorter curve γ from p_n to q_n whose image in any triangle of the triangulation of P_n is a line segment, and moreover, the endpoints of these line segments lie on β . It follows that $f_n \circ \gamma$ is a broken line in \mathbb{R}^2 with at most $M(n)$ edges, whose vertices we denote, in order, by

$$f_n(p_n) = z_0, z_1, \dots, z_k = f_n(q_n).$$

Note that $k \leq M(n)$ and each z_i lies on the curve $f_n \circ \beta$. Therefore,

$$\begin{aligned} |p_n - q_n|_{P_n} &\leq \text{length } \gamma \\ &= \ell_{M(n)}(f_n \circ \gamma) \\ \textcircled{5} \quad &= |z_0 - z_1| + \dots + |z_{k-1} - z_k| \\ &\leq \ell_{M(n)}(f_n \circ \beta). \end{aligned}$$

Fix a curve α from p to q in \mathbb{S}^2 . By Exercise 4.5 and $\textcircled{2}$, for all n we have

$$\textcircled{6} \quad |\ell_{M(n)}(f \circ \alpha) - \ell_{M(n)}(f_n \circ \psi_n \circ \alpha)| \leq 2 \cdot M(n) \cdot \varepsilon_n = \varepsilon_{n-1}.$$

Given $\varepsilon > 0$, we can choose n so large that $\varepsilon_{n-1} \leq \frac{\varepsilon}{2}$ and

$$|p - q|_{\mathbb{S}^2} - |p_n - q_n|_{P_n} \leq \frac{\varepsilon}{2},$$

which can be arranged by ❶. Applying ❺ for $\beta = \psi_n \circ \alpha$ and ❻, we see that

$$\begin{aligned} \text{length}(f \circ \alpha) &\geq \ell_{M^{(n)}}(f \circ \alpha) \\ &\geq \ell_{M^{(n)}}(f_n \circ \psi_n \circ \alpha) - \varepsilon_{n-1} \\ &\geq |p_n - q_n|_{P_n} - \varepsilon_{n-1} \\ &\geq |p - q|_{\mathbb{S}^2} - \frac{\varepsilon}{2} - \varepsilon_{n-1} \\ &\geq |p - q|_{\mathbb{S}^2} - \varepsilon. \end{aligned}$$

Since $\varepsilon > 0$ was arbitrary,

$$\text{length}(f \circ \alpha) \geq |p - q|_{\mathbb{S}^2}.$$

Hence ❷ follows. □


§5. ARNOLD'S PROBLEM ON PAPER FOLDING

This lecture is meant to be entertaining. Here we discuss the following problem posed by V. Arnold in 1956 [5, Problem 1956-1].

5.1. Problem. *Is it possible to fold a square on the plane so that the resulting figure will have a longer perimeter?*


The answer to this problem depends on the meaning of the word “fold”.

For example, one can consider a sequence of *foldings* in which all layers are folded simultaneously along a line. By the following exercise, the perimeter can never increase under a folding of this type.



5.2. Exercise. *Show that each folding described above indeed decreases the perimeter. (Note that, in general, the intersection of the line q with the polygon M in the picture may be a union of line segments.)*

Using only the *foldings* described above makes it impossible to unfold a layer which lies on top of another layer, as shown in the following picture.




Note that the *unfold* increases the perimeter, although not beyond the perimeter of the original square. It is still unknown if it is possible to increase perimeter by a sequence of such “folds” and “unfolds”.


Japanese crane. Now let us consider a more general definition of folding. Imagine that we mark in advance the lines of folding and start to fold the paper in such a way that each domain between folds remains flat all the time.

If you understand “folding” this way, then the answer to the problem is “yes”. In some sense, this problem was solved by origami practitioners well before it was even posed. The possibility to increase the perimeter slightly can be seen in the base for the

crane. This was known by origami masters for centuries⁶, but mathematicians learned this answer only in 1998.



The base for the crane has four long ends and one short end. Two ends are used for wings and the other two have to be thinned, as one is used for the head and the other for the tail. Thinning twice each of the long ends makes it possible to produce a base which can then be folded into the plane to obtain a figure with larger perimeter.




On the picture above, you can see the net of folds, the base, and the base with opened out ends. On the net of folds, you can see the number of the layer in the base. The dashed lines are the folds that appear at the opening out. The perimeter increases by about 0.5%, and there are 80 layers in the end. A similar example was described in Lang’s book [18]. We do not know of a way to increase the perimeter with a smaller number of layers.


If a is the side of the original square, then it takes a bit less than a to go around each of four needles and it takes about $(\sqrt{2} - 1) \cdot a$ to go around the short end, resulting in a longer perimeter. Thinning the ends many times makes it possible to increase the perimeter by a value arbitrarily close to $(\sqrt{2} - 1) \cdot a$.

The following picture describes another way to increase the perimeter, based on an idea of Yaschenko [25]. It can be obtained by recursive application of one simple move. If one repeats this move sufficiently many times, we obtain a figure with a longer perimeter. Since each iteration adds two layers near the concave corner, the total number of layers in this model is much larger than in the crane base.

⁶It appears in the oldest known book on origami, “Senbazuru Orikata,” dated 1797; but for sure it was known much earlier.




The sea urchin and the comb. It turns out that the perimeter can be made arbitrarily large. This can be seen in the origami model for a sea urchin constructed by Robert Lang in 1987, see [19]. In 2004, a complete solution was discovered independently by Alexei Tarasov [24]. Tarasov constructed a folding of a “comb”, which is shown in the picture on the right. More importantly, he proved that the comb can be folded in a true way, in particular without starching and crooking the paper, as is often done in origami.



The online version of this lectures [22] also contains three movies that describe Tarasov’s solution.

Foldings in 4-dimensional space. One can define a “folding” as a piecewise distance preserving map from the square to the plane. These foldings are yet more general than those considered above. The following exercise shows that it is not always possible to realize such a map by folding a paper model.



5.3. Exercise. Consider the part of regular tessellation in the square \square as in the picture.

Show that there is a map $f: \square \rightarrow \mathbb{R}^2$ that is distance preserving on each polygon in the tessellation and is such that it only reverses the orientation of the gray polygons.⁷

Show that it is not possible to make a paper folding model for f .⁸

The obstructions described in the above exercise disappear in \mathbb{R}^4 ; i.e., one can regard piecewise distance preserving maps as paper folding in 4-dimensional space. Moreover, one can actually fold the square in \mathbb{R}^4 , as prescribed by a given piecewise distance preserving map.

By this we mean that one can construct a continuous one-parameter family of piecewise distance preserving maps $f_t: \square \rightarrow \mathbb{R}^4$, $t \in [0, 1]$, with fixed triangulation, say \mathcal{T} , such that

⁷Less formally, you need to “fold” along each segment in this tessellation.

⁸More formally, we need to think that the plane lies in the space \mathbb{R}^3 , and we need to show that the map f cannot be approximated by injective continuous maps $\square \rightarrow \mathbb{R}^3$.

- ◊ f_0 is a distance preserving map from \square to the coordinate plane $\mathbb{R}^2 \times \{0\}$ in $\mathbb{R}^4 = \mathbb{R}^2 \times \mathbb{R}^2$;
- ◊ the map f_1 is our given piecewise distance preserving map to the same coordinate plane;
- ◊ the map f_t is injective for any $t \neq 1$.

The proof of the last statement is based on Exercise 2.4. Let a_1, \dots, a_k be the vertices of \mathcal{T} and b_1, \dots, b_k the corresponding images for the piecewise distance preserving map. Set $f_t(a_i) \in \mathbb{R}^2 \times \mathbb{R}^2$ to be

$$f_t(a_i) = \left(\frac{a_i + b_i}{2} + \cos(\pi \cdot t) \cdot \frac{a_i - b_i}{2}, \sin(\pi \cdot t) \cdot \frac{a_i - b_i}{2} \right);$$

so $f_0(a_i) = (a_i, 0)$ and $f_1(a_i) = (b_i, 0)$ for any i . We can extend f_t linearly to each triangle of \mathcal{T} . Direct calculation shows that $\ell_{i,j}(t) = |f_t(a_i) - f_t(a_j)|$ is monotone in t ; in particular, if $|a_i - a_j| = |b_i - b_j|$, then $\ell_{i,j}(t)$ is constant. This proves that f_t is piecewise distance preserving.

Further, direct calculations show that for any $x, y \in \square$ we have

$$|f_t(x) - f_t(y)|^2 = p - q \cdot \cos(\pi \cdot t)$$


for some constants p and q . Therefore, if $x \neq y$, then $|f_t(x) - f_t(y)| > 0$ for any $t \neq 1$. In other words, f_t is injective for any $t \neq 1$.

We see that for the paper folding in \mathbb{R}^4 , the existence of perimeter increasing folds follows from Brehm’s theorem. It suffices to construct a distance nonexpanding map f from the square to the plane so that the perimeter of its image is sufficiently long. Then applying Brehm’s theorem for a sufficiently dense finite set of points in the square, we get a piecewise distance preserving map h that is arbitrarily close to f . In particular, we can arrange it so that the perimeter of the image $f(\square)$ is still sufficiently long.

The required map can be constructed as follows. Fix a large n and divide the square \square into n^2 squares with side length $\frac{a}{n}$. Let $d(x)$ denote the distance from a point $x \in \square$ to the boundary of the small square that contains x . The function $d: \square \rightarrow \mathbb{R}$ takes values in $[0, \frac{a}{2n}]$. Further, we enumerate the squares by integers from 1 to n^2 . Given $x \in \square$, denote by $i(x)$ the (say, minimal) number in this enumeration of a small square that contains x .

Now for each $i \in \{1, \dots, n^2\}$ choose a unit vector $u_i \in \mathbb{R}^2$ so that $u_i \neq u_j$ if $i \neq j$. Consider the map $f: \square \rightarrow \mathbb{R}^2$ defined by

$$f(x) = d(x) \cdot u_{i(x)}.$$



It is straightforward to check that the obtained map is distance nonexpanding. The image $f(\square)$ consists of n^2 segments of length $\frac{a}{2n}$ that start at the origin. Thus, the perimeter of $f(\square)$ is equal to $2 \cdot n^2 \cdot \frac{a}{2n} = a \cdot n$. So, by taking n sufficiently large, one can make the perimeter of the image $f(\square)$ arbitrarily large. (The picture shows the case where $n = 4$. In this case the perimeter of $f(\square)$ is $4 \cdot a$, which is the same as the perimeter of the original square. However for $n > 4$, it gets larger.)

FINAL REMARKS

Zalgaller’s folding theorem. Zalgaller’s theorem holds true in all dimensions: any m -dimensional polyhedral space P admits a piecewise distance preserving map to \mathbb{R}^m .

In [27], Zalgaller proved this statement for $m \leq 4$. The trick described in the “proof with no cheating” makes the proof work in all dimensions. This trick first appeared in Krat’s thesis [17].

One may call a higher dimensional simplex *acute* if it contains its own circumcenter — this provides a natural generalization of an acute triangle to higher dimensions. The existence of acute triangulations in higher dimensions seems to be unlikely, but as far as we can see, nothing is known about these triangulations.

Brehm’s extension theorem. This was proved by Brehm in [9] and rediscovered independently many years later by Akopyan and Tarasov in [2]. The proofs are based on the same idea.

Brehm’s extension theorem holds in all dimensions and it can be proved along the same lines.

Kirszbraun theorem. This remarkable theorem was proved by Kirszbraun in his thesis, defended in 1930. A few years later he published the result in [16]. Independently the same result was reproved later by Valentine [26].

The paper of Danzer, Grünbaum, and Klee [11], which is delightful to read, gives a very nice proof of this theorem, based on Helly’s theorem on the intersection of convex sets.

Akopyan’s approximation theorem. This theorem also admits direct generalizations to higher dimensions. Moreover, the condition that f is piecewise linear is redundant. This is because any distance nonexpanding map from a polyhedral space to \mathbb{R}^m can be approximated by piecewise linear distance nonexpanding maps.

The 2-dimensional case was proved by Krat in her thesis [17]. In [1], Akopyan noticed that Brehm’s extension theorem simplifies the proof and also makes it possible to prove the higher dimensional case.

Much earlier, an analogous question was considered by Burago and Zalgaller. They proved that any piecewise linear embedding of a 2-dimensional polyhedral surface in \mathbb{R}^3 can be approximated by a piecewise distance preserving embedding; see [7] and [8].

Rumpling the sphere. Theorem 4.1 admits the following generalization, which can be proved along the same lines.

5.4. Gromov’s rumpling theorem. *Let M be an m -dimensional Riemannian manifold. Then any distance nonexpanding map $f: M \rightarrow \mathbb{R}^m$ can be approximated by length-preserving maps. More precisely, given $\varepsilon > 0$, there is a length-preserving map $f_\varepsilon: M \rightarrow \mathbb{R}^m$ such that*

$$|f_\varepsilon(x) - f(x)| < \varepsilon$$

for any $x \in M$.

This result is a partial case of Gromov’s theorem in [13, 2.4.11]. The proof presented here is based on [21]; Gromov’s original proof is different. The proof in [21] also makes it possible to construct surprising examples of spaces that admit length-preserving maps to \mathbb{R}^m , such as sub-Riemannian manifolds.

Gromov’s theorem roughly states that length-preserving maps have no nontrivial global properties. This is a typical “local to global” problem. Here the length-preserving property is “local” and the only “global” consequence is trivial: it is the distance nonexpanding property.

For such “local to global” problems the answer “no nontrivial global properties” is the most common, but it does not mean that it is easy to prove. There is a machinery invented by Gromov which makes it possible to prove this for many local to global problems. This machinery is called the “ h -principle” (homotopy principle). The h -principle is not a theorem, it is a property which often holds for different geometric structures. There are a few methods to prove the h -principle, including the one described in the proof

of Theorem 4.1⁹. Gromov’s rumpling theorem is one of the simplest examples. Other examples include

- ◊ The cone eversion theorem, which states that there is a continuously varying one-parameter family of smooth functions $f_t(x, y)$, $t \in [0, 1]$, without critical points in the punctured plane $\mathbb{R}^2 \setminus \{0\}$, such that $f_0(x, y) = -\sqrt{x^2 + y^2}$ and $f_1(x, y) = \sqrt{x^2 + y^2}$. See [12, Lecture 27] and read the whole book; it is nice.
- ◊ The Nash–Kuiper theorem, which in particular implies the existence of C^1 -smooth length-preserving maps $\mathbb{S}^2 \rightarrow \mathbb{R}^3$ whose image has arbitrarily small diameter.
- ◊ Smale’s sphere eversion paradox, which states that there is a continuous one-parameter family of smooth immersions $f_t: \mathbb{S}^2 \rightarrow \mathbb{R}^3$, $t \in [0, 1]$, such that $f_0: \mathbb{S}^2 \rightarrow \mathbb{R}^3$ is the standard inclusion and $f_1(x) = -f_0(x)$ for all $x \in \mathbb{S}^2$.
- ◊ Combining the techniques of Smale and Nash–Kuiper, one can make sphere eversions f_t in the class of C^1 -smooth length-preserving maps.

For further reading we suggest the comprehensive introduction to the h -principle by Eliashberg and Mishachev [15].

Paper folding. The aspects of paper folding related to geometric constructions are discussed in [14]; that paper is very entertaining. Interesting aspects of paper folding in the 3-dimensional space are covered in [12, Lecture 15].

Acknowledgments. We would like to thank Arseniy Akopyan, Robert Lang, Alexei Tarasov for their help. Also we would like to thank all the students in our class for their participation and true interest.

REFERENCES

- [1] A. V. Akopyan, *PL-analog of Nash–Kuiper theorem*, <http://www.moebiuscontest.ru/files/2007/akopyan.pdf>.
- [2] A. V. Akopyan and A. S. Tarasov, *A constructive proof of Kirszbraun’s theorem*, *Mat. Zametki* **84** (2008), no. 5, 781–784; English transl., *Math. Notes* **84** (2008), no. 5–6, 725–728. MR2500644 (2010c:47123)
- [3] A. D. Alexandrov, *Convex polyhedra*, Gosudarstv. Izdat. Tehn.-Teor. Lit., Moscow–Leningrad, 1950; English transl., Springer Monogr. Math., Springer-Verlag, Berlin, 2005. MR2127379 (2005j:52002)
- [4] R. Alexander, *Lipschitzian mappings and total mean curvature of polyhedral surfaces*. I, *Trans. Amer. Math. Soc.* **288** (1985), no. 2, 661–678. MR776397 (86c:52004)
- [5] V. I. Arnol’d, *Arnold problems*, FAZIS, Moscow, 2000; English transl., Springer-Verlag, Berlin, 2004. MR2078115 (2005c:58001)
- [6] K. Bezdek and R. Connelly, *Pushing disks apart — the Kneser–Poulsen conjecture in the plane*, *J. Reine Angew. Math.* **553** (2002), 221–236. MR1944813 (2003m:52001)
- [7] Yu. D. Burago and V. A. Zalgaller, *Polyhedral embedding of a net*, *Vestn. Leningrad. Univ. Ser. Mat. Mekh. Astronom.* **1960**, vyp. 7, 66–80. (Russian) MR0116317 (22:7112)
- [8] ———, *Isometric piecewise-linear embeddings of two-dimensional manifolds with a polyhedral metric into \mathbb{R}^3* , *Algebra i Analiz* **7** (1995), no. 3, 76–95; English transl., *St. Petersburg Math. J.* **7** (1996), no. 3, 369–385. MR1353490 (96g:53091)
- [9] U. Brehm, *Extensions of distance reducing mappings to piecewise congruent mappings on \mathbb{R}^m* , *J. Geom.* **16** (1981), no. 2, 187–193. MR642266 (83b:51020)
- [10] D. Yu. Burago, Yu. D. Burago, and S. V. Ivanov, *A course in metric geometry*, Graduate Stud. Math., vol. 33, Amer. Math. Soc., Providence, RI, 2001. MR1835418 (2002e:53053)
- [11] L. Danzer, B. Grünbaum, and V. Klee, *Helly’s theorem and its applications*, *Proc. Sympos. Pure Math.*, vol. 7, Amer. Math. Soc., Providence, RI, 1963, pp. 101–180. MR0157289 (28:524)
- [12] D. B. Fuchs and S. L. Tabachnikov, *Mathematical omnibus, Thirty lectures on classic mathematics*. Amer. Math. Soc., Providence, RI, 2007. MR2350979 (2008h:00002)
- [13] M. Gromov, *Partial differential relations*, *Ergeb. Math. Grenzgeb.* (3), vol. 9, Springer-Verlag, Berlin, 1986. MR864505 (90a:58201)

⁹Usually, the h -principle is formulated in terms of partial differential equations, but one may think of “length-preserving maps” as weak solutions of a particular partial differential equation.

- [14] T. C. Hull, *Solving cubics with creases: the work of Beloch and Lill*, Amer. Math. Monthly **118** (2011), no. 4, 307–315. MR2800341 (2012e:52069)
- [15] Ya. M. Eliashberg and N. M. Mishachev, *Introduction to the h-principle*, MCNMO, Moscow, 2004.
- [16] M. D. Kirszbraun, *Über die zusammenziehende und Lipschitzsche Transformationen*, Fund. Math. **22** (1934), 77–108.
- [17] S. Krat, *Approximation problems in length geometry*, Thesis, Penn. State. Univ., 2005. MR2706488
- [18] R. J. Lang, *Origami design secrets. Mathematical methods for an ancient art*, CRC Press, Boca Raton, FL, 2012. MR2841394
- [19] J. Montroll and R. J. Lang, *Origami sea life*, Dover Publ., New York, 1991.
- [20] A. M. Petrunin, *Arnold's problem on paper folding*, Tasks St. Petersburg School Olympiad on Mathematics, Nevskii Dialect, St. Petersburg, 2008, pp. 116–125, arXiv: 1004.0545. (Russian)
- [21] ———, *Intrinsic isometries in Euclidean space*, Algebra i Analiz **22** (2010), no. 5, 140–153; English transl., St. Petersburg Math. J. **22** (2011), no. 5, 803–812. MR2828830 (2012e:54037)
- [22] A. Petrunin and A. Yashinski, *Lectures on piecewise distance preserving maps*, arXiv:1405.6606.
- [23] S. Saraf, *Acute and nonobtuse triangulations of polyhedral surfaces*, European J. Combin. **30** (2009), no. 4, 833–840. MR2504642 (2010e:52028)
- [24] A. S. Tarasov, *Solution of Arnold's "folded ruble" problem*, Chebyshevskii Sb. **5** (2004), no. 1, 174–187. (Russian) MR2098977 (2005e:52005)
- [25] I. Yaschenko, *Make your dollar bigger now!!!*, Math. Intelligencer **20** (1998), no. 2, 38–40.
- [26] F. A. Valentine, *A Lipschitz condition preserving extension for a vector function*, Amer. J. Math. **67** (1945), no. 1, 83–93. MR0011702 (6:203e)
- [27] V. A. Zalgaller, *Isometric imbedding of polyhedra*, Dokl. Akad. Nauk SSSR **123** (1958), no. 4, 599–601. (Russian) MR0103511 (21:2279)

MATHEMATICS DEPARTMENT, PENN STATE UNIVERSITY, UNIVERSITY PARK, PENNSYLVANIA 16802
E-mail address: anton.petrinin@gmail.com

MATHEMATICS DEPARTMENT, PENN STATE UNIVERSITY, UNIVERSITY PARK, PENNSYLVANIA 16802
E-mail address: allan@math.hawaii.edu

Received 25/MAY/2014

Translated by THE AUTHORS