

ON CONFORMAL MAPPING OF INFINITE STRIPS

BY

S. E. WARSCHAWSKI

INTRODUCTION

Let S be the strip in the plane of the complex variable $w = u + iv$ defined by the relations

$$\phi_-(u) < v < \phi_+(u), \quad -\infty < u < +\infty,$$

where $\phi_-(u), \phi_+(u)$ are continuous for $-\infty < u < +\infty$. Let $\theta(u) \equiv \phi_+(u) - \phi_-(u)$ and $\psi(u) \equiv \frac{1}{2}[\phi_+(u) + \phi_-(u)]$. S can be mapped conformally onto the strip $|y| < \pi/2$ of the z -plane, $z = x + iy$, by means of an analytic function $z = Z(w) = X(w) + iY(w)$ in such a manner that $\lim_{u \rightarrow +\infty} X(w) = +\infty$. The principal object of this paper is to obtain asymptotic expressions for $Z(w)$ and its derivative $Z'(w)$ as $u \rightarrow +\infty$. For this purpose two inequalities concerning the difference $X(w_2) - X(w_1)$ ($w_1 = u_1 + iv_1, w_2 = u_2 + iv_2$ in S) are established which are similar to certain inequalities of Ahlfors⁽¹⁾, but which, due to some assumptions regarding the smoothness of the boundary of S , yield sharper estimates for large values of u_1 and u_2 .

We say that S is an L -strip⁽²⁾ with the boundary inclination γ at $u = +\infty$, $|\gamma| < \pi/2$, if, for $u_2 > u_1$,

$$\frac{\phi_+(u_2) - \phi_+(u_1)}{u_2 - u_1}, \quad \frac{\phi_-(u_2) - \phi_-(u_1)}{u_2 - u_1}$$

approach the same limit, $\tan \gamma$, as u_1 and $u_2 \rightarrow +\infty$ simultaneously. The two inequalities in question (the "basic inequalities") are then as follows:

I. If S is an L -strip with the boundary inclination $\gamma = 0$ at $u = +\infty$, then

$$X(w_2) - X(w_1) \leq \pi \int_{u_1}^{u_2} \frac{1 + \psi'^2(u)}{\theta(u)} du + \frac{\pi}{12} \int_{u_1}^{u_2} \frac{\theta'^2(u)}{\theta(u)} du + o(1),$$

where $o(1) \rightarrow 0$ as $u_1, u_2 \rightarrow +\infty$, uniformly with respect to v_1 and v_2 .

II. If S is an L -strip as in I and if, in addition, $\phi'_+(u)$ and $\phi'_-(u)$ are continuous and of bounded variation for $u_0 \leq u \leq +\infty$, then

$$X(w_2) - X(w_1) \geq \pi \int_{u_1}^{u_2} \frac{1 + \psi'^2(u)}{\theta(u)} du - \frac{\pi}{4} \int_{u_1}^{u_2} \frac{\theta'^2(u)}{\theta(u)} du + o(1).$$

Presented to the Society, January 1, 1941; received by the editors January 23, 1941.

⁽¹⁾ Ahlfors [1, p. 10 and p. 16]. The number in the brackets refers to the author's paper quoted in the bibliography.

⁽²⁾ For a justification of this notation see §1 (b).

If now the integral $\int_{u_0}^{\infty} [\theta'^2(u)/\theta(u)] du$ converges, I and II together yield an asymptotic expression for the difference $X(w_2) - X(w_1)$:

$$X(w_2) - X(w_1) = \pi \int_{u_1}^{u_2} \frac{1 + \psi'^2(u)}{\theta(u)} du + o(1) \quad \text{as } u_1, u_2 \rightarrow + \infty.$$

Combining this with a result on $Y(w)$ established in this paper under the same hypothesis as in I, we obtain the following asymptotic representation for $Z(w)$:

$$Z(w) = \lambda + \pi \int_{u_0}^u \frac{1 + \psi'^2(t)}{\theta(t)} dt + i\pi \frac{v - \psi(u)}{\theta(u)} + o(1), \quad \text{as } u \rightarrow + \infty,$$

uniformly with respect to v . Here λ is a real constant.

As to $Z'(w)$, we find under the same hypothesis as in I,

$$Z'(w) \sim \frac{\pi}{\theta(u)} \quad \text{as } u \rightarrow + \infty,$$

uniformly in any subregion $S_\beta: \{(|v - \psi(u)|)/\theta(u) \leq \beta/\pi\}$ where $0 < \beta < \pi/2$. The "approach of u to $+\infty$ in any S_β " is the analogue of the "approach within any angle" in the case of a finite boundary point at which the boundary curve possesses a tangent. In this connection we obtain an extension of Carathéodory's well known theorem which states that the map of a region bounded by a Jordan curve onto a circle is quasi-conformal at a boundary point which is the vertex of a corner (cf. §16).

Similar expressions for $Z(w)$ and $Z'(w)$ are obtained when $\gamma \neq 0$, but $|\gamma| < \pi/2$. Further theorems are derived as corollaries from the above mentioned results.

An important part in the proof of some of these results is played by a theorem of A. Qstrowski (cf. §2) which deals with the argument of the derivative of the mapping function in a neighborhood of a point at which the boundary curve has a cusp. In the present paper a new proof of this theorem is given.

By the use of suitable transformations these results can be applied to the study of the mapping function in a neighborhood of a boundary point for various boundary configurations. Let Γ be a closed Jordan curve in the ω -plane, R the interior of Γ , let $\zeta = \zeta(\omega)$ map R conformally onto the circle $|\zeta - 1| < 1$, let ω_0 be a point on Γ and let $\zeta(\omega_0) = 0$. By means of simple logarithmic transformations, asymptotic expressions for $\zeta(\omega)$ and $\zeta'(\omega)$ as $\omega \rightarrow \omega_0$ are derived from the above stated results. In particular, when Γ possesses a tangent at ω_0 , the expression for $\zeta(\omega)$ yields a new criterion for the existence of the derivative of $\zeta(\omega)$ at ω_0 (i.e., $\lim_{\omega \rightarrow \omega_0} \zeta(\omega)/(\omega - \omega_0)$ for unrestricted approach). Another case is that in which Γ has a cusp at ω_0 . It was partly for the purpose of treating this case that the present investigation was started.

In addition to the asymptotic expressions for $\zeta(\omega)$ and $\zeta'(\omega)$ as $\omega \rightarrow \omega_0$ we obtain some extensions of earlier results on the cusp which are due to Ostrowski.

Other applications include the study of $\zeta(\omega)$ for a region R whose boundary contains two "concurrent" spirals. These spirals may both approach a single "asymptotic" point ω_0 , or else the set of their limiting points might be a more general point set which forms a prime end of R .

I. PRELIMINARIES

1. **Definitions.** We begin with a few definitions which will be used in the paper.

(a) **SIMPLE JORDAN STRIP.** Let C_+ and C_- be two curves in the w -plane ($w = u + iv$) represented by continuous functions

$$v = \phi_+(u), \quad v = \phi_-(u), \quad u \geq u_0, \quad \phi_+(u) > \phi_-(u),$$

respectively, and let C_1 be a Jordan arc⁽³⁾ which lies in the half-plane $u \leq u_0$ and connects the finite end points of C_+ and C_- . The curve C , consisting of C_+ , C_- and C_1 decomposes the complex plane into two regions (by Jordan's theorem). Let S be the one which contains the region

$$\phi_-(u) < v < \phi_+(u), \quad u_0 < u < +\infty.$$

We call S a simple Jordan strip.

We set $\theta(u) = \phi_+(u) - \phi_-(u)$ and denote by θ_u the segment $\{\Re w = u, \phi_-(u) \leq v \leq \phi_+(u)\}$.

(b) **L-TANGENT AT $u = +\infty$; L-STRIP.** Let C be a curve represented by the equation $v = \phi(u)$ where $\phi(u)$ is continuous for $u_0 \leq u < +\infty$. We say that C has an L-tangent with the angle of inclination γ , $-\pi/2 \leq \gamma \leq \pi/2$ at $u = +\infty$, if the angle of inclination of any chord w_1w_2 of C ($w_1 = u_1 + iv_1, w_2 = u_2 + iv_2$), $u_1 < u_2$, approaches the limit γ when u_1 and u_2 approach $+\infty$ simultaneously, or, in other words, if for every $\epsilon > 0$ there exists an $R(\epsilon) > 0$ such that for the principal value of the argument,

$$|\arg(w_2 - w_1) - \gamma| < \epsilon \quad \text{when } u_2 > u_1 > R(\epsilon).$$

Let S be a simple Jordan strip whose boundary curves C_+ and C_- have both L-tangents at $u = +\infty$ with the angles of inclination γ_+ and γ_- , $|\gamma_+| < \pi/2$, $|\gamma_-| < \pi/2$, respectively. Then, for sufficiently large u , say $u \geq u_1 \geq u_0$, $\theta(u)$ has bounded difference quotients and hence $\theta'(u)$ exists for $u \geq u_1$, except possibly for a set of measure zero. Furthermore, $\theta'(u)$ is bounded and $\theta(b) - \theta(c) = \int_c^b \theta'(u) du$, $b \geq c \geq u_1$.

If, in particular, the L-tangents of C_+ and C_- at $u = +\infty$ have the same angle of inclination γ , then we call S an L-strip with the boundary inclination γ at $u = +\infty$.

(3) C_1 may pass through the infinite point (" $u = -\infty$ ").

The definition of an L -tangent at $u = +\infty$ is patterned after that of an L -tangent to a curve at a finite point⁽⁴⁾. Let β be a Jordan arc which possesses a tangent at one of its end points, P . If the angle of inclination of every chord P_1P_2 of β ($P_1 \neq P_2$), approaches that of the tangent at P as P_1 and P_2 approach P simultaneously, then we say that β has an L -tangent at P .

2. A theorem of Ostrowski on L -cusps. Let Γ be a closed Jordan curve. If, in a neighborhood of a point P of Γ , Γ consists of two branches Γ_+ and Γ_- each of which possesses a tangent at P , and if the interior angle made by these two tangents is θ , $0 < \theta \leq 2\pi$, then we shall say that Γ has a *corner of measure θ* at P . The limiting case where $\theta = 0$ will be called a *cusp*. If both tangents at P are L -tangents then we shall say that Γ has an *L -corner* ($\theta > 0$) or an *L -cusp* ($\theta = 0$) at P .

The following theorem is due to A. Ostrowski⁽⁵⁾.

THEOREM I. (Ostrowski.) *Let Γ be a closed Jordan curve in the ω -plane which has an L -cusp at the point P of Γ , and let $\omega = f(\zeta)$ map the circle $|\zeta| < 1$ conformally onto the interior of Γ in such a manner that $\zeta = 1$ corresponds to P . Then, for any determination of the argument in $|\zeta| < 1$, $\lim_{\zeta \rightarrow 1} \arg [f'(\zeta)(\zeta - 1)]$ exists for unrestricted approach.*

This theorem is an extension of a theorem by Lindelöf⁽⁶⁾ on $\arg f'(\zeta)$ in the case that Γ has an L -corner of measure $\theta = \pi$ at P , and was first stated and proved by Ostrowski⁽⁵⁾. We give here a proof of this theorem, which is different from the one by Ostrowski. It follows to some extent the ideas which Lindelöf used in proving his theorem.

Proof of Theorem I. (i) It may be assumed that P is the point $\omega = 0$, that Γ_- follows Γ_+ when Γ is traversed in the mathematically positive direction and that the L -tangents to Γ_+ and Γ_- at $\omega = 0$ fall on the negative real axis. Then we have, for a proper choice of the determination of the argument for ω_1 and ω_2 on Γ_- ,

$$(2.1) \quad \lim_{\omega_1, \omega_2 \rightarrow 0} \arg (\omega_2 - \omega_1) = \pi \quad (\omega_1 \text{ between } \omega = 0 \text{ and } \omega_2)$$

and for ω_1 and ω_2 on Γ_+ ,

$$(2.2) \quad \lim_{\omega_1, \omega_2 \rightarrow 0} \arg (\omega_2 - \omega_1) = 2\pi \quad (\omega_2 \text{ between } \omega_1 \text{ and } \omega = 0).$$

Let γ_+ and γ_- be subarcs of Γ_+ and Γ_- , respectively, such that both have $\omega = 0$ as an end point and that for all ω_1, ω_2 on γ_-

$$(2.3) \quad \left| \arg (\omega_2 - \omega_1) - \pi \right| < \pi/8 \quad (\omega_1 \text{ between } \omega_2 \text{ and } \omega = 0)$$

⁽⁴⁾ The idea of an L -tangent at a finite point was introduced by Lindelöf, [1, pp. 89-91], the term " L -tangent" by Ostrowski [1, p. 93].

⁽⁵⁾ Ostrowski [1, pp. 181-183].

⁽⁶⁾ Lindelöf [1, pp. 89-91].

and for all ω_1, ω_2 on γ_+ ,

$$(2.4) \quad \left| \arg (\omega_2 - \omega_1) - 2\pi \right| < \pi/8 \quad (\omega_2 \text{ between } \omega = 0 \text{ and } \omega_1).$$

It follows from (2.3) and (2.4) that γ_- and γ_+ can be represented in the form $(\omega = \xi + i\eta)$:

$$\eta = g_-(\xi), \quad \eta = g_+(\xi),$$

respectively. Both of these functions are continuous and have derivatives almost everywhere in a certain interval $\xi_0 \leq \xi \leq 0$ ($\xi_0 < 0$). We can assume both functions have a derivative at ξ_0 , so that γ_- and γ_+ have tangents at the points $P_1 (\xi_0, g_-(\xi_0))$ and $P_2 (\xi_0, g_+(\xi_0))$, respectively. Denote the arcs OP_1 and OP_2 by Γ'_- and Γ'_+ , respectively.

Let γ be an arc with continuously turning tangent which joins P_1 and P_2 , which has the same tangents at P_1 and P_2 as Γ , and which lies in the interior R of Γ except for its end points. Call Γ_1 the closed Jordan curve formed by γ , Γ'_- and Γ'_+ . We shall prove the theorem first for the function $\omega = f_1(\zeta)$ which maps the circle $|\zeta| < 1$ onto the interior R_1 of Γ_1 , and for which $f_1(1) = 0$.

(ii) We establish first an auxiliary inequality. Let a be a point on Γ'_+ and b a point on Γ'_- (so that $\Re a \geq \xi_0, \Re b \geq \xi_0$). If that determination of the argument is chosen which lies between $-\pi/2$ and $3\pi/2$ (inclusive),

$$(2.5) \quad -\frac{\pi}{8} \leq \arg (b - a) \leq \pi + \frac{\pi}{8}.$$

Assume first that $\Re b > \Re a$. Then only the left-hand side of (2.5) needs proof, since then $\arg (b - a) \leq \pi/2$. Draw the line $\xi = \Re a$. This line intersects Γ'_- in a point a' whose ordinate is greater than that of a . By use of elementary properties of the angles of a triangle, it is easily seen that

$$\arg (b - a) \geq \arg (b - a'), \quad \arg (b - a') \geq -\pi/8,$$

because of (2.3), since b and a' are both on Γ'_- and b is between a' and 0. The case $\Re b < \Re a$ is treated similarly. If $\Re a = \Re b$, $\arg (b - a) = \pi/2$ and (2.5) is true.

(iii) Let $f_1(e^{i\theta_1})$ and $f_1(e^{i\theta_2})$ be interior points of Γ'_- and Γ'_+ , respectively, $0 < \theta_1 < \theta_2 < 2\pi$, and let δ_0 be a positive number which is less than $\min \{ \theta_1, 2\pi - \theta_2 \}$ and also so small that (2.3) and (2.4) are satisfied on the arcs of Γ_1 corresponding to $0 \leq \theta \leq \theta_1 + \delta_0$ and $\theta_2 - \delta_0 \leq \theta \leq 2\pi$ of $|\zeta| = 1$, respectively. Since $\omega = f_1(\zeta)$ is continuous and univalent on $|\zeta| = 1$,

$$Q(\theta; \delta) \equiv \arg [f_1(e^{i(\theta+\delta)}) - f_1(e^{i\theta})], \quad 0 < \delta \leq \delta_0, \delta \text{ fixed},$$

is continuous for all real θ , once the determination for one value, say $\theta = 0$, is selected. Since (2.3) holds on Γ'_- , $Q(\theta; \delta)$ can be determined by the condition $|Q(0; \delta) - \pi| < \pi/8$. We have then also

$$(2.6) \quad \left| Q(\theta; \delta) - \pi \right| < \pi/8 \quad \text{for } 0 \leq \theta \leq \theta_1.$$

We assert now that

$$(2.7) \quad |Q(\theta; \delta) - 2\pi| < \pi/8 \quad \text{for } \theta_2 - \delta \leq \theta \leq 2\pi - \delta.$$

To prove this it is sufficient to show that

$$(2.8) \quad |Q(\theta_2; \delta) - 2\pi| < \pi/8.$$

For since (2.4) holds on the arc of Γ_1 which corresponds to the arc $\theta_2 - \delta_0 \leq \theta \leq 2\pi$ of $|\zeta| = 1$ and $Q(\theta; \delta)$ is continuous in θ , (2.7) will then follow.

To prove (2.8) we observe first that for any θ ,

$$(2.9) \quad Q(\theta + 2\pi; \delta) - Q(\theta; \delta) = 2\pi.$$

This is an application of the principle of the argument, since the function $f_1(\zeta e^{i\theta}) - f_1(\zeta)$ is regular in $|\zeta| < 1$, continuous in $|\zeta| \leq 1$, and has exactly one zero, $\zeta = 0$, in $|\zeta| \leq 1$.

Now let $Q(\theta_2; \delta) = h_2 + 2k\pi$ where k is an integer and $|h_2| < \pi/8$ (because of (2.4)). Since $Q(\theta; \delta)$ is continuous, it follows from (2.4) and (2.5) that for $\theta_2 \leq \theta \leq 2\pi$

$$(2.10) \quad 2k\pi - \pi/8 \leq Q(\theta; \delta) \leq 2k\pi + \pi + \pi/8.$$

Let $Q(0; \delta) = h_1 + \pi$, $|h_1| < \pi/8$ by (2.6). Then by (2.9)

$$Q(2\pi; \delta) = h_1 + 3\pi$$

and hence by (2.10) for $\theta = 2\pi$

$$2k\pi - \pi/8 \leq h_1 + 3\pi \leq 2k\pi + \pi + \pi/8.$$

The left-hand side of this inequality implies $k \leq 1$, the right-hand side $k \geq 1$, so that $k = 1$ and (2.8) is proved.

If we set $\omega_1 = f_1(e^{i\theta})$, $\omega_2 = f_1(e^{i(\theta+\delta)})$ then (2.6) shows that $Q(\theta; \delta)$ for $0 \leq \theta \leq \theta_1$ is identical with the determination of $\arg(\omega_2 - \omega_1)$ chosen in (2.1). Hence, by (2.1), $Q(\theta; \delta) \rightarrow \pi$ as θ and δ approach 0 simultaneously. Similarly, we infer from (2.7) and (2.2) that $Q(\theta; \delta) \rightarrow 2\pi$ as $\theta \rightarrow 2\pi$ and $\delta \rightarrow 0$ with the restriction that $\theta \leq 2\pi - \delta$.

Finally we note that there exists an $M > 0$ such that for all $0 \leq \theta \leq 2\pi$ and all $0 < \delta \leq \delta_0$

$$(2.11) \quad |Q(\theta; \delta)| \leq M.$$

For $0 \leq \theta \leq \theta_1$ this follows from (2.6) for $\theta_2 - \delta \leq \theta \leq 2\pi - \delta$ from (2.7), for $2\pi - \delta \leq \theta \leq 2\pi$ from (2.10) with $k = 1$, and for $\theta_1 \leq \theta \leq \theta_2 - \delta$ from (2.3), (2.4) and the fact that γ has a continuously turning tangent.

(iv) For fixed $\delta \leq \delta_0$, the function

$$g(\zeta; \delta) \equiv \frac{f_1(\zeta e^{i\delta}) - f_1(\zeta)}{\zeta(e^{i\delta} - 1)} \quad \text{for } \zeta \neq 0, \quad g(0; \delta) = f_1'(0),$$

is regular in $|\zeta| < 1$, continuous in $|\zeta| \leq 1$, and never vanishes because $f_1(\zeta)$ is univalent in $|\zeta| \leq 1$. Therefore, any branch of $\arg g(\zeta; \delta)$ is harmonic in $|\zeta| < 1$ and continuous in $|\zeta| \leq 1$. We choose that branch of $\arg g(\zeta; \delta)$ which reduces to

$$Q(\theta; \delta) - \delta/2 - \theta - \pi/2, \quad 0 \leq \theta \leq 2\pi,$$

for $|\zeta| = 1$. Consider now, for $|\zeta| < 1$, the harmonic function

$$(2.12) \quad P(\zeta; \delta) = \arg [g(\zeta; \delta)(\zeta - 1)] = \arg g(\zeta; \delta) + \arg (\zeta - 1),$$

where $\arg (\zeta - 1)$ is determined by the condition that it reduces to π for $\zeta = 0$. $P(\zeta; \delta)$ is bounded in $|\zeta| < 1$ and it has continuous boundary values on $|\zeta| = 1$ except at $\zeta = 1$. Hence it can be represented by the Poisson integral

$$P(\zeta; \delta) = \frac{1}{2\pi} \int_0^{2\pi} P(e^{i\theta}; \delta) \frac{(1 - \rho^2)d\theta}{1 + \rho^2 - 2\rho \cos(\theta - \alpha)}, \quad \zeta = \rho e^{i\alpha}, \quad \rho < 1.$$

(v) Let $\epsilon > 0$ be given. Then by our statement at the end of section (iii) there exists an $\eta = \eta(\epsilon) < \epsilon/4$ such that, for all positive $\delta \leq \min(\delta_0, \eta)$:

$$(2.13) \quad |Q(\theta; \delta) - \pi| < \epsilon/4 \quad \text{for } 0 \leq \theta \leq \eta$$

and

$$(2.14) \quad |Q(\theta; \delta) - 2\pi| < \epsilon/4 \quad \text{for } 2\pi - \eta \leq \theta \leq 2\pi - \delta.$$

Observing that, for the determination of $\arg (\zeta - 1)$ selected in (2.12)

$$\arg (e^{i\theta} - 1) = \frac{\theta}{2} + \frac{\pi}{2}, \quad 0 < \theta < 2\pi,$$

we find, for $0 < \theta < \eta$,

$$\begin{aligned} |P(e^{i\theta}; \delta) - \pi| &= \left| Q(\theta; \delta) - \frac{\delta}{2} - \theta - \frac{\pi}{2} + \left(\frac{\theta}{2} + \frac{\pi}{2} \right) - \pi \right| \\ &\leq |Q(\theta; \delta) - \pi| + \frac{\delta}{2} + \frac{|\theta|}{2} \\ &\leq |Q(\theta; \delta) - \pi| + \frac{\delta}{2} + \frac{\eta}{2}, \end{aligned}$$

and for $2\pi - \eta \leq \theta \leq 2\pi - \delta$,

$$\begin{aligned} |P(e^{i\theta}; \delta) - \pi| &\leq |Q(\theta; \delta) - 2\pi| + \frac{\delta}{2} + \left| \frac{\theta}{2} - \pi \right| \\ &\leq |Q(\theta; \delta) - 2\pi| + \frac{\delta}{2} + \frac{\eta}{2}. \end{aligned}$$

Because of (2.13) and (2.14), we have therefore

$$(2.15) \quad |P(e^{i\theta}; \delta) - \pi| < \frac{\epsilon}{4} + \frac{\epsilon}{8} + \frac{\epsilon}{8} = \frac{\epsilon}{2}$$

for $0 \leq \theta \leq \eta$ and $2\pi - \eta \leq \theta \leq 2\pi - \delta$.

To estimate the difference $|P(\zeta; \delta) - \pi|$ we decompose the Poisson integral as follows:

$$P(\zeta; \delta) = \frac{1}{2\pi} \left\{ \int_0^\eta + \int_\eta^{2\pi-\eta} + \int_{2\pi-\eta}^{2\pi-\delta} + \int_{2\pi-\delta}^{2\pi} \right\}.$$

Then we find, from (2.15), (2.12), and (2.11),

$$\begin{aligned} |P(\zeta; \delta) - \pi| \leq & \frac{1}{2\pi} \frac{\epsilon}{2} \int_0^\eta \frac{(1 - \rho^2)d\theta}{1 + \rho^2 - 2\rho \cos(\theta - \alpha)} \\ & + \frac{M + 2\pi}{2\pi} \frac{2\pi(1 - \rho^2)}{1 + \rho^2 - 2\rho \cos(\eta - |\alpha|)} \\ & + \frac{1}{2\pi} \frac{\epsilon}{2} \int_{2\pi-\eta}^{2\pi-\delta} \frac{(1 - \rho^2)d\theta}{1 + \rho^2 - 2\rho \cos(\theta - \alpha)} \\ & + \frac{M + 2\pi}{2\pi} \frac{\delta(1 - \rho^2)}{(1 - \rho^2)}. \end{aligned}$$

Now we keep ζ fixed and let $\delta \rightarrow 0$. Since M is independent of δ , we find

$$|\arg [f'_1(\zeta)(\zeta - 1)] - \pi| \leq \epsilon + (M + 2\pi) \frac{1 - \rho^2}{1 + \rho^2 - 2\rho \cos(\eta - |\alpha|)}.$$

Hence, as $\zeta \rightarrow 1$,

$$(2.16) \quad \limsup |\arg [f'_1(\zeta)(\zeta - 1)] - \pi| \leq \epsilon,$$

q.e.d.

(vi) It remains to prove the theorem for the original function $f(\zeta)$. The inverse function of $\omega = f(\zeta)$ maps R_1 onto a region E of the circle $|\zeta| < 1$ and the arc P_1OP_2 of Γ_1 onto an arc of the circumference $|\zeta| = 1$ which contains $\zeta = 1$ as an interior point. If $\zeta = \zeta(s)$, $\zeta(1) = 1$, is a suitably chosen function which maps the circle $|s| < 1$ onto E , then $f_1(s) = f(\zeta(s))$. By Schwarz's reflection principle, $\zeta(s)$ is regular at $\zeta = 1$ and $\zeta'(1) \neq 0$. Now we have, in a sufficiently small neighborhood of $\zeta = 1$, $|\zeta| < 1$,

$$\arg f'(\zeta) = \arg f'_1(s) - \arg \zeta'(s)$$

and

$$\arg [f'(\zeta)(\zeta - 1)] = \arg [f'_1(s)(s - 1)] - \arg \zeta'(s) + \arg \frac{\zeta - 1}{s - 1}.$$

Letting $\zeta \rightarrow 1$ in $|\zeta| < 1$ and consequently $s \rightarrow 1$ in $|s| < 1$, we find the desired result from (2.16).

3. **An application.** As an application of Theorem I we prove the following

THEOREM II. *Let S be an L -strip with the boundary inclination $\gamma = 0$ at $u = +\infty$. Let $w = W(z)$, $z = x + iy$, map the strip $|y| < \pi/2$ conformally onto S in such a manner that $\Re W(x + iy) \rightarrow +\infty$ as $x \rightarrow +\infty$. Then*

(a) *For a suitable choice of the branch of the argument, $\lim_{z \rightarrow +\infty} \arg W'(z) = 0$, for unrestricted approach.*

(b) *For z_1 and z_2 in any fixed strip $|y| \leq \beta < \pi/2$ which satisfy the condition $|z_2 - z_1| \leq M$ (M a constant), $\lim [W'(z_2)/W'(z_1)] = 1$ as $x_1 = \Re z_1$ (and hence $x_2 = \Re z_2$) approaches $+\infty$, uniformly in $|y| \leq \beta$ (τ).*

Proof. (a) The transformation $\omega = 1/w$ maps S onto a region R bounded by a closed Jordan curve⁽⁸⁾ Γ in such a manner that $u = +\infty$ corresponds to $\omega = 0$. Let w_1 and w_2 be two points on one of the boundary curves C_+ or C_- of S and $\omega_1 = 1/w_1$ and $\omega_2 = 1/w_2$ their images on Γ . Since S is an L -strip with the boundary inclination $\gamma = 0$ at $u = +\infty$, the principal value of $\arg w$ ($-\pi < \arg w \leq \pi$) is single-valued in S if $u = \Re w$ is sufficiently large, and $\arg w_1$ and $\arg w_2$ approach 0 as w_1 and w_2 approach ∞ along C_+ or C_- . Hence, it follows from the relation $\arg(\omega_2 - \omega_1) = \arg(w_1 - w_2) - \arg w_1 - \arg w_2$, which holds when the principal value is taken for each of the arguments, provided $\Re w_1 > \Re w_2$ and both are sufficiently large, that Γ has an L -cusp at $\omega = 0$.

Let now, for $|\zeta| < 1$, $z = \log [(1 + \zeta)/(1 - \zeta)]$ be the branch of the logarithm which is 0 when $\zeta = 0$. The function

$$\omega = f(\zeta) = \frac{1}{W(z)}, \quad z = \log \frac{1 + \zeta}{1 - \zeta}$$

maps the circle $|\zeta| < 1$ conformally onto R . It can be defined as a continuous function in $|\zeta| \leq 1$, and then $f(1) = 0$. Hence by Theorem I,

$$\lim_{\zeta \rightarrow 1} \arg [f'(\zeta)(\zeta - 1)] = l$$

exists, or

$$\lim_{z \rightarrow +\infty} \arg W'(z) = \lim_{\zeta \rightarrow 1} \arg \left[(W(z))^2 f'(\zeta) \frac{\zeta^2 - 1}{2} \right] = l$$

exists. That this limit is 0 for a suitable choice of the branch of the argument can be seen in the following way. The image L in the w -plane of the real axis

(⁷) Part (b) is due to Ostrowski [1, p. 185, relation (68.6), and p. 177, relation (64.2)].

(⁸) If the arc C_1 of the boundary curve of S passes through $w = \infty$, Γ will have a double point at $\omega = 0$. In this case, however, Γ will be a Jordan curve on the Riemann surface of $(\omega - a)^{1/2}$ for a suitable choice of the point a ($a \neq 0$).

in the z -plane by means of $w = W(z)$ has a tangent at the point $W(x)$ which forms the angle $\arg W'(x)$ with the positive u -axis. If $l \neq 0$, we may assume that l is also not a multiple of 2π . Since L lies in S , the slope of any chord through two points $W(x_0)$ and $W(x_1)$, x_0 fixed, will approach 0 as $x_1 \rightarrow +\infty$, no matter how large x_0 might be chosen. On the other hand, $\lim_{x \rightarrow +\infty} \arg W'(z) = l$ implies that L has an L -tangent with the direction l at $u = +\infty$, which is impossible unless $l = 0$ or a multiple of 2π .

(b) To prove part (b) we note first: If $F(z) = U(z) + iV(z)$ is regular for $|y| < \pi/2$ and if $\lim_{x \rightarrow +\infty} V(z) = V_0$ in $|y| < \pi/2$, then uniformly in any fixed sub-strip $|y| \leq \beta < \pi/2$, $\lim_{x \rightarrow +\infty} F'(z) = 0$ ⁽⁹⁾.

This statement follows immediately from the integral representation⁽¹⁰⁾:

$$F'(z) = \frac{i}{\pi r} \int_0^{2\pi} V(z + re^{i\theta}) e^{-i\theta} d\theta = \frac{i}{\pi r} \int_0^{2\pi} [V(z + re^{i\theta}) - V_0] e^{-i\theta} d\theta,$$

where $0 < r < \pi/2 - \beta$, $z = x + iy$, $|y| \leq \beta$.

Now, if z_1 and z_2 are points in $|y| \leq \beta < \pi/2$, and $|z_2 - z_1| \leq M$,

$$|F(z_2) - F(z_1)| = \left| \int_{z_1}^{z_2} F'(\zeta) d\zeta \right| = o(|z_2 - z_1|) = o(1)$$

as $x_1 \rightarrow \infty$, uniformly in $|y| \leq \beta$. Applying this result to

$$F(z) \equiv \log W'(z) = \log |W'(z)| + i \arg W'(z)$$

and using the result of part (a), we have, for $|z_2 - z_1| \leq M$,

$$\lim_{x_1 \rightarrow +\infty} [\log W'(z_2) - \log W'(z_1)] = 0$$

or

$$\lim_{x_1 \rightarrow +\infty} \frac{W'(z_2)}{W'(z_1)} = 1,$$

q.e.d.

4. Some auxiliary results. As an application of Theorem II we prove the following

LEMMA 1. *Let S be an L -strip in the w -plane with the boundary inclination 0 at $u = +\infty$. For all u exceeding a certain number u_1 , let l_u denote a line segment within S which joins the point $w = u + i\phi_-(u)$ of C_- with some point of C_+ and which forms the angle $\gamma(u)$, $\lim_{u \rightarrow +\infty} \gamma(u) = 0$, with the positive v -axis⁽¹¹⁾. Let*

⁽⁹⁾ This result is well known; see for example, Wolff [1, p. 221, §6], Ostrowski [2, p. 23, Theorem V, Part 3]. For the last part of this proof compare Ostrowski [2, p. 31, Theorem VII].

⁽¹⁰⁾ Copson [1, p. 88, Example 1].

⁽¹¹⁾ The angle γ which a line forms with the positive v -axis is the smaller of the two angles between them (if there be a smaller one), and it is considered as positive if the direction of rotation from the positive v -axis to the line is counterclockwise.

$z = Z(w) = X(w) + iY(w)$ map the strip S conformally onto the strip $|y| < \pi/2$ in such a manner that $\lim_{u \rightarrow +\infty} X(w) = +\infty$. If

$$x^* = \max_{w \in l_u} X(w), \quad x_* = \min_{w \in l_u} X(w), \quad m(u) = \max_{w \in l_u} |\arg Z'(w)|,$$

then, for all sufficiently large u ,

$$x^* - x_* \leq 2\pi \tan \{m(u) + |\gamma(u)|\} \rightarrow 0 \quad \text{as } u \rightarrow +\infty.$$

Proof. By Theorem I, a suitable branch of $\arg Z'(w)$ approaches 0 uniformly in S as $u \rightarrow +\infty$. Let, for $u \geq u_2 \geq u_1$, $|\arg Z'(w)| < \pi/8$, $|\gamma(u)| < \pi/8$. The image of l_u in the z -plane by means of $z = Z(w)$, is an arc λ_u joining $z^* = Z(u + i\phi_-(u))$ on $y = -\pi/2$ with a point on $y = \pi/2$. If $w \in l_u$ (w in the interior of S), then λ_u has a tangent at $z = Z(w)$ which forms the angle $\tau = \arg Z'(w) + \gamma(u)$ with the positive y -axis. A simple application of the mean value theorem shows that no point of λ_u is in the exterior of the isosceles triangle whose top is at z^* , whose base is on $y = \pi/2$ and whose angle at z^* is $2[m(u) + |\gamma(u)|]$. The base of this triangle is $2\pi \tan [m(u) + |\gamma(u)|]$, and therefore $x^* - x_* \leq 2\pi \tan [m(u) + |\gamma(u)|]$.

THEOREM III. Let S be a simple Jordan strip and let $z = Z(w) = X(w) + iY(w)$ map S conformally onto the strip $|y| < \pi/2$ in such a manner that $\lim_{u \rightarrow +\infty} X(w) = +\infty$. Let $w_1 = u_1 + iv_1$, $w_2 = u_2 + iv_2$, $u_0 \leq u_1 \leq u_2$, $x_1 = X(w_1)$, $x_2 = X(w_2)$. Then

(a) the integral

$$(4.1) \quad \pi \int_{u_1}^{u_2} \frac{du}{\theta(u)} \leq x_2 - x_1 + 4\pi;$$

(b) if S is an L -strip with the boundary inclination 0 at $u = +\infty$,

$$(4.2) \quad \pi \int_{u_1}^{u_2} \frac{du}{\theta(u)} \leq x_2 - x_1 + o(1) \quad \text{as } u \rightarrow +\infty,$$

uniformly with respect to v_1, v_2 .

Proof. (a) Part (a) is a theorem of Ahlfors (see Ahlfors [1, p. 10] or Nevanlinna [1, p. 92]).

(b) In the proof of (4.1) an inequality (Ahlfors [1, p. 8, relation (3) or Nevanlinna [1, p. 90, relation (35)]) is first derived which contains the following

$$\pi \int_{u_1}^{u_2} \frac{du}{\theta(u)} \leq x_2^* - x_{*1},$$

where $x_{*1} = \min_{w \in \theta_{u_1}} X(w)$, $x_2^* = \max_{w \in \theta_{u_2}} X(w)$. Now, by Lemma 1, (applied with $l_u \equiv \theta_u$), $x_2^* - x_2 \rightarrow 0$ and $x_1 - x_{*1} \rightarrow 0$ as $u_1 \rightarrow +\infty$. Hence, if we write $x_2^* = x_2 + (x_2^* - x_2)$, $x_{*1} = x_1 + (x_{*1} - x_1)$, the result (4.2) follows immediately.

II. THE FIRST BASIC INEQUALITY

5. **Statement of Theorem IV.** The content of Theorem IV (b) will be referred to as the first basic inequality.

THEOREM IV. Let S be a simple Jordan strip and let the functions $v = \phi_+(u)$ and $v = \phi_-(u)$ representing the boundary curves C_+ and C_- of S have uniformly bounded difference quotients for $u \geq u_0$. Let

$$\theta(u) = \phi_+(u) - \phi_-(u), \quad \psi(u) = \frac{1}{2}[\phi_+(u) + \phi_-(u)].$$

Suppose that the function $w = W(z) = U(z) + iV(z)$, $z = x + iy$, maps the strip $|y| < \pi/2$ conformally onto S in such a manner that $x = +\infty$ corresponds to $u = +\infty$. Let $z_1 = x_1 + iy_1$ and $z_2 = x_2 + iy_2$ be two points in $|y| < \pi/2$, $x_1 < x_2$, and let $u_1 = U(z_1)$, $u_2 = U(z_2)$.

(a) If $|\phi'_+(u)|$ and $|\phi'_-(u)| \leq m$ for $u \geq u_0$, there is an x_0 , depending on u_0 , such that for $x_0 \leq x_1 \leq x_2$

$$(5.1) \quad x_2 - x_1 \leq \pi \int_{u_1}^{u_2} \frac{1 + \psi'^2(u)}{\theta(u)} du + \frac{\pi}{12} \int_{u_1}^{u_2} \frac{\theta'^2(u)}{\theta(u)} du + 8\pi(1 + \frac{4}{3}m^2).$$

(b) If S is an L -strip with the boundary inclination $\gamma = 0$ at $u = +\infty$, then

$$(5.2) \quad x_2 - x_1 \leq \pi \int_{u_1}^{u_2} \frac{1 + \psi'^2(u)}{\theta(u)} du + \frac{\pi}{12} \int_{u_1}^{u_2} \frac{\theta'^2(u)}{\theta(u)} du + o(1)$$

as $x_1 \rightarrow +\infty$, uniformly with respect to y_1, y_2 .

REMARK. Ahlfors proves [1, pp. 12-16] the following theorem. Let S be a region represented in the form $-\theta(u)/2 < v < \theta(u)/2$, $-\infty < u < +\infty$, where $\theta(u)$ is of bounded variation in any finite interval and $0 < \theta(u) < L$ for all u . Then, if x_1, x_2, u_1, u_2 have the same meaning as in Theorem IV,

$$(5.3) \quad x_2 - x_1 \leq \pi \int_{u_1}^{u_2} \frac{du}{\theta(u)} + 8\pi + 16\pi L^2 \frac{\theta_m^2 + V}{\theta_m^4},$$

where θ_m is the infimum of $\theta(u)$ for $u_1 - 4L \leq u \leq u_2 + 4L$ and V is the total variation of $\theta^2(u)$ in this interval. (It should be noted that $\psi(u) \equiv 0$ here because S is symmetrical with respect to the real axis.) While the hypotheses of Theorem IV regarding the smoothness of the boundary of S are more stringent than those of Ahlfors' theorem, Theorem IV contains no restriction as to the symmetry of S or boundedness of $\theta(u)$. Moreover, since $V = 2 \int_{u=u_1-4L}^{u_2+4L} \theta(u) |d\theta(u)|$, for continuous $\theta(u)$,

$$(5.4) \quad 16\pi L^2 \frac{\theta_m^2 + V}{\theta_m^4} \geq 16\pi \frac{L^2}{\theta_m^2} \left\{ 1 + 2 \int_{u=u_1-4L}^{u_2+4L} \frac{|d\theta(u)|}{\theta(u)} \right\},$$

and this shows that under the hypotheses of Theorem IV (b) the “remainder” term $\pi \int_{u_1}^{u_2} (\theta'^2(u)/\theta(u)) du + o(1)$ is smaller than the one in (5.3) if u_1 is sufficiently large. This becomes important in the case when $\liminf_{u \rightarrow +\infty} \theta(u) = 0$, since then (5.4) is not bounded as $u_2 \rightarrow +\infty$ (u_1 fixed), while $\int_{u_1}^{\infty} (\theta'^2(u)/\theta(u)) du$ converges for very general classes of functions $\theta(u)$.—In the proof of Theorem IV (as well as in that of Theorem VI below) we make use of Ahlfors’ method of relating area and arc length employed in the proof of his inequalities.

6. **Proof of Theorem IV.** 1. Let for $x_1 \leq x \leq x_2$, $a < U(z) < b$. We assume first that the functions $\phi_+(u)$ and $\phi_-(u)$ have two continuous derivatives for $a \leq u \leq b$.

The functions

$$(6.1) \quad \bar{u} = h(u) = \int_{u_0}^u \frac{\eta dt}{(\theta(t)(1+t^2))^{1/2}}, \quad \bar{v} = \frac{v - \psi(u)}{\theta(u)}, \quad \eta > 0, \text{ a constant,}$$

map the domain $S(a, b): \{\phi_-(u) \leq v \leq \phi_+(u), a \leq u \leq b\}$ in a one-to-one and continuous manner onto a rectangle: $\{\bar{a} \leq \bar{u} \leq \bar{b}, |\bar{v}| \leq 1\}$ of the \bar{w} -plane, $\bar{w} = \bar{u} + i\bar{v}$. The function $W(z)$ maps the line segment $\{x, |y| \leq \pi/2\}$, $x_1 \leq x \leq x_2$, onto a rectifiable arc l_x in S and the transformation (6.1) carries l_x into a rectifiable arc \bar{l}_x within the strip $|\bar{v}| \leq 1$ of the w -plane. If the variable arc lengths of l_x and \bar{l}_x are denoted by s and \bar{s} , respectively, the

$$\text{length of } \bar{l}_x = \int_{l_x} \frac{d\bar{s}}{ds} ds = \int_{-\pi/2}^{\pi/2} \frac{d\bar{s}}{ds} |W'(x + iy)| dy.$$

Since \bar{l}_x connects a point on the line $\bar{v} = +1$ with another on $\bar{v} = -1$, its length is greater than or equal to 1. Hence,

$$(6.2) \quad 1 \leq \left\{ \int_{-\pi/2}^{\pi/2} \frac{d\bar{s}}{ds} |W'(x + iy)| dy \right\}^2 \leq \pi \int_{-\pi/2}^{\pi/2} \left(\frac{d\bar{s}}{ds} \right)^2 |W'(x + iy)|^2 dy.$$

Now, if $u(s), v(s)$ denote the parametric representation of l_x in terms of s ,

$$\begin{aligned} \left(\frac{d\bar{s}}{ds} \right)^2 &= \left(\frac{d\bar{u}}{du} \frac{du}{ds} \right)^2 + \left(\frac{\partial \bar{v}}{\partial u} \frac{du}{ds} + \frac{\partial \bar{v}}{\partial v} \frac{dv}{ds} \right)^2 \\ &= \left[h'^2(u) + \left(\frac{\partial \bar{v}}{\partial u} \right)^2 \right] \left(\frac{du}{ds} \right)^2 + 2 \frac{\partial \bar{v}}{\partial u} \frac{\partial \bar{v}}{\partial v} \frac{du}{ds} \frac{dv}{ds} + \left(\frac{\partial \bar{v}}{\partial v} \right)^2 \left(\frac{dv}{ds} \right)^2 \\ &\leq \left[h'^2(u) + \left(\frac{\partial \bar{v}}{\partial u} \right)^2 \right] \left(\frac{du}{ds} \right)^2 \\ &\quad + 2 \left| \frac{\partial \bar{v}}{\partial v} \right| \left| \frac{du}{ds} \right| \left| \frac{dv}{ds} \right| \left(h'^2(u) + \left(\frac{\partial \bar{v}}{\partial u} \right)^2 \right)^{1/2} + \left(\frac{\partial \bar{v}}{\partial v} \right)^2 \left(\frac{dv}{ds} \right)^2 \\ &= \left\{ \left(h'^2(u) + \left(\frac{\partial \bar{v}}{\partial u} \right)^2 \right)^{1/2} \frac{du}{ds} + \left| \frac{\partial \bar{v}}{\partial v} \right| \left| \frac{dv}{ds} \right| \right\}^2. \end{aligned}$$

Hence, by Schwarz's inequality,

$$\begin{aligned} \left(\frac{d\bar{s}}{ds}\right)^2 &\leq \left[h'^2(u) + \left(\frac{\partial\bar{v}}{\partial u}\right)^2 + \left(\frac{\partial\bar{v}}{\partial v}\right)^2 \right] \left[\left(\frac{du}{ds}\right)^2 + \left(\frac{dv}{ds}\right)^2 \right] \\ &= h'^2(u) + \left(\frac{\partial\bar{v}}{\partial u}\right)^2 + \left(\frac{\partial\bar{v}}{\partial v}\right)^2, \end{aligned}$$

since $(du/ds)^2 + (dv/ds)^2 = 1$. Observing that (we leave off the argument u)

$$\frac{\partial\bar{v}}{\partial u} = -\frac{\theta\psi' + (v - \psi)\theta'}{\theta^2}, \quad \frac{\partial\bar{v}}{\partial v} = \frac{1}{\theta}$$

(ψ', θ' are derivatives with respect to u), we find at every point (u, v) of l_x

$$(6.3) \quad \left(\frac{d\bar{s}}{ds}\right)^2 \leq \frac{\eta^2}{\theta \cdot (u^2 + 1)} + \frac{\theta^2\psi'^2 + 2(v - \psi)\theta\theta'\psi' + (v - \psi)^2\theta'^2}{\theta^4} + \frac{1}{\theta^2} \equiv f(u, v).$$

Hence, from (6.2)

$$1 \leq \pi \int_{-\pi/2}^{\pi/2} f(U(z), V(z)) |W'(x + iy)|^2 dy.$$

Integrating this inequality between the limits x_1, x_2 with respect to x , we find

$$(x_2 - x_1) \leq \pi \int_{x_1}^{x_2} dx \int_{-\pi/2}^{\pi/2} f(U(z), V(z)) |W'(x + iy)|^2 dy.$$

Now, introducing the variables (u, v) in place of (x, y) by means of the transformation $w = W(x + iy)$, we obtain

$$x_2 - x_1 \leq \pi \iint_{(T)} f(u, v) dudv$$

where T is the image of the rectangle $x_1 < x < x_2, |y| < \pi/2$, in the w -plane. If $u_{*1} = \min_{|v| \leq \pi/2} U(x_1 + iy)$ and $u_2^* = \max_{|v| \leq \pi/2} U(x_2 + iy)$, it is clear that T is contained in the region $\phi_-(u) < v < \phi_+(u), u_{*1} < u < u_2^*$. Using this fact and substituting the value in (6.3) for $f(u, v)$, we find

$$\begin{aligned} x_2 - x_1 &\leq \pi \int_{u_{*1}}^{u_2^*} \int_{\phi_-(u)}^{\phi_+(u)} \frac{1 + \psi'^2(u)}{\theta^2(u)} dvdu + \pi\eta^2 \int_{u_{*1}}^{u_2^*} \int_{\phi_-(u)}^{\phi_+(u)} \frac{dvdu}{\theta(u)(u^2 + 1)} \\ &\quad + \pi \int_{u_{*1}}^{u_2^*} \int_{\phi_-(u)}^{\phi_+(u)} \frac{2(v - \psi)\theta\theta'\psi' + (v - \psi)^2\theta'^2}{\theta^4} dvdu. \end{aligned}$$

The integration with respect to v can be easily carried through. The first two integrals yield the result

$$\pi \int_{u_{*1}}^{u_2^*} \frac{1 + \psi'^2(u)}{\theta(u)} du + \pi \eta^2 \int_{u_{*1}}^{u_2^*} \frac{du}{1 + u^2}.$$

The third integral is equal to

$$2\pi \int_{u_{*1}}^{u_2^*} 0 \cdot \frac{\theta' \psi'}{\theta^3} du + \pi \int_{u_{*1}}^{u_2^*} \frac{2}{3} \left(\frac{\theta}{2}\right)^3 \frac{\theta'^2}{\theta^4} du = \frac{\pi}{12} \int_{u_{*1}}^{u_2^*} \frac{\theta'^2(u)}{\theta(u)} du.$$

Hence we finally obtain

$$x_2 - x_1 \leq \pi \int_{u_{*1}}^{u_2^*} \frac{1 + \psi'^2(u)}{\theta(u)} du + \frac{\pi}{12} \int_{u_{*1}}^{u_2^*} \frac{\theta'^2(u)}{\theta(u)} du + \pi \eta^2 \int_{-\infty}^{\infty} \frac{du}{1 + u^2}.$$

This inequality is true for every η . Keeping x_1, x_2 fixed and letting $\eta \rightarrow 0$ we find

$$(6.4) \quad x_2 - x_1 \leq \pi \int_{u_{*1}}^{u_2^*} \frac{1 + \psi'^2(u)}{\theta(u)} du + \frac{\pi}{12} \int_{u_{*1}}^{u_2^*} \frac{\theta'^2(u)}{\theta(u)} du.$$

2. If the hypothesis made at the beginning of part 1 of this proof (that $\phi_+(u)$ and $\phi_-(u)$ have continuous second derivatives for $a \leq u \leq b$) is not satisfied, we replace the arcs $\beta_+ : v = \phi_+(u)$ and $\beta_- : v = \phi_-(u)$, $a \leq u \leq b'$ ($b' > b$) by certain arcs $\beta_+^{(n)}$ and $\beta_-^{(n)}$ for which this assumption is true and which converge to β_+ and β_- , respectively, as $n \rightarrow +\infty$. We proceed as follows:

Since $\phi_+(u)$ and $\phi_-(u)$ have bounded difference quotients, $\phi'_+(u)$ and $\phi'_-(u)$ exist almost everywhere in $a \leq u \leq b'$ and are, in absolute value, less than or equal to m . There exist therefore two sequences of continuous functions, $\phi'_+(u; n)$ and $\phi'_-(u; n)$, $a \leq u \leq b$, such that

$$(6.5.1) \quad \int_a^b |\phi'_+(u; n) - \phi'_+(u)| du \rightarrow 0, \quad \int_a^b |\phi'_-(u; n) - \phi'_-(u)| du \rightarrow 0 \text{ as } n \rightarrow \infty,$$

$$(6.5.2) \quad |\phi'_+(u; n)| \leq 2m, \quad |\phi'_-(u; n)| \leq 2m, \quad a \leq u \leq b, \quad n = 1, 2, 3, \dots,$$

$$(6.5.3) \quad \frac{d}{du} \phi'_+(u; n), \quad \frac{d}{du} \phi'_-(u; n) \quad \text{exist and are continuous for } a \leq u \leq b.$$

Now we define

$$(6.6) \quad \phi_+(u; n) = \phi_+(a) + \int_a^u \phi'_+(t; n) dt, \quad \phi_-(u; n) = \phi_-(a) + \int_a^u \phi'_-(t; n) dt, \\ a \leq u \leq b.$$

Furthermore, we may assume that $\phi_+(b) \neq 0$ and $\phi_-(b) \neq 0$ and set for $b \leq u \leq b'$

$$q_+(u; n) = q_n + \frac{1 - q_n}{b' - b} (u - b) \quad \text{where } q_n = \frac{\phi_+(b; n)}{\phi_+(b)},$$

$$q_-(u; n) = p_n + \frac{1 - p_n}{b' - b} (u - b) \text{ where } p_n = \frac{\phi_-(b; n)}{\phi_-(b)},$$

and define

$$(6.7) \quad \phi_+(u; n) = \phi_+(u)q_+(u; n), \quad \phi_-(u; n) = \phi_-(u)q_-(u; n) \quad \text{for } b \leq u \leq b'.$$

Then it is evident from (6.5.1), (6.6) and (6.7) that the arcs

$$\beta_+^{(n)}: v = \phi_+(u; n) \quad \text{and} \quad \beta_-^{(n)}: v = \phi_-(u; n), \quad a \leq u \leq b',$$

converge to the arcs β_+ and β_- , respectively, in the sense that

$$(6.8) \quad \lim_{n \rightarrow \infty} \phi_+(u; n) = \phi_+(u), \quad \lim_{n \rightarrow \infty} \phi_-(u; n) = \phi_-(u), \text{ uniformly for } a \leq u \leq b'.$$

If now, for sufficiently large n , the arcs β_+ and β_- are replaced by $\beta_+^{(n)}$ and $\beta_-^{(n)}$, respectively, we obtain a Jordan strip S_n whose boundary curves satisfy the hypotheses stated in the theorem and in the beginning of part 1 of this proof. Let $W_n(z)$ be the function which maps the strip $|y| < \pi/2$ onto S_n in such a manner that $x = +\infty$ corresponds to $u = +\infty$ and that $W_n(0) = W(0)$. Then we may define $u_{*1}^{(n)}$ and $u_2^{*(n)}$ for $W_n(z)$ in the same way as u_{*1} and u_2^* are defined for $W(z)$. Since $\beta_+^{(n)}$ and $\beta_-^{(n)}$ converge to β_+ and β_- , respectively, in the above specified manner,

$$(6.9) \quad \lim_{n \rightarrow \infty} u_{*1}^{(n)} = u_{*1}, \quad \lim_{n \rightarrow \infty} u_2^{*(n)} = u_2^*(12).$$

Let $\theta_n(u)$ and $\psi_n(u)$ have the same meaning for S_n as $\theta(u)$ and $\psi(u)$ for S . We now apply (6.4) to the region S_n , where n is sufficiently large, and obtain an analogous inequality in which $\theta, \psi', \theta', u_{*1}$, and u_2^* are replaced by $\theta_n, \psi_n', \theta_n', u_{*1}^{(n)}, u_2^{*(n)}$, respectively. Here we let $n \rightarrow \infty$, and it follows from (6.5.1) and (6.5.2), that the inequality holds for the original region S without the assumption of the existence of the second derivatives of $\phi_+(u)$ and $\phi_-(u)$.

(12) To see this, we may assume that $w=0$ is neither in S nor on the boundary of S and transform S and S_n by means of the function $\omega = 1/w$ into limited regions R and R_n , respectively. R and R_n are bounded by closed curves Γ and Γ_n , respectively, which coincide except for the arcs $\gamma_+^{(n)}$ and $\gamma_-^{(n)}$ of Γ_n , the images of $\beta_+^{(n)}$ and $\beta_-^{(n)}$, and γ_+ and γ_- of Γ , the images of β_+ and β_- . From (6.8) it follows that the curves Γ_n converge to Γ in the sense that their Fréchet distance approaches 0. (The Fréchet distance d of two Jordan curves C and C' is defined as follows: For any continuous one-to-one transformation of C onto C' , the distance of corresponding points has a maximum. The greatest lower bound of these maxima for all possible transformations is d .) If $\omega_n(\zeta)$ and $\omega(\zeta)$ map $|\zeta| < 1$ conformally onto R_n and R , respectively, and if $\omega_n(0) = \omega(0)$ and $\omega_n(1) = \omega(1) = 0$ (the image of $u = +\infty$), then it follows from a theorem of Radó [1, pp. 180-186] that $\omega_n(\zeta) \rightarrow \omega(\zeta)$ uniformly in $|\zeta| \leq 1$ as $n \rightarrow \infty$. Now the functions $W_n(z)$ and $W(z)$ of the text are

$$W_n(z) = [\omega_n(\zeta)]^{-1}, \quad W(z) = [\omega(\zeta)]^{-1}, \quad \text{where } z = \log \frac{1 + \zeta}{1 - \zeta},$$

and it follows therefore that $W_n(z) \rightarrow W(z)$ uniformly in any fixed rectangle $\xi_1 \leq x \leq \xi_2, |y| \leq \pi/2$. This implies (6.9).

3. Now the proof is easily completed. We treat the cases (a) and (b) of our theorem separately.

(a) By Theorem III (a),

$$\pi \int_{u_{*1}}^{u_1} \frac{du}{\theta(u)} \leq x_1 - x_1 + 4\pi = 4\pi, \quad \pi \int_{u_2}^{u_2^*} \frac{du}{\theta(u)} \leq 4\pi.$$

Moreover, since

$$1 + \psi'^2(u) \leq 1 + m^2, \quad \theta'^2(u) \leq 4m^2,$$

we find (5.1) from (6.4).

(b) In the case (b), we have by Theorem III (b),

$$\int_{u_{*1}}^{u_1} \frac{du}{\theta(u)} = o(1), \quad \int_{u_2}^{u_2^*} \frac{du}{\theta(u)} = o(1)$$

as $u_1 \rightarrow +\infty$, and by hypothesis

$$\lim_{u \rightarrow +\infty} \psi'(u) = \lim_{u \rightarrow +\infty} \theta'(u) = 0.$$

From these facts (5.2) follows immediately.

7. A corollary of Theorem IV. *If S is an L -strip with the boundary inclination $\gamma = 0$ at $u = +\infty$, for which the integrals*

$$(7.1) \quad \int_{u_0}^{\infty} \frac{\phi_+^{\prime 2}(u)}{\theta(u)} du \quad \text{and} \quad \int_{u_0}^{\infty} \frac{\phi_-^{\prime 2}(u)}{\theta(u)} du \quad \text{converge,}$$

and if $z = Z(w) = X(w) + iY(w)$ maps S conformally onto the strip $|y| < \pi/2$, in such a manner that $\lim_{u \rightarrow +\infty} X(w) = +\infty$, then there exists a constant λ such that, for $w = u + iv \in S$,

$$(7.2) \quad X(w) = \lambda + \pi \int_{u_0}^u \frac{dt}{\theta(t)} + o(1) \quad \text{as } u \rightarrow +\infty, \text{ uniformly with respect to } v.$$

Proof. First we note that (7.1) implies the convergence of $\int_{u_0}^{\infty} (\theta'^2(u)/\theta(u)) du$ and $\int_{u_0}^{\infty} (\psi'^2(u)/\theta(u)) du$. Applying now Theorems III (b) and IV (b) we find that, for $w_1 = u_1 + iv_1, w_2 = u_2 + iv_2$ in $S, u_1 \leq u_2$,

$$(7.3) \quad \pi \int_{u_1}^{u_2} \frac{dt}{\theta(t)} + o(1) \leq X(w_2) - X(w_1) \leq \pi \int_{u_1}^{u_2} \frac{dt}{\theta(t)} + o(1),$$

where $o(1) \rightarrow 0$ as $u_1, u_2 \rightarrow +\infty$, uniformly with respect to v_1, v_2 . If we set $A(w) \equiv X(w) - \pi \int_{u_0}^u [\theta(t)]^{-1} dt$, then (7.3) means that for every ϵ there exists an $N(\epsilon)$ such that

$$|A(w_2) - A(w_1)| < \epsilon \quad \text{if } u_2 \geq u_1 > N(\epsilon).$$

Hence (7.2) is true by Cauchy's convergence principle.

The condition (7.1) is very restrictive, since $\theta(u)$ might be of smaller order of magnitude than $\phi_+{}^{1/2}(u)$ and $\phi_-{}^{1/2}(u)$. Our next aim is, therefore, to establish, in place of Theorem III (b), a lower bound for the difference $x_2 - x_1$ in terms of the integral $\int_{u_1}^{u_2} \{ (1 + \psi'^2(u)) / \theta(u) \} du$ and of a suitable "remainder" term. This inequality, combined with Theorem IV, will yield an asymptotic representation for $X(w)$ which will require only the existence of the integral $\int_{u_0}^{\infty} (\theta'^2(u) / \theta(u)) du$ in place of (7.1).

III. THE SECOND BASIC INEQUALITY

8. Preliminaries. In order to derive the inequality indicated at the end of §7 we shall establish several preliminary results.

(a) Let S be an L -strip in the w -plane with the boundary inclination γ , $-\pi/2 \leq \gamma \leq \pi/2$, at $u = +\infty$. Moreover, let its boundary curves $C_+ : v = \phi_+(u)$ and $C_- : v = \phi_-(u)$, $u \geq u_0$, satisfy the hypothesis that $\phi_+{}'(u)$ and $\phi_-{}'(u)$ are continuous and of bounded variation in any finite interval contained in $u \geq u_0$.

(b) Let s denote the variable arc length of C_- measured from $u = u_0$ and $w = w(s)$ the parametric representation of C_- by means of s as parameter. Let $\alpha(s)$, $-\pi/2 < \alpha(s) < \pi/2$, be the angle of inclination of the tangent to C_- at the point $w(s)$.

(c) Since the boundary inclination of S at $u = +\infty$ is γ , there exists a half-plane $H_c : u \geq c$, such that the angle of inclination of a tangent at any point of C_+ or C_- in H_c satisfies the relation

$$|\alpha - \gamma| < \frac{\pi}{8}.$$

Let C_+^* and C_-^* denote the parts of C_+ and C_- , respectively, which lie in H_c . Application of the mean value theorem shows then that any straight line with the angle of inclination β , where

$$\left| \beta - \left(\gamma + \frac{\pi}{2} \right) \right| < \frac{\pi}{8},$$

can intersect C_+^* and C_-^* in but one point each. Hence, there exists a number $\sigma > 0$ such that any normal to C_- at $w(s)$, where $s \geq \sigma$, will intersect C_+^* and C_-^* in exactly one point each. The segment of this normal which lies in S will be denoted by Δ_s , its length by $\Delta(s)$. Evidently $\Delta(s)$ is a continuous function for $s \geq \sigma$.

Denote the point $w(s)$ by A , the other end point of Δ_s (on C_+) by B and the point $u + i\phi_+(u)$ by C ($u = \Re w(s)$). Since, in the triangle ABC , the angle A approaches $|\gamma|$ and B approaches $\pi/2$ as $s \rightarrow \infty$, it follows by the law of sines that

$$(8.1) \quad \lim_{u \rightarrow +\infty} \frac{\Delta(s)}{\theta(u)} = \cos \gamma, \quad u = \mathcal{R}w(s).$$

(d) For the following two lemmas we assume that hypothesis (a) is satisfied.

LEMMA 2. *If $\alpha'(s)$ is continuous for $\sigma \leq a \leq s \leq b$, and if $\Delta(s)\alpha'(s) < 1$ in the open interval $a < s < b$, then no two normals $\Delta_s, a \leq s \leq b$, intersect in $\bar{S}^{(13)}$.*

Proof. We show first that no two normals Δ_s of any closed subarc $a < \alpha \leq s \leq \beta < b$ of the arc $a < s < b$, intersect. If this assertion were not true, there would exist a point $A_1 (s = a_1)$ and a point $B_1 (s = b_1)$ on C_- , $\alpha \leq a_1 < b_1 \leq \beta$, such that Δ_{a_1} and Δ_{b_1} would intersect at some point $D_1 \in \bar{S}^{(13)}$. Let I_1 be the arc $a_1 \leq s \leq b_1$. The curvilinear triangle $A_1B_1D_1$ lies entirely within \bar{S} . Let $s' = (1/2)(a_1 + b_1)$. Then $\Delta_{s'}$ enters this triangle at the point $s = s'$ of C_- and will intersect either A_1D_1 or B_1D_1 or both at some point $D_2 \in \bar{S}$. If D_2 lies on A_1D_1 , we let I_2 be the arc $a_2 \leq s \leq s'$ and denote its end points by $A_2 (s = a_2)$ and $B_2 (s = b_2)$ where $a_2 = a_1, b_2 = s'$. Then $\Delta_{s''}$ where $s'' = (1/2)(a_2 + b_2)$, will intersect either A_2D_2 or B_2D_2 or both at some point D_3 . If D_3 does not lie on A_1D_1 , let I_2 be the arc $s' \leq s \leq b$ where now $a_2 = s'$ and $b_2 = b_1$ and the end points of I_2 are again denoted by A_2 and B_2 . Everything said about $\Delta_{s''}$ holds then for this second choice of I_2 . Continuing in the manner indicated, we obtain a sequence of intervals $I_n (n = 1, 2, 3, \dots)$ whose end points we shall denote by $A_n (s = a_n)$ and $B_n (s = b_n)$, $\alpha \leq a_n < b_n \leq \beta$, such that $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = c$ exist and that Δ_{a_n} and Δ_{b_n} intersect at some point $D_n \in \bar{S}$. By the law of sines, we obtain from the triangle $A_nB_nD_n$:

$$\frac{\sin D_n}{A_nB_n} = \frac{\sin A_n}{B_nD_n}.$$

Since the angle A_n approaches $\pi/2$ as $n \rightarrow \infty$ and the angle $D_n = \alpha(b_n) - \alpha(a_n) \geq 0$, and since $B_nD_n \leq \Delta(b_n)$ we find

$$\lim_{n \rightarrow \infty} \frac{\alpha(b_n) - \alpha(a_n)}{b_n - a_n} \geq \lim_{n \rightarrow \infty} \frac{\sin A_n}{\Delta(b_n)} = \frac{1}{\Delta(c)}$$

or

$$\alpha'(c)\Delta(c) \geq 1,$$

which contradicts the hypothesis.

To complete the proof of the lemma we must show that no $\Delta_s, a < s < b$, intersects Δ_a or Δ_b . Suppose $\Delta_{s_0}, a < s_0 < b$, intersected Δ_b at some point D_0 . Then any Δ_{s_1} with $s_0 < s_1 < b$ must intersect Δ_b , since it must intersect Δ_{s_0} or Δ_b and it cannot intersect Δ_{s_0} (by the part already proved). Let B denote the point $s = b$ on C_- . The point D_1 at which Δ_{s_1} intersects Δ_b must lie on the seg-

⁽¹³⁾ The closure of a point set M in the complex plane will be denoted by \bar{M} .

ment BD_0 of Δ_b (for otherwise Δ_{s_1} would meet Δ_{s_0}) and hence $BD_1 < BD_0 \leq \Delta(b)$.

Let now $\{s_n\}$, $n = 1, 2, 3, \dots$, $s_0 < s_n < s_{n+1} < b$, denote a sequence which converges to b as $n \rightarrow \infty$, and let Δ_{s_n} intersect Δ_b at D_n . For all n , $BD_n \leq BD_1 < \Delta(b)$. Call A_n the point $s = s_n$ on C_- . Then, by the law of sines,

$$\frac{\sin D_n}{A_n B} = \frac{\sin A_n}{BD_n} \geq \frac{\sin A_n}{BD_1}.$$

Since the angle $A_n \rightarrow \pi/2$ as $n \rightarrow \infty$ and since $D_n = \alpha(b) - \alpha(s_n)$,

$$\lim_{n \rightarrow \infty} \frac{\alpha(b) - \alpha(s_n)}{b - s_n} \geq \frac{1}{BD_1} > \frac{1}{\Delta(b)}, \quad \text{or} \quad \Delta(b)\alpha'(b) > 1,$$

while the continuity of $\Delta(s) \cdot \alpha'(s)$ implies that $\Delta(b)\alpha'(b) \leq 1$. A similar argument applies to the point $s = a$.

LEMMA 3. Let Δ_a and Δ_b , $\sigma \leq a < b$, intersect at a point C in \bar{S} . Suppose that any of the angles, which a chord of C_- or of C_+ through the end points⁽¹⁴⁾ of Δ_s and $\Delta_{s'}$, $a \leq s < s' \leq b$, forms with Δ_s and $\Delta_{s'}$, differs from $\pi/2$ in absolute value by less than ϵ , $0 < \epsilon < \pi/8$ ⁽¹⁵⁾. Then

$$(8.2) \quad \int_a^b \frac{ds}{\Delta(s)} \leq \frac{\alpha(b) - \alpha(a)}{\cos^3(2\epsilon)}.$$

Proof. Call the points $s = a$ and $s = b$ on C_- , A and B , respectively. Draw Δ_s , $a < s < b$, and denote its end points on C_- and C_+ by D and E , respectively. DE will intersect either AC or BC or both. Suppose it intersects BC at a point F . Call C' the end point of Δ_b on C_+ . From the two triangles BDF and $C'EF$ we find by the law of sines

$$DF = BF \frac{\sin B}{\sin D}, \quad FE = FC' \frac{\sin C'}{\sin E}.$$

Since, by hypothesis, $\sin B > \sin(\pi/2 - \epsilon) = \cos \epsilon$, $\sin C' > \cos \epsilon$, we have

$$(8.3) \quad \Delta(s) = DF + FE > BC' \cos \epsilon = \Delta(b) \cos \epsilon.$$

If Δ_s intersects Δ_a we obtain in a similar way

$$(8.4) \quad \Delta(s) \geq \Delta(a) \cos \epsilon.$$

Now, by the mean value theorem, $\int_a^b [\Delta(s)]^{-1} ds = (b - a) / \Delta(\bar{s})$ where $a < \bar{s} < b$. Hence, by (8.3) or (8.4)

⁽¹⁴⁾ If the end points of Δ_s and $\Delta_{s'}$ on C_+ coincide at a point P , then a chord through these end points on C_+ means the tangent to C_+ at P .

⁽¹⁵⁾ This condition will always be satisfied if a is sufficiently large (because of the hypothesis in §8 (a)).

$$\int_a^b \frac{ds}{\Delta(s)} \leq (b - a) \frac{1}{\Delta(a) \cos \epsilon}$$

or

$$\int_a^b \frac{ds}{\Delta(s)} \leq (b - a) \frac{1}{\Delta(b) \cos \epsilon},$$

respectively.

From the triangle ABC we find

$$\Delta(a) \geq AC = AB \frac{\sin B}{\sin C} \geq AB \frac{\cos \epsilon}{\sin C}, \quad \Delta(b) \geq AB \frac{\cos \epsilon}{\sin C},$$

so that

$$\int_a^b \frac{ds}{\Delta(s)} \leq \frac{\sin C}{\cos^2 \epsilon} \frac{b - a}{AB}.$$

Observing that the hypothesis regarding the chords of C_- implies that⁽¹⁶⁾

$$b - a \leq \frac{AB}{\cos(2\epsilon)}$$

and that the angle $C = \alpha(b) - \alpha(a)$, we obtain (8.2).

9. "Invariant" formulation of the second basic inequality. We prove now

THEOREM V. *Let S be an L -strip which satisfies the hypothesis of §8 (a) and let $\alpha(s)$, Δ_s , and $\Delta(s)$ be defined as in §8 (b) and (c). Suppose that $z = Z(w) = X(w) + iY(w)$ maps S conformally onto the strip $|y| < \pi/2$ in such a manner that $X(w) \rightarrow +\infty$ as $u \rightarrow +\infty$. Let w_1 and w_2 be points in \bar{S} and Δ_{s_1} and Δ_{s_2} normals to C_- which pass through w_1 and w_2 , respectively⁽¹⁷⁾. Let $x_1 = X(w_1)$, $x_2 = X(w_2)$. Then, if $s_2 > s_1$,*

$$(9.1) \quad x_2 - x_1 \geq \pi \int_{s_1}^{s_2} \frac{ds}{\Delta(s)} - \pi(1 + o(1)) \int_{s=s_1}^{s_2} |d\alpha(s)| + o(1)$$

where the second integral on the right-hand side is taken in the sense of Stieltjes and $o(1) \rightarrow 0$ as $s_1 \rightarrow +\infty$, uniformly in w_1 and w_2 .

⁽¹⁶⁾ The tangents to any two points of the arc $AB: a \leq s \leq b$, of C form an angle not exceeding 2ϵ . By the mean value theorem it follows therefore that the chord AB forms an angle of measure less than 2ϵ with the tangent to any point of the arc AB . Hence, if the arc AB is represented in the form $y=f(x)$, AB being the x -axis, A the origin and B the point $(l, 0)$, we have

$$b - a = \int_0^l (1 + [f'(x)]^2)^{1/2} dx \leq l(1 + \tan^2(2\epsilon))^{1/2} = \frac{AB}{\cos(2\epsilon)}.$$

⁽¹⁷⁾ If $\mathcal{R}w_1$ and $\mathcal{R}w_2$ are sufficiently large, there always exist normals Δ_{s_1} and Δ_{s_2} passing through w_1 and w_2 , respectively.

Proof. 1. It is sufficient to prove this theorem under the assumption that $\gamma=0$ since the statement of the theorem is invariant with respect to a rotation of the coordinate system in the w -plane through the angle γ in the positive direction. Since S is an L -strip with the boundary inclination $\gamma=0$ at $u=+\infty$, there exists for every $\epsilon>0$, $\epsilon<\pi/8$, a half-plane $H_c: u\geq c$, such that the angle of inclination α , $|\alpha|<\pi/2$, of the tangent at any point of C_+ or C_- in H_c satisfies the condition

$$|\alpha| < \epsilon/2.$$

Let an ϵ be fixed. We assume s_1 so large that all Δ_s with $s\geq s_1$ lie entirely in H_c , say $s_1>\sigma_1$.

2. We assume first that $\alpha'(s)$ exists and is continuous for $s_1\leq s\leq s_2$. Let

$$x_2^* = \max_{w\in\Delta_{s_2}} X(w), \quad x_{*1} = \min_{w\in\Delta_{s_1}} X(w).$$

Then we prove that

$$(9.2) \quad x_2^* - x_{*1} \geq \pi \int_{s_1}^{s_2} \frac{ds}{\Delta(s)} - \frac{\pi}{\cos^3(2\epsilon)} \int_{s_1}^{s_2} |\alpha'(s)| ds.$$

Let $I_1: a_1 < s < b_1$, be a largest open arc of $I: s_1 < s < s_2$, where

$$(9.3) \quad \Delta(s)\alpha'(s) < 1.$$

If there is no such arc we have on I

$$(9.4) \quad \Delta(s)\alpha'(s) \geq 1$$

and hence

$$\pi \int_{s_1}^{s_2} \frac{ds}{\Delta(s)} - \pi \int_{s_1}^{s_2} \alpha'(s) ds \leq 0 \leq x_2^* - x_{*1},$$

so that (9.2) is evidently true. If there are several such arcs, let I_1 be one of them. By Lemma 2, no two normals Δ_s of the closed arc $\bar{I}_1: a_1 \leq s \leq b_1$, will intersect. If there are any normals Δ_s with $b_1 < s \leq s_2$ which intersect Δ_{b_1} , let $\Delta_{b'_1}$ be the one for which b'_1 is as large as possible. Otherwise we set $b'_1 = b_1$. If there are any Δ_s with $s_1 \leq s < a_1$ which intersect Δ_{a_1} , let $\Delta_{a'_1}$ be the one for which a'_1 is as small as possible. Otherwise we set $a'_1 = a_1$.

Call I'_1 the arc $a'_1 < s < b'_1$. No normal Δ_s of the arcs $(\bar{I} - \bar{I}'_1)$ will intersect any of the Δ_s of I_1 .

If $(\bar{I} - \bar{I}'_1)$ is void we do not proceed any further. In case $(\bar{I} - \bar{I}'_1)$ is not void, either (9.4) holds for all $s \in (\bar{I} - \bar{I}'_1)$, or else there exists a largest open arc $I_2: a_2 < s < b_2$, $I_2 \subset (\bar{I} - \bar{I}'_1)$, where (9.3) is true. By Lemma 2 again no two normals Δ_s of I_2 will intersect. If there are any normals Δ_s , $s \in (\bar{I} - \bar{I}'_1)$, with $s > b_2$ which intersect Δ_{b_2} , let $\Delta_{b'_2}$ be the one for which b'_2 is as large as possible.

Otherwise let $b'_2 = b_2$. Similarly, if there are any $\Delta_s, s \in (\bar{I} - I'_1)$, with $s < a_2$ which intersect Δ_{a_2} , let $\Delta'_{a'_2}$ be the one for which a'_2 is as small as possible. Otherwise let $a'_2 = a_2$. Call I'_2 the arc $a'_2 < s < b'_2$. I'_1 and I'_2 have no points in common, except possibly an end point. Moreover, no normal $\Delta_s, s \in \{\bar{I} - (\bar{I}'_1 + \bar{I}'_2)\}$, will intersect any of the normals of \bar{I}_1 and \bar{I}_2 , and none of the normals of I_1 will intersect any of the normals of I_2 .

Continuing this construction in the manner indicated as long as possible we obtain a finite or infinite sequence of arcs $I_n: a_n < s < b_n$, in which (9.3) holds, I_n being a largest open arc of the set $(\bar{I} - \sum_{k=1}^{n-1} I'_k)$ for which (9.3) is true. With each I_n we obtain an arc $I'_n: a'_n < s < b'_n$ containing I_n which is defined in the following manner: If any $\Delta_s, s \in (\bar{I} - \sum_{k=1}^{n-1} I'_k)$, with $s > b_n$ intersects Δ_{b_n} , let $\Delta_{b'_n}$ be the one for which $s = b'_n$ is a maximum, and if any $\Delta_s, s \in (\bar{I} - \sum_{k=1}^{n-1} I'_k)$, with $s < a_n$ intersects Δ_{a_n} , let $\Delta_{a'_n}$ be the one for which $s = a'_n$ is a minimum. We set $b'_n = b_n$ or $a'_n = a_n$ if there are no such Δ_s for $s > b_n$ or $s < a_n$ respectively. Then no normal $\Delta_s, s \in (\bar{I} - \sum_{k=1}^n I'_k)$, intersects any of the normals Δ_s with $s \in I_k, (k = 1, 2, 3, \dots, n)$. This implies that the normals of I_n and $I_m, n > m$, do not intersect. For no $\Delta_s, s \in (\bar{I} - \sum_{k=1}^m I'_k)$ intersects any of the $\Delta_s, s \in I_k (k = 1, 2, \dots, m)$, and $I_n \subset (\bar{I} - \sum_{k=1}^m I'_k)$, since $n > m$. Moreover, since, for $s \in \bar{I}$, any Δ_s with $s < a'_k$ cannot intersect any of the Δ_s with $s > b'_k (k = 1, 2, 3, \dots)$, it is easily seen that any two arcs I'_n and $I'_m, n \neq m$ have no points in common except possibly an end point. Hence, the same is true for $(I'_n - I_n)$ and $(I'_m - I_m)$ if $m \neq n$. Let

$$A = \sum_{k=1}^{\infty} I_k, \quad B = \bar{I} - \sum_{k=1}^{\infty} I'_k, \quad C = \sum_{k=1}^{\infty} (\bar{I}'_k - I_k),$$

so that $\bar{I} = A + B + C$. By our construction, we have

$$(9.5) \quad \text{for } s \in A: \Delta(s)\alpha'(s) < 1.$$

If B is not void, we have

$$(9.6) \quad \text{for } s \in B: \Delta(s)\alpha'(s) \geq 1.$$

If (9.6) were not true, there would be for at least one

$$(9.7) \quad s_0 \in B: \Delta(s_0)\alpha'(s_0) < 1.$$

Then there exists an *open* arc $J \subset I$ containing s_0 such that (9.3) holds for $s \in J$. We show first that no arc $I'_k \subset J$. Such an arc $I'_k (a'_k, b'_k)$ must necessarily coincide with its subarc $I_k (a_k, b_k)$ on which (9.3) holds, for otherwise $\Delta_{a'_k}$ and $\Delta_{a'_k}$ or $\Delta_{b'_k}$ and $\Delta_{b'_k}$ would intersect and that is impossible by Lemma 2 since a'_k, a_k, b_k, b'_k are all points of J , where (9.3) holds. Let I'_n be the first arc in the sequence $\{I'_m\}$ (in the order in which the I'_m were constructed) which is entirely contained in J . Since s_0 is not in any I'_m, \bar{I}'_n lies either to the right or to the left of s_0 . In either case it follows that, in choosing I_n , we did

not select the largest available interval in which (9.3) holds, since in the first case at least the arc (s_0, a_n) and in the second case the arc (b_n, s_0) should be part of I_n . Thus, no I'_m can be entirely contained in J .

Therefore, there can be in J at most one left end point $a'_\nu > s_0$ of some arc I'_ν and one right end point $b'_\mu < s_0$ of some I'_μ and the subarc $b'_\mu < s < a'_\nu$ of J must remain free of any points belonging to any arcs I'_m ($m = 1, 2, 3, \dots$). But this is impossible: The set of arcs I_m is at most denumerable and, if infinite, the lengths of the I_m approach 0 as $m \rightarrow \infty$. Since at each step of our construction a largest possible arc in which (9.3) holds is being taken for an I_m , the arc $b'_\mu < s < a'_\nu$ would have to be included at some step in the sequence $\{I_m\}$. If there is no end point a'_ν or no end point b'_μ in J , an even larger subarc of J remains free of points of any I'_m , and this again is impossible. The assumption (9.7) thus leads to a contradiction, and hence (9.6) is true. We obtain therefore

$$(9.8) \quad \int_{(B)} \frac{ds}{\Delta(s)} \leq \int_{(B)} \alpha'(s) ds = \int_{(B)} |\alpha'(s)| ds.$$

Finally, by Lemma 3, on each of the two arcs of $I'_m - I_m$, $a'_m \leq s \leq a_m$ and $b_m \leq s \leq b'_m$, we have

$$\begin{aligned} \int_{a'_m}^{a_m} \frac{ds}{\Delta(s)} &\leq \frac{1}{\cos^3(2\epsilon)} |\alpha(a_m) - \alpha(a'_m)|, \\ \int_{b_m}^{b'_m} \frac{ds}{\Delta(s)} &\leq \frac{1}{\cos^3(2\epsilon)} |\alpha(b'_m) - \alpha(b_m)|. \end{aligned}$$

Hence,

$$(9.9) \quad \int_{(C)} \frac{ds}{\Delta(s)} \leq \frac{1}{\cos^3(2\epsilon)} \int_{(C)} |\alpha'(s)| ds.$$

3. Consider now an arc $I_n: a_n < s < b_n$ of A . The points on a normal Δ_s are given by the equation

$$(9.10) \quad w = w(s) + ite^{i\alpha(s)}, \quad 0 \leq t \leq \Delta(s).$$

The integral

$$\int_0^{\Delta(s)} |Z'(w(s) + ite^{i\alpha(s)})| dt \equiv \int_{\Delta_s} |Z'(w)| dt \geq \pi,$$

since it represents the length of the image of Δ_s in the z -plane. By Schwarz's inequality,

$$\frac{\pi^2}{\Delta(s)} \leq \int_{\Delta_s} |Z'(w)|^2 dt.$$

If $s \in I_n$ and $\alpha'(s) \geq 0$, we have, because of (9.5),

$$0 \leq 1 - \alpha'(s)\Delta(s) \leq 1 - \alpha'(s)t \quad \text{for } 0 \leq t \leq \Delta(s);$$

and if $\alpha'(s) < 0$, we have

$$1 \leq 1 - \alpha'(s)t \quad \text{for } 0 \leq t \leq \Delta(s).$$

Hence, in either case

$$-\pi^2 \int_{a_n}^{b_n} |\alpha'(s)| ds + \pi^2 \int_{a_n}^{b_n} \frac{ds}{\Delta(s)} \leq \int_{a_n}^{b_n} ds \int_0^{\Delta(s)} |Z'(w)|^2 (1 - \alpha'(s)t) dt.$$

By what was said in part 2, no two normals Δ_s , $a_n \leq s \leq b_n$, will intersect, and the transformation (9.10) maps therefore the region $\{0 < t < \Delta(s), a_n < s < b_n\}$, of an (s, t) -plane in a one-to-one manner onto the (limited) sub-region T_n of S which is bounded by C_-, C_+, Δ_{a_n} and Δ_{b_n} . The Jacobian of this transformation is $(1 - \alpha'(s)t)$. Hence, the last double integral may be written as

$$\iint_{(T_n)} |Z'(u + iv)|^2 du dv.$$

Summation over n gives

$$(9.11) \quad -\pi^2 \int_{(A)} |\alpha'(s)| ds + \pi^2 \int_{(A)} \frac{ds}{\Delta(s)} \leq \sum_{n=1}^{\infty} \iint_{(T_n)} |Z'(u + iv)|^2 du dv.$$

Since, again by the discussion of part 2, the normals Δ_s of any I_n do not intersect those of any other I_m ($n \neq m$), the regions T_n for $n = 1, 2, 3, \dots$ do not overlap. Each T_n is mapped by $z = Z(w)$ onto a subregion of the strip $|y| < \pi/2$ whose area is given by $\iint_{(T_n)} |Z'(u + iv)|^2 du dv$. Since the T_n do not overlap, their images in the z -plane do not overlap either, and the total area which these images cover is therefore less than or equal to $\pi(x_2^* - x_{*1})$. Hence, from (9.11)

$$(9.12) \quad -\pi \int_{(A)} |\alpha'(s)| ds + \pi \int_{(A)} \frac{ds}{\Delta(s)} \leq x_2^* - x_{*1}.$$

Combining (9.12) with (9.8) and (9.9) we find (9.2).

4. In order to prove now the general case of the theorem (where $\alpha'(s)$ is not necessarily continuous) we approximate the arc $\beta: w = w(s), s_1 \leq s \leq s_2$ of C_- by a sequence of arcs $\beta_n: w = w_n(s), s_1 \leq s \leq s_2$ ($n = 1, 2, \dots$), with the following properties⁽¹⁸⁾:

(i) $w_n(s), w_n'(s), w_n''(s)$ are continuous for $s_1 \leq s \leq s_2$, $w_n(s_k) = w(s_k), w_n'(s_k) = w'(s_k)$, ($k = 1, 2$), and β_n does not intersect any part of the boundary of S except β .

⁽¹⁸⁾ The functions $w_n(s)$ may be found as follows: Let $\alpha(s; h) = (1/h) \int_s^{s+h} \alpha(t) dt, s \geq s_1, h > 0$. Then $\alpha(s; h) \rightarrow \alpha(s)$ as $h \rightarrow 0$, uniformly for $s_1 \leq s \leq s_2$. Let $\alpha(s) = \alpha_1(s) - \alpha_2(s)$, where $\alpha_1(s), \alpha_2(s)$ are continuous, non-decreasing functions. The total variation of $\alpha(s; h)$ in $s_1 \leq s \leq s_2$ is

(ii) $w_n(s) \rightarrow w(s)$ and $w'_n(s) \rightarrow w'(s)$ as $n \rightarrow \infty$, uniformly for $s_1 \leq s \leq s_2$. This condition implies that if $\sigma = \sigma_n(s)$ is the variable arc length of β_n , measured from $s = s_1$, and if $\alpha_n(s)$ is the angle ($|\alpha_n(s)| < \pi/2$) which the tangent to β_n at $w_n(s)$ forms with the positive real axis, then

$$(9.13) \quad \sigma_n(s) \rightarrow s - s_1, \quad \alpha_n(s) \rightarrow \alpha(s) \quad \text{as } n \rightarrow \infty,$$

uniformly for all $s_1 \leq s \leq s_2$.

(iii) The total variation

$$(9.14) \quad \int_{s=s_1}^{s_2} |d\alpha_n(s)| \rightarrow \int_{s=s_1}^{s_2} |d\alpha(s)| \quad \text{as } n \rightarrow \infty.$$

Denote by S_n the region obtained from S when the arc β is replaced by β_n . For sufficiently large n and $s \geq s_1$, $\Delta_n(s)$ can be defined for S_n as $\Delta(s)$ is defined for S (see part 1 and §8 (c)). Condition (ii) implies that, as $n \rightarrow \infty$,

$$(9.15) \quad \Delta_n(s) \rightarrow \Delta(s) \quad \text{uniformly for } s_1 \leq s \leq s_2.$$

By a well known theorem, it follows from (9.13) and (9.15) that

$$(9.16) \quad \int_{s=s_1}^{s_2} \frac{d\sigma_n(s)}{\Delta_n(s)} \rightarrow \int_{s_1}^{s_2} \frac{ds}{\Delta(s)}.$$

Finally let S_n be mapped onto $|y| < \pi/2$ so that $u = +\infty$ and $x = +\infty$ correspond to each other and that a point w_0 which is in S and in all S_n is carried

$$\int_{s=s_1}^{s_2} |d\alpha(s; h)| \leq \frac{1}{h} \int_{s=s_1}^{s_2} d \left[\int_s^{s+h} \alpha_1(t) dt \right] + \frac{1}{h} \int_{s=s_1}^{s_2} d \left[\int_s^{s+h} \alpha_2(t) dt \right].$$

Since

$$\begin{aligned} \frac{1}{h} \int_{s=s_1}^{s_2} d \left[\int_s^{s+h} \alpha_i(t) dt \right] &= \frac{1}{h} \int_{s_1}^{s_2} [\alpha_i(s+h) - \alpha_i(s)] ds = \frac{1}{h} \int_{s_1}^{s_2} ds \int_{t=s}^{s+h} d\alpha_i(t) \\ &\leq \frac{1}{h} \int_{s=s_1}^{s_2+h} d\alpha_i(t) \int_{t-h}^t ds = \int_{s=s_1}^{s_2+h} d\alpha_i(t), \end{aligned}$$

we have

$$\int_{s=s_1}^{s_2} |d\alpha(s; h)| \leq \int_{t=s_1}^{s_2+h} d\alpha_1(t) + \int_{t=s_1}^{s_2+h} d\alpha_2(t).$$

Thus, for $h \leq h_0$, the total variation of $\alpha(s; h)$ in $s_1 \leq s \leq s_2$ is uniformly bounded, and there exists, therefore, by a theorem of E. Helly, a sequence $h_n \rightarrow 0$ as $n \rightarrow \infty$, such that, for $\bar{\alpha}_n(s) \equiv \alpha(s; h_n)$:

$$\int_{s_1}^{s_2} |\bar{\alpha}'_n(s)| ds = \int_{s=s_1}^{s_2} |d\bar{\alpha}_n(s)| \rightarrow \int_{s=s_1}^{s_2} |d\alpha(s)| \quad \text{as } n \rightarrow \infty.$$

Let $w_1^* = w(s_1)$, $w_2^* = w(s_2)$. Then we set

$$w_n(s) = w_1^* + \frac{w_2^* - w_1^*}{g_n} \int_{s_1}^s e^{i\bar{\alpha}_n(t)} dt + (a_n s + b_n)(s - s_1)(s - s_2),$$

where $g_n = \int_{s_1}^{s_2} e^{i\bar{\alpha}_n(t)} dt$ and a_n and b_n are so determined that $w'_n(s_1) = w'(s_1)$, $w'_n(s_2) = w'(s_2)$. It is easily seen that $\lim_{n \rightarrow +\infty} g_n = w_2^* - w_1^*$, $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = 0$ and that the $w_n(s)$ satisfy the three conditions (i), (ii), (iii) of the text.

into $Z(w_0)$. Let $x_{*1}^{(n)}$ and $x_2^{*(n)}$ be defined for S_n as x_{*1} and x_2^* are for S . Then⁽¹⁹⁾

$$(9.17) \quad \lim_{n \rightarrow \infty} x_{*1}^{(n)} = x_{*1} \quad \text{and} \quad \lim_{n \rightarrow \infty} x_2^{*(n)} = x_2^*.$$

For sufficiently large n , S_n will satisfy the condition stated at the end of part 1 of this proof, since S satisfies it. Moreover, since β_n has continuous curvature, we have by (9.2)

$$-\frac{\pi}{\cos^3(2\epsilon)} \int_{s=s_1}^{s_2} |d\alpha_n(s)| + \pi \int_{s_1}^{s_2} \frac{d\sigma_n(s)}{\Delta_n(s)} \leq x_2^{*(n)} - x_{*1}^{(n)}.$$

Letting $n \rightarrow \infty$ we find because of (9.14), (9.16), and (9.17),

$$-\frac{\pi}{\cos^3(2\epsilon)} \int_{s=s_1}^{s_2} |d\alpha(s)| + \pi \int_{s_1}^{s_2} \frac{ds}{\Delta(s)} \leq x_2^* - x_{*1}.$$

Finally, to prove (9.1) we need only observe that by Lemma 1 $x_2^* - x_2 \rightarrow 0$ and $x_1 - x_{*1} \rightarrow 0$ as $s_1 \rightarrow \infty$ and that ϵ may be taken arbitrarily small, provided s_1 is chosen sufficiently large. This completes the proof of Theorem V.

10. A definition and further lemmas. Theorem V will enable us to obtain a lower bound for the difference $x_2 - x_1$ of the type mentioned in §7. We introduce first the following definition.

DEFINITION. Let S be an L -strip in the w -plane with the boundary inclination γ , $|\gamma| < \pi/2$, at $u = +\infty$. Let $v = \phi_+(u)$ and $v = \phi_-(u)$ represent the boundary curves C_+ and C_- of S , respectively. We shall say that S is a strip with finite boundary turning at $u = +\infty$ if $\phi'_+(u)$ and $\phi'_-(u)$ are continuous for all sufficiently large u , say $u \geq u_1$, and if the integrals

$$\int_{u=u_1}^{\infty} |d\phi'_+(u)|, \quad \int_{u=u_1}^{\infty} |d\phi'_-(u)|$$

converge.

Our object will be accomplished by the following lemma.

LEMMA 4. Let S be an L -strip in the w -plane with the boundary inclination γ , $|\gamma| < \pi/2$, and with finite boundary turning at $u = +\infty$. If $\Delta(s)$ and $w(s)$ are defined as in §8, then for $u = \mathcal{R}w(s)$ and $s_1 < s_2$ ($u_1 = \mathcal{R}w(s_1)$, $u_2 = \mathcal{R}w(s_2)$)

$$(10.1) \quad \int_{s_1}^{s_2} \frac{ds}{\Delta(s)} = \int_{u_1}^{u_2} \frac{1 + \psi'^2(u)}{\theta(u)} du - \frac{1}{4} \int_{u_1}^{u_2} \frac{\theta'^2(u)}{\theta(u)} du + o(1) \quad \text{as } s_1 \rightarrow \infty.$$

Proof. Denote the point $w(s)$ on C_- by A , the other end point of Δ_s by B and the point $u + i\phi_+(u)$, where $u = \mathcal{R}w(s)$, by C . Suppose, for the present, that B lies in the half-plane to the left of AC . In the triangle ABC the angle

⁽¹⁹⁾ This follows from the theorem of Radó used in the proof of (6.9); see (12).

$A = \alpha = \alpha(s)$. To find the angle C , observe that by the mean value theorem there exists a point $w^* = u^* + i\phi_+(u^*)$ on the arc BC of C_+ such that the tangent to C_+ at w^* is parallel to the chord BC . If the angle of inclination of this tangent is β^* , $|\beta^*| < \pi/2$, then the angle $C = \pi/2 - \beta^*$. Let β , $|\beta| < \pi/2$, denote the angle of inclination of the tangent to C_+ at C . By the law of sines we obtain from the triangle ABC :

$$(10.2) \quad \frac{AC}{AB} = \frac{\theta(u)}{\Delta(s)} = \frac{\sin(A + C)}{\sin C} = \frac{\cos(\beta^* - \alpha)}{\cos \beta^*} = \frac{\cos[(\beta - \alpha) + (\beta^* - \beta)]}{\cos[\beta + (\beta^* - \beta)]}.$$

Since α, β, β^* approach γ as $s \rightarrow \infty$ and since $|\gamma| < \pi/2$, we may write (10.2) in the form

$$\frac{\theta(u)}{\Delta(s)} = \frac{\cos(\beta - \alpha)}{\cos \beta} (1 + O(|\beta^* - \beta|)) \quad \text{as } s \rightarrow \infty.$$

Now

$$|\beta^* - \beta| \leq |\tan \beta^* - \tan \beta| \leq \left| \int_{t=u}^{u^*} |d\phi'_+(t)| \right|.$$

Moreover,

$$|u - u^*| \leq BC = \frac{\sin A}{\sin B} \theta(u) < \theta(u)$$

for sufficiently large u , since $B \rightarrow \pi/2$ and $A \rightarrow |\gamma|$ as $u \rightarrow +\infty$. Hence

$$(10.3) \quad |\beta^* - \beta| \leq \Phi_+(u) \equiv \int_{t=u-\theta(u)}^{u+\theta(u)} |d\phi'_+(t)|.$$

The quotient

$$\frac{\cos(\beta - \alpha)}{\cos \beta \cos \alpha} = 1 + \tan \alpha \tan \beta = 1 + \phi'_+(u)\phi'_-(u) = 1 + \psi'^2(u) - \left[\frac{\theta'(u)}{2} \right]^2,$$

so that

$$\frac{1}{\Delta(s)} = \frac{\cos \alpha}{\theta(u)} (1 + \psi'^2(u)) - \frac{\cos \alpha}{\theta(u)} \left[\frac{\theta'(u)}{2} \right]^2 + O\left(\frac{\Phi_+(u)}{\theta(u)} \cos \alpha \right)$$

as $u \rightarrow +\infty$. This relation is also easily verified if B lies to the right of AC . Observing now that $\cos \alpha \cdot ds = du$, we obtain

$$\int_{s_1}^{s_2} \frac{ds}{\Delta(s)} = \int_{u_1}^{u_2} \frac{1 + \psi'^2(u)}{\theta(u)} du - \frac{1}{4} \int_{u_1}^{u_2} \frac{\theta'^2(u)}{\theta(u)} du + O\left\{ \int_{u_1}^{u_2} \frac{\Phi_+(u)}{\theta(u)} du \right\}.$$

The result (10.1) follows now immediately from

LEMMA 5. *Under the hypotheses of Lemma 4 the integral*

$$(10.4) \quad \int^{\infty} \frac{\Phi_+(u)}{\theta(u)} du \text{ converges, } \Phi_+(u) \equiv \int_{t=u-\theta(u)}^{u+\theta(u)} |d\phi_+(t)|.$$

Proof. We have for $b < c$:

$$I \equiv \int_b^c \frac{\Phi_+(u)}{\theta(u)} du = \int_b^c \frac{1}{\theta(u)} \left[\int_{t=u}^{u+\theta(u)} |d\phi_+(t)| \right] du + \int_b^c \frac{1}{\theta(u)} \left[\int_{t=u-\theta(u)}^u |d\phi_+(t)| \right] du = I_1 + I_2.$$

Interchanging of the order of integration in I_1 gives

$$I_1 = \int_{t=b}^c |d\phi_+(t)| \int_{g(t)}^t \frac{du}{\theta(u)} + \int_{t=c}^{c+\theta(c)} |d\phi_+(t)| \int_{g(t)}^c \frac{du}{\theta(u)} \leq \int_{t=b}^{c+\theta(c)} |d\phi_+(t)| \int_{g(t)}^t \frac{du}{\theta(u)},$$

where $u = g(t)$ is the inverse function⁽²⁰⁾ of $t = u + \theta(u)$ for $b + \theta(b) \leq t \leq c + \theta(c)$, $u = g(t) \equiv b$ for $b \leq t \leq b + \theta(b)$. Now $0 \leq t - g(t) \leq \theta(g(t))$ for $b \leq t \leq c + \theta(c)$. Hence, by Lemma 7 (which is proved in §15),

$$\theta(u) \geq \frac{1}{2}\theta(g(t)) \quad \text{for } g(t) \leq u \leq t,$$

provided only b is sufficiently large. Thus

$$\int_{g(t)}^t du/\theta(u) \leq 2[(t - g(t))/\theta(g(t))] \leq 2,$$

and hence

$$I_1 \leq 2 \int_{t=b}^{c+\theta(c)} |d\phi_+(t)|.$$

Similarly,

$$I_2 \leq 2 \int_{t=b-\theta(b)}^c |d\phi_+(t)|,$$

if b is sufficiently large, so that

$$I \leq 4 \int_{t=b-\theta(b)}^{c+\theta(c)} |d\phi_+(t)|.$$

Hence (10.4) follows from the hypothesis that S has finite boundary turning at $u = +\infty$.

11. The second basic inequality. We are now in the position to prove our second basic inequality.

⁽²⁰⁾ The inverse function of $t = u + \theta(u)$ exists if b is sufficiently large, since $dt/du = 1 + \theta'(u)$, and therefore approaches 1 as $u \rightarrow +\infty$.

THEOREM VI. *Let S be an L -strip in the w -plane with the boundary inclination $\gamma=0$ and with finite boundary turning at $u=+\infty$. Suppose that $z=Z(w)=X(w)+iY(w)$ maps S conformally onto the strip $|y|<\pi/2$ and that $\lim_{u\rightarrow+\infty} X(w)=+\infty$. Let $w_1=u_1+iv_1, w_2=u_2+iv_2, u_1<u_2$, be two points in \bar{S} and let $x_1=X(w_1), x_2=X(w_2)$. Then*

$$(11.1) \quad x_2 - x_1 \geq \pi \int_{u_1}^{u_2} \frac{1 + \psi'^2(u)}{\theta(u)} du - \frac{\pi}{4} \int_{u_1}^{u_2} \frac{\theta'^2(u)}{\theta(u)} du + o(1)$$

as u_1 and hence u_2 approach $+\infty$, uniformly in v_1 and v_2 .

Proof. Let $w(s)$ be defined as in §8 (b), and let $\Re w(s_k) = u_k, X(u_k + i\phi_-(u_k)) = x'_k$ ($k=1, 2$). By Theorem V we have, since $\int_{t=u_1}^{\infty} |d\phi'_-(t)| < \infty$,

$$(11.2) \quad x'_2 - x'_1 \geq \pi \int_{s_1}^{s_2} \frac{ds}{\Delta(s)} + o(1) \text{ as } s_1 \text{ and hence } s_2 \rightarrow \infty.$$

By Lemma 1, $\lim_{u_2 \rightarrow +\infty} (x_2 - x'_2) = 0$ and $\lim_{u_1 \rightarrow +\infty} (x_1 - x'_1) = 0$ so that the difference $x'_2 - x'_1$ in (11.2) may be replaced by $x_2 - x_1 + o(1)$. The result (11.1) follows now immediately from Lemma 4.

IV. A DISTORTION THEOREM

12. A lemma. *If in Theorems IV and VI it is assumed that the integral*

$$\int_{u_1}^{\infty} \frac{\theta'^2(u)}{\theta(u)} du \text{ converges,}$$

then their combination yields an asymptotic expression for $x_2 - x_1$,

$$(12.1) \quad x_2 - x_1 = \int_{u_1}^{u_2} \frac{1 + \psi'^2(u)}{\theta(u)} du + o(1) \text{ as } u_1 \text{ and } u_2 \rightarrow +\infty.$$

One of the hypotheses under which this formula is obtained is that the boundary inclination of S at $u=+\infty$ is $\gamma=0$. It is easy to modify it so as to obtain a result which holds for any $\gamma, |\gamma|<\pi/2$. For this purpose we prove first the following lemma.

LEMMA 6. *Let S be an L -strip in the w -plane ($w=u+iv$) with the boundary inclination $\gamma, |\gamma|<\pi/2$, and with finite boundary turning at $u=+\infty$. Let $v=\phi_+(u)$ and $v=\phi_-(u)$ represent its boundary curves C_+ and C_- , respectively. Suppose that a new set of coordinates \bar{u}, \bar{v} ($\bar{u}+i\bar{v}=\bar{w}$) is introduced by means of the rotation $\bar{w}=we^{-i\gamma}$ and that, for sufficiently large \bar{u} , C_+ and C_- are represented in the new coordinate system by $\bar{v}=\bar{\phi}_+(\bar{u})$ and $\bar{v}=\bar{\phi}_-(\bar{u})$, respectively. Set*

$$\theta(u) = \phi_+(u) - \phi_-(u), \quad \bar{\theta}(\bar{u}) = \bar{\phi}_+(\bar{u}) - \bar{\phi}_-(\bar{u}).$$

If u and \bar{u} are connected by the relation

$$(12.2) \quad \bar{u} = u \cos \gamma + \phi_-(u) \sin \gamma,$$

then, for sufficiently large u ,

$$(12.3) \quad |\bar{\theta}'(\bar{u})| \leq 2 \{ |\theta'(u)| + \Phi_+(u) \}, \quad \Phi_+(u) \equiv \int_{t=u-\theta(u)}^{u+\theta(u)} |d\phi'_+(t)|.$$

Thus, the convergence of the integral

$$\int^\infty \frac{\theta'^2(u)}{\theta(u)} du$$

implies that of

$$\int^\infty \frac{\bar{\theta}'^2(\bar{u})}{\bar{\theta}(\bar{u})} d\bar{u}$$

(by (8.1) and Lemma 5).

Proof. Let u and \bar{u} satisfy (12.2). We denote the points $\bar{u} + i\bar{\phi}_-(\bar{u})$ and $\bar{u} + i\bar{\phi}_+(\bar{u})$ of the \bar{w} -plane by A and B , respectively. The coordinates of A and B in the w -plane are then $u + i\phi_-(u)$ and $u_1 + i\phi_+(u_1)$, respectively (where u_1 is determined by the relation $e^{-i\gamma}(u_1 + i\phi_+(u_1)) = \bar{u} + i\bar{\phi}_+(\bar{u})$). Finally, we call C the point $u + i\phi_+(u)$ of the w -plane. In the triangle ABC , $AB = \bar{\theta}(\bar{u})$, $AC = \theta(u)$ and the angle $A = |\gamma|$.

Let $\alpha_+(u)$ and $\alpha_-(u)$ denote in the w -plane the angles of inclination of the tangents to C_+ and to C_- , respectively, at a point with the abscissa u . Then

$$\begin{aligned} \phi'_-(u) &= \tan \alpha_-(u), & \phi'_+(u) &= \tan \alpha_+(u); \\ \bar{\phi}'_-(\bar{u}) &= \tan \{ \alpha_-(u) - \gamma \}, & \bar{\phi}'_+(\bar{u}) &= \tan \{ \alpha_+(u_1) - \gamma \}. \end{aligned}$$

Hence,

$$\bar{\theta}'(\bar{u}) = \tan \{ \alpha_+(u_1) - \alpha_-(u) \} \cdot \{ 1 + \bar{\phi}'_-(\bar{u})\bar{\phi}'_+(\bar{u}) \}.$$

Since $\bar{\phi}'_-(\bar{u})$ and $\bar{\phi}'_+(\bar{u})$ approach 0 as $\bar{u} \rightarrow +\infty$, we may write:

$$(12.4) \quad |\bar{\theta}'(\bar{u})| \leq \{ |\theta'(u)| + |\phi'_+(u_1) - \phi'_+(u)| \} (1 + o(1))$$

as $u \rightarrow +\infty$. Now

$$|u_1 - u| \leq BC = \frac{\sin A}{\sin B} \theta(u) \leq \theta(u)$$

for sufficiently large u , since the angle $B \rightarrow \pi/2$ and $A = |\gamma| < \pi/2$. Hence, $|\phi'_+(u_1) - \phi'_+(u)| \leq \left| \int_{t=u}^{u_1} |d\phi'_+(t)| \right| \leq \Phi_+(u)$, and (12.3) follows from (12.4).

13. **First form of the distortion theorem.** We prove now

THEOREM VII. *Let S be an L -strip with the boundary inclination γ , $|\gamma| < \pi/2$ and with finite boundary turning at $u = +\infty$. Moreover let the integral*

$$(13.1) \quad \int^{\infty} \frac{\theta'^2(u)}{\theta(u)} du \text{ be convergent}^{(21)}.$$

Suppose that $z = Z(w) = X(w) + iY(w)$ maps S conformally onto the strip $|y| < \pi/2$ in such a manner that $\lim_{u \rightarrow +\infty} X(w) = +\infty$. Let Δ_s , $\Delta(s)$ and $w(s)$ be defined as in §8. Then, if $s_1 < s_2$, $w_1 \in \Delta_{s_1}$, $w_2 \in \Delta_{s_2}$,

$$(13.2) \quad X(w_2) - X(w_1) = \pi \int_{s_1}^{s_2} \frac{ds}{\Delta(s)} + o(1) \quad \text{as } s_1 \text{ and } s_2 \rightarrow \infty,$$

uniformly with respect to w_1 and w_2 .

Proof. As in Lemma 6, let there be introduced a new set of axes (\bar{u}, \bar{v}) , $\bar{u} + i\bar{v} = \bar{w}$, by means of the rotation $\bar{w} = e^{-i\gamma}w$. Let $\bar{\phi}_+(u)$, $\bar{\phi}_-(u)$, $\bar{\theta}(\bar{u})$ be defined as in Lemma 6 and let u and \bar{u} be two numbers related by equation (12.2). Then, by Lemma 6, (13.1) implies that

$$(13.3) \quad \int^{\infty} \frac{\bar{\theta}'^2(\bar{u})}{\bar{\theta}(\bar{u})} d\bar{u}$$

converges. Let \bar{u}_1 and \bar{u}_2 be the abscissas of the points $w(s_1)$ and $w(s_2)$ (on C_-) in the (\bar{u}, \bar{v}) -system, and let $x'_1 = X(w(s_1))$, $x'_2 = X(w(s_2))$. If $\bar{\psi}(\bar{u}) = \frac{1}{2}[\bar{\phi}_+(\bar{u}) + \bar{\phi}_-(\bar{u})]$, we have by Theorem IV,

$$x'_2 - x'_1 \leq \pi \int_{\bar{u}_1}^{\bar{u}_2} \frac{1 + \bar{\psi}'^2(\bar{u})}{\bar{\theta}(\bar{u})} d\bar{u} + \frac{\pi}{12} \int_{\bar{u}_1}^{\bar{u}_2} \frac{\bar{\theta}'^2(\bar{u})}{\bar{\theta}(\bar{u})} d\bar{u} + o(1),$$

as s_1 and $s_2 \rightarrow \infty$. By Lemma 4 and by (13.3) we obtain

$$(13.4) \quad x'_2 - x'_1 \leq \int_{s_1}^{s_2} \frac{ds}{\Delta(s)} + o(1) \quad \text{as } s_1, s_2 \rightarrow \infty.$$

Finally, application of Lemma 1 shows that $\lim_{s_1 \rightarrow \infty} (x_1 - x'_1) = \lim_{s_1 \rightarrow \infty} (x_2 - x'_2) = 0$ ($x_k = X(w_k)$), so that the difference $x'_2 - x'_1$ in (13.4) may be replaced by $x_2 - x_1$. Combining this result with Theorem V, we obtain (13.2).

14. Second form of the distortion theorem. For some applications the following form of Theorem VII will be more convenient.

THEOREM VIII. *Let S be an L -strip satisfying all the hypotheses of Theorem VII, and let $z = Z(w)$ be defined as in that theorem. Then for $w_1 = u_1 + iv_1$ and $w_2 = u_2 + iv_2$ in \bar{S} , $u_1 < u_2$,*

⁽²¹⁾ Condition (13.1) is automatically satisfied for L -strips with finite boundary turning at $u = +\infty$ for which $0 < c_1 \leq \theta(u) \leq c_2$ (c_1, c_2 constants). For

$$\int_a^b \frac{\theta'^2(u)}{\theta(u)} du = \theta'(u) \log \theta(u) \Big|_a^b - \int_{u=a}^b \log \theta(u) d\theta'(u).$$

Since $|\log \theta(u)|$ is bounded and $\int_{u=a}^{\infty} |d\theta'(u)|$ exists, (13.1) follows.

$$X(w_2) - X(w_1) = \pi \int_{u_1}^{u_2} \frac{1 + \psi'^2(u)}{\theta(u)} du + \pi \left[\frac{v_2 - \psi(u_2)}{\theta(u_2)} - \frac{v_1 - \psi(u_1)}{\theta(u_1)} \right] \tan \gamma + o(1),$$

as u_1 and $u_2 \rightarrow +\infty$, uniformly with respect to v_1 and v_2 .

Proof. Let Δ_s , $\Delta(s)$ and $w(s)$ be defined as in §8. Let Δ_{s_1} and Δ_{s_2} be two normals of C_- which pass through w_1 and w_2 , respectively. By Theorem VII,

$$X(w_2) - X(w_1) = \pi \int_{s_1}^{s_2} \frac{ds}{\Delta(s)} + o(1) \quad \text{as } s_1, s_2 \rightarrow \infty.$$

Let $\mathcal{R}w(s_1) = u'_1$, $\mathcal{R}w(s_2) = u'_2$. By Lemma 4 and (13.1) this equals

$$\pi \int_{u'_1}^{u'_2} \frac{1 + \psi'^2(u)}{\theta(u)} du + o(1), \quad \text{as } u_1, u_2 \rightarrow +\infty,$$

or

$$(14.1) \quad X(w_2) - X(w_1) = \pi \int_{u'_1}^{u_1} + \pi \int_{u_1}^{u_2} + \pi \int_{u_2}^{u'_2} + o(1).$$

We estimate the first and third of the integrals in (14.1). Assume, for the present, that $\gamma > 0$. Denote the points $u_1 + i\phi_-(u_1)$, $w(s_1)$ and w_1 by A , B , and C , respectively. In the triangle ABC the side $AC = v_1 - \phi_-(u_1)$, the angle $C \rightarrow \gamma$ and the angle $B \rightarrow \pi/2$ as $u_1 \rightarrow +\infty$. These limits, as well as all following in this proof (taken as $u_1 \rightarrow +\infty$) exist uniformly with respect to the position of w_1 on θ_{u_1} . Now

$$u'_1 - u_1 = BC \sin(\gamma + \epsilon_1), \quad \lim_{u_1 \rightarrow +\infty} \epsilon_1 = 0,$$

$$BC = AC \frac{\sin A}{\sin B} = AC \frac{\cos(\gamma + \epsilon_2)}{\sin B}, \quad \lim_{u_1 \rightarrow +\infty} \epsilon_2 = 0,$$

and hence

$$(14.2) \quad u'_1 - u_1 = [v_1 - \phi_-(u_1)] \sin(\gamma + \epsilon_1) \cos(\gamma + \epsilon_2) \frac{1}{\sin B}.$$

This result is also easily verified when $\gamma \leq 0$.

Now, by the law of the mean

$$\int_{u_1}^{u'_1} \frac{1 + \psi'^2(u)}{\theta(u)} du = (u'_1 - u_1) \frac{1 + \psi'^2(u^*)}{\theta(u^*)} \quad (u^* \text{ between } u_1 \text{ and } u'_1).$$

Since, by (14.2), $|u'_1 - u_1| \leq \theta(u_1)$ for sufficiently large u_1 , it follows from Lemma 7 (which is proved in §15) that $(\theta(u^*)/\theta(u_1)) \rightarrow 1$ as $u_1 \rightarrow +\infty$. Furthermore, as $u_1 \rightarrow +\infty$, $1 + \psi'^2(u^*) \rightarrow 1/\cos^2 \gamma$. Hence

$$\int_{u_1}^{u'} \frac{1 + \psi'^2(u)}{\theta(u)} du = \frac{v_1 - \phi_-(u_1)}{\theta(u_1)} \tan \gamma + o(1)$$

$$= \left[\frac{v_1 - \psi(u_1)}{\theta(u_1)} + \frac{1}{2} \right] \tan \gamma + o(1)$$

as $u_1 \rightarrow +\infty$. An analogous result is obtained for the third integral in (14.1). Substitution of these expressions into (14.1) proves the theorem.

COROLLARY OF THEOREM VIII. *Under the hypotheses of the theorem, for $w = u + iv \in \bar{S}$,*

$$(14.3) \quad \lim_{u \rightarrow +\infty} \left\{ X(w) - \left[\pi \int_{u_0}^u \frac{1 + \psi'^2(t)}{\theta(t)} dt + \pi \frac{v - \psi(u)}{\theta(u)} \tan \gamma \right] \right\} = \lambda$$

exists uniformly in v and is finite.

For, if the difference within the braces in (14.3) is denoted by $A(w)$, Theorem VIII states that for any given $\epsilon > 0$ there exists an $N(\epsilon)$ such that for all w_1 and w_2 in S for which $\Re w_2 > \Re w_1 > N(\epsilon)$:

$$|A(w_2) - A(w_1)| < \epsilon.$$

THEOREM IX. *Let S be an L -strip satisfying the hypotheses of Theorem VII and let $z = Z(w)$ be defined as in that theorem. Then, for $w = u + iv \in \bar{S}$,*

$$Z(w) = \lambda + \pi \int_{u_0}^u \frac{1 + \psi'^2(t)}{\theta(t)} dt + \pi \frac{v - \psi(u)}{\theta(u)} \tan \gamma + i\pi \frac{v - \psi(u)}{\theta(u)} + o(1),$$

as $u \rightarrow +\infty$, uniformly in v . λ is a real constant.

This result follows immediately by combining (14.3) with the result (18.1) of Corollary 1 of Theorem X (which is proved in §18).

V. ASYMPTOTIC BEHAVIOR OF THE MAPPING FUNCTION OF AN L -STRIP AND OF ITS DERIVATIVE

15. Preliminary remarks and lemmas. We shall establish now asymptotic expressions for $Z(w)$ and $Z'(w)$ under the mere assumption that S is an L -strip. These results will be less sharp than those of Part IV. While in Part IV (Theorem IX) we obtained, under more restrictive assumptions, an expression $f(w)$ such that the difference $Z(w) - f(w)$ approaches a finite limit, as $u \rightarrow +\infty$, we shall find here expressions for $Z(w)$ and $Z'(w)$ which represent these functions merely in the sense of *asymptotic equivalence* (that is, the *quotient* of the function in question and its asymptotic expression approaches 1 as $u \rightarrow +\infty$). None of the results of Parts III and IV will be used here.

Throughout this part we assume S to be an L -strip in the w -plane with the boundary inclination γ , $|\gamma| < \pi/2$ at $u = +\infty$, and $Z(w) = X(w) + iY(w)$

a function which maps S conformally onto the strip $|y| < \pi/2$ in such a manner that $\lim_{u \rightarrow +\infty} X(w) = +\infty$. The inverse of $z = Z(w)$ will be denoted by $w = W(z) = U(z) + iV(z)$.

We shall make use of the following simple lemma.

LEMMA 7. *Let, for $u_0 < u_1 < u_2$,*

$$\theta^*(u_1, u_2) = \max_{u_1 \leq u \leq u_2} \theta(u), \quad \theta_*(u_1, u_2) = \min_{u_1 \leq u \leq u_2} \theta(u)$$

and let

$$(15.1) \quad u_2 - u_1 \leq k\theta^*(u_1, u_2), \quad k \text{ a constant.}$$

Then, uniformly for all u_2 ,

$$\lim_{u_1 \rightarrow +\infty} \frac{\theta_*(u_1, u_2)}{\theta^*(u_1, u_2)} = 1.$$

Proof. Let $\theta^*(u_1, u_2) = \theta(b)$ and $\theta_*(u_1, u_2) = \theta(c)$, $u_1 \leq b$, $c \leq u_2$. If u_1 is sufficiently large, we have

$$|\theta(b) - \theta(c)| = \left| \int_c^b \theta'(u) du \right| \leq \int_{u_1}^{u_2} |\theta'(u)| du \leq (u_2 - u_1) \sup_{u_1 \leq u \leq u_2} |\theta'(u)|,$$

(where $\sup_{u_1 \leq u \leq u_2} |\theta'(u)|$ denotes the least upper bound of $|\theta'(u)|$ in $u_1 \leq u \leq u_2$). Hence by use of (15.1):

$$0 \leq 1 - \frac{\theta_*(u_1, u_2)}{\theta^*(u_1, u_2)} \leq \frac{u_2 - u_1}{\theta^*(u_1, u_2)} \sup_{u_1 \leq u \leq u_2} |\theta'(u)| \leq k \sup_{u_1 \leq u \leq u_2} |\theta'(u)| \rightarrow 0,$$

as $u_1 \rightarrow +\infty$, by hypothesis. This proves the lemma.

COROLLARY. *Let*

$$(15.2) \quad 0 \leq x_2 - x_1 \leq c, \quad c > 0, \text{ a constant,}$$

and $u_1 = \min_{|y| \leq \pi/2} U(x_1 + iy)$, $u_2 = \max_{|y| \leq \pi/2} U(x_2 + iy)$. Then, uniformly for all x_2 , satisfying (15.2),

$$\lim_{x_1 \rightarrow +\infty} \frac{\theta^*(u_1, u_2)}{\theta_*(u_1, u_2)} = 1.$$

Proof. From Theorem III (a) we have

$$\pi \int_{u_1}^{u_2} \frac{du}{\theta(u)} \leq x_2 - x_1 + 4\pi \leq c + 4\pi$$

by (15.2). On the other hand, since $u_2 \geq u_1$,

$$\int_{u_1}^{u_2} \frac{du}{\theta(u)} \geq \frac{u_2 - u_1}{\theta^*(u_1, u_2)},$$

and hence $0 \leq u_2 - u_1 \leq \{(4\pi + c)/\pi\} \theta^*(u_1, u_2)$. Thus the result follows from Lemma 7

16. **Asymptotic expressions for $Z(w)$ and $Z'(w)$.** We prove now the following

THEOREM X. *If S is an L -strip with boundary inclination 0 at $u = +\infty$, we have*

(i) *For $w = u + iv$ in S , uniformly in v ,*

$$\lim_{u \rightarrow +\infty} \frac{Z(w)}{\int_{u_0}^u \frac{dt}{\theta(t)}} = \pi, \quad u_0 \text{ defined as in } \S 1 \text{ (a)}.$$

(ii) *Uniformly in any fixed strip $|y| \leq \beta < \pi/2$ ($z = x + iy$)⁽²²⁾,*

$$\lim_{x \rightarrow +\infty} \frac{|W'(z)|}{\theta(u)} = \frac{1}{\pi} \quad (u = U(z)), \quad \lim_{x \rightarrow +\infty} \frac{\log |W'(z)|}{|z|} = 0.$$

(iii) *The straight line $\Lambda_y: y = \text{const.}, |y| < \pi/2$, is mapped by $Z(w)$ onto a curve L_y which for sufficiently large u is represented by an equation of the form*

$$(16.1) \quad v = f_y(u) \equiv \psi(u) + \frac{\theta(u)}{\pi} y + o[\theta(u)] \quad \text{as } u \rightarrow +\infty,$$

uniformly in $|y| < \pi/2$ ⁽²³⁾.

(iv) *Let the image of the strip*

$$\Sigma_\alpha: \{ -\infty < x < \infty, |y| \leq \alpha \}, \quad 0 < \alpha < \pi/2,$$

by means of $w = W(z)$ be T_α , and let S_β be the region

$$\left\{ u > u_0, \left| \frac{v - \psi(u)}{\theta(u)} \right| < \frac{\beta}{\pi} \right\}, \quad 0 < \beta < \frac{\pi}{2}.$$

If $0 < \alpha \pm \epsilon < \pi/2$, $\epsilon > 0$, then there exists an $N = N(\epsilon; \alpha)$ such that the part of T_α which lies in $u \geq N$, contains that part of $S_{\alpha-\epsilon}$ which is in $u \geq N$ and is contained in $S_{\alpha+\epsilon}$.

⁽²²⁾ Since $\arg W'(z) \rightarrow 0$ as $x \rightarrow +\infty$, the first of these relations implies that even

$$\lim_{x \rightarrow +\infty} \frac{W'(z)}{\theta(u)} = \frac{1}{\pi}.$$

⁽²³⁾ This implies the following fact: If Λ is a curve in the strip $|y| < \pi/2$ which approaches the line Λ_t as an asymptote, as $x \rightarrow +\infty$, then, for any point (u, v) on the image L of Λ in the w -plane, $v = \psi(u) + (\theta(u)/\pi)t + o[\theta(u)]$ as $u \rightarrow +\infty$. (L need not be representable in the form $v = f(u)$.) For, if $\epsilon > 0$ is given, there exists an x_1 such that the part of Λ for $x \geq x_1$ lies "between" the lines $\Lambda_{t-\epsilon}$ and $\Lambda_{t+\epsilon}$ (if $|t \pm \epsilon| < \pi/2$; in case $t = \pi/2$ or $t = -\pi/2$, it lies "between" $\Lambda_{(\pi/2)-\epsilon}$ and $\Lambda_{\pi/2}$ or "between" $\Lambda_{-(\pi/2)}$ and $\Lambda_{-(\pi/2)+\epsilon}$, respectively). The image of that part of Λ lies in a certain half-plane $u \geq u_1$ "between" the curves $L_{t-\epsilon}$ and $L_{t+\epsilon}$ (or $L_{(\pi/2)-\epsilon}$, $L_{\pi/2}$ or $L_{-\pi/2}$, $L_{-(\pi/2)+\epsilon}$, respectively). Since ϵ may be taken arbitrarily small, this proves the above assertion.

(v) In any region S_β , $0 < \beta < \pi/2$, β fixed,

$$(16.2) \quad \lim_{u \rightarrow +\infty} [|Z'(w)| \theta(u)] = \pi, \quad w = u + iv, w \in S_\beta.$$

REMARK. Part (iii) may be considered as an extension in a certain direction of Carathéodory's well known result, which states that the map of the interior of a closed Jordan curve Γ onto the circle $|\zeta| < 1$ is "quasi-conformal" at a boundary point ω_0 with a corner of measure $\beta > 0$, i.e., that the angles at corresponding boundary points ω_0 and ζ_0 are transformed proportionally⁽²⁴⁾. Transformation of the interior of Γ onto a strip S by means of the function $w = \log [1/(\omega - \omega_0)]$ and of $|\zeta| < 1$ onto the strip $|y| < \pi/2$, in such a way that $\zeta = \zeta_0$ corresponds to $x = +\infty$, leads to the following statement of Carathéodory's theorem: Let C be a closed Jordan curve through $w = \infty$ which, in a neighborhood of $w = \infty$, consists of two branches C_+ and C_- having the lines $v = \phi_+$ and $v = \phi_-$, respectively, for asymptotes as $u \rightarrow +\infty$ ($\phi_+ - \phi_- = \theta > 0$). If $W(z) = U(z) + iV(z)$ maps $|y| < \pi/2$ onto the interior S of C and if $\lim_{x \rightarrow +\infty} U(z) = +\infty$, then

$$(16.3) \quad V(z) = \frac{1}{2}[\phi_+ + \phi_-] + \frac{\theta}{\pi} y + o(1) \quad \text{as } x \rightarrow +\infty,$$

uniformly in $|y| < \pi/2$.

If S is a simple Jordan strip, the hypothesis regarding the asymptotes of C_+ and C_- means that

$$\lim_{u \rightarrow +\infty} \phi_+(u) = \phi_+ \quad \text{and} \quad \lim_{u \rightarrow +\infty} \phi_-(u) = \phi_- \quad \text{exist.}$$

Similarly, Part (ii) can be considered as an extension in a certain direction of a theorem of Ostrowski which (formulated for infinite strips) states: Under the hypothesis of Carathéodory's theorem

$$\lim_{x \rightarrow +\infty} |W'(z)| = \frac{\theta}{\pi}, \quad \lim_{x \rightarrow +\infty} \left[\frac{\log |W'(z)|}{|z|} \right] = 0,$$

uniformly in any strip $|y| \leq \beta < \pi/2$. Here θ may be greater than or equal to 0⁽²⁵⁾.

⁽²⁴⁾ Carathéodory [1, pp. 40-41] and [2, pp. 19-93]. See also Lindelöf [1, p. 87]. Carathéodory's theorem has been generalized in two other directions: 1. The assumption that C_+ and C_- approach distinct asymptotes has been replaced by the condition that C_+ and C_- "oscillate" within the strips $\phi_+ - k_+ \leq v \leq \phi_+ + k_+$ and $\phi_- - k_- \leq v \leq \phi_- + k_-$ respectively (k_+ , k_- constants). 2. It has been extended to the case where C_+ and C_- may be arbitrary continua forming the boundary of S (and not necessarily Jordan curves). For these extensions see Gross, [1, p. 278], Ostrowski [1, pp. 172-174], Wolff [2, p. 42] and [3, p. 46], Warschawski [2, p. 674]. Ostrowski [2, p. 77] gives a necessary and sufficient condition in order that the mapping function of a simply-connected region onto a half-plane preserve angles at an accessible boundary point.

⁽²⁵⁾ Ostrowski [1]; the first of these equations is his relation (11.3) of page 101, the second follows, for $\theta > 0$, by combination of his relations (62.2) and (62.1) of page 174, and, for $\theta = 0$, by combination of his relations (68.7) of page 185 and (62.1) of page 174.

Both of these theorems do not require that C_+ and C_- have an L -tangent at $u = +\infty$, as our Theorem X does. Our extension, however, refers to the fact that (iii) might be substituted for (16.3) when C_+ and C_- have *no* asymptotes and that in (ii) the *asymptotic behavior* of $|W'(z)|$ at $x = +\infty$ is given, whether $\theta(u) = \phi_+(u) - \phi_-(u)$ approaches a limit as $u \rightarrow +\infty$ or not.

17. **Proof of Theorem X.** (i) By hypothesis, $\psi'(u)$ and $\theta'(u)$ approach 0 as $u \rightarrow +\infty$. Hence it follows from Theorem IV (b) that there exists, for every $\epsilon > 0$, an $N_1(\epsilon) \geq u_0$ such that for all $w = u + iv$ and $w_1 = u_1 + iv_1$ in S for which $u \geq u_1 \geq N_1(\epsilon)$:

$$X(w) - X(w_1) \leq \pi(1 + \epsilon) \int_{u_1}^u \frac{dt}{\theta(t)} + \epsilon \leq \pi(1 + \epsilon) \int_{u_0}^u \frac{dt}{\theta(t)} + \epsilon.$$

Hence,

$$\frac{X(w) - X(w_1)}{\int_{u_0}^u \frac{dt}{\theta(t)}} \leq \pi(1 + \epsilon) + \frac{\epsilon}{\int_{u_0}^u \frac{dt}{\theta(t)}}.$$

Keeping here w_1 fixed and letting $u \rightarrow +\infty$ we find, since $\int_{u_0}^u [\theta(t)]^{-1} dt \rightarrow \infty$ as $u \rightarrow +\infty$,

$$\limsup_{u \rightarrow +\infty} \frac{X(w)}{\int_{u_0}^u \frac{dt}{\theta(t)}} \leq \pi(1 + \epsilon), \quad \text{uniformly with respect to } v.$$

Since the left-hand side of this inequality is *independent* of ϵ , we may let $\epsilon \rightarrow 0$ and find

$$(17.1) \quad \limsup_{u \rightarrow +\infty} \frac{X(w)}{\int_{u_0}^u \frac{dt}{\theta(t)}} \leq \pi.$$

On the other hand, by Theorem III (a), uniformly with respect to v ,

$$(17.2) \quad \liminf_{u \rightarrow +\infty} \frac{X(w)}{\int_{u_0}^u \frac{dt}{\theta(t)}} \geq \pi,$$

and (17.1) and (17.2) together imply that, uniformly in v ,

$$(17.3) \quad \lim_{u \rightarrow \infty} \frac{X(w)}{\int_{u_0}^u \frac{dt}{\theta(t)}} = \pi.$$

The result (i) follows now from (17.3), if we observe that $Z(w) = X(w) + iY(w)$ and that $|Y(w)| \leq \pi/2$, while $\int_{u_0}^u [\theta(t)]^{-1} dt \rightarrow \infty$ as $u \rightarrow +\infty$.

(ii) Let L denote the image in the w -plane of the real axis of the z -plane by means of $w = W(z)$. Let $z_n = 3n$ ($n = 1, 2, 3, \dots$) and $w_n = W(z_n) = u_n + iv_n$. Observing that $\lim_{u \rightarrow +\infty} \psi'(u) = \lim_{u \rightarrow +\infty} \theta'(u) = 0$, we have, by Theorem IV (b), for all sufficiently large n ,

$$3 = 3(n+1) - 3n \leq 2\pi \int_{u_n}^{u_{n+1}} \frac{du}{\theta(u)} + 1,$$

or

$$(17.4) \quad \pi \int_{u_n}^{u_{n+1}} \frac{du}{\theta(u)} \geq 1.$$

Let $\epsilon > 0$ be given. By Theorem IV (b) there exists an $n_0(\epsilon)$ such that for $n \geq n_0(\epsilon)$

$$(17.5) \quad \begin{aligned} X(w_{n+1}) - X(w_n) &\leq \pi \left(1 + \frac{\epsilon}{2}\right) \int_{u_n}^{u_{n+1}} \frac{du}{\theta(u)} + \frac{\epsilon}{2} \\ &\leq \pi(1 + \epsilon) \int_{u_n}^{u_{n+1}} \frac{du}{\theta(u)}, \end{aligned}$$

because of (17.4). Similarly, by Theorem III (b), there is an $n_1(\epsilon) \geq n_0(\epsilon)$ such that for $n \geq n_1(\epsilon)$:

$$(17.6) \quad X(w_{n+1}) - X(w_n) \geq \pi \int_{u_n}^{u_{n+1}} \frac{du}{\theta(u)} - \frac{\epsilon}{1 + \epsilon} \geq \frac{\pi}{1 + \epsilon} \int_{u_n}^{u_{n+1}} \frac{du}{\theta(u)}.$$

Since $\arg W'(z) \rightarrow 0$ as $x \rightarrow +\infty$, the curve L can be represented for sufficiently large u in the form $v = f(u)$ where $f'(u)$ exists and approaches 0 as $u \rightarrow +\infty$. Evidently, for all sufficiently large n , $w_n = u_n + if(u_n)$. The functions $X(u + if(u))$ and $\int_{u_n}^u [\theta(t)]^{-1} dt$ are differentiable for $u_n \leq u \leq u_{n+1}$, and $d/du \int_{u_n}^u [\theta(t)]^{-1} dt > 0$ for $u \geq u_n$. Hence, by the extended mean value theorem, for sufficiently large n , say $n \geq n_2 \geq n_1$:

$$\frac{X(w_{n+1}) - X(w_n)}{\int_{u_n}^{u_{n+1}} \frac{du}{\theta(u)}} = \theta(u'_n) \left[\frac{\partial X}{\partial u} \Big|_{w'_n} + \frac{\partial X}{\partial v} \Big|_{w'_n} f'(u'_n) \right],$$

where $u_n < u'_n < u_{n+1}$ and $w'_n = u'_n + if(u'_n)$. Thus, we obtain from (17.5) and (17.6):

$$\frac{\pi}{1 + \epsilon} \leq \theta(u'_n) \left[\frac{\partial X}{\partial u} \Big|_{w'_n} + \frac{\partial X}{\partial v} \Big|_{w'_n} f'(u'_n) \right] \leq \pi(1 + \epsilon).$$

Since

$$\frac{\partial X}{\partial u} = |Z'(w)| \cos(\arg Z'(w)), \quad \frac{\partial X}{\partial v} = -|Z'(w)| \sin(\arg Z'(w))$$

and

$$\lim_{u \rightarrow +\infty} [\arg Z'(w)] = \lim_{u \rightarrow +\infty} f'(u) = 0,$$

we have, for all sufficiently large n , say $n \geq n_3 \geq n_2$:

$$\frac{\pi}{(1 + \epsilon)^2} \leq \theta(u'_n) |Z'(w'_n)| \leq \pi(1 + \epsilon)^2.$$

Let $z'_n = Z(w'_n)$. Evidently $3n \leq z'_n \leq 3(n+1)$. Since $Z'(w'_n) = 1/W'(z'_n)$ we have also

$$\frac{1}{\pi(1 + \epsilon)^2} \leq \frac{|W'(z'_n)|}{\theta(u'_n)} \leq \frac{(1 + \epsilon)^2}{\pi}.$$

Now let $z = x + iy$ be a point in the strip $|y| \leq \beta$ with $x \geq 3n_3$, and let n be such that $3n \leq x < 3(n+1)$. Since then $|z'_n - x| \leq 3$, we infer from Theorem II (b) that, uniformly for $|y| \leq \beta$,

$$\lim_{x \rightarrow \infty} \left| \frac{W'(z)}{W'(z'_n)} \right| = 1.$$

Moreover, by the corollary of Lemma 7, we have for $u = U(z)$, $3n \leq x < 3(n+1)$,

$$\lim_{u \rightarrow +\infty} \frac{\theta(u)}{\theta(u'_n)} = 1.$$

Hence, if x is sufficiently large,

$$\frac{1}{\pi(1 + \epsilon)^3} \leq \frac{|W'(z)|}{\theta(u)} = \left| \frac{W'(z)}{W'(z'_n)} \right| \frac{|W'(z'_n)|}{\theta(u'_n)} \frac{\theta(u'_n)}{\theta(u)} \leq \frac{(1 + \epsilon)^3}{\pi},$$

and this proves the first relation of part (ii) of the theorem.

To prove the second relation we use the first one:

$$(17.7) \quad \log |W'(z)| = \log \theta(u) - \log \pi + \log(1 + \delta), \quad \lim_{x \rightarrow +\infty} \delta = 0,$$

uniformly in $|y| \leq \beta$. Now, by part (i), uniformly in $|y| < \pi/2$, as $x \rightarrow +\infty$,

$$(17.8) \quad \frac{\log \theta(u)}{|z|} \sim \frac{\int_{u_1}^u \frac{\theta'(t)}{\theta(t)} dt + \log \theta(u_1)}{\pi \int_{u_1}^u \frac{dt}{\theta(t)}}.$$

Here u_1 may be chosen arbitrarily large, but fixed. Let $\epsilon > 0$ be given. We choose u_1 such that $|\theta'(t)| < \epsilon$ for $t \geq u_1$. Then by the mean value theorem

$$\frac{\left| \int_{u_1}^u \frac{\theta'(t)}{\theta(t)} dt \right|}{\int_{u_1}^u \frac{dt}{\theta(t)}} = |\theta'(\xi)| < \epsilon, \quad u_1 < \xi < u.$$

Keeping u_1 fixed and letting $u \rightarrow +\infty$, and hence $x \rightarrow +\infty$, we find from (17.8)

$$\limsup_{x \rightarrow +\infty} \left| \frac{\log \theta(u)}{z} \right| \leq \epsilon,$$

and the result follows from (17.7).

(iii) a. We shall first prove the *weaker* statement that (16.1) holds uniformly for all y with $|y| \leq \beta$, β being any fixed positive number less than $\pi/2$.

The image L_y of Λ_y is represented in parametric form by the equations $u = U(x+iy)$, $v = V(x+iy)$, where x is the parameter, $-\infty < x < \infty$, and y is fixed. Evidently

$$(17.9) \quad V(x+iy) = V(x) + \int_0^y \frac{\partial V(x+i\eta)}{\partial \eta} d\eta.$$

For $\zeta = x+i\eta$, $|\eta| \leq |y|$,

$$\frac{\partial V(x+i\eta)}{\partial \eta} = \Re \left\{ \frac{dW(\zeta)}{d\zeta} \right\} = |W'(\zeta)| \cos(\arg W'(\zeta)),$$

and by Theorem II (a), the first of the relations (ii), and the corollary of Lemma 7, we have for $u = U(x)$

$$\frac{\partial V(\zeta)}{\partial \eta} = \frac{\theta(u)}{\pi} (1 + o(1)) \quad \text{as } u \rightarrow +\infty,$$

uniformly for $|\eta| \leq \beta$. Therefore,

$$(17.10) \quad V(x+iy) = V(x) + \frac{\theta(u)}{\pi} y + o[\theta(u)], \quad \text{as } x \rightarrow +\infty,$$

uniformly for $|y| \leq \beta$.

Since $\arg W'(z) \rightarrow 0$ as $x \rightarrow +\infty$, uniformly in $|y| < \pi/2$, there exists an N' such that for $u \geq N'$ every curve L_y , $|y| < \pi/2$, is representable in the form $v = f_y(u)$, where $df_y(u)/du$ is continuous and approaches 0 as $u \rightarrow +\infty$, uniformly for $|y| < \pi/2$. Let $u = U(x)$, $u' = U(x+iy)$ and let x be so large that $u, u' \geq N'' \geq N'$, where N'' is so chosen that $|df_y(u)/du| < 1$ for $u \geq N''$. Hence⁽²⁶⁾

(26) $\theta^*(u, u') = \max \theta(t)$ for t between u and u' .

$$|f_v(u) - V(x + iy)| = |f_v(u) - f_v(u')| < |u - u'| \leq \theta^*(u, u') \left| \int_u^{u'} \frac{dt}{\theta(t)} \right|.$$

By Theorem III (b), $\int_u^{u'} [\theta(t)]^{-1} dt = o(1)$ and, by the Corollary of Lemma 7, $\theta^*(u, u')/\theta(u) \rightarrow 1$ as $u \rightarrow +\infty$. We find therefore

$$|f_v(u) - V(x + iy)| \leq \theta(u) o(1), \quad \text{as } u \rightarrow +\infty,$$

uniformly in $|y| < \pi/2$. Hence we obtain from (17.10)

$$(17.11) \quad f_v(u) = V(x) + \frac{\theta(u)}{\pi} y + o[\theta(u)] \quad \text{as } u \rightarrow +\infty,$$

uniformly for $|y| \leq \beta$.

Let now an ϵ , $0 < \epsilon < \pi/8$, be assigned. We may apply (17.11) for the particular value of $\beta = \pi/2 - \epsilon$. Let the curves L_v for $y = \pi/2 - \epsilon$ and $y = -\pi/2 + \epsilon$, for sufficiently large u , be represented by the equations $v = f_+(u)$ and $v = f_-(u)$, respectively. Then it follows from (17.11) that

$$f_+(u) = V(x) + \theta(u) \left[\frac{1}{2} - \frac{\epsilon}{\pi} \right] + o[\theta(u)],$$

$$f_-(u) = V(x) + \theta(u) \left[\frac{\epsilon}{\pi} - \frac{1}{2} \right] + o[\theta(u)]$$

as $u \rightarrow +\infty$. Addition and subtraction of these expressions give

$$(17.12) \quad V(x) = \frac{f_+(u) + f_-(u)}{2} + o[\theta(u)],$$

$$f_+(u) - f_-(u) = \theta(u) \left[1 - \frac{2\epsilon}{\pi} \right] + o[\theta(u)],$$

respectively. Substituting $\theta(u) = \phi_+(u) - \phi_-(u)$ into the second expression we find

$$[\phi_+(u) - f_+(u)] + [f_-(u) - \phi_-(u)] = \frac{2\epsilon}{\pi} \theta(u) + o[\theta(u)].$$

Since each of the summands on the left-hand side is greater than or equal to 0,

$$0 \leq \phi_+(u) - f_+(u) \leq \left[\frac{2\epsilon}{\pi} + o(1) \right] \theta(u),$$

$$0 \leq f_-(u) - \phi_-(u) \leq \left[\frac{2\epsilon}{\pi} + o(1) \right] \theta(u).$$

Hence we obtain from the first of the relations (17.12)

$$(17.13) \quad V(x) = \frac{\phi_+(u) + \phi_-(u)}{2} + \left[\lambda \frac{2\epsilon}{\pi} + o(1) \right] \theta(u)$$

for a suitable λ , $|\lambda| \leq 1$. Substitution of (17.13) into (17.11) gives

$$f_v(u) = \frac{1}{2}(\phi_+(u) + \phi_-(u)) + \frac{\theta(u)}{\pi} \cdot y + \left[\frac{2\epsilon}{\pi} \lambda + o(1) \right] \theta(u) \quad \text{as } u \rightarrow +\infty,$$

uniformly for $|y| \leq \beta$. This proves our result in the above stated weaker form, since it shows that, for all $|y| \leq \beta$,

$$(17.14) \quad \left| f_v(u) - \psi(u) - \frac{\theta(u)}{\pi} y \right| \leq \left[\frac{2\epsilon}{\pi} + o(1) \right] \theta(u) < \epsilon \theta(u),$$

for all sufficiently large u , say $u \geq N = N(\epsilon; \beta) \geq N''$.

b. Using this (weaker) result *only* we shall prove part (iv) of our theorem below. Anticipating here part (iv), we can readily prove (16.1) in the complete form, namely that $o[\theta(u)]$ in (16.1) holds uniformly for $|y| < \pi/2$.

Let ϵ , $0 < \epsilon < \pi/8$, be given. Choose $\beta = \pi/2 - \epsilon/2$ and determine the index $N(\epsilon; \pi/2 - \epsilon/2)$ so that (17.14) holds. Let $\pi/2 - \epsilon/2 < y < \pi/2$. Then, by part (iv), there exists an $N_1(\epsilon) \geq N$ such that all points $w = u + if_v(u)$ with $u \geq N_1$ lie in the exterior of $S_{\beta - \epsilon/2}$, i.e., either

$$\psi(u) + \frac{\theta(u)}{\pi} \left(\frac{\pi}{2} - \epsilon \right) < f_v(u) < \phi_+(u)$$

or

$$\phi_-(u) < f_v(u) < \psi(u) + \frac{\theta(u)}{\pi} \left(\epsilon - \frac{\pi}{2} \right).$$

Since we assumed $\pi/2 - \epsilon/2 < y < \pi/2$ and $\lim_{y \rightarrow \pi/2} f_v(u) = \phi_+(u)$, the first inequality holds. Similarly, if $-\pi/2 < y < -\pi/2 + \epsilon/2$, the second inequality holds. Hence, for all $u \geq N_1(\epsilon)$,

$$\left| f_v(u) - \psi(u) - \frac{\theta(u)}{\pi} y \right| < \epsilon \theta(u) \quad \text{for all } |y| < \pi/2.$$

(iv) Let ϵ be given as stated in the theorem. By (17.14) there exists an $N = N(\epsilon/\pi; \alpha)$ such that all points $w \in T_\alpha$ with $\Re w \geq N$ are contained in the region

$$\psi(u) - \frac{\theta(u)}{\pi} \alpha - \frac{\theta(u)}{\pi} \epsilon < v < \psi(u) + \frac{\theta(u)}{\pi} \alpha + \frac{\theta(u)}{\pi} \epsilon, \quad u \geq N.$$

This proves that the part of T_α which lies in $u \geq N$ is contained in $S_{\alpha+\epsilon}$. In a similar manner it may be seen that it contains the part of $S_{\alpha-\epsilon}$ within $u \geq N$.

(v) Relation (16.2) follows immediately from the first of the relations (ii) and part (iv) if we observe that, for $w = W(z)$, $Z'(w) = 1/W'(z)$.

18. **Corollaries of Theorem X.** We now prove the following corollaries.

COROLLARY 1. *If S is an L -strip with boundary inclination γ , $|\gamma| < \pi/2$ at $u = +\infty$, then parts (iii) and (iv) of Theorem X remain unchanged and parts (ii) and (v) are to be replaced by the relations*

(ii*) $\lim_{z \rightarrow +\infty} |W'(z)|/\theta(u) = (\cos \gamma)/\pi$, $\lim_{z \rightarrow +\infty} [\log |W'(z)|]/|z| = 0$, uniformly in y ,

(v*) $\lim_{u \rightarrow +\infty} [|Z'(w)|\theta(u)] = \pi/\cos \gamma$, uniformly for $w = u + iv \in S_\beta$, $0 < \beta < \pi/2$, respectively. Moreover, uniformly for $w \in S$,

$$(18.1) \quad Y(w) = \pi \frac{v - \psi(u)}{\theta(u)} + o(1) \quad \text{as } u \rightarrow +\infty.$$

To prove the corollary we rotate the coordinate system through the angle γ in the positive direction, thus obtaining a new set of coordinates (\bar{u}, \bar{v}) , $\bar{u} + i\bar{v} = \bar{w}$, where $\bar{w} = e^{-i\gamma}w$. Let, for sufficiently large \bar{u} , $\bar{v} = \bar{\phi}_+(\bar{u})$ and $\bar{v} = \bar{\phi}_-(\bar{u})$ represent C_+ and C_- , respectively. Consider a point $C: w = u + iv \in S$ and let $\bar{w} = e^{-i\gamma}w$. Denote the points $u + i\phi_-(u)$, $u + i\phi_+(u)$, in the w -plane, and $\bar{u} + i\bar{\phi}_-(\bar{u})$, $\bar{u} + i\bar{\phi}_+(\bar{u})$ in the \bar{w} -plane by A, B, \bar{A}, \bar{B} respectively. In the triangle $\bar{A}AC$, the angles \bar{A} and A approach $\pi/2$ and $\pi/2 - |\gamma|$ respectively, and in $\bar{B}BC$, the angles \bar{B} and B approach $\pi/2$ and $\pi/2 - |\gamma|$ respectively as $u \rightarrow +\infty$, uniformly with respect to the position of C on AB . Hence

$$(18.2) \quad \frac{AC}{\bar{AC}} = \frac{\sin \bar{A}}{\sin A} \rightarrow \frac{1}{\cos \gamma}, \quad \frac{BC}{\bar{BC}} \rightarrow \frac{1}{\cos \gamma} \quad \text{as } u \rightarrow +\infty.$$

and therefore $(\bar{\theta}(\bar{u}) = \bar{\phi}_+(\bar{u}) - \bar{\phi}_-(\bar{u}))$

$$(18.3) \quad \theta(u) = AB = AC + BC = \frac{\bar{A}\bar{B}}{\cos \gamma} + o(\bar{A}\bar{B}) = \frac{\bar{\theta}(\bar{u})}{\cos \gamma} (1 + o(1))$$

as $u \rightarrow +\infty$,

the convergence in (18.2) and (18.3) being uniform with respect to the position of C on AB .

Now (ii*) follows immediately if we observe that for $\bar{W}(z) = e^{-i\gamma}W(z)$ by Theorem X (ii) and subsequently by (18.3), uniformly for $|y| \leq \beta$:

$$\frac{1}{\pi} = \lim_{z \rightarrow +\infty} \frac{|\bar{W}'(z)|}{\bar{\theta}(\bar{u})} = \lim_{z \rightarrow +\infty} \frac{|W'(z)|}{\theta(u) \cos \gamma}, \quad \bar{u} = \Re \bar{W}(z); u = \Re W(z),$$

and that

$$\frac{\log |\bar{W}'(z)|}{|z|} = \frac{\log |W'(z)|}{|z|}.$$

To prove now that part (iii) of Theorem X remains unchanged, we note that in the figure which was used above, $AC = v - \phi_-(u)$, $BC = \phi_+(u) - v$,

$\overline{AC} = \bar{v} - \bar{\phi}_-(\bar{u})$, $\overline{BC} = \bar{\phi}_+(\bar{u}) - \bar{v}$. From (18.2) we obtain therefore (leaving off the arguments u , \bar{u})

$$(18.4) \quad v - \phi_- = (\bar{v} - \bar{\phi}_-) \frac{1}{\cos \gamma} + o(\bar{\theta}), \quad \phi_+ - v = (\bar{\phi}_+ - \bar{v}) \frac{1}{\cos \gamma} + o(\bar{\theta}),$$

as $u \rightarrow +\infty$, uniformly with respect to the position of C on AB . By Theorem X (iii), there exists an \bar{N} (independent of γ), such that any line $\Lambda_y: y = \text{const.}$ ($|y| < \pi/2$) is mapped by $\bar{w} = \bar{W}(z)$ onto a curve L_y which, for all $\bar{u} \geq \bar{N}$, can be represented in the form $(\bar{\psi} = \frac{1}{2}[\bar{\phi}_+ + \bar{\phi}_-])$:

$$\bar{v} = \bar{\psi} + \frac{\bar{\theta}}{\pi} y + o(\bar{\theta}) \quad \text{as } \bar{u} \rightarrow +\infty, \text{ uniformly for } |y| < \pi/2.$$

Hence, for $\bar{w} = \bar{u} + \bar{v}i$ on L_y ($\bar{u} \geq \bar{N}$), uniformly for $|y| < \pi/2$,

$$\bar{v} - \bar{\phi}_- = \frac{\bar{\theta}}{\pi} \left[\frac{\pi}{2} + y + o(1) \right], \quad \bar{\phi}_+ - \bar{v} = \frac{\bar{\theta}}{\pi} \left[\frac{\pi}{2} - y + o(1) \right]$$

as $u \rightarrow +\infty$. Substituting these values for $\bar{v} - \bar{\phi}_-$ and $\bar{\phi}_+ - \bar{v}$ into the first and second equations of (18.4), respectively and using (18.3) we find that there exists an N (independent of γ) such that any point $w = u + iv$ on L_y with $u \geq N$ satisfies the relations

$$(18.5) \quad v - \phi_- = \frac{\theta}{\pi} \left[\frac{\pi}{2} + y + o(1) \right], \quad \phi_+ - v = \frac{\theta}{\pi} \left[\frac{\pi}{2} - y + o(1) \right],$$

as $u \rightarrow +\infty$, uniformly in $|y| < \pi/2$. Subtraction of the second equation in (18.5) from the first gives

$$(18.6) \quad v = \psi + \frac{\theta}{\pi} y + o(\theta) \quad \text{as } u \rightarrow +\infty, \text{ uniformly in } |y| < \pi/2.$$

It may be shown now by use of (18.6) (as in the proof of part (iv) of Theorem X) that the statement of part (iv) remains unchanged in the case of any γ , $|\gamma| < \pi/2$. Part (v*) of the corollary follows then from part (ii*) in the same manner as part (v) of the theorem follows from part (ii).

Finally, (18.1) is obtained by solving (18.6) for y , since for $w = u + iv$ on L_t : $Y(w) = t$.

COROLLARY 2. Let S be an L -strip with the boundary inclination γ , $|\gamma| < \pi/2$, at $u = +\infty$.

(i) If $\gamma = 0$, then for $z_1 = x_1 + iy_1$, $z_2 = x_2 + iy_2$, $|x_2 - x_1| \leq M$ ($M = \text{const.}$),

$$(18.7) \quad U(z_2) - U(z_1) = \frac{\theta(u)}{\pi} (x_2 - x_1) + o[\theta(u)]$$

uniformly for $|y| < \pi/2$ as $x_1, x_2 \rightarrow +\infty$. Here u may be taken equal to any number between u_1 and u_2 .

In particular, if $\lim_{u \rightarrow +\infty} \theta(u) = \theta$ exists, then (18.7) becomes

$$(18.7^*) \quad U(z_2) - U(z_1) = \frac{\theta}{\pi} (x_2 - x_1) + o(1).$$

(ii) If $\limsup_{u \rightarrow +\infty} \theta(u) = \theta^*$, $\liminf_{u \rightarrow +\infty} \theta(u) = \theta_*$, then, uniformly in $|y| < \pi/2$,

$$(18.8) \quad \frac{\theta_* \cos^2 \gamma}{\pi} \leq \liminf_{x \rightarrow +\infty} \frac{U(z)}{x} \leq \limsup_{x \rightarrow +\infty} \frac{U(z)}{x} \leq \frac{\theta^* \cos^2 \gamma}{\pi}.$$

Thus, if $\lim_{u \rightarrow +\infty} \theta(u) = \theta$ exists:

$$(18.8^*) \quad \lim_{x \rightarrow +\infty} \frac{U(z)}{x} = \frac{\theta \cos^2 \gamma}{\pi}, \quad \lim_{z \rightarrow +\infty} \frac{W(z)}{z} = \frac{\theta \cos \gamma}{\pi} e^{i\gamma}.$$

REMARK. In the case that

$$(18.9) \quad \lim_{u \rightarrow +\infty} \phi_+(u) = \phi_+, \quad \lim_{u \rightarrow +\infty} \phi_-(u) = \phi_- \text{ exist,}$$

both of these parts are known theorems⁽²⁷⁾.

Proof. (i) Let $u_1 = U(z_1)$, $u_2 = U(z_2)$, $x_2 \geq x_1$. By Theorems III (b) and IV (b)

$$(18.10) \quad -\epsilon_1 + \pi \int_{u_1}^{u_2} \frac{du}{\theta(u)} \leq x_2 - x_1 \leq \pi(1 + \epsilon_2) \int_{u_1}^{u_2} \frac{du}{\theta(u)} + \epsilon_3,$$

$$\lim_{x_1 \rightarrow +\infty} \epsilon_k = 0, \quad k = 1, 2, 3.$$

By the corollary of Lemma 7, for any u between u_1 and u_2 :

$$\int_{u_1}^{u_2} \frac{dt}{\theta(t)} = \frac{1}{\theta(u)} (u_2 - u_1)(1 + \epsilon_4), \quad \lim_{x_1 \rightarrow +\infty} \epsilon_4 = 0.$$

Substituting this value of $\int_{u_1}^{u_2} [\theta(t)]^{-1} dt$ into (18.10) and observing that $|x_2 - x_1| \leq M$, we find (18.7).

⁽²⁷⁾ In the case that $\phi_+ - \phi_- = \theta > 0$, (18.7*) has been proved under the less restrictive hypothesis that S is a region as described in the statement of Carathéodory's theorem in the remark of §16, whose boundary curves C_+ and C_- (which approach the asymptotes $v = \phi_+$ and $v = \phi_-$ respectively) satisfy an *additional condition* regarding their "oscillation" ("Reguläre Unbewalltheit"). See Wolff [1, p. 217], Warschawski [1, p. 326], Ostrowski [1, p. 117, relation (21.2)]. For $\theta = 0$, Ostrowski [1, p. 177] proved (18.7*) under the assumption that S is an L -strip satisfying (18.9). Compare also Ostrowski's extension of the first case ($\theta > 0$) to regions with general boundaries [2, pp. 88, 95].—Part (ii) of Corollary 2, under the assumption (18.9) is due to Ostrowski [1, p. 174, relation (62.1)].

(ii) Assume first that $\gamma=0$. Then it is sufficient to prove that

$$(18.11) \quad \frac{\theta_*}{\pi} \leq \liminf_{u \rightarrow +\infty} \frac{u}{X(w)} \leq \limsup_{u \rightarrow +\infty} \frac{u}{X(w)} \leq \frac{\theta^*}{\pi}, \quad \text{uniformly for } w \in S.$$

Let $\epsilon > 0$ be given. Take u_1 so large that

$$\theta_* - \epsilon \leq \theta(u) \leq \theta^* + \epsilon \quad \text{for } u \geq u_1.$$

By Theorem X (i), uniformly for $w \in S$, as $u \rightarrow +\infty$,

$$\frac{u}{X(w)} \sim \frac{u}{\pi \int_{u_1}^u \frac{dt}{\theta(t)}} = \frac{u}{u - u_1} \cdot \frac{\theta(\xi)}{\pi}, \quad u_1 < \xi < u.$$

Hence, keeping u_1 fixed, we have

$$\frac{\theta_* - \epsilon}{\pi} \leq \liminf_{u \rightarrow +\infty} \frac{u}{X(w)} \leq \limsup_{u \rightarrow +\infty} \frac{u}{X(w)} \leq \frac{\theta^* + \epsilon}{\pi}.$$

Since ϵ is arbitrary this proves (18.11).

If $\gamma \neq 0$, let $\bar{w} = e^{-i\gamma}w$, $\bar{W}(z) = \bar{U}(z) + i\bar{V}(z) = e^{-i\gamma}W(z)$. Moreover, let $\bar{\theta}(\bar{u})$ be defined in the \bar{w} -plane as $\theta(u)$ is in the w -plane. Then, by the part just proved,

$$(18.12) \quad \frac{1}{\pi} \liminf_{\bar{u} \rightarrow +\infty} \bar{\theta}(\bar{u}) \leq \liminf_{x \rightarrow +\infty} \frac{\bar{U}(z)}{x} \leq \limsup_{x \rightarrow +\infty} \frac{\bar{U}(z)}{x} \leq \frac{1}{\pi} \limsup_{\bar{u} \rightarrow +\infty} \bar{\theta}(\bar{u}).$$

Now

$$U(z) = \bar{U}(z) \cos \gamma - \bar{V}(z) \sin \gamma = \bar{U}(z) \cos \gamma \left\{ 1 - \frac{\bar{V}(z)}{\bar{U}(z)} \tan \gamma \right\}.$$

Here $\bar{V}(z)/\bar{U}(z) \rightarrow 0$ as $x \rightarrow +\infty$, uniformly for $|y| < \pi/2$. Using this relation in connection with (18.3) we obtain (18.8) from (18.12).

VI. APPLICATIONS

19. The general result. We shall now apply our results to the study of the mapping function in a neighborhood of a finite boundary point ω_0 of a region bounded by a closed Jordan curve. First we derive from our results a theorem which deals with a certain general boundary configuration and then apply it to various special cases.

(a) Let R be the interior of a closed Jordan curve Γ in the ω -plane and let $\omega=0$ be on Γ . Suppose that in a neighborhood of $\omega=0$, say $|\omega| \leq a$, Γ consists of two arcs Γ_+ and Γ_- which are represented in polar coordinates in the form

$$\begin{aligned} \phi &= \Phi_+(\rho), & \phi &= \Phi_-(\rho), \\ 0 < \rho &\leq a, & \Phi_+(\rho) < \Phi_-(\rho), \end{aligned}$$

respectively, the functions $\Phi_+(\rho)$, $\Phi_-(\rho)$ being continuous in the interval $0 < \rho \leq a$. The region

$$0 < \rho < a, \quad \Phi_+(\rho) < \phi < \Phi_-(\rho)$$

is contained in R . We set $\Theta(\rho) \equiv \Phi_-(\rho) - \Phi_+(\rho)$ and $\Psi(\rho) = \frac{1}{2} [\Phi_-(\rho) + \Phi_+(\rho)]$.

Suppose that $\Phi_+(\rho)$ and $\Phi_-(\rho)$ are absolutely continuous in any closed interval within $0 < \rho \leq a$ and that $\rho [d\Phi_+(\rho)/d\rho]$ and $\rho [d\Phi_-(\rho)/d\rho]$, which exist for $0 < \rho \leq a$ except possibly for a set of measure 0, approach the same limit, $\tan \gamma$, $|\gamma| < \pi/2$, as $\rho \rightarrow 0$.

Finally, let $\zeta = \zeta(\omega)$ map R conformally onto the circle $|\zeta - 1| < 1$ in such a manner that $\omega = 0$ corresponds to $\zeta = 0$ and let $\omega = \omega(\zeta)$ denote its inverse function.

(b) Logarithmic transformation of R by means of the function⁽²⁸⁾ $w = \log(1/\omega)$ and of the circle $|\zeta - 1| < 1$ by means of $z = \log [(2 - \zeta)/\zeta]$ gives at once the following results:

THEOREM XI(A). *Under the above stated hypotheses we have⁽²⁹⁾:*

(i) *If $\gamma = 0$, then for any branch of $\log \zeta(\omega)$, $|\omega| = \rho$, uniformly in R ,*

$$\lim_{\rho \rightarrow 0} \frac{\log \zeta(\omega)}{\int_{\rho}^a \frac{dr}{r\Theta(r)}} = -\pi.$$

(ii) *Uniformly in any angle $|\arg \zeta| \leq \beta < \pi/2$, as $\zeta \rightarrow 0$,*

$$\frac{\left| \frac{\omega'(\zeta)}{\omega(\zeta)} \right|}{\left| \frac{\zeta}{\omega(\zeta)} \right|} \sim \frac{\Theta(\rho)}{\pi} \cos \gamma, \quad |\omega| = \rho.$$

(iii) *Any circular arc λ_t , t fixed, $|t| < \pi/2$: $\{\arg \{(2 - \zeta)/\zeta\} = t, |\zeta - 1| < 1\}$ is mapped by $\omega(\zeta)$ onto a curve l_t which in a neighborhood of $\omega = 0$ is representable in the form*

$$\phi = \Psi(\rho) - \frac{\Theta(\rho)}{\pi} t + o[\Theta(\rho)] \quad \text{as } \rho \rightarrow 0,$$

uniformly for $|t| < \pi/2$.

⁽²⁸⁾ R is transformed by $w = \log 1/\omega$ into an L -strip S whose boundary curves C_+ and C_- are given by the equations $v = \phi_+(u) = -\Phi_+(e^{-u})$, $v = \phi_-(u) = -\Phi_-(e^{-u})$, respectively. To see that C_+ (and similarly C_-) has an L -tangent at $u = +\infty$, one only has to note that

$$\begin{aligned} \frac{\phi_+(u_2) - \phi_+(u_1)}{u_2 - u_1} &= -\frac{1}{u_2 - u_1} \int_{\rho_1}^{\rho_2} \frac{d\Phi_+}{d\rho} d\rho = -(1 + \epsilon) \frac{\tan \gamma}{u_2 - u_1} \int_{\rho_1}^{\rho_2} \frac{d\rho}{\rho} \\ &= -(1 + \epsilon) \tan \gamma \frac{\log \rho_2 - \log \rho_1}{u_2 - u_1} = (1 + \epsilon) \tan \gamma, \end{aligned}$$

where $\epsilon \rightarrow 0$ as $\rho_1, \rho_2 \rightarrow 0$.

⁽²⁹⁾ Part (A) follows from Theorem X and Corollary 1 of this theorem.

(iv) Let σ_α denote the region within $|\zeta - 1| < 1$ which is bounded by the two circular arcs $\arg \{(2 - \zeta)/\zeta\} = \alpha$ and $\arg \{(2 - \zeta)/\zeta\} = -\alpha$, $0 < \alpha < \pi/2$, let τ_α be its image in the ω -plane by means of $\omega(\zeta)$, and let s_β ($0 < \beta < \pi/2$) denote the region $\{0 < \rho < a, |\phi - \Psi(\rho)| \leq (\beta/\pi)\Theta(\rho)\}$.

If $0 < \alpha \pm \epsilon < \pi/2$, $\epsilon > 0$, then there exists an $r = r(\epsilon; \alpha)$ such that the part of τ_α which lies in $0 < \rho \leq r$ contains that part of $s_{\alpha-\epsilon}$ which is in $\rho \leq r$ and is contained in $s_{\alpha+\epsilon}$.

(v) Uniformly in any region s_β , $0 < \beta < \pi/2$,

$$\frac{|\zeta'(\omega)|}{\left| \frac{\zeta(\omega)}{\omega} \right|} \sim \frac{\pi}{\Theta(\rho) \cos \gamma} \quad \text{as } |\omega| = \rho \rightarrow 0.$$

(vi) Uniformly in R , for $\omega = \rho e^{i\phi}$,

$$(19.1) \quad \arg \zeta(\omega) = \pi \frac{\phi - \Psi(\rho)}{\Theta(\rho)} + o(1) \quad \text{as } \omega \rightarrow 0.$$

THEOREM XI (B). If in addition to the hypotheses stated in §19 (a), $\rho[d\Phi_+(\rho)/d\rho]$ and $\rho[d\Phi_-(\rho)/d\rho]$ are continuous for $0 \leq \rho \leq a$ and the integrals

$$(19.2) \quad \int_{\rho=0}^a |d(\rho\Phi'_+(\rho))|, \quad \int_{\rho=0}^a |d(\rho\Phi'_-(\rho))|, \quad \int_0^a \frac{\Theta'^2(\rho)}{\Theta(\rho)} d\rho \text{ converge,}$$

then there exists a constant $c > 0$ such that, for $\omega = \rho e^{i\phi}$,

$$(19.3) \quad |\zeta(\omega)| = c \exp \left\{ -\pi \int_\rho^a \frac{1 + (r\Psi'(r))^2}{r\Theta(r)} dr + \pi \frac{\phi - \Psi(\rho)}{\Theta(\rho)} \tan \gamma + o(1) \right\}$$

as $\omega \rightarrow 0$ in any way at all in $R^{(30)}$.

(c) REMARK. If Γ is a curve as described in §19 (a) and if, in addition the integrals

$$(19.4) \quad \int_0^a \rho \left(\frac{d\Phi_+}{d\rho} \right)^2 \frac{d\rho}{\Theta(\rho)} \quad \text{and} \quad \int_0^a \rho \left(\frac{d\Phi_-}{d\rho} \right)^2 \frac{d\rho}{\Theta(\rho)} \text{ converge,}$$

then $\gamma = 0^{(31)}$ and the integral $\int_0^a (\rho\Psi'^2(\rho)/\Theta(\rho))d\rho$ exists, and therefore by the corollary of Theorem IV (§7), (19.3) reduces to

$$(19.5) \quad |\zeta(\omega)| = c' \exp \left\{ -\pi \int_\rho^a \frac{dr}{r\Theta(r)} + o(1) \right\} \text{ as } \omega \rightarrow 0 \text{ in } R \text{ (} c' = \text{const.)}$$

(d) COROLLARY. Let R be a region in the ω -plane satisfying the hypotheses

⁽³⁰⁾ Part (B) follows from the corollary to Theorem VIII (§14).

⁽³¹⁾ If γ were not 0, the convergence of either of the integrals (19.4) would imply that of $\int_0^a [r\Theta(r)]^{-1}dr$, and this integral diverges.

of §19 (a). If in addition, $\lim_{\rho \rightarrow 0} \Theta(\rho) = \theta$ exists, then we have⁽³²⁾:

(i) If $\gamma = 0$, for ζ_1 and ζ_2 in $|\zeta - 1| < 1$, for which $0 < c_1 < |\zeta_1/\zeta_2| < c_2$ (c_1, c_2 consts.),

$$\left| \frac{\omega(\zeta_2)}{\omega(\zeta_1)} \right| = \left| \frac{\zeta_2}{\zeta_1} \right|^{\theta/\pi} (1 + o(1)) \quad \text{as } \zeta_1, \zeta_2 \rightarrow 0,$$

uniformly in the circle $|\zeta - 1| < 1$.

(ii) As $\zeta \rightarrow 0$ in $|\zeta - 1| < 1$ in any way at all,

$$(19.6) \quad \frac{\log |\omega(\zeta)|}{\log |\zeta|} \rightarrow \frac{\theta}{\pi} \cos^2 \gamma, \quad \frac{\log \omega(\zeta)}{\log \zeta} \rightarrow \frac{\theta}{\pi} e^{i\gamma} \cos \gamma.$$

(iii) As $\zeta \rightarrow 0$ in any angle $|\arg \zeta| \leq \alpha < \pi/2$ ⁽³³⁾,

$$(19.7) \quad \frac{\log |\omega'(\zeta)|}{\log |\zeta|} \rightarrow \frac{\theta}{\pi} \cos^2 \gamma - 1.$$

We now apply these results to various special cases.

20. Boundary "elements" with bounded argument oscillation. Let Γ be a curve as described in §19 (a) for which $\Phi_+(\rho)$ and $\Phi_-(\rho)$ are bounded for $0 < \rho \leq a$. Then, necessarily, $\gamma = 0$ ⁽³⁴⁾, and the results of §19 hold with γ replaced by 0. We consider the case where

$$(20.1) \quad \lim_{\rho \rightarrow 0} \Phi_+(\rho) = \phi_+ \quad \text{and} \quad \lim_{\rho \rightarrow 0} \Phi_-(\rho) = \phi_- \quad \text{exist.}$$

(a) *Corners.* If $\theta \equiv \phi_- - \phi_+ > 0$, then Γ has an L -corner of measure θ at $\omega = 0$ (see §2). In this case parts (ii), (iii) and (vi) of Theorem XI (A) are known, (ii) has been proved by Ostrowski, even under the weaker hypothesis that Γ has a corner at $\omega = 0$ (and not necessarily an L -corner), and (iii) and (vi) express the fact that the map is quasi-conformal at $\omega = 0$. Furthermore, the results of the corollary for this case as well as those for $\theta = 0$ are due to Ostrowski (see (27)).

Formulas (19.3) and (19.5) give expressions for the order of magnitude of $|\zeta(\omega)|$. One may ask here, when, in particular, $\zeta(\omega) \sim c\omega^{\pi/\theta}$ as $\omega \rightarrow 0$ in $R + \Gamma$ ($c = \text{constant} \neq 0$). This question has received much attention in the recent lit-

⁽³²⁾ For parts (i) and (ii) see Corollary 2 of Theorem X (§18, (18.7*) and (18.8*)).

⁽³³⁾ The relation (19.7) follows from Theorem XI (A), (ii) and (19.6) if one observes that

$$\log |\omega'(\zeta)| = \log |\omega(\zeta)| - \log |\zeta| + \log \left[\frac{\Theta(\rho)}{\pi} \cos \gamma \right] + o(1)$$

as $\zeta \rightarrow 0$ in $|\arg \zeta| \leq \alpha < \pi/2$.

⁽³⁴⁾ If γ were not equal to 0, we would have since $(d\Phi_+/d\rho)\rho \rightarrow \tan \gamma$, ($\rho_1 < \rho_2$),

$$|\Phi_+(\rho_2) - \Phi_+(\rho_1)| = \left| \int_{\rho_1}^{\rho_2} \frac{d\Phi_+}{d\rho} d\rho \right| \geq \frac{|\tan \gamma|}{2} \int_{\rho_1}^{\rho_2} \frac{d\rho}{\rho}$$

for all sufficiently small ρ_1 and ρ_2 . Keeping ρ_2 fixed and letting $\rho_1 \rightarrow 0$, we would find that $\Phi_+(\rho)$ is not bounded.

erature, especially for the case $\theta = \pi$ where this means that $\zeta(\omega)$ possesses a *nonvanishing derivative at $\omega = 0$* . Since by (19.1) $\lim_{\omega \rightarrow 0} \arg \{\zeta(\omega)/\omega^{\pi/\theta}\}$ exists, it is sufficient to consider $\lim_{\omega \rightarrow 0} |\zeta(\omega)/\omega^{\pi/\theta}|$. The remark of §19 (c) yields the following result *which does not even presuppose any assumption regarding the existence of $\lim_{\rho \rightarrow 0} \theta(\rho)$ or that of the limits (20.1)*.

THEOREM XII. *If Γ is a curve as described in §19 (a) and if the integrals (19.4) converge, then a necessary and sufficient condition that, for some positive θ , $\lim_{\omega \rightarrow 0} |\zeta(\omega)/\omega^{\pi/\theta}|$ exist for unrestricted approach and be different from 0 is that*

$$(20.2) \quad \int_0^a \frac{\theta - \Theta(\rho)}{\rho\Theta(\rho)} d\rho \text{ converge.}$$

For, we have, uniformly in R ,

$$\begin{aligned} \left| \frac{\zeta(\omega)}{\omega^{\pi/\theta}} \right| &= c' \exp \left\{ -\pi \int_{\rho}^a \frac{dr}{r\Theta(r)} + \frac{\pi}{\theta} \log \frac{a}{\rho} - \frac{\pi}{\theta} \log a + o(1) \right\} \\ &= c'' \exp \left\{ -\pi \int_{\rho}^a \frac{\theta - \Theta(r)}{r\Theta(r)} dr + o(1) \right\} \end{aligned}$$

as $|\omega| = \rho \rightarrow 0$ (c'' is a constant different from 0).

Combining this result with (19.1) we find:

THEOREM XIII. *If Γ is a curve as described in §19 (a) and if the integrals (19.4) converge, then a necessary and sufficient condition in order that for some $\theta > 0$, $\lim_{\omega \rightarrow 0} (\zeta(\omega)/\omega^{\pi/\theta})$ exist for unrestricted approach and be different from 0 is that the conditions (20.1) and (20.2) be satisfied and that $\phi_- - \phi_+ = \theta$.*

The sufficiency of the conditions stated is clear. That (20.2) is necessary follows from Theorem XII. That (20.1) and the relation $\phi_- - \phi_+ = \theta$ are necessary is immediately seen if we observe that $\arg \{\zeta/[\omega(\zeta)]^{\pi/\theta}\}$ approaches a limit as $\zeta \rightarrow 0$ in $|\zeta - 1| \leq 1$, and then let $\zeta \rightarrow 0$, first along the upper and then along the lower semi-circle of $|\zeta - 1| = 1$.

For $\theta = \pi$ this theorem gives a criterion for the existence of the derivative of $\zeta(\omega)$ at a boundary point. Several criteria (sufficient conditions) for the existence of the angular derivative (i.e., $\lim_{\zeta \rightarrow 0} (\omega(\zeta)/\zeta)$ in any fixed angle $|\arg \zeta| \leq \alpha < \pi/2$) are known which apply to even more general types of regions than those bounded by Jordan curves. (For sufficiently smooth boundaries the existence of the derivatives for unrestricted approach can be inferred from that of the angular derivative.) The sharpest of these criteria to date is due to Ahlfors⁽³⁵⁾. Since our theorem refers to a smaller class of regions it obviously does not contain that of Ahlfors⁽³⁶⁾. On the other hand, the follow-

⁽³⁵⁾ Ahlfors [1, p. 36].

⁽³⁶⁾ However, by use of Theorem XII (B) and a modification of Ahlfors' proof of his criterion, one may obtain a sharper criterion for the existence of the angular derivative.

ing example shows that it also is not contained in Ahlfors' result: Let Γ be a closed Jordan curve through $\omega=0$ and let, in a neighborhood of $\omega=0$ ($0 \leq \rho \leq a$), Γ consist of the two branches Γ_+ and Γ_- represented by

$$\phi = \Phi_+(\rho) \equiv -\frac{\pi}{2} + \delta_1(\rho), \quad \phi = \Phi_-(\rho) \equiv \frac{\pi}{2} + \delta_2(\rho), \quad \delta_i(\rho) > 0, \quad \lim_{\rho \rightarrow 0} \delta_i(\rho) = 0,$$

where $\int_0^a [\delta_i(\rho)/\rho] d\rho$ is divergent ($i=1, 2$), $\int_0^a \{|\delta_1(\rho) - \delta_2(\rho)|/\rho\} d\rho$ convergent, $\rho \delta'_i(\rho)$ continuous, $\rho(\delta'_i(\rho))^2$ integrable, $0 \leq \rho \leq a$. It is easily seen that the hypotheses of Theorem XIII are satisfied and that therefore $\lim_{\omega \rightarrow 0} [\zeta(\omega)/\omega]$ exists and does not equal 0. However, one of the conditions of Ahlfors' criterion is not satisfied. Suppose Γ is mapped by the function $w = \log(1/\omega)$ onto a strip. The images C_+ and C_- of Γ_+ and Γ_- are represented by the equations

$$v = \phi_+(u) \equiv \frac{\pi}{2} - \delta_1(e^{-u}), \quad v = \phi_-(u) \equiv -\frac{\pi}{2} - \delta_2(e^{-u}),$$

respectively. Then (in Ahlfors' notation [1, p. 36]) the series $\sum_{\nu=1}^{\infty} m_\nu$ does not converge. For, if $\nu k \leq u \leq (\nu+1)k$ are the intervals for which the m_ν are formed, then

$$k \sum_{\nu=1}^n m_\nu \geq \sum_{k=1}^n \int_{\nu k}^{(\nu+1)k} \delta_1(e^{-u}) du = \int_k^{(n+1)k} \delta_1(e^{-u}) du \rightarrow +\infty$$

as $n \rightarrow \infty$. However, the convergence of this series is one of the conditions of his criterion.

A corollary of Theorem XIII is the following

THEOREM XIV. *Let Γ be a closed Jordan curve through $\omega=0$ which has an L-corner of measure θ at $\omega=0$, formed by the two branches Γ_+ and Γ_- . Suppose that the angles of inclination of the tangents to Γ_+ and Γ_- , considered as functions of ρ , are of bounded variation in a neighborhood of $\omega=0$, $0 \leq \rho \leq a$. Let $\zeta(\omega)$ be defined as in §19 (a). Then, a necessary and sufficient condition that $\lim_{\omega \rightarrow 0} (\zeta(\omega)/\omega^{\tau/\theta})$ exist and be different from zero is that the integral (20.2) converge. ($\Phi_+(\rho)$, $\Phi_-(\rho)$ and $\Theta(\rho)$ in (20.2) are defined as in §19 (a).)*

Proof. We represent Γ_+ and Γ_- in the form $\phi = \Phi_+(\rho)$ and $\phi = \Phi_-(\rho)$ respectively, $0 \leq \rho \leq a$, and show first that the integrals (19.4) converge. Let $\tau_+(\rho)$ denote the angle of inclination of the tangent to Γ_+ at $\omega = \rho e^{i\Phi_+(\rho)}$ (if the tangent exists), $\tau_+(\rho)$ being so chosen that $\lim_{\rho \rightarrow 0} \tau_+(\rho) = \phi_+$ (see (20.1)). As is well known,

$$\rho \frac{d\Phi_+(\rho)}{d\rho} = \tan [\tau_+(\rho) - \Phi_+(\rho)].$$

To prove the convergence of the first integral in (19.4) it is sufficient to show that

$$(20.3) \quad \int_0^a [\tau_+(\rho) - \Phi_+(\rho)] \frac{d\Phi_+(\rho)}{d\rho} d\rho \quad \text{converges.}$$

Integration by parts gives ($0 \leq \epsilon < a$)

$$\begin{aligned} \int_{\epsilon}^a [\tau_+ - \Phi_+] \frac{d\Phi_+}{d\rho} d\rho &= - \int_{\rho=\epsilon}^a (\Phi_+(\rho) - \phi_+) d[\tau_+(\rho) - \Phi_+(\rho)] \\ &\quad + (\Phi_+(\rho) - \phi_+) (\tau_+(\rho) - \Phi_+(\rho)) \Big|_{\epsilon}^a \\ &= - \int_{\rho=\epsilon}^a (\Phi_+(\rho) - \phi_+) d\tau_+(\rho) \\ &\quad + \int_{\rho=\epsilon}^a (\Phi_+(\rho) - \phi_+) d[\Phi_+(\rho) - \phi_+] + M(\epsilon, a), \end{aligned}$$

where $M(\epsilon, a)$ is continuous at $\epsilon=0$. Since $\tau_+(\rho)$ is of bounded variation and $\Phi_+(\rho)$ is continuous for $0 \leq \rho \leq a$, the first of the last two integrals converges as $\epsilon \rightarrow 0$. The second of these integrals has the value $[\frac{1}{2}(\Phi_+(\rho) - \phi_+)^2]_{\epsilon}^a$ and it approaches, therefore, a finite limit as $\epsilon \rightarrow 0$. This proves (20.3).

Similarly it is shown that the second integral in (19.4) converges and Theorem XIV follows now from Theorem XIII.

(b) *Cusps*. Suppose now that Γ has an L -cusp at $\omega=0$. Our main results here are the formulas (19.3), with $\gamma=0$, and (19.5) which give *asymptotic expressions for* $|\zeta(\omega)|$. Parts (iii) and (vi) of Theorem XI (A) take the place of the quasi-conformality in the case of a corner. Part (ii) of this theorem (with $\gamma=0$) is an extension of Ostrowski's result which states that under the weaker assumption that Γ has a cusp at $\omega=0$ (and not necessarily an L -cusp), $\lim_{\zeta \rightarrow 0} [\omega'(\zeta)/(\omega(\zeta)/\zeta)] = 0$, in any fixed angle $|\arg \zeta| \leq \beta < \pi/2$ (cf. the remark to Theorem X, §16). Similarly, part (i) of Theorem XI (A) can be considered as a sharper form of (19.6), since it gives the order of magnitude of $\log |\zeta(\omega)|$ while (19.6) merely states that $(\log |\zeta(\omega)| / \log |\omega|) \rightarrow +\infty$ as $\omega \rightarrow 0$. The results of the corollary are due to Ostrowski (see (27)).

EXAMPLE. It might be of some interest to apply our results to a cusp formed by two arcs Γ_+ and Γ_- which have "finite order of contact." Γ_+ and Γ_- are represented in Cartesian coordinates (ξ, η) by the equations

$$(20.4) \quad \eta = [a + \epsilon(\xi)]\xi^n, \quad \eta = [b + \delta(\xi)]\xi^m, \quad 0 \leq \xi \leq \xi_0.$$

Here a, b, n, m are any real numbers, $n \geq m > 1$; if $n > m$ then $b > 0$, if $n = m$, $b > a$; $\epsilon(\xi), \delta(\xi)$ have continuous first derivatives, $0 \leq \xi \leq \xi_0$ and approach 0 with ξ . It is clear that Γ_+ and Γ_- form an L -cusp so that Theorem XI (A) holds. We examine, therefore, the possibility of applying part (B).

We introduce polar coordinates about $\omega=0$. Using ρ as parameter and writing $\phi = \Phi_+(\rho)$ on Γ_+ and $\phi = \Phi_-(\rho)$ on Γ_- we obtain from (20.4), $\rho > 0$,

$$(20.5) \quad \rho \sin \Phi_+(\rho) = [a + \epsilon(\xi)]\rho^n \cos^n \Phi_+(\rho), \quad \rho \sin \Phi_-(\rho) = [b + \delta(\xi)]\rho^m \cos^m \Phi_-(\rho).$$

Hence

$$\sin [\Phi_-(\rho) - \Phi_+(\rho)] = \rho^{m-1} \{ (b + \delta) \cos^m \Phi_- \cos \Phi_+ - (a + \epsilon)\rho^{n-m} \cos^n \Phi_+ \cos \Phi_- \}.$$

Thus we find

$$\Theta(\rho) = \Phi_- - \Phi_+ \sim b\rho^{m-1} \quad \text{if } n > m; \quad \Theta(\rho) \sim (b - a)\rho^{m-1} \quad \text{if } n = m,$$

as $\rho \rightarrow 0$. Furthermore, differentiation of the first relation of (20.5) with respect to ρ gives

$$\cos \Phi_+(\rho) \frac{d\Phi_+}{d\rho} = (a + \epsilon)(n - 1)\rho^{n-2} \{ 1 + o(1) \} \quad \text{as } \rho \rightarrow 0,$$

so that

$$\frac{\rho}{\Theta(\rho)} \left(\frac{d\Phi_+}{d\rho} \right)^2 = O(\rho^{n-2}) \quad \text{as } \rho \rightarrow 0,$$

and similarly

$$\frac{\rho}{\Theta(\rho)} \left(\frac{d\Phi_-}{d\rho} \right)^2 = O(\rho^{m-2}).$$

Hence the conditions (19.4) of the remark of §19 (c) are satisfied and therefore (19.5) holds.

21. Boundary "elements" with unbounded argument oscillation. Suppose that Γ is a curve as described in §19 (a) and that $\lim_{\rho \rightarrow 0} \Phi_+(\rho)$ and $\lim_{\rho \rightarrow 0} \Phi_-(\rho)$ are both $+\infty$ (or $-\infty$). In this case Γ_+ and Γ_- are two "concurrent" spirals having $\omega=0$ as an asymptotic point. Our results of §19 described the behavior of $\zeta(\omega)$ and $\zeta'(\omega)$ as ω approaches the asymptotic point.

However, our methods still apply to a case not included in §19, in which $\rho(d\Phi_+/d\rho)$ and $\rho(d\Phi_-/d\rho)$ both approach $+\infty$ (or $-\infty$). This case is contained in the following more general configuration.

Let R be a simply-connected (single-sheeted) region whose boundary consists of the spirals

$$\Gamma_+: \rho = \rho_+(\phi), \quad \Gamma_-: \rho = \rho_-(\phi), \quad \phi_0 \leq \phi < \infty,$$

and of a Jordan arc connecting the end points $(\phi_0, \rho_+(\phi_0))$ of Γ_+ and $(\phi_0, \rho_-(\phi_0))$ of Γ_- . It is assumed that $\rho_+(\phi)$ and $\rho_-(\phi)$ are positive and absolutely continuous in any interval $\phi_0 \leq \phi \leq \phi_1 < \infty$, and that $(1/\rho_+(\phi))d\rho_+/d\phi$ and $(1/\rho_-(\phi))d\rho_-/d\phi$ which exist for almost all $\phi \geq \phi_0$, both approach the limit $-\tan \gamma$, $|\gamma| < \pi/2$ as $\phi \rightarrow \infty$. The region

$$(21.1) \quad \phi_0 < \phi < \infty, \quad \rho_+(\phi) < \rho < \rho_-(\phi)$$

is contained in R . We set

$$\Theta(\phi) \equiv \log \frac{1}{\rho_+(\phi)} - \log \frac{1}{\rho_-(\phi)}, \quad \Psi(\rho) \equiv \frac{1}{2} \left[\log \frac{1}{\rho_+(\phi)} + \log \frac{1}{\rho_-(\phi)} \right].$$

There exists a function $\zeta(\omega)$ which maps R conformally onto the circle $|\zeta - 1| < 1$ in such a manner that $\lim \zeta(\omega) = 0$ as $\arg \omega \rightarrow +\infty$, ω being in the region (21.1). The inverse function of $\zeta(\omega)$ will again be denoted by $\omega(\zeta)$.

The function $w = u + iv = i \log(1/\omega) = i \log(1/\rho) + \phi$ maps R onto an L -strip S in the w -plane with the boundary inclination γ at $u = +\infty$. We map the circle $|\zeta - 1| < 1$ onto the strip $|y| < \pi/2$ by means of the function $z = \log \{(2 - \zeta)/\zeta\}$ ($z = 0$, when $\zeta = 1$). In this way we have again reduced our problem to that of a strip. We can carry over all our theorems on L -strips thus obtaining results on $\zeta(\omega)$ and $\omega(\zeta)$ as $\arg \omega \rightarrow +\infty$ for ω in (21.1) and $\zeta \rightarrow 0$ in $|\zeta - 1| < 1$.

To apply Theorem IX and to obtain the analogue of Theorem XI (B) we assume here that the integrals

$$\int_{\phi=\phi_0}^{\infty} \left| d \left(\frac{\rho'_+(\phi)}{\rho_+(\phi)} \right) \right|, \quad \int_{\phi=\phi_0}^{\infty} \left| d \left(\frac{\rho'_-(\phi)}{\rho_-(\phi)} \right) \right|, \quad \int_{\phi_0}^{\infty} \frac{\Theta'^2(\phi)}{\Theta(\phi)} d\phi$$

converge. There is no difficulty in actually writing out the results obtained for this case.

BIBLIOGRAPHY

L. AHLFORS

1. *Untersuchungen zur Theorie der konformen Abbildung und der ganzen Funktionen*, Acta Societatis Scientiarum Fennicae, (n. s.), A, vol. 1 (1930).

C. CARATHÉODORY

1. *Elementarer Beweis für den Fundamentalsatz der konformen Abbildung*, Schwarz-Festschrift, 1914, pp. 19–41.
2. *Conformal Representation*, Cambridge Tracts, 1932.

E. T. COPSON

1. *Introduction to the Theory of Functions of a Complex Variable*, Oxford University Press, 1935.

W. GROSS

1. *Zum Verhalten analytischer Funktionen in der Umgebung singulärer Stellen*, Mathematische Zeitschrift, vol. 2 (1918), pp. 241–294.

E. LINDELÖF

1. *Sur la représentation conforme d'une aire simplement connexe sur l'aire d'un cercle*, Comptes Rendu du Quatrième Congrès des Mathématiciens Scandinaves, Stockholm, (1916), pp. 59–90.

R. NEVANLINNA

1. *Eindeutige analytische Funktionen*, Berlin, 1936.

A. OSTROWSKI

1. *Über den Habitus der konformen Abbildung am Rande des Abbildungsbereiches*, Acta Mathematica, vol. 64 (1934), pp. 81–185.
2. *Zur Randverzerrung bei konformer Abbildung*, Prace Matematyczne Fizyczne, vol. 44 (1937), pp. 371–471.

T. RADÓ

1. *Sur la représentation conforme de domaines variables*, Acta Litterarum ac Scientiarum Universitatis Franciscio Josephinae, vol. 1 (1923), pp. 180–186.

S. E. WARSZAWSKI

1. *Zur Randverzerrung bei konformer Abbildung*, Compositio Mathematica, vol. 1 (1935), pp. 314–343.
2. *On the preservation of angles at a boundary point in conformal mapping*, Bulletin of the American Mathematical Society, vol. 42 (1936), pp. 674–680.

J. WOLFF

1. *Sur la représentation conforme des bandes*, Compositio Mathematica, vol. 1 (1934), pp. 217–222.
2. *Sur la conservation des angles dans la représentation conforme d'un domaine au voisinage d'un point frontière*, Comptes Rendus de l'Académie des Sciences, Paris, vol. 200 (1935), pp. 42–43.
3. *Démonstration d'un théorème sur la conservation des angles dans la représentation conforme d'un domaine au voisinage d'un point frontière*, Proceedings, Akademie van Wetenschappen, Amsterdam, vol. 38 (1935), pp. 46–50.

WASHINGTON UNIVERSITY,
ST. LOUIS, MO.