

A GENERALIZATION OF K. T. CHEN'S INVARIANTS FOR PATHS UNDER TRANSFORMATION GROUPS

BY
H. H. JOHNSON⁽¹⁾

In a series of papers [1; 2; 3; 4], K. T. Chen introduced and studied certain infinite series of numbers associated with paths in Euclidean n -space. These numbers were invariants under translations, and in [4] he proved that they uniquely characterize paths under translations.

This paper's purpose is investigating similar path invariants for other transformation groups. Integral invariants are studied using prolongations in the context of C. Ehresmann's "jets" [5].

A general discussion of the ideas involved is given in §1. §2 is devoted to differential equations and some illuminating examples. We find in §3 a sufficient condition that a group's invariants uniquely characterize paths. The theory also applies to pseudo groups having sufficiently regular behavior. All structures assumed C^∞ (infinitely differentiable).

1. INVARIANTS

1.1. Prolongations. Let M be a real C^∞ -manifold. Let G be a Lie group acting effectively as a transformation group of C^∞ -homeomorphisms on M . If $g \in G$, then $g : M \rightarrow M$. Let $\Theta = \{\theta\}$ be the Lie algebra of vector fields on M induced by the Lie algebra of G .

R is the real line, $t \in R$, $x \in M$. A k -jet $j_t^k(\tau)$ with source $t = \alpha(j_t^k(\tau))$ and target $x = \beta(j_t^k(\tau))$ is the equivalence class of all C^∞ -functions τ mapping a neighborhood of t into M , which satisfy $\tau(t) = x$ and have the first same k derivatives at t . If (x^1, \dots, x^n) are local coordinates on a neighborhood U of x , this k -jet will be specified by numbers $t, x^i, x_1^i, x_2^i, \dots, x_k^i$; $i = 1, \dots, n$, where

$$x_h^i = \frac{d^h x^i(\tau(t))}{dt^h}.$$

$J^k(M)$ will denote the k -jets of M . Then $(t, x^1, \dots, x^n, x_1^1, \dots, x_1^n, \dots, x_k^1, \dots, x_k^n)$ are local coordinates for $J^k(M)$ on $\beta^{-1}(U)$. There are natural C^∞ -projections $\rho_k^{k+r} : J^{k+r}(M) \rightarrow J^k(M)$, namely, $\rho_k^{k+r}(j_t^{k+r}(\tau)) = j_t^k(\tau)$.

If $g : M \rightarrow M$, there is a natural prolongation $p^k(g) : J^k(M) \rightarrow J^k(M)$, namely, $p^k(g)(j_t^k(\tau)) = j_t^k(g \circ \tau)$. Hence G can be prolonged to a transformation group on

Received by the editors July 13, 1961.

(1) Supported by OOR Contract DA-36-ORD-2164.

$J^k(M)$. Vector fields $\theta = (dg_t/dt)_{t=0}$ on M are prolonged to $p^k(\theta) = (dp^k(g_t)/dt)_{t=0}$ on $J^k(M)$. Let $p^k(\Theta) = \{p^k(\theta) \mid \theta \in \Theta\}$.

Similarly, if $h : R \rightarrow R$, there is a prolongation $p^k(h)(j_t^k(\tau)) = j_{h(t)}^k(\tau \circ h^{-1})$. This corresponds to changing the variable t . Vector fields ϕ on R may thus be prolonged to vector fields $p^k(\phi)$ on $J^k(M)$.

Later, exact expressions in local coordinates will be found for these prolongations. If $g : M \rightarrow M$ and $h : R \rightarrow R$, then $p^k(g) \circ p^k(h) = p^k(h) \circ p^k(g)$. Hence if θ and ϕ are vector fields on M and R respectively, then $[p^k(\theta), p^k(\phi)] = 0$. ($[\]$ denotes the Lie bracket.)

1.2. Paths. A path in M is a C^∞ map $\tau : [a, b] \rightarrow M$. If $\lambda : [c, d] \rightarrow [a, b]$ is C^∞ and increasing, $\tau \circ \lambda$ is considered equivalent to τ . There is a natural prolongation of τ to $p^k(\tau) : [a, b] \rightarrow J^k(M)$ defined by $p^k(\tau)(t) = j_t^k(\tau)$.

If t denotes the coordinate on R , $\alpha^*(dt)$ on $J^k(M)$ will be denoted more shortly by dt .

DEFINITION 1.1. A 1-form on an open set of $J^k(M)$ of the type Adt is *invariant* under G when for every $\theta \in \Theta$ or vector field ϕ on R , the Lie derivative of Adt with respect to $p^k(\theta)$ and $p^k(\phi)$ is zero. Note that if ψ is a vector field on $J^k(M)$, the Lie derivative of Adt with respect to ψ is

$$[\psi, Adt] = \langle \psi, dA \rangle dt + Ad(\langle \psi, dt \rangle).$$

This implies that $p^k(g) * (Adt) = Adt$ for all $g \in G$.

1.3. Chen constants. The *Chen constants* for a path $\tau : [a, b] \rightarrow M$ with respect to 1-forms $A_1 dt, A_2 dt, \dots, A_r dt$ on $J^k(M)$ are defined to be the infinite set of numbers

$$\int_\tau A_{i_1} dt \cdots A_{i_j} dt, \quad 1 \leq i_j \leq r; \quad j = 1, 2, \dots$$

These integrals are defined inductively by

$$\int_\tau A_i dt = \int_a^b (p^k(\tau)) * (A_i dt) = \int_a^b A_i(j_t^k(\tau)) dt,$$

and if $\tau_t = \tau \mid [a, t]$,

$$\int_\tau A_{i_1} dt \cdots A_{i_j} dt = \int_a^b \left[\int_{\tau_t} A_{i_1} dt \cdots A_{i_{j-1}} dt \right] (p^k(\tau)) * (A_{i_j} dt).$$

Following Chen [4], these $A_1 dt, \dots, A_r dt$ define an "exponential" mapping θ on paths to the space of formal, noncommutative power series in indeterminants X_1, \dots, X_r , over R :

$$\theta(\tau) = 1 + \sum_{j=1}^\infty \sum \left(\int_\tau A_{i_1} dt \cdots A_{i_j} dt \right) X_{i_1} \cdots X_{i_j}.$$

(These integrals are of course defined only when $p^k(\tau)([a,b])$ is contained in the domains of $A_1 dt, \dots, A_r dt$.)

THEOREM 1.1. *Let $A_1 dt, \dots, A_r dt$ be invariant under G . If τ is any path in M , g any element of G , then the paths τ and $g \circ \tau$ have the same Chen constants.*

Proof.

$$\begin{aligned} \int_{g \circ \tau} A_j dt &= \int_a^b A_j(j_i^k(g \circ \tau)) dt = \int_a^b A_j \circ p^k(g)(j_i^k(\tau)) dt \\ &= \int_a^b A_j(j_i^k(\tau)) dt = \int_{\tau} A_j dt. \end{aligned}$$

Since $(g \circ \tau)_t = g \circ \tau_t$, equality for the other Chen constants also follows.

2. DIFFERENTIAL EQUATIONS

Next we study the partial differential equations satisfied by invariant 1-forms $A dt$. All functions and forms are still assumed to be infinitely differentiable.

2.1. Formal derivatives. The following construction is due to M. Kuranishi. A 1-form ω on an open subset of $J^k(M)$ is *canonical* when $(p^k(\tau)) * (\omega) = 0$ for all paths τ . Let Ω^k denote the set of canonical 1-forms. Take local coordinates (x^1, \dots, x^n) on the open set U in M . Then on $\beta^{-1}(U)$, $J^k(M)$ has local coordinates $(t, x^i, x_1^i, \dots, x_k^i)$, $i = 1, \dots, n$. If $\tau : (t) \rightarrow (x^i(t))$, then $p^k(\tau) : (t) \rightarrow (t, x^i(t), dx^i(t)/dt, \dots, d^k x^i(t)/dt^k)$.

Consider the 1-forms $\pi_h^i = dx_h^i - x_{h+1}^i dt$; $h = 1, \dots, k - 1$, and $\pi_0^i = dx^i - x_1^i dt$; $i = 1, \dots, n$, on $\beta^{-1}(U)$.

$$(p^k(\tau)) * (\pi_h^i) = d \left(\frac{d^h x^i}{dt^h} \right) - \frac{d^{h+1} x^i}{dt^{h+1}} dt = 0.$$

Hence $\pi_h^i \in \Omega^k$ for $h = 0, 1, \dots, k - 1$. In fact, it may be shown that any $\omega \in \Omega^k|_U$ is $\equiv 0$ modulo $\{\pi_h^i\}$ over the real-valued functions.

If $g \in G$ and $\omega \in \Omega^k$, and τ is any path in M , then $(p^k(\tau)) * ((p^k(g)) * (\omega)) = (p^k(g \circ \tau)) * (\omega) = 0$, since $g \circ \tau$ is a path in M . Hence, Ω^k is invariant.

It follows that if $\theta \in \Theta$ is defined on an open subset of U , then $[p^k(\theta), \pi_h^i] \equiv 0$ modulo $\{\pi_h^i\}$. Similar reasoning shows that if ϕ is a vector field on R , then $[p^k(\phi), \pi_h^i] \equiv 0$ modulo $\{\pi_h^i\}$.

If F is a real-valued C^∞ -function on an open subset V of $J^k(M)$, then on $(\rho_k^{k+1})^{-1}(V)$ we define ∂F by

$$dF \equiv (\partial F) dt \text{ modulo } (\Omega^{k+1}).$$

When $V \subset \beta^{-1}(U)$,

$$\partial F = \frac{\partial F}{\partial t} + \frac{\partial F}{\partial x^j} x_1^j + \dots + \frac{\partial F}{\partial x_k^j} x_{k+1}^j.$$

The following lemmas may be proved by induction using the canonical forms $\{\pi_h^i\}$.

LEMMA 2.1. *If $\theta = \theta^j(\partial/\partial x^j)$ is a vector field on U , then*

$$p^k(\theta) = \theta + \sum_{h=1}^k \partial^h \theta^j \frac{\partial}{\partial x_h^j}.$$

LEMMA 2.2. *If $\phi = \xi(t)(\partial/\partial t)$ is a vector field on R , then on $\beta^{-1}(U)$,*

$$p^k(\phi) = \phi - \sum_{h=1}^k \frac{d^h \xi}{dt^h} \left[\sum_{m=h}^k \binom{m}{h} x_{m-h+1}^j \frac{\partial}{\partial x_m^j} \right].$$

2.2. **Differential equations for integral invariants.** Let Adt be an invariant 1-form on an open subset of $\beta^{-1}(U)$. If the Lie algebra Θ is generated by $\theta_\lambda = \theta_\lambda^j(\partial/\partial x^j)$, $\lambda = 1, \dots, \lambda_0$, A must satisfy

$$0 = [p^k(\theta_\lambda), Adt] = \langle p^k(\theta_\lambda), dA \rangle dt + Ad(\langle p^k(\theta_\lambda), dt \rangle),$$

or,

$$(E_1) \quad \theta_\lambda^j \frac{\partial A}{\partial x^j} + \sum_{h=1}^k \partial^h \theta_\lambda^j \frac{\partial A}{\partial x_h^j} = 0, \quad \lambda = 1, \dots, \lambda_0.$$

Similarly, for every choice of $\xi(t)$ on R , A must satisfy

$$0 = \xi \frac{\partial A}{\partial t} - \sum_{h=1}^k \frac{d^h \xi}{dt^h} \left[\sum_{m=h}^k \binom{m}{h} x_{m-h+1}^j \frac{\partial A}{\partial x_m^j} \right] + A \frac{d\xi}{dt}.$$

Taking successively $\xi = 1, t, t^2, \dots$, one obtains

$$(E_2) \quad \begin{aligned} \frac{\partial A}{\partial t} &= 0, \\ A &= \sum_{m=1}^k m x_m^j \frac{\partial A}{\partial x_m^j}, \\ \sum_{m=h}^k \binom{m}{h} x_{m-h+1}^j \frac{\partial A}{\partial x_m^j} &= 0, \quad h = 2, \dots, k. \end{aligned}$$

Thus, the following theorem is true.

THEOREM 2.1. *The 1-form Adt defined on an open subset of the coordinate neighborhood $\beta^{-1}(U)$ is invariant under G if and only if equations (E_1) and (E_2) hold.*

It is natural to ask whether more information could be gained by studying more general invariant 1-forms on $J^k(M)$. The following may be proved by induction on k .

PROPOSITION 2.1. *If*

$$\omega = A dt + A_j dx^j + \sum_{h=1}^k A_j^h dx_h^j$$

is invariant on $\beta^{-1}(U)$ in $J^k(M)$, then

$$\omega' = \left(A + A_j x_1^j + \sum_{h=1}^k A_j^h x_{h+1}^j \right) dt$$

is invariant on $J^{k+1}(M)$.

2.3. **Complete integrability.** Instead of solving equations (E₁) and (E₂) it is customary [6, §75] to introduce a new variable V and a new system (E') with V as an unknown function of $A, t, x^i, x_1^i, \dots, x_k^i; i = 1, \dots, n$; namely,

$$\begin{aligned} \theta_\lambda^j \frac{\partial V}{\partial x^j} + \sum_{h=1}^k \partial^h \theta_\lambda^j \frac{\partial V}{\partial x_h^j} &= 0, & \lambda = 1, \dots, \lambda_0, \\ \frac{\partial V}{\partial t} &= 0, \\ \text{(E')} \quad A \frac{\partial V}{\partial A} + \sum_{h=1}^k h x_h^j \frac{\partial V}{\partial x_h^j} &= 0, \\ \sum_{m=h}^k \binom{m}{h} x_{m-h+1}^j \frac{\partial V}{\partial x_m^j} &= 0, & h = 2, \dots, k. \end{aligned}$$

THEOREM 2.2. *The system (E') is completely integrable.*

Proof. The following lemmas may be proved by induction arguments, after which the theorem can be proved by direct computation.

Let F be a real-valued C^∞ -function on an open subset of U .

LEMMA 2.3. *If $h \leq k$,*

$$h \partial^h F = \sum_{m=1}^k m x_m^i \frac{\partial (\partial^h F)}{\partial x_m^i}.$$

LEMMA 2.4.

$$\partial^h F = \frac{\partial F}{\partial x_h^j} x_h^j + G,$$

where G is a function on $J^{h-1}(M)$.

LEMMA 2.5.

$$\sum_{q=m}^k \binom{q}{m} x_{q-m+1}^j \frac{\partial (\partial^h F)}{\partial x_q^j} = \begin{cases} 0 & \text{if } h < m, \\ \binom{h}{m} \partial^{h-m+1} F & \text{if } h \geq m. \end{cases}$$

2.4. **Examples.** (a) *Translations.* Here, $M = E^n =$ Euclidean n -space. Θ is generated by $\{\partial/\partial x^i \mid i = 1, \dots, n\}$. $p^k(\Theta) = \Theta$. When $k = 1$, (E') becomes

$$\frac{\partial V}{\partial t} = \frac{\partial V}{\partial x^i} = 0; \quad i = 1, \dots, n,$$

$$A \frac{\partial V}{\partial A} + x_1^j \frac{\partial V}{\partial x_1^j} = 0,$$

having solutions $V^j = x_1^j/A$; $j = 1, \dots, n$. Hence the invariant 1-forms on $J^1(M)$ are $x_1^j dt$. The Chen constants of paths with respect to these invariant 1-forms are the original invariants found by Chen.

(b) *Motions.* Let $M = E^2 = \{(x^1, x^2)\}$. Θ is generated by $\partial/\partial x^1, \partial/\partial x^2$, and $x^2(\partial/\partial x^1) - x^1(\partial/\partial x^2)$. $p^2(\Theta)$ is generated by $\partial/\partial x^1, \partial/\partial x^2$, and $x^2(\partial/\partial x^1) - x^1(\partial/\partial x^2) + x_1^2(\partial/\partial x_1^1) - x_1^1(\partial/\partial x_1^2) + x_2^2(\partial/\partial x_2^1) - x_2^1(\partial/\partial x_2^2)$. For $k=2$ equations (E') yield two independent solutions,

$$\frac{((x_1^1)^2 + (x_1^2)^2)^{1/2}}{A}, \quad \frac{x_1^1 x_2^2 - x_1^2 x_2^1}{A[(x_1^1)^2 + (x_1^2)^2]}.$$

Hence if ds denotes arc length and $1/\rho$ denotes curvature, the invariant 1-forms on $J^2(M)$ are generated by $((x_1^1)^2 + (x_1^2)^2 dt)^{1/2} = ds$ and

$$\{(x_1^1 x_2^2 - x_1^2 x_2^1) / [(x_1^1)^2 + (x_1^2)^2]\} dt = 1/\rho ds.$$

In [4] Chen proved that the invariant 1-forms of example (a) are *complete*, i.e., they essentially characterize paths under translations. This cannot be true in general, especially if one allows the paths to have isolated discontinuities, as Chen does. For example,

$$\tau_1(t) = \begin{cases} (t, 0), & 0 \leq t \leq 1/2, \\ (1-t, 0), & 1/2 \leq t \leq 1 \end{cases} \quad \text{and} \quad \tau_2(t) = \{(t, 0) \mid 0 \leq t \leq 1\}$$

have the same Chen constants with respect to ds and $(1/\rho)ds$, but cannot be obtained from one another by motions. To show that this completeness may fail no matter how large k may be taken, consider the next example.

(c) *Special linear group.* $M = E^2, \Theta$ is generated by

$$\frac{\partial}{\partial x^1}, \frac{\partial}{\partial x^2}, x^2 \frac{\partial}{\partial x^1}, x^1 \frac{\partial}{\partial x^2}, x^1 \frac{\partial}{\partial x^1} - x^2 \frac{\partial}{\partial x^2}.$$

For any k , equations (E_1) and (E_2) contain the equations

$$\frac{\partial A}{\partial x^1} = \frac{\partial A}{\partial x^2} = x^2 \frac{\partial A}{\partial x^1} + \sum_{h=1}^k x_h^2 \frac{\partial A}{\partial x_h^1} = 0,$$

$$x^1 \frac{\partial A}{\partial x^2} + \sum_{h=1}^k x_h^1 \frac{\partial A}{\partial x_h^2} = 0,$$

$$A = \sum_{h=1}^k h \left(x_h^1 \frac{\partial A}{\partial x_h^1} + x_h^2 \frac{\partial A}{\partial x_h^2} \right).$$

Suppose $A dt$ is invariant. Let $\tau: (t) \rightarrow (t, t), 0 \leq t \leq 1$. Then $x_h^1(p^k(\tau)) = x_h^2(p^k(\tau)) = 0$ if $h > 1$ and $x_1^1(p^k(\tau)) = x_1^2(p^k(\tau)) = 1$.

One finds from the above equations that $A(p^k(\tau)) = 0$. This means all the Chen constants for τ are the same as for the constant path. Since the special linear group is a subgroup of the general projective group, the same conclusion holds for that case.

(d) *The existence of invariants depends on how the group acts.* Let $G = \{g = (a, b) \mid a \in R, b \in R\}$ and $(a, b) \cdot (a', b') = (aa', a'b + b')$.

If G acts on E^2 by $g(x^1, x^2) = (x^1, ax^2 + b)$, then Θ is generated by $\partial/\partial x^2$ and $x^2(\partial/\partial x^2)$. As in example (c) the path $\tau: (t) \rightarrow (0, t), 0 \leq t \leq 1$, has $A(p^k(\tau)) = 0$ for any invariant 1-form $A dt$ on $J^k(M)$.

However, if G acts by $g(x^1, x^2) = (ax^1 + b, ax^2)$, then Θ is generated by $\partial/\partial x^1$ and $x^1(\partial/\partial x^1) + x^2(\partial/\partial x^2)$. Hence $(x_1^1/x^2)dt$ and $(x_1^2/x^2)dt$ are invariant on $J^1(M)$ where $x^2 \neq 0$. The only path (not passing through $x^2 = 0$) on which these invariants are identically zero is the constant path.

3. COMPLETENESS

3.1. Algebraic theory. Chen's invariants for translation groups possess remarkable algebraic properties. Chen [4] considered paths which were piecewise "regular," i.e., had nonvanishing tangent vectors at all but a finite number of points and a finite number of jump discontinuities. Then the "product" $\tau \cdot \sigma$ of two paths τ and σ could be defined as well as "the inverse" τ^{-1} .

Such a course could be imitated here, defining "piecewise C^∞ -paths" in the obvious way. As in [4], one can prove that the Chen constants for any invariant 1-forms under a group G satisfy $\theta(\tau \cdot \sigma) = \theta(\tau) \cdot \theta(\sigma)$. However, $\theta(\tau \cdot \tau^{-1})$ need not be 1 (cf. example (b) above). Hence $\theta(\tau^{-1})$ need not be $\theta(\tau)^{-1}$.

3.2. Characterization. Chen's invariants in [4] characterize paths under translations: $\theta(\tau) = \theta(\sigma)$ implies τ and σ are obtained from each other by translations and such operations as multiplication by $\gamma \cdot \gamma^{-1}$ which do not change θ . Example (c) shows that this property fails in general. Next, a criterion will be given under which it does hold. (Recall that our paths are C^∞ , hence such paths as $\gamma \cdot \gamma^{-1}$ do not enter into the considerations.)

DEFINITION 3.1. Let $A_1 dt, \dots, A_r dt$ be invariant 1-forms under G on an open set $U \subset J^k(M)$. Assume that for all sets f_1, \dots, f_r of r real-valued C^∞ -functions defined on an open set $V \subset R$, the set

$$\mathcal{U} = \{X \in \alpha^{-1}(V) \cap U \mid A_1(X) = f_1(\alpha(X)), \dots, A_r(X) = f_r(\alpha(X))\}$$

satisfies the following three conditions:

- (1) \mathcal{U} is a submanifold of $\alpha^{-1}(V) \cap U$,
- (2) $p^k(G)$ acts transitively on \mathcal{U} ,
- (3) Ω^k , the canonical 1-forms on $J^k(M)$, satisfy $\dim \Omega^k|_{\mathcal{U}} = \dim \mathcal{U} - 1$ at all points of \mathcal{U} .

Then G is called *complete of order k with respect to $A_1 dt, \dots, A_r dt$ on U* .

THEOREM 3.1. *Let G be complete of order k with respect to $A_1 dt, \dots, A_r dt$ on U . Let σ and τ be two C^∞ -paths on M such that $p^k(\sigma)$ and $p^k(\tau)$ lie in U . If the Chen constants of σ and τ with respect to $A_1 dt, \dots, A_r dt$ are identical, then $\tau = g \circ \sigma$ for some $g \in G$.*

Proof. Suppose σ and τ have domains $[a, b]$ and $[c, d]$, respectively. Define C^∞ -paths σ' and τ' in $E^r = \{(y^1, \dots, y^r)\}$ by

$$\sigma'(t) = \left(\int_a^t A_1 \circ p_s^k(\sigma) ds, \dots, \int_a^t A_r \circ p_s^k(\sigma) ds \right),$$

$a \leq t \leq b$; and similarly for τ' .

Now, the Chen constants for σ' under the translation group on E^r coincide with those for σ with respect to $A_1 dt, \dots, A_r dt$:

$$\begin{aligned} \int_{\sigma'} dy^{i_1} \dots dy^{i_j} &= \int_a^b A_{i_1} \circ p_s^k(\sigma) ds \dots A_{i_j} \circ p_s^k(\sigma) ds \\ &= \int_{\sigma} A_{i_1} dt \dots A_{i_j} dt; \end{aligned}$$

similarly for τ' and τ . By hypothesis, these constants for σ' and τ' are the same. Since σ' and τ' are C^∞ -paths, they are irreducible, and it follows from [4, Theorem 4.1] that τ' can be obtained from σ' by a translation. However, $\sigma'(a) = (0, \dots, 0) = \tau'(c)$. Hence σ' and τ' coincide. Hence, one can change parameters so that σ' and τ' are both defined on $[a, b]$ and agree completely: $\sigma'(t) = \tau'(t)$.

After this change, let $f_1(t) = A_1 \circ p_t^k(\sigma) = A_1 \circ p_t^k(\tau), \dots, f_r(t) = A_r \circ p_t^k(\sigma) = A_r \circ p_t^k(\tau)$, for $a < t < b$. Then

$$\mathcal{U} = \{X \in \alpha^{-1}((a, b)) \mid A_1(X) = f_1(\alpha(X)), \dots, A_r(X) = f_r(\alpha(X))\}$$

is a submanifold of $\alpha^{-1}((a, b))$ containing $j_t^k(\sigma)$ and $j_t^k(\tau)$ for $a < t < b$. Since $p^k(G)$ is transitive on \mathcal{U} , there exists $g \in G$ such that for any fixed $t_0 \in (a, b)$, $j_{t_0}^k(\tau) = j_{t_0}^k(g \circ \sigma)$.

Condition (3) of Definition 3.1 implies that the system of exterior differential forms

$$\begin{cases} dx^i - x_1^i dt, & i = 1, \dots, n, \\ \vdots \\ dx_{k-1}^i - x_k^i dt, & i = 1, \dots, n, \end{cases}$$

restricted to \mathcal{U} , is completely integrable and has unique solutions of degree 1 through each point of \mathcal{U} . $j_i^k(g \circ \sigma)$ and $j_i^k(\tau)$ are two such solutions through t_0 . Therefore, $j_i^k(g \circ \sigma) = j_i^k(\tau)$ in a neighborhood of t_0 . Hence $g \circ \sigma(t) = \beta(j_i^k(g \circ \sigma)) = \beta(j_i^k(\tau)) = \tau(t)$ in a neighborhood of t_0 . Since t_0 is arbitrary, this equality holds on (a, b) , and by continuity on $[a, b]$.

COROLLARY. Let $\tau_i : [a_i, b_i] \rightarrow M, i = 1, 2$, be two C^∞ -paths in the plane for which $(x_1^1 \circ p_i^1(\tau_i))^2 + (x_2^1 \circ p_i^1(\tau_i))^2 \neq 0$ for $t \in [a_i, b_i]$. Let $s_i(t)$ denote arc length of τ_i measured from a_i to t , $K_i(t)$ the curvature of τ_i at t . Then if

$$F_{i1} = \frac{ds_i}{dt}, \quad F_{i2} = K_i \frac{ds_i}{dt}, \quad i = 1, 2,$$

satisfy

$$\int_{a_1}^{b_1} F_{1i_1} dt \cdots F_{1i_j} dt = \int_{a_2}^{b_2} F_{2i_1} dt \cdots F_{2i_j} dt$$

for all $i_1, \dots, i_j = 1, 2; j = 1, 2, 3, \dots$, then τ_1 may be obtained from τ_2 by a motion.

Proof. It suffices to observe that the group of motions is complete of order 2 with respect to $A_1 dt = ((x_1^1)^2 + (x_2^1)^2)^{1/2} dt$ and

$$A_2 dt = \{(x_1^1 x_2^2 - x_2^1 x_1^2) / [(x_1^1)^2 + (x_2^1)^2]\} dt$$

on $U = \{X \in J^2(M) \mid (x_1^1)^2 + (x_2^1)^2 \neq 0\}$.

BIBLIOGRAPHY

1. K. T. Chen, *Integration in free groups*, Ann. of Math. **54** (1951), 147-162.
2. ———, *Iterated integrals and exponential homomorphisms*, Proc. London Math. Soc. (3) **4** (1954), 502-512.
3. ———, *Integration of paths — geometric invariants and a generalized Baker-Housdorff formula*, Ann. of Math. **65** (1957), 163-178.
4. ———, *Integration of paths — a faithful representation of paths by noncommutative formal power series*, Trans. Amer. Math. Soc. **89** (1958), 395-407.
5. C. Ehresmann, *Introduction à la théorie des structures infinitésimales et des pseudo-groupes de Lie*, Colloque International de Géométrie Différentielle du C.N.R.S., Paris, 1953.
6. E. Goursat, *A course in mathematical analysis*, Vol. II, Part 2, Trans. by E. R. Hedrick and O. Dunkel, Dover, N. Y., 1917.

PRINCETON UNIVERSITY,
 PRINCETON, NEW JERSEY
 UNIVERSITY OF WASHINGTON,
 SEATTLE, WASHINGTON