

ON THE NUMBER OF SINGULARITIES OF AN ANALYTIC CURVE

BY
HENRY B. LAUFER⁽¹⁾

I. Introduction. Burau [1] and Zariski [7] showed that algebroid singularities can be classified by the knot type obtained by looking at that part of the curve lying on the boundary of a small polydisc (or ball) centered at the singularity. In this paper, by examining the isotopy class of the link obtained by looking at that part of an analytic curve lying on the boundary of an unbounded polydisc, we will obtain information about the genus and the number of singularities of that part of the curve lying within the polydisc. Thus, we get global instead of local results.

As an application to algebraic geometry, we give a topological proof of a Plücker formula. Also, for many proper maps of Riemann surfaces into \mathbf{C}^2 , we can count the number of points where the map fails to be regular or one-one.

The complex analysis and the knot theory are in different sections, with independent proofs.

II. Counting singularities. Let S be a one-dimensional analytic subvariety in a neighborhood of $B_R = \{(z_1, z_2) \in \mathbf{C}^2 \mid |z_1| \leq R\}$ such that $S \cap B_R$ is compact. More generally, since we are not concerned with S outside of B_R , S can be the image of a continuous map $F = (f_1, f_2): M \rightarrow B_R$, where M is a compact complex 1-manifold with boundary, $F(\partial M) \subset \partial B_R$, F is holomorphic on the interior of M , and F is one-one on ∂M , the topological boundary of M . Topologically ∂B_R is the product of a 1-sphere S^1 and an open 2-cell E^2 . We shall coordinatize ∂B_R by z_2 and the argument of z_1 . F restricted to ∂M gives a link $L_R: \partial M \rightarrow \partial B_R$. We shall use L_R to refer to either the map L_R or to its image. We want to count the singularities of F using topological properties of L_R .

All links in this paper will have a finite number of components.

To see what L_R looks like, we approximate L_R by $L_{R-\varepsilon}$, which is F restricted to $F^{-1}(\partial B_{R-\varepsilon}) = \partial M_{R-\varepsilon}$. Choose ε so that $f'_1(z) \neq 0$ on $\partial M_{R-\varepsilon}$. As we traverse a component of $\partial M_{R-\varepsilon}$ in the positive direction, $\arg z_1$ in the image is strictly monotone increasing since f_1 is orientation preserving and a local homeomorphism. $H_1(\partial B_{R-\varepsilon}; \mathbf{Z}) \approx \mathbf{Z}$ and we choose a generator which has an increasing z_1 argument. Then the homology class of $L_{R-\varepsilon}$ is, by the argument principle, equal in number to the number of zeroes of f_1 . Since as $\varepsilon \rightarrow 0$, $L_{R-\varepsilon} \rightarrow L_R$, for small enough ε (see

Received by the editors December 4, 1967.

(¹) Research sponsored by the Air Force Office of Scientific Research, Office of Aerospace Research, under Contract No. F44620-67-6-0029.

Lemma 2 for the case of arbitrary links) the links $L_{R-\varepsilon}$ are all isotopic via homeomorphisms which have strictly monotone increasing z_1 arguments.

DEFINITION 1. Let $L = (f_1, f_2): \bigcup S^1_i \rightarrow \partial B_R$ be a link with a monotone increasing z_1 argument. Approximate L by differentiable links \bar{L}_ε having positive \bar{f}_1 derivatives. The self-winding $sw(L)$ of L is

$$(1) \quad \lim_{\varepsilon \rightarrow 0} sw(\bar{L}_\varepsilon) = \lim_{\varepsilon \rightarrow 0} \frac{1}{2\pi} \int_{\bigcup S^1_i} \sum_{z'} d(\arg(\bar{f}_2(z') - \bar{f}_2(z)),$$

where \bar{f}_1 and \bar{f}_2 are the coordinates for \bar{L}_ε and where, for fixed z , the sum is over all z' such that $\bar{f}_1(z') = \bar{f}_1(z)$ and $z' \neq z$.

Thus $sw(L)$ looks at the change in argument between points having the same z_1 coordinate and is integer valued. Since $sw(\bar{L}_\varepsilon)$ depends continuously on \bar{L}_ε and is integer valued, $sw(L)$ is well defined and is unchanged if we deform L through links with monotone increasing z_1 argument. In the next section, we shall see that $sw(L)$ is an isotopy invariant that can be defined for any link in $S^1 \times E^2$.

DEFINITION 2. Let L and L' be links in ∂B_R with monotone increasing z_1 arguments. Approximate L and L' by differentiable links \bar{L}_ε and \bar{L}'_ε with positive \bar{f}_1 derivatives.

$$(2) \quad l(L, L') = \lim_{\varepsilon \rightarrow 0} l(\bar{L}_\varepsilon, \bar{L}'_\varepsilon) = \lim_{\varepsilon \rightarrow 0} \frac{1}{2\pi} \int \sum_{z'} d(\arg \bar{f}_2(z') - \bar{f}_2(z)),$$

where \bar{f}_1 and \bar{f}_2 are the coordinates and where, for fixed $z \in \bar{L}_\varepsilon$, the sum is over z' such that $\bar{f}_1(z') = \bar{f}_1(z)$ and $z' \in \bar{L}'_\varepsilon$. The integration is over z in \bar{L}_ε .

$l(L, L')$ is just the usual linking number if we embed $S^1 \times E^2$ in S^3 in the usual manner. In our case, this corresponds to replacing ∂B_R by

$$S^3 = \{(z_1, z_2) \mid \sup |z_1|, |z_2|/N = R\}$$

where N is chosen sufficiently large so that L and L' lie in S^3 .

We must assign multiplicities to the singularities of F . Nodes (i.e. points in the image of F where the image is locally two intersecting submanifolds with different tangent planes) receive multiplicity one. Any singularity can be approximated by some fixed positive number of nodes (see Corollary 2), which is its multiplicity.

Statement (4) below is a weaker result than one of Martinelli [4], who shows that B_R can be quite arbitrary.

THEOREM 1. Let $F = (f_1, f_2): M \rightarrow B_R$ be a continuous map from M , a compact Riemann surface with boundary to $B_R = \{(z_1, z_2) \in \mathbf{C}^2 \mid |z_1| \leq R\}$ such that $F(\partial M) \subset \partial B_R$, F is holomorphic on the interior of M , and F is one-one on ∂M . F restricted to ∂M defines a link L . Then

$$(3) \quad sw(L) = (h-1) + (\mu-1) + 2g + 2d,$$

where h is the homology class of L , μ is the number of components of L , g is the genus of M , and d is the number of singularities of F .

If F_1 and F_2 are two such maps, then

(4) $l(L_1, L_2)$ is the number of intersections of $F(M_1)$ with $F(M_2)$.

Proof. We can first approximate F arbitrarily closely on the interior of M so that the only singularities of the map are nodes and so that $f'_1(z)$ has only simple zeroes. $L_{R-\epsilon}$ is deformed only slightly so that $\text{sw}(L_{R-\epsilon})$ is unchanged. Singularities are replaced by the appropriate number of nodes. Thus we may assume that the only singularities of F are nodes, where we must replace B_R by $B_{R-\epsilon}$. There can be no singularities near ∂M since as $\epsilon \rightarrow 0$, $\text{sw}(L_{R-\epsilon})$ is constant and thus d on the right side of (3) cannot increase.

On M , form the function $p(z) = \prod_{z'} f_2(z') - f_2(z)$ where $z' \neq z$ and $f_1(z') = f_1(z)$. $p(z)$ is holomorphic except possibly at a z such that $f'_1(z') = 0$ for some z' with $f_1(z') = f_1(z)$. Since $p(z)$ is locally bounded, it must have a removable singularity at such a z .

$2\pi \text{sw}(L)$ is nothing more than the total change in argument for $p(z)$ as z traverses ∂M . Thus $\text{sw}(L)$ equals the number of zeroes of $p(z)$. $p(z) = 0$ can occur only when $f_2(z) = f_2(z')$, $f_1(z) = f_1(z')$, $z \neq z'$, i.e. at a node, or when $z' \rightarrow z$, i.e. where $f'_1(z) = 0$. Every node yields two zeroes, one for each of the points which are not separated. For the other possibility, $f'_1(z) = 0$ is a simple zero and $f'_2(z) \neq 0$ by our approximation assumption. If $f_1(z)$ is thus locally z^2 , then $z' = -z$ and $p(z)$ has a simple zero.

Look at $F^{-1}(\partial B_r) = \{z \in M \mid |f_1(z)| = r\}$. When $r = 0$ we get h points and for small r , h circles (we may assume that $f_1(z)$ and $f'_1(z)$ have disjoint zeroes). Let r increase to $R - \epsilon$. The number of components of $F^{-1}(\partial B_r)$ increases or decreases by one each time we go through a z such that $f'_1(z) = 0$. The number of components cannot change at other z since they can only combine or divide when f_1 is not a local homeomorphism, can only, by the minimum principle, appear at points where $f_1(z) = 0$, and, by the maximum principle, cannot contract to points and disappear. We start with h components, end with μ components and have g handles. Since M is connected, there must be $(h-1) + 2g + (\mu-1)$ places where $f'_1(z) = 0$ ($(h-1)$ points to join the h components, 2 points for each of the handles, and $(\mu-1)$ points to end with μ components, although generally not in that order). Hence (3) is proved:

(4) is easier to prove. Let, for $z \in M_1$, $g(z) = \prod_{z'} f_2(z') - f_2(z)$ where $z' \in M_2$ and $f_1(z) = f_1(z')$. Now argue as before.

If we have a singularity at the origin having one branch and $z_2 = 0$ gives the tangent line, then the knot corresponding to the singularity necessarily lies in ∂B_ϵ , for small ϵ . We shall see in the next section that generalizations of $\text{sw}(\cdot)$ serve to classify such knots in $S^1 \times E^2$.

COROLLARY 2. Let $K \subset \partial B_\epsilon$ be the knot associated to a singularity with one branch. Then $2d = \text{sw}(K) - (h-1)$ where d is the multiplicity of the singularity and h is the homology class of the knot.

Proof. Locally the singularity is given by $F=(z^h, f_2): D \rightarrow B_\varepsilon$, where D is a disc. There is one component to the knot and D is simply connected.

COROLLARY 3 (PLÜCKER). *Let C be an irreducible algebraic curve of degree m and genus g having only cusps and nodes as singularities. Then*

$$g = \frac{1}{2}(m-1)(m-2) - s - n,$$

where s is the number of cusps and n is the number of nodes.

Proof. Since a cusp may be parametrized by (z^2, z^3) and has as corresponding knot a torus knot of type $(2, 3)$, the multiplicity of a cusp is one.

Choose the line l at infinity so that $C \cap l$ is m distinct points, having homogeneous coordinates $(1, p_i, 0)$, $1 \leq i \leq m$. Then as $R = |z_1| \rightarrow \infty$, looking at $\partial B_R \cap C$, $z_2/z_1 \rightarrow p_i$ and we get m distinct circles, each with homology class one and each pair having linking number one. There are $m(m-1)$ ordered pairs of circles, so by (3)

$$m(m-1) = (m-1) + (m-1) + 2g + 2n + 2s.$$

We can get some information about trying to embed Riemann surfaces to \mathbf{C}^2 . See [3, pp. 220–221] for the construction of proper maps of the needed type.

COROLLARY 4. *Let $F=(f_1, f_2): S \rightarrow \mathbf{C}^2$ be a proper holomorphic map of S , an open Riemann surface of finite genus g with μ boundary components, such that f_1 is an almost proper map. Let L_R be links corresponding to components M_R of*

$$\{z \in S \mid |f_1(z)| \leq R\}$$

such that $M_R \subset M_{R+1}$ and $\bigcup M_R = S$. Then F is an embedding if and only if for all sufficiently large R

$$(5) \quad \text{sw}(L_R) = (h_R - 1) + (\mu - 1) + 2g.$$

Proof. The maximum principle shows that M_R is, for sufficiently large R , a Riemann surface with boundary of genus g and with μ boundary components. Since every singularity of F has multiplicity at least one, (5) is just the condition that there be no singularities.

III. Isotopy invariance. If we deform a link L in $T = S^1 \times E^2$, (1) will not, in general, suffice to define $\text{sw}(L)$ since $\arg z_1$ need not be increasing and singularities may occur in the integral, even assuming that L remains differentiable. We shall define, in the topological category, i.e. for all continuous embeddings $L: \bigcup_i S^1_i \rightarrow T$, an isotopy invariant $\text{sw}(L)$ which coincides with (1) when L has an increasing z_1 argument. L , or more precisely the image of L , need not be connected, although stronger invariants may be obtained by looking at each component of L separately. We retain our parametrization of

$$T = B_1 = \{(z_1, z_2) \in \mathbf{C}^2 \mid |z_1| = 1\}.$$

Let $\{L_i\}$ be the components of L . Let h_i be the homology class of L_i . To every point (z_1, z_2) of L_i assign $\arg z_1$ which, after starting somewhere and varying the argument continuously, is well defined modulo $2\pi h_i$. To generalize $sw(L)$ from (1), start at $p \in L$ and $p' \in L$ such that $\arg p \equiv \arg p'$, mod 2π and p and p' are in different components or $\arg p \not\equiv \arg p'$, mod $2\pi h_i$. Look at the argument of the z_2 coordinate of $p' - p$ and follow the change in argument as p varies around L . When the derivative of $\arg p'$ changes sign we reverse the direction in which we are traversing L . Eventually we return to our starting pair. $sw(L)$ is the sum of all changes in the z_2 argument which occur in this manner. Thus,

DEFINITION 3. Let L be a polygonal link, $L = (f_1, f_2) : \bigcup_i S_i^1 \rightarrow T$ such that f_1 is not constant on any segment of L . Let h_i be the homology class of L_i , a component of L . Modulo $2\pi h_i$, assign continuously z_1 arguments to the points of L_i for all i . Let s be arclength on $\bigcup_i S_i^1$. Then

$$(6) \quad sw(L) = \frac{1}{2\pi} \int_{\bigcup_i S_i^1} \sum_{s'} \varepsilon(s') \frac{d(\arg(f_2(s') - f_2(s)))}{ds} ds$$

where the sum is over all s' such that $f_1(s') = f_1(s)$ and s and s' are in different components or $\arg f_1(s') \not\equiv \arg f_1(s)$ mod $2\pi h_i$. $\varepsilon(s')$ is the sign of the derivative of the z_1 argument at s' .

$sw(L)$ is necessarily an integer, for look at the pairs (s, s') . They form a 1-complex and every 0-cell, occurring when s or s' is a vertex, is on the boundary of an even number of 1-cells, and in fact is on the boundary of two 1-cells unless both s and s' are vertices and the derivatives of both $\arg(f_1(s))$ and $\arg(f_1(s'))$ change sign at s and s' . In this last case, (s, s') is on the boundary of four 1-cells. Thus as we follow $\arg(f_2(s') - f_2(s))$ around the 1-complex, we eventually return to our starting point, without retracing paths. (6) shows that $sw(L)$ depends continuously on f_2 . Hence $sw(L)$ is invariant under homotopies which keep the z_1 -coordinates fixed and which have no self-intersections at s and s' such that s and s' are in different components or $\arg f_1(s) \not\equiv \arg f_1(s')$, mod $2\pi h_i$.

$T = S^1 \times E^1 \times E^1$. Projecting onto $S^1 \times E^1$, we get a link diagram for L . By arbitrarily small deformations of L we may get a link whose projection is in general position (see [2, p. 7]). To each double point, or crossing, c we assign $\eta(c) = 1$ if the overpass crosses the underpass from right to left and $\eta(c) = -1$ if the overpass crosses from left to right.

LEMMA 1. Let L be a polygonal link, $L = (f_1, f_2) : \bigcup_i S_i^1 \rightarrow T$ such that f_1 is not constant on any segment of L and such that the link diagram of L is in general position. Then

$$(7) \quad sw(L) = \sum \eta(c)$$

where the sum is over those crossings whose under- and overpass are in different components of L or have different arguments, modulo $2\pi h_i$ where h_i is the homology class of L_i , the component of L containing the under- and overpass.

Proof. Without changing f_1 or the link diagram we deform L so that it lies in the plane of $S^1 \times E^1$ except near the crossings. This does not change $\text{sw}(L)$.

$$\frac{d(\arg f_2(s') - \arg f_2(s))}{ds} = 0$$

except near the crossings. One checks the cases, i.e. $\eta = \pm 1$ and the under- and overpass have increasing or decreasing z_1 arguments and one sees that (7) is correct.

In T , measure distance in S^1 by arclength in S^1 .

LEMMA 2. *Let L and \bar{L} be polygonal links*

$$L = (f_1, f_2): \bigcup_i S_i^1 \rightarrow T, \quad \bar{L} = (\bar{f}_1, \bar{f}_2): \bigcup_i S_i^1 \rightarrow T$$

such that f_1 and \bar{f}_1 are not constant on any segment and such that the link diagrams of L and \bar{L} are in general position. Let $\delta > 0$ be such that

$$|\arg f_1(s') - \arg f_1(s)| < \frac{2\pi}{9} \quad \text{if } |s - s'| < \delta.$$

Let

$$m = \inf [\max |f_1(s') - f_1(s)|, |f_2(s') - f_2(s)|]$$

with \inf taken over $\{s, s' \mid |s - s'| \geq \delta \text{ or } s \text{ and } s' \text{ are in different components of } L\}$.

Let \bar{L} satisfy the following conditions:

(i) *The preimages $\{v_j\}$ of the vertices of \bar{L} coincide with the preimages of the vertices of L .*

(ii) *Let Δ be a triangle with vertices*

$$\{L(v_j), \bar{L}(v_j), L(v_{j+1})\} \quad \text{or} \quad \{L(v_{j+1}), \bar{L}(v_j), \bar{L}(v_{j+1})\}$$

and let Δ' be another such triangle. Then Δ and its interior intersect the sides of Δ' only in isolated points and sides in common.

(iii) *In their link diagrams, the vertices of L and \bar{L} are all distinct.*

(iv) $|f_1(s) - \bar{f}_1(s)| < \min(2\pi/9, m/6)$ and $|f_2(s) - \bar{f}_2(s)| < m/6$ for all s .

Then $\text{sw}(L) = \text{sw}(\bar{L})$.

Proof. $m > 0$ by compactness.

We shall use the triangles Δ to construct a suitable homotopy between L and \bar{L} and then look at the changes in the link diagram. We may add vertices to L and \bar{L} so that $|v_j - v_{j+1}| < \delta'$, with δ' such that $\delta' \leq \delta$ and $|f_1(s') - f_1(s)| < m/6$ and $|f_2(s') - f_2(s)| < m/6$ if $|s' - s| < \delta'$, and not change the hypotheses of the lemma.

Replace the segment with end points $L(v_j)$ and $L(v_{j+1})$ by two segments, $(L(v_j), \bar{L}(v_j))$ and $(\bar{L}(v_j), L(v_{j+1}))$. Then replace $(\bar{L}(v_j), L(v_{j+1}))$ by $(L(v_{j+1}), \bar{L}(v_{j+1}))$ and $(\bar{L}(v_j), \bar{L}(v_{j+1}))$. This moved $(L(v_j), L(v_{j+1}))$ to $(\bar{L}(v_j), \bar{L}(v_{j+1}))$. Move $(L(v_{j+1}), L(v_{j+2}))$ to $(\bar{L}(v_{j+1}), \bar{L}(v_{j+2}))$ in a similar fashion (replacing also $(L(v_{j+1}), \bar{L}(v_{j+1}))$ by $(L(v_{j+2}), \bar{L}(v_{j+2}))$). Eventually we get a homotopy H (requiring a small change of the parametrization of \bar{L}), $H: \bigcup S_i^1 \times I \rightarrow T$.

Use H to assign a continuously varying argument, modulo $2\pi h_i$, on each component in the image of H . The replacement of segments as defined in the previous paragraph is just like [6, p. 8] except that there may be some intersections. Since (iii) prevents pathologies, an intersection corresponds to changing an underpass to an overpass. We wish first to show that at any such crossing the z_1 arguments of the under- and overpasses, which must be in the same component, are equal. Hence $\text{sw}(L)$, by Lemma 1, will be unchanged.

Suppose $f_1(s, t) = f_1(s', t)$ and $f_2(s, t) = f_2(s', t)$.

$$|\arg f_1(s, t) - \arg f_1(s)| < \frac{2\pi}{3} \quad \text{and} \quad |\arg f_1(s', t) - \arg f_1(s')| < \frac{2\pi}{3}$$

from considering the possible variation from $\arg f_1(v_j)$ or $\arg f_1(v_{j+1})$ in Δ . Similarly,

$$\begin{aligned} |f_1(s, t) - f_1(s)| &< m/2, & |f_1(s', t) - f_1(s')| &< m/2, \\ |f_2(s, t) - f_2(s)| &< m/2, & |f_2(s', t) - f_2(s')| &< m/2. \end{aligned}$$

$|s - s'| \geq \delta$ or s and s' in different components is impossible for then either

$$|f_1(s') - f_1(s)| \geq m \quad \text{or} \quad |f_2(s') - f_2(s)| \geq m.$$

If $|s - s'| < \delta$, then $|\arg f_1(s') - \arg f_1(s)| < 2\pi/9$ and indeed for all s' such that $|s - s'| < \delta$. From

$$|\arg f_1(s, t) - \arg f_1(s)| < 2\pi/3 \quad \text{and} \quad |\arg f_1(s', t) - \arg f_1(s')| < 2\pi/3,$$

which hold for all t , we see that $|\arg f_1(s', t) - \arg f_1(s, t)| < 2\pi$ and hence $\arg f_1(s', t) = \arg f_1(s)$ as desired.

Thus the intersections in H have no effect on $\text{sw}(L_t)$. Otherwise H is an isotopy and we need only observe the effect of the operations Ω_1 , Ω_2 and Ω_3 [6, pp. 7–8] on $\text{sw}(L_t)$, calculated from Lemma 1. But Ω_1 , Ω_2 and Ω_3 do indeed leave $\text{sw}(L)$ invariant.

If L is any continuous link and we approximate L by polygonal links L' and L'' such that f_1 is not constant on any segment and the link diagrams of L' and L'' are in general position, Lemma 2 indicates how to construct \bar{L} such that $\text{sw}(L') = \text{sw}(\bar{L}) = \text{sw}(L'')$, if L' and L'' are sufficiently close to L .

DEFINITION 4. Let $L: \bigcup_i S_i^1 \rightarrow T$ be a link. $\text{sw}(L) = \text{sw}(\bar{L})$ where \bar{L} is any polygonal link such that \bar{f}_1 is not constant on any segment of \bar{L} , the link diagram of \bar{L} is in general position, and \bar{L} is sufficiently close to L .

THEOREM 5. Let $H = (f_1, f_2): (\bigcup_i S_i^1) \times I \rightarrow T$ be an isotopy between the links $L_0 = (f_1, f_2, 0)$ and $L_1 = (f_1, f_2, 1)$. Then $\text{sw}(L_0) = \text{sw}(L_1)$.

Proof. Use H to assign continuously z_1 arguments to the image of H . Since $(\bigcup_i S_i^1) \times I$ is compact, $m(t) = m(f_1, f_2, t)$ of Lemma 2 is bounded away from 0. Thus we may use Lemma 2 to find polygonal links \bar{L}_t which approximate (f_1, f_2, t) and which have the same $\text{sw}(\bar{L}_t)$ for all t . Thus $\text{sw}(L_0) = \text{sw}(L_1)$.

Besides $\text{sw}(L)$, we may define isotopy invariants $\text{sw}_i(L)$ as follows. For simplicity, we shall assume that L is connected, i is an integer.

DEFINITION 5. Let $K: S^1 \rightarrow T$ be a knot. Approximate K by polygonal knots \bar{K}_ϵ such that \bar{f}_1 is not constant on any segment of \bar{K}_ϵ . Let h be the homology class of K . For $i \not\equiv 0 \pmod{h}$, let

$$\text{sw}_i(\bar{K}_\epsilon) = \frac{1}{2\pi} \int_{S^1} \sum_{s'} \epsilon(s') \frac{d(\arg(f_2(s') - f_2(s)))}{ds} ds,$$

where the sum is over all s' such that $\arg f_1(s') - \arg f_1(s) \equiv 2\pi i, \pmod{2\pi h}$.

$$\text{sw}_i(K) = \lim_{\epsilon \rightarrow 0} \text{sw}_i(\bar{K}_\epsilon).$$

THEOREM 6. Let $K: S^1 \rightarrow T$ be a knot. $\text{sw}_i(K)$ is independent of the choice of the approximating knots and is an isotopy invariant. If h is the homology class of K ,

$$\text{sw}(K) = \sum_{i=1}^{h-1} \text{sw}_i(K).$$

Theorem 6 is proved, using Lemma 2, in the same manner as Theorem 5 was proved.

If an algebroid singularity has a Puiseaux expansion [5, p. 163–164]

$$(8) \quad y = a_1 x^{n_1/m_1} + a_2 x^{n_2/m_1 m_2} + a_3 x^{n_3/m_1 m_2 m_3} + \dots$$

with $m_1 \leq n_1$, $n_1 m_2 < n_2$, $n_2 m_3 < n_3$, ... and $(m_1, n_1) = 1$, $(m_2, n_2) = 1$, $(m_3, n_3) = 1$, ... then the associated knot is a tubular knot A_k given by the symbol

$$\{(m_1, n_1), (m_2, n_2), \dots, (m_k, n_k)\}.$$

Without changing the singularity or the knot, we may assume that $m_i \neq 1$, $1 \leq i \leq k$. $m = m_1 \cdots m_k$ is the homology class of A_k . A_k is constructed inductively as follows. Let $A_0 = \{(1, 1)\}$ be the circle $z_2 = 0$, oriented positively. A_k winds m_k times about A_{k-1} in the direction of the parallels and n_k times about A_{k-1} in the direction of the meridians, where A_k must lie on a small tubular surface centered at A_{k-1} .

THEOREM 7. Let K be the knot in ∂B_ϵ associated to a singularity at the origin with one branch given by (8). Then the m_j and the n_j are determined by the homology class of K and the $\text{sw}_i(K)$.

Proof. We shall show by induction that

$$(9) \quad \begin{aligned} \text{sw}_i &= m_k \cdots m_2 n_1 && \text{if } i \not\equiv 0 \pmod{m_1} \\ \text{sw}_i &= m_k \cdots m_3 n_2 && \text{if } i \equiv 0 \pmod{m_1}, i \not\equiv 0 \pmod{m_1 m_2} \\ &\vdots && \\ \text{sw}_i &= m_k \cdots m_{j+1} n_j && \text{if } i \equiv 0 \pmod{m_1 \cdots m_{j-1}}, i \not\equiv 0 \pmod{m_1 \cdots m_j} \\ &\vdots && \\ \text{sw}_i &= n_k && \text{if } i \equiv 0 \pmod{m_1 \cdots m_{k-1}}. \end{aligned}$$

Since $n_{i+1} > n_i m_{i+1}$, the sw_i are listed in (9) with strictly increasing values, so that the m_j are all determined, using $m = m_1 \cdots m_k$ to determine m_k from the previous m_j and m , the homology class of K . Then the n_j are also all determined.

To prove (9), when $k=1$, we have a torus knot of type (m_1, n_1) and (9) is true. We now proceed by induction.

In general A_k winds m_k times about A_{k-1} in the direction of the parallels. Thus if $i \equiv 0 \pmod{m_1 \cdots m_{j-1}}$ and $i \not\equiv 0 \pmod{m_1 \cdots m_j}$ and we look at s' and s with $\arg s' - \arg s \equiv 2\pi i, \pmod{2\pi m}$, after s winds about S^1 $m_1 \cdots m_{k-1}$ times, we see that $f_2(s') - f_2(s)$ changes argument by $m_{k-1} \cdots m_{j+1} n_j 2\pi$. But instead of completing a full circuit around A_k , we have only gone around A_{k-1} and A_k winds m_k times around A_{k-1} in this direction. Thus $\text{sw}_i = m_k m_{k-1} \cdots m_{j+1} n_j$.

Finally if $i \equiv 0 \pmod{m_1 \cdots m_{k-1}}$, we have that A_k winds n_k times around A_{k-1} in the direction of the meridians, so that $\text{sw}_i(K)$ is the same as for the knot $\{(m_k, n_k)\}$ or just n_k , as desired.

REFERENCES

1. W. Burau, *Kennzeichnung der Schlauchknoten*, Abh. Math. Sem. Univ. Hamburg **9** (1932), 125–133.
2. R. H. Crowell and R. H. Fox, *Introduction to knot theory*, Ginn, New York, 1963.
3. R. C. Gunning and H. Rossi, *Analytic functions of several complex variables*, Prentice-Hall, Englewood Cliffs, N. J., 1965.
4. E. Martinelli, *Sulle intersezioni delle curve analitiche complesse*, Rend. Mat. e Appl. (5) **14** (1955), 422–430.
5. J. E. Reeve, *A summary of results in the topological classification of plane algebroid singularities*, Univ. e Politec. Torino Rend. Sem. Mat. **14** (1955), 159–187.
6. K. Reidemeister, *Knotentheorie*, Springer, Berlin, 1932.
7. O. Zariski, *On the topology of algebroid singularities*, Amer. J. Math. **54** (1932), 453–465.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY,
CAMBRIDGE, MASSACHUSETTS