

# UNIFORM APPROXIMATION ON A REAL-ANALYTIC MANIFOLD

BY

MICHAEL FREEMAN<sup>(1)</sup>

**1. Introduction.** Let  $M$  be a compact subset of a real-analytic manifold of dimension  $n$  and  $F$  a set of real-analytic complex-valued functions on  $M$  which separates  $M$ , meaning that for each pair  $p, q$  of distinct points in  $M$  there is a function  $f$  in  $F$  with  $f(p) \neq f(q)$ . We wish to study the Banach algebra  $A$  obtained as the closure in the norm of uniform convergence on  $M$  of the algebra of all polynomials in the functions of  $F$  (including constants). Thus  $A$  is the smallest closed subalgebra of  $C(M)$  with identity which contains  $F$ , where  $C(M)$  is all continuous complex-valued functions on  $M$ . Closed, separating subalgebras of  $C(M)$  with identity are frequently called function algebras, and the term is used elsewhere in much more general circumstances, where arbitrary compact Hausdorff spaces are admitted for  $M$ .

As we note below, the study of this type of function algebra includes the classical problem of uniform polynomial approximation on certain polynomially convex subsets of complex Euclidean space  $C^n$ .

Our study of these algebras continues a program initiated by Wermer [12], [13], and treated by the author [2], Wells [11], Nirenberg and Wells [9], [10], and very recently by Hörmander and Wermer [6]. In all of this work it has been shown that the set

$$E = \{p \in M : df_1 \wedge \cdots \wedge df_n(p) = 0 \text{ for all } n\text{-tuples } \{f_1, \dots, f_n\} \text{ of functions in } F\}$$

plays a major role in determining the structure of  $A$ . Our principal object in this paper is to prove a result announced earlier [3].

**THEOREM 1.** *If  $M_A = M$  then  $A$  contains the ideal of all continuous complex-valued functions which vanish on  $E$ .*

Here  $M_A$  is the spectrum or maximal ideal space of the Banach algebra  $A$ , and consists of all algebra homomorphisms of  $A$  onto  $C$ . Each point  $p$  of  $M$  provides such a homomorphism, defined by sending a function  $f$  in  $A$  into  $f(p)$ . The hypothesis  $M_A = M$  means that all homomorphisms arise in this manner. It is a necessary condition for the conclusion when  $E$  is empty and in certain other cases [2]. We refer to [2] for further properties of  $E$ .

---

Received by the editors May 23, 1968 and, in revised form, February 17, 1969.

<sup>(1)</sup> This work was supported in part by Army Grant DA-ARO-D-31-124-G866 while the author was at Brandeis University.

The theorem says nothing about the behavior of  $A$  on  $E$ , a problem of great interest since the theorem shows it equivalent to the problem of describing  $A$  (it is easy to see that a continuous function  $f$  on  $M$  is in  $A$  if and only if its restriction to  $E$  coincides with that of some function in  $A$ ). In the case where  $M$  is contained in a real-analytic submanifold of  $C^n$ , and with the usual coordinate functions comprising  $F$ ,  $A$  is the algebra of all continuous functions on  $M$  which can be approximated uniformly by polynomials. Here the condition  $M_A = M$  is equivalent to the assertion that  $M$  is polynomially convex [5], [14]. In this case Hörmander and Wermer [6], and earlier Wermer [13] in a special case, have shown that  $A$  is the set of all continuous functions on  $M$  which admit uniform approximation on  $E$  by functions holomorphic in a neighborhood of  $E$ . Their result of course contains Theorem 1 in this case. Wermer [13] also gives a differential description of  $A$  in a special case. A description of this type is conjectured in [3] for the more general situation treated here, but we have not proved it.

The author proved Theorem 1 when  $n=2$  in [2], by extending techniques first used by Wermer [13]. The same basic ideas are used here, but a number of modifications and extensions have been necessary to adapt the earlier proof to manifolds of dimension greater than two.

A simple example which verifies Theorem 1 is obtained when

$$M = \{(x, y, t) \in \mathbf{R}^3 : x^2 + y^2 + t^2 \leq 1\}$$

is the closed unit ball in real Euclidean space  $\mathbf{R}^3$  and  $F = \{f, g, h\}$  where

$$f(x, y, t) = x + iy = z, \quad g(x, y, t) = t\bar{z}, \quad \text{and} \quad h(x, y, t) = t.$$

Then the function  $(f, g, h): M \rightarrow C^3$  with the indicated coordinates maps  $M$  homeomorphically onto a compact polynomially convex [4], [5] subset of  $C^3$ , so [5]  $M_A = M$ . Clearly,  $E = \{(x, y, t) : t = 0\}$ . Moreover,  $A$  contains  $fh$ ,  $g$ , and  $h$ , which together separate  $M - E$ , which have no common zero there, and which generate an algebra closed under complex conjugation. An application of the Stone-Weierstrass theorem now shows that  $A$  contains every continuous function which vanishes on  $E$ .

**2. Proof of Theorem 1.** The argument is similar and in some places identical to that used before [2]. We show again that every bounded regular complex Borel measure which annihilates  $A$  also annihilates every continuous function which vanishes on  $E$ . This follows from Theorem 2 below in exactly the same way as it did in [2]. Theorem 2 reduces the study of such a measure  $\mu$  to the study of the family of bounded, regular, compactly supported Borel measures  $f_*\mu$  induced on the plane from  $\mu$  by certain functions in  $A$ . These measures are defined for each Borel set  $E$  by  $f_*\mu(E) = \mu(f^{-1}(E))$  and their relevant elementary properties are listed in [2].

We write  $\mu \perp f$  if  $\int f d\mu = 0$  and write  $\mu \perp A$  if this holds for all functions in  $A$ . If  $f_1, \dots, f_k$  are functions on  $M$  we denote by

$$(f_1, \dots, f_k): M \rightarrow C^k$$

the map with these functions as coordinates.

The following well-known facts are collected as lemmas for future use.

LEMMA 1. *If  $\nu$  is a bounded regular complex Borel measure with compact support in  $C$ , then*

(1)  $\int d|\nu|(\lambda)/|\lambda-z|$  is finite for almost all  $z$  in  $C$  (in the sense of Lebesgue measure), and

(2) if  $K$  is compact in  $C$  and  $\int d\nu(\lambda)/(\lambda-z)=0$  for almost all  $z$  in  $C-K$  (in the same sense), then support  $\nu \subset K$ .

In particular, if (1) holds for almost all  $z$  in  $C$ , then it follows from (2) that  $\nu=0$ . For a proof, the reader is referred to [14].

LEMMA 2. *If  $M$  is a compact subset of a complex manifold and  $G$  is a holomorphic map of a neighborhood of  $M$  into  $C^n$  which is injective and nonsingular on  $M$ , then  $G$  is injective and nonsingular on some neighborhood of  $M$ .*

LEMMA 3. *Let  $X$  be a compact Hausdorff space and  $F$  a subset of  $C(X)$  satisfying*

(3)  *$F$  separates  $X$ , and*

(4) *for each  $x$  in  $X$  there exists a finite subset of  $F$  which separates some neighborhood of  $x$ .*

*Then there exists a finite subset of  $F$  which separates  $X$ .*

This type of result has been used by Narasimhan [8], and was brought to my attention by H. Rossi.

**Proof.** Property (4) and compactness yield open sets  $U_1, \dots, U_k$  and a finite subset  $\{g_1, \dots, g_l\}$  of  $F$  which separates each  $U_i$ . Property (3) and the compactness of  $X \times X$  provide open sets  $V_1, \dots, V_p, W_1, \dots, W_p$  and functions  $g_{l+1}, \dots, g_{l+p}$  in  $F$  such that

$$X \times X - \bigcup_{i=1}^k U_i \times U_i \subset \bigcup_{i=1}^p V_i \times W_i,$$

and

$$g_{l+j}(V_j) \cap g_{l+j}(W_j) = \emptyset, \quad j = 1, \dots, p.$$

It follows easily that  $\{g_1, \dots, g_{l+p}\}$  separates  $X$ .

LEMMA 4. *Let  $M$  be a compact Hausdorff space and  $F$  a separating subset of  $C(M)$ . If  $U$  is open in  $M$  and  $\{f_1, \dots, f_k\}$  is a subset of  $F$  which separates  $U$ , then for each compact subset  $K$  of  $U$  there exist functions  $f_{k+1}, \dots, f_m$  in  $F$  such that  $(f_1, \dots, f_m)(K)$  is disjoint from  $(f_1, \dots, f_m)(M-K)$ .*

**Proof.** Since  $F$  separates  $M$ , a standard compactness argument shows that there exist functions  $f_{k+1}, \dots, f_m$  in  $F$  such that

$$(f_{k+1}, \dots, f_m)(K) \text{ is disjoint from } (f_{k+1}, \dots, f_m)(M-U).$$

It is straightforward to verify that  $\{f_1, \dots, f_m\}$  has the stated property.

THEOREM 2. If  $M_A = M$ ,  $\mu \perp A$ , and  $f$  is a polynomial in the functions of  $F$ , then

$$\int_C \frac{d(f_*\mu)(\lambda)}{\lambda - a} = 0$$

for almost all points  $a$  in  $C - f(E)$ . Thus by Lemma 1, support  $f_*\mu \subset f(E)$ .

**Proof.** As in [2], it will suffice to show that  $\int d\mu/(f - a) = 0$  for all points  $a$  in  $C - f(E)$  for which the integral is absolutely convergent. For each such  $a$ , we construct a sequence  $\{f_n\}$  of functions in  $A$  such that

$$f_n \rightarrow 1/(f - a) \text{ a.e. } |\mu|, \text{ and } |f_n| \leq 2/|f - a| \text{ a.e. } |\mu|.$$

These functions are obtained as before from the solution of a certain Cousin Problem I on a domain in  $C^n$ . This problem has the same basic structure as it did in [2], but is somewhat more complicated because of the higher dimension. To set it up, we appeal to a result of Whitney and Bruhat [15], which states that there exists a complex manifold  $\tilde{M}$  in which the ambient manifold of  $M$  can be imbedded as a real-analytic submanifold in such a way that every real-analytic function on  $M$  can be extended to a holomorphic function on an  $\tilde{M}$ -neighborhood of  $M$ . While some of the constructions below could be executed on  $\tilde{M}$ , it is more convenient to transfer immediately to complex Euclidean space. The basic idea of the proof is clearest when  $F$  is finite, so we present that case first. Technical modifications required to handle the general situation are given afterwards.

*Case of finite  $F$ .* Here the functions in  $F$  comprise the coordinates of a map

$$H = (f_1, \dots, f_p): M \rightarrow C^p.$$

By our assumption,  $f = q \circ H$  for some polynomial  $q$  in  $p$  variables. Given a point  $a$  in  $C - f(E)$  we claim that:

*There exists an open set  $W \supset H(M) \cap q^{-1}(a)$  and a function  $k$  holomorphic on  $W$  such that  $k|_{H(M) \cap W} = (q - a)^*|_{H(M) \cap W}$ . Here the  $*$  denotes complex conjugation.*

There is an  $\tilde{M}$ -open set  $\tilde{V}$  which contains  $f^{-1}(a)$  and to which  $H$  has a holomorphic extension  $\tilde{H} = (\tilde{f}_1, \dots, \tilde{f}_p)$ . Since  $a$  is not in  $f(E)$  the  $p$ -form  $df_1 \wedge \dots \wedge df_p$  has no zeros on  $f^{-1}(a)$  so the same is true of  $d\tilde{f}_1 \wedge \dots \wedge d\tilde{f}_p$ . Since  $\tilde{H}$  is injective on  $M$  we can use Lemma 2 to choose  $\tilde{V}$  small enough so that  $\tilde{H}$  imbeds  $\tilde{V}$  as a complex submanifold  $V$  of  $C^p$ .

We may assume that  $(f - a)^*$  has a holomorphic extension to  $\tilde{V}$ . Now  $V$  is a closed submanifold of some open set  $U$  in  $C^p$  (for instance, a union of ambient coordinate neighborhoods whose slices define  $V$  locally), and since  $H(M \cap \tilde{V})$  is disjoint from  $H(M - \tilde{V})$  we can remove the compact set  $H(M - \tilde{V})$  from  $U$  to obtain  $V$  as a closed submanifold of the open set  $U$  and satisfying

$$(5) \quad U \cap H(M) = V \cap H(M) = H(M \cap \tilde{V}).$$

The set  $H(M) \cap q^{-1}(a)$  is polynomially convex [5], as a consequence [14] of our assumption that  $M_A = M$ . Therefore there exists [5] a domain of holomorphy  $W$  such that  $H(M) \cap q^{-1}(a) \subset W \subset U$ . We have an extension of  $(f-a)^* = (q-a)^* \circ H$  to a holomorphic function on  $\tilde{V}$ . Because of (5), composition of this extension with  $(\tilde{H}|\tilde{V})^{-1}: V \rightarrow \tilde{V}$  yields a holomorphic extension to  $V$  of  $(q-a)^*|H(M) \cap V$ . The theorem of Grauert and Docquier [4, Theorem 8, pp. 257-258], which says that  $W \cap V$  is a holomorphic retraction of  $W$ , finally yields the desired extension  $k$  on  $W$ .

We wish now to proceed along lines similar to [2], and construct functions  $h$  and  $h_1$  in  $A$  such that  $h = (f-a)h_1$  and  $h(M) \subset \{w : |w-1| > 1\} \cup \{0\}$ . A "local" solution to this problem is given by  $h_1 = -k \circ H$  and  $h = (f-a)h_1$ . However  $k \circ H$  is not in  $A$ , and we wish to use  $k$  to obtain a holomorphic function  $\psi$  with appropriate divisibility properties in a neighborhood of  $H(M)$ . Since  $H(M)$  is polynomially convex, the Oka-Weil theorem [5, Theorem 2.7.7, p. 55] will imply that  $h = -\psi \circ H$  is in  $A$ . Let  $g = (q-a)k$ , a function holomorphic on  $W$ .

*Then there exists a function  $\psi$  holomorphic on a neighborhood of  $H(M)$  such that*

(6)  $\psi$  has no zeros on  $H(M) - q^{-1}(a)$ , and

(7)  $\psi$  is divisible by  $g$  in a neighborhood of  $H(M) \cap q^{-1}(a)$  and the holomorphic quotient  $\psi/g$  has the value 1 everywhere on  $H(M) \cap q^{-1}(a)$ .

This function  $\psi$  is exhibited as the solution to a Cousin Problem I [5] with data determined from  $g$  as follows. Since  $g|H(M) \cap W = |q-a|^2|H(M) \cap W$ , it follows that

$$H(M) \cap \{\text{Re } g = 0\} = H(M) \cap q^{-1}(a).$$

Therefore by shrinking  $W$  if necessary, it can be assumed that  $\{\text{Re } g = 0\}^-$  is disjoint from  $H(M) - W$  (here the superscript bar denotes closure).

Thus  $C^p - q^{-1}(a) - \{\text{Re } g = 0\}^-$  is an open set containing  $H(M) - W$ , so that  $W$  and this set constitute an open cover of  $H(M)$ . Since  $H(M)$  is polynomially convex, there exists an open domain of holomorphy  $S$  in  $C^p$  such that

$$H(M) \subset S \subset W \cup [C^p - q^{-1}(a) - \{\text{Re } g = 0\}^-].$$

Replacing  $W$  by its intersection with  $S$ , we have  $W \subset S$ . Since  $\{\text{Re } g = 0\}$  is closed in  $S$ , the set  $T = S - q^{-1}(a) - \{\text{Re } g = 0\}$  is open and  $S = W \cup T$ .

We have designed  $W, T$ , and  $g$  so that  $g$  has a holomorphic logarithm  $\log g$  on

$$W \cap T = W - q^{-1}(a) - \{\text{Re } g = 0\},$$

and so that  $(\log g)/(q-a)$  is holomorphic on  $W \cap T$ . Therefore the Cousin Problem I defined on  $S$  for the covering  $\{W, T\}$  by  $(\log g)/(q-a)$  has a solution [5]; that is, there exist functions  $g_1$  holomorphic on  $T$  and  $g_2$  holomorphic on  $W$  such that

$$g_1 - g_2 = (\log g)/(q-a) \text{ on } W \cap T.$$

Thus  $\log g + (q-a)g_2 = (q-a)g_1$  on  $W \cap T$  so the holomorphic functions

$$g \exp((q-a)g_2) \text{ on } W \text{ and } \exp((q-a)g_1) \text{ on } T$$

coincide on  $W \cap T$ . Hence they define a holomorphic function  $\psi$  on  $S$  with the desired properties.

This result is used to construct functions  $h$  and  $h_1$  in  $A$  such that

$$(8) \ h \text{ has no zeros on } H(M) - q^{-1}(a),$$

$$(9) \ h = (f - a)h_1, \text{ and}$$

$$(10) \ h(M) \subset \{w : |w - 1| > 1\} \cup \{0\}.$$

From (6) and (7) it follows that  $\psi_1 = \psi/(q - a)$  is holomorphic in a neighborhood of  $H(M)$ . By (7) there exists an  $H(M)$ -neighborhood  $P$  of  $H(M) \cap q^{-1}(a)$  on which  $\operatorname{Re}(\psi/g) > 0$ . It follows from this and the positivity of  $g$  on  $H(M) - q^{-1}(a)$  that  $\operatorname{Re} \psi > 0$  on  $P - q^{-1}(a)$ . The function  $\psi$  has no zeros on the compact set  $H(M) - P$ , so after multiplication of  $\psi$  by a suitable positive constant, it will satisfy  $|\psi| \geq 2$  on  $H(M) - P$ . We have already noted that  $h = -\psi \circ H$  and  $h_1 = -\psi_1 \circ H$  are in  $A$ , and they clearly have the properties (8), (9) and (10).

These functions are used exactly as in [2, p. 54] to construct the sequence  $\{f_n\}$ . There we defined the rational functions  $\phi_n$  by

$$\phi_n(w) = \frac{1}{w} \left( 1 - \frac{1}{(w-1)^{2n}} \right), \quad n = 1, 2, \dots$$

and showed easily that the sequence  $\{f_n\}$  of functions in  $A$  defined by  $f_n = (\phi_n \circ h)h_1$ ,  $n = 1, 2, \dots$  has the properties set forth at the beginning of the proof. Theorem 2 is thereby proved when  $F$  is finite.

**Proof of Theorem 2 for arbitrary  $F$ .** The proof has the same basic structure as before, but it must surmount two additional difficulties. Since we cannot expect to find any finite subset of  $F$  to provide the coordinates of a homeomorphism of  $M$  into a complex Euclidean space, the separation arguments made at the beginning are somewhat more involved. A more serious problem is that the image of  $M$  under a map  $G = (f_1, \dots, f_m)$  with the  $f_j$ 's in  $F$  will not necessarily be polynomially convex. Because of this, domains of holomorphy corresponding to  $W$  and  $S$  will be harder to find. To construct them, we will use a standard technique due to Arens and Calderón [1], [5].

We again choose a point  $a$  in  $C - f(E)$  and claim that *there exist functions  $\{f_1, \dots, f_m\}$  in  $F$ , an open set  $U$  in  $C^m$ , and a closed complex submanifold  $V$  of  $U$  such that if  $G = (f_1, \dots, f_m)$  then*

$$(11) \ f = q \circ G \text{ for some polynomial } q \text{ in } m \text{ variables,}$$

$$(12) \ V \supset G(M) \cap q^{-1}(a),$$

$$(13) \ U \cap G(M) = V \cap G(M) = G(M \cap \tilde{V}), \text{ and}$$

$$(14) \ (q - a)^* \text{ extends from } G(M) \cap V \text{ to a function } k \text{ holomorphic on } V.$$

To begin the construction of  $G$ ,  $U$ , and  $V$  we note that for each point of  $f^{-1}(a)$  there can be found functions  $f_1, \dots, f_n$  in  $F$  and a neighborhood of the point on which  $df_1 \wedge \dots \wedge df_n$  has no zeros. These functions extend to holomorphic functions  $\tilde{f}_1, \dots, \tilde{f}_n$  on an  $\tilde{M}$ -neighborhood of the point on which the extended form  $d\tilde{f}_1 \wedge \dots \wedge d\tilde{f}_n$  has no zeros. Since  $f^{-1}(a)$  is compact it can be covered by finitely

many such neighborhoods with the result that there exist functions  $f_1, \dots, f_i$  in  $F$  whose holomorphic extensions provide a map  $(\tilde{f}_1, \dots, \tilde{f}_i)$  with maximum rank  $n$  on some  $\tilde{M}$ -neighborhood of  $f^{-1}(a)$ . Since  $F$  separates  $M$ , Lemma 3 applied to  $F$  and  $f^{-1}(a)$  yields the existence of functions  $f_{i+1}, \dots, f_m$  in  $F$  whose adjunction provides a map

$$G = (f_1, \dots, f_m): M \rightarrow \mathbb{C}^m$$

which is injective and of rank  $n$  in an  $M$ -neighborhood of  $f^{-1}(a)$ . By Lemma 2, the map  $(\tilde{f}_1, \dots, \tilde{f}_m)$  imbeds an open  $\tilde{M}$ -neighborhood  $\tilde{V}$  of  $f^{-1}(a)$  as a complex submanifold  $V$  of  $\mathbb{C}^m$ . This property is clearly unaffected by the adjunction of more coordinate functions, so we may assume that (11) is true. Statement (12) is then immediate.

By applying Lemma 4 to  $M$  and  $M \cap \tilde{V}$  and passing from  $\tilde{V}$  to a relatively compact subset of it which contains  $f^{-1}(a)$ , we can assume that  $G(M \cap \tilde{V})$  and  $G(M - \tilde{V})$  are disjoint. Just as before we can arrange that  $V$  is a closed submanifold of an open set  $U$  in  $\mathbb{C}^m$  which satisfies (13), and find an extension  $k$  verifying (14).

However,  $U$  need not contain a domain of holomorphy containing  $G(M) \cap q^{-1}(a)$ , since the latter set is not necessarily polynomially convex. Hence the theorem of Grauert and Docquier cannot yet be used to extend  $k$  to an open set in  $\mathbb{C}^m$ .

To effect this extension and thus prepare the way for the Cousin I construction above, we shall use a technique of Arens and Calderón [1], [5]. In fact, we assert that *there are functions  $f_{m+1}, \dots, f_p$  in  $F$  such that if  $H = (f_1, \dots, f_p)$  and  $B$  is the closed subalgebra of  $A$  with identity generated by  $\{f_1, \dots, f_p\}$ , then there exists a domain of holomorphy  $W$  in  $\mathbb{C}^p$  such that  $M_B \cap q^{-1}(a) \subset W \subset U \times \mathbb{C}^{p-m}$  and  $k$  extends to a holomorphic function on  $W$ .* Here we have made the usual identification of the maximal ideal space  $M_B$  of  $B$  with the polynomially convex hull of  $H(M)$  in  $\mathbb{C}^p$ , and the functions  $k$  and  $q$  are transferred in the obvious way to functions on  $V \times \mathbb{C}^{p-m}$  and  $\mathbb{C}^p$ , respectively.

These additional functions are obtained by means of the Lemma of Arens and Calderón [5], which says that there can be found  $f_{m+1}, \dots, f_p$  in  $F$  such that with  $H$  and  $B$  as defined above and  $\sigma_B(f_1, \dots, f_m)$  the joint spectrum [5] of the indicated functions relative to  $B$ , we have

$$(15) \quad \sigma_B(f_1, \dots, f_m) \subset U \cup [\mathbb{C}^m - q^{-1}(a)].$$

Their lemma is applicable because  $U \supset G(M) \cap q^{-1}(a)$ , so the right side of (15) is an open neighborhood of  $G(M)$ .

Now since  $\sigma_B(f_1, \dots, f_m)$  is the projection of  $M_B$  on  $\mathbb{C}^m$ , we have

$$M_B \subset [U \times \mathbb{C}^{p-m}] \cup [\mathbb{C}^p - q^{-1}(a)], \quad \text{so} \quad M_B \cap q^{-1}(a) \subset U \times \mathbb{C}^{p-m}.$$

Moreover,  $M_B \cap q^{-1}(a)$  is polynomially convex, since  $M_B$  has this property. Thus there exists a domain of holomorphy  $W$  in  $\mathbb{C}^p$  with

$$M_B \cap q^{-1}(a) \subset W \subset U \times \mathbb{C}^{p-m}.$$

Finally,  $(V \times C^{p-m}) \cap W$  is a closed submanifold of  $W$ , and the function  $k$  extends to  $W$  exactly as it did before.

We can now construct a function  $\psi$  holomorphic in a neighborhood of  $M_B$  with no zeros on  $M_B - q^{-1}(a)$ , divisible by  $g = (p-a)k$  in a neighborhood of  $M_B \cap q^{-1}(a)$  and such that the holomorphic quotient  $\psi/g$  has the value 1 everywhere on  $H(M) \cap q^{-1}(a)$ .

For we have again that

$$\{\operatorname{Re} g = 0\} \cap H(M) = q^{-1}(a) \cap H(M),$$

and we may therefore assume that  $\{\operatorname{Re} g = 0\}^-$  is disjoint from  $H(M) - W$ . However, it may not be the case that  $\{\operatorname{Re} g = 0\}^-$  is disjoint from  $M_B - W$ . If not, this separation may be achieved by another application of the Arens-Calderón Lemma, noting that

$$W \cup [C^p - q^{-1}(a) - \{\operatorname{Re} g = 0\}^-]$$

is an open neighborhood of  $H(M)$  and proceeding as above. In other words, we can assume that

$$M_B \subset W \cup [C^p - q^{-1}(a) - \{\operatorname{Re} g = 0\}^-],$$

which enables the construction of  $\psi$  as the solution to the same Cousin Problem I that we have already described. The proof is then completed exactly as it was when  $F$  is finite.

### 3. Some conditions for $A = C(M)$ .

**COROLLARY 1.** *If  $M_A = M$  and  $E$  is totally disconnected, then  $A = C(M)$ .*

This result appears in [2]. Since it depends solely on the conclusion of Theorem 1 and not on the dimension of the ambient manifold of  $M$ , the proof given there holds without modification.

In [2] we also deduced for the two-dimensional case that  $A = C(M)$  if  $M_A = M$  and  $E$  has Lebesgue measure zero. The example presented in §1 shows that this result fails in higher dimensions. In this example,  $E$  has three-dimensional Lebesgue measure zero but every function in  $A$  is a uniform limit on  $E$  of polynomials in  $f$ , with  $f(x, y, t) = x + iy$ . Thus each function in  $A$  is holomorphic on  $E$  (in the obvious sense), so that  $A \neq C(M)$ . However, a stronger measure-theoretic restriction on  $E$  will still yield the same result:

**COROLLARY 2.** *If  $M_A = M$  and  $E$  has two-dimensional Hausdorff measure zero, then  $A = C(M)$ .*

**Proof.** It is easily seen that the image by a continuously differentiable map of a set of two-dimensional Hausdorff measure zero also has this property. Because of the relation [7, p. 104] between Hausdorff two-dimensional measure and plane Lebesgue measure, we then have for any polynomial  $f$  in the functions of  $F$  that

$f(E)$  has measure zero. From Theorem 2 it follows that  $f_*\mu=0$ , which implies that  $\mu=0$  [2, p. 56].

We remark that it is clear how, by adjoining more coordinates, an example of the type presented in §1 can be constructed where  $M$  has any dimension greater than two but  $E$  has Hausdorff three-dimensional measure zero and  $A \neq C(M)$ .

#### REFERENCES

1. R. Arens and A. P. Calderón, *Analytic functions of several Banach algebra elements*, Ann. of Math. **62** (1955), 204–216.
2. M. Freeman, "Some conditions for uniform approximation on a manifold" in *Function algebras*, Scott, Foresman and Co., Chicago, Ill., 1966.
3. ———, *Function algebras on a manifold*, Proc. Rice Univ. Conf. on Complex Analysis, April 1967, Rice Univ. Studies **54** (1968), 15–19.
4. R. Gunning and H. Rossi, *Analytic functions of several complex variables*, Prentice-Hall, Englewood Cliffs, N. J., 1965.
5. L. Hörmander, *An introduction to complex analysis in several variables*, Van Nostrand, Princeton, N. J., 1966.
6. L. Hörmander and J. Wermer, *Uniform approximation on compact sets in  $C^n$* , Math. Scand. (to appear).
7. W. Hurewicz and H. Wallman, *Dimension theory*, Princeton Univ. Press, Princeton, N. J., 1941.
8. R. Narasimhan, *Imbedding of holomorphically complete complex spaces*, Amer. J. Math. **82** (1960), 917–934.
9. R. Nirenberg and R. O. Wells, Jr., *Holomorphic approximation on real submanifolds of a complex manifold*, Bull. Amer. Math. Soc. **73** (1967), 378–381. (This is an announcement of some results in reference [10].)
10. ———, *Approximation theorems on differentiable submanifolds of a complex manifold*, Trans. Amer. Math. Soc. **142** (1969), 15–35.
11. R. O. Wells, Jr., *Holomorphic approximation on real-analytic submanifolds of a complex manifold*, Proc. Amer. Math. Soc. **17** (1966), 1272–1275.
12. J. Wermer, *Approximation on a disk*, Math. Ann. **155** (1964), 331–333.
13. ———, *Polynomially convex disks*, Math. Ann. **158** (1965), 6–10.
14. ———, *Banach algebras and analytic functions*, Advances in Math. **1** (1961), 51–102.
15. H. Whitney and F. Bruhat, *Quelques propriétés fondamentales des ensembles analytiques-réels*, Comment. Math. Helv. **33** (1959), 132–160.

RICE UNIVERSITY,  
HOUSTON, TEXAS