

INVOLUTIONS ON HOMOTOPY SPHERES AND THEIR GLUING DIFFEOMORPHISMS

BY

CHAO-CHU LIANG

ABSTRACT. Let $hS(P^{2n+1})$ denote the set of equivalence classes of smooth fixed-point free involutions on $(2n + 1)$ -dimensional homotopy spheres. Browder and Livesay defined an invariant $\sigma(\Sigma^{2n+1}, T)$ for each $(\Sigma^{2n+1}, T) \in hS(P^{2n+1})$, where $\sigma \in \mathbb{Z}$ if n is odd, $\sigma \in \mathbb{Z}_2$ if n is even. They showed that for $n \geq 3$, $\sigma(\Sigma^{2n+1}, T) = 0$ if and only if (Σ^{2n+1}, T) admits a codim 1 invariant sphere. For any (Σ^{2n+1}, T) , there exists an A -equivariant diffeomorphism f of $S^n \times S^n$ such that $(\Sigma^{2n+1}, T) = (S^n \times D^{n+1}, A) \cup_f (D^{n+1} \times S^n, A)$, where A denotes the antipodal map. Let $\beta(f) = \sigma(\Sigma^{2n+1}, T)$. In the case n is odd, we can show that the Browder-Livesay invariant is additive: " $\beta(fg) = \beta(f) + \beta(g)$ ". But if n is even, then there exists f and g such that $\beta(gf) = \beta(g) + \beta(f) \neq \beta(fg)$. Let $D_0(S^n \times S^n, A)$ be the group of concordance classes of A -equivariant diffeomorphisms which are homotopic to the identity map of $S^n \times S^n$. We can prove that "For $n \equiv 0, 1, 2 \pmod{4}$, $hS(P^{2n+1})$ is in 1-1 correspondence with a subgroup of $D_0(S^n \times S^n, A)$ ". As an application of these theorems, we demonstrated that "Let Σ_0^{8k+3} denote the generator of bP_{8k+4} . Then the number of (Σ_0^{8k+3}, T) 's with $\sigma(\Sigma_0^{8k+3}, T) = 0$ is either 0 or equal to the number of (S^{8k+3}, T) 's with $\sigma(S^{8k+3}, T) = 0$, where S^{8k+3} denotes the standard sphere".

0. Introduction. In [7], [8], Browder and Livesay studied differentiable fixed-point free involutions on homotopy spheres. They defined the Browder-Livesay desuspension invariant σ for each free involution (Σ^m, T) : $\sigma(\Sigma^m, T) = 0$, for m even; $\sigma(\Sigma^m, T) \in \mathbb{Z}$, for $m = 4k + 3$; $\sigma(\Sigma^m, T) \in \mathbb{Z}_2$, for $m = 4k + 1$. For $m \geq 6$, they proved that $\sigma(\Sigma^m, T) = 0$ if and only if (Σ^m, T) admits a codim 1 invariant subsphere $(S^{m-1}, T|_{S^{m-1}})$ embedded in it. It was shown by several people that all these desuspension invariants can be realized, [2], [6], [21], [33], and [34] etc.

Livesay and Thomas, [20], showed that any (Σ^{2n+1}, T) can be obtained by gluing $(S^n \times D^{n+1}, A)$ and $(D^{n+1} \times S^n, A)$ together by an A -equivariant diffeomorphism f of their boundaries, where A is the antipodal map. We shall denote this (Σ^{2n+1}, T) by (Σ_f, T_f) . The purpose of this paper is to investigate the rela-

Received by the editors August 7, 1974.

AMS (MOS) subject classifications (1970). Primary 57E25, 57E30, 57D65, 55E05.

Key words and phrases. Free differentiable involutions, Browder-Livesay invariant, equivariant diffeomorphism, concordance group of diffeomorphisms, curious involutions.

Copyright © 1976, American Mathematical Society

tion between the free involutions on odd dimensional homotopy spheres and their gluing diffeomorphisms.

Let A denote the antipodal map on $S^n \times S^n$, defined by $A(x, y) = (-x, -y)$ for $(x, y) \in S^n \times S^n$. An A -equivariant map $f: S^n \times S^n \rightarrow S^n \times S^n$ is a map such that $fA = Af$, we shall call f an A -map. An A -map f induces a map $f': S^n \times S^n/A \rightarrow S^n \times S^n/A$, where $S^n \times S^n/A$ denotes the orbit space of $S^n \times S^n$ under the action of A . If f' is a diffeomorphism (or a homotopy equivalence), we will call f an A -diffeomorphism (or an A -homotopy equivalence). Considering the action of $A \times \text{identity}$ on $S^n \times S^n \times [0, 1]$, we have the notion of A -homotopy and A -concordance etc. (see §2 below).

Let $D(S^n \times S^n, A)$ denote the group of A -concordance classes of A -diffeomorphisms of $S^n \times S^n$. Define $J_{2n+1} = \{f \in D(S^n \times S^n, A) \mid f \text{ is homotopic to identity}\}$, (but f might not be A -homotopic to identity). J_{2n+1} is a subgroup of $D(S^n \times S^n, A)$. We will show that the gluing diffeomorphism can always be chosen from J_{2n+1} , (§4).

Two involutions (Σ, T) and (Σ', T') are called equivalent, $(\Sigma, T) = (\Sigma', T')$, if there exists an orientation-preserving diffeomorphism $f: \Sigma \rightarrow \Sigma'$ such that $f \circ T = T' \circ f$. Let $hS(P^{2n+1})$ denote the set of homotopy smoothings of P^{2n+1} , [34], which is also the set of equivalence classes of differentiable free involutions on $(2n+1)$ -homotopy spheres, [21]. In §5 below, we will prove that for $n \equiv 0, 1, 2 \pmod{4}$, $hS(P^{2n+1})$ is in 1-1 correspondence with a subgroup G_{2n+1} of J_{2n+1} . Thus, in these cases, $hS(P^{2n+1})$ forms a group by carrying over the composition law of diffeomorphisms in G_{2n+1} .

Also, we will show in Theorem 6.15 below that the Browder-Livesay index invariant is additive. For $m = 4k + 3$, f and $g \in J_m$, we have $\sigma(\Sigma_f, T_f) + \sigma(\Sigma_g, T_g) = \sigma(\Sigma_{fg}, T_{fg})$.

From Theorems 5.2 and 6.15, we can deduce the following theorem concerning the curious involutions in the sense of [13]. Let Σ_0^n denote the generator of bP_{n+1} , [17]. S^n denotes the standard sphere.

THEOREM 8.2. *For $n = 8k + 3$, the number of curious involutions (Σ_0^n, T) with $\sigma(\Sigma_0^n, T) = 0$ is either 0 or equal to the number of involutions (S^n, T) with $\sigma(S^n, T) = 0$.*

Everything considered here is assumed to be in the smooth category.

This paper is essentially the author's doctoral thesis written at Cornell University. I wish to thank my thesis advisor, Professor G. R. Livesay, for his constant help and encouragement. I also wish to thank Professors I. Berstein, P. J. Kahn, and H. C. Wang for many useful discussions.

1. Livesay-Thomas decomposition theorem. We have the following theorem from [20].

THEOREM 1.1. *For any free involution on a homotopy sphere (Σ^{2n+1}, T) , $n \geq 3$, there exists an A -diffeomorphism g of $S^n \times S^n$ such that $(\Sigma^{2n+1}, T) = (S^n \times D^{n+1}, A) \cup_g (D^{n+1} \times S^n, A)$, denoted by (Σ_g, T_g) .*

Note. $M \cup_g N$ denotes a manifold obtained by gluing two manifolds M and N together by a diffeomorphism $g: M_0 \rightarrow N_0$, where $M_0 \subseteq \partial M$ and $N_0 \subseteq \partial N$.

We shall prove the following proposition in §4 below.

PROPOSITION 1.2. *For any (Σ^{2n+1}, T) , where $n \neq 3, 7$, there exists an A -diffeomorphism g of $S^n \times S^n$ such that g is homotopic to the identity and $(\Sigma^{2n+1}, T) = (\Sigma_g, T_g)$.*

Notation. Let “ \sim ” denote homotopic, and “ \sim^A ” denote A -homotopic.

LEMMA 1.3. *If g is an A -diffeomorphism of $S^n \times S^n$ such that $g \sim \text{Id}$ (the identity), then there exists a pair of A -homotopy equivalences f_1, f_2 of $S^n \times S^n$ such that $f_1 \sim f_2 \sim \text{Id}$, where f_1 (f_2) extends to an A -homotopy equivalence h_1 (h_2) of $S^n \times D^{n+1}$ ($D^{n+1} \times S^n$); and $g \sim^A f_2 f_1$.*

PROOF. Let p_j denote the projection of $S^n \times S^n$ onto the j th factor S_j^n , $j = 1$ or 2 . p_j is A -equivariant: $p_j(Ax, Ay) = A(p_j(x, y))$. Let $g_j = p_j \circ g$. We define f_1 by $f_1(x, y) = (x, g_2(x, y))$. g^{-1} is also an A -diffeomorphism. Let $k_j = p_j \circ g^{-1}$. We define f_3 by $f_3(x, y) = (k_1(x, y), y)$.

$$\begin{aligned} f_3 \circ g(x, y) &= f_3(g_1(x, y), g_2(x, y)) = (k_1(g_1(x, y), g_2(x, y)), g_2(x, y)) \\ &= (x, g_2(x, y)) = f_1(x, y). \end{aligned}$$

f_1 and f_3 are obviously A -maps.

Since $g \sim \text{Id}$, $(x_0, y) \rightarrow (x_0, g_2(x_0, y))$ is a degree 1 map of $x_0 \times S^n$ to itself for each $x_0 \in S^n$. Hence $f_1|_{x_0 \times S^n} \rightarrow x_0 \times S^n$ is a homotopy equivalence for each $x_0 \in S^n$. We have a locally trivial fibre bundle $S_2^n \rightarrow S^n \times S^n/A \rightarrow S_1^n/A$, with base space S_1^n/A and fibre S_2^n . The map $f'_1: S^n \times S^n/A \rightarrow S^n \times S^n/A$, induced by f_1 , is fibre preserving. The restriction of f'_1 to each fibre is a homotopy equivalence. Hence f'_1 is a fibre homotopy equivalence by a theorem of Dold [10], and so f_1 is an A -homotopy equivalence. Similarly, we can show that f_3 induces a fibre homotopy equivalence f'_3 of the bundle $S_1^n \rightarrow S^n \times S^n/A \rightarrow S_2^n/A$. Let f'_2 be the fibre homotopy inverse of f'_3 , and write f_2 for the double cover of f'_2 such that f_2 is the A -homotopy inverse for f_3 . Now since $f_3 \circ g = f_1$, it follows that $g \sim^A f_2 f_1$.

Since the A -map $g_2: S^n \times S^n \rightarrow S_2^n$ extends to an A -map $\bar{g}_2: S^n \times D^{n+1} \rightarrow D^{n+1}$ by radial extension, we define an A -homotopy equivalence h_1 of $S^n \times D^{n+1}$ by $h_1(x, y) = (x, \bar{g}_2(x, y))$. An A -homotopy equivalence h_2 of $D^{n+1} \times S^n$ can be defined similarly. $f_1 \sim f_2 \sim \text{Id}$ follows from [19, 2.5]. Q.E.D.

LEMMA 1.4. Suppose $(\Sigma^{2n+1}, T) = (\Sigma_g, T_g) = S^n \times D^{n+1} \cup_{\text{Id}} S^n \times S^n \times I \cup_g D^{n+1} \times S^n$ for an A -diffeomorphism g as in (1.2). Then there exists an equivariant homotopy equivalence $F: (\Sigma^{2n+1}, T) \rightarrow (S^{2n+1}, A) = S^n \times D^{n+1} \cup_{\text{Id}} S^n \times S^n \times I \cup_{\text{Id}} D^{n+1} \times S^n$ such that each summand is mapped into the corresponding one by an A -homotopy equivalence.

PROOF. For such an A -diffeomorphism g , there exists f_1, f_2, h_1, h_2 as in (1.3). Write f_2^{-1}, h_2^{-1} for A -homotopy inverses of f_2, h_2 . Let H be an A -homotopy between $f_2^{-1} \circ g$ and f_1 with $H(x, 0) = f_1(x)$. We then define $F = h, H, h_2^{-1}$ on each summand as follows:

$$\begin{array}{ccccccc} (\Sigma^{2n+1}, T) & = & S^n \times D^{n+1} & \cup_{\text{Id}} & S^n \times S^n \times I & \cup_g & D^{n+1} \times S^n \\ \downarrow F & & \downarrow h_1 & & \downarrow H & & \downarrow h_2^{-1} \\ (S^{2n+1}, A) & = & S^n \times D^{n+1} & \cup_{\text{Id}} & S^n \times S^n \times I & \cup_{\text{Id}} & D^{n+1} \times S^n \quad \text{Q.E.D.} \end{array}$$

An invariant m -manifold for (Σ^k, T) is an embedded m -manifold $M^m \subseteq \Sigma^k$ which is invariant under T . An invariant M^m for (Σ^k, T) is called characteristic if there is an equivariant map $F: (\Sigma^k, T) \rightarrow (S^N, A), N \geq k$, such that F is transverse to $S^{N+m-k} \subseteq S^N$ and $F^{-1}(S^{N+m-k}) = M^m$.

PROPOSITION 1.5. Let $(\Sigma^{2n+1}, T) = (\Sigma_g, T_g)$ for an A -diffeomorphism g as in (1.2). If one of f_1, f_2 corresponding to g in (1.3) is A -homotopic to Id , then (Σ^{2n+1}, T) admits S^m , where $m = 1, \dots, n$, as characteristic spheres, such that $(S^m, T|S^m)$ is conjugate to (S^m, A) .

PROOF. Suppose $f_1 \sim^A \text{Id}$. We take H in (1.4) to be an A -homotopy between $f_2^{-1} \circ g$ and Id , and $h_1 = \text{Id}$. Let $S^m = S^m \times 0 \subseteq S^n \times D^{n+1}$. From (1.4), we see that $F^{-1}(S^m) = h_1^{-1}(S^m) = S^m$. Since $h_1 = \text{Id}$, $(S^m, T|S^m)$ is equivalent to (S^m, A) . Q.E.D.

REMARK 1.6. In [6], Browder showed that there exists a smooth involution (Σ_0^{4k+1}, T_0) which admits no m -dimensional homotopy sphere, $m \neq 4l+1$, as characteristic manifold. Hence, any A -diffeomorphism g of $S^n \times S^n$ such that $(\Sigma_g, T_g) = (\Sigma_0^{4k+1}, T_0)$ is not A -homotopic to the identity by (1.5).

2. Nonuniqueness of the decomposition. The decomposition for (Σ^{2n+1}, T) in (1.1) is not unique: we may have different A -diffeomorphisms f and g such that $(\Sigma^{2n+1}, T) = (\Sigma_f, T_f) = (\Sigma_g, T_g)$. But we have the following

PROPOSITION 2.1 [20]. For $n \geq 3$, $(\Sigma^{2n+1}, T) = (\Sigma_f, T_f) = (\Sigma_g, T_g)$ iff there exist A -diffeomorphisms $H: S^n \times D^{n+1} \rightarrow S^n \times D^{n+1}$ and $K: D^{n+1} \times S^n \rightarrow D^{n+1} \times S^n$ such that, when we restrict our attention to the boundary, $g = KfH$.

Two diffeomorphisms f and g of a manifold M are called concordant, if there exists a diffeomorphism $H: M \times [0, 1] \rightarrow M \times [0, 1]$ such that $H(x, 0) =$

$(f(x), 0), H(x, 1) = (g(x), 1)$. Similarly, we have the notion of A -concordance between two A -diffeomorphisms. If f and g are A -concordant diffeomorphisms of $S^n \times S^n$, then $(\Sigma_f, T_f) = (\Sigma_g, T_g)$, which can be seen by constructing an equivariant diffeomorphism F between them as follows:

$$\begin{array}{ccccccc} (\Sigma_f, T_f) = S^n \times D^{n+1} & \cup_f & S^n \times S^n \times I & \cup_{\text{Id}} & D^{n+1} \times S^n \\ \downarrow F & & \downarrow \text{Id} & & \downarrow H & & \downarrow \text{Id} \\ (\Sigma_g, T_g) = S^n \times D^{n+1} & \cup_{\text{Id}} & S^n \times S^n \times I & \cup_g & D^{n+1} \times S^n \end{array}$$

where H^{-1} is an A -concordance between f and g . Q.E.D.

Now, we are going to determine the A -diffeomorphisms H and K in (2.1) within A -concordance classes.

DEFINITION. A bundle map f_a for $S^n \times S^n$ over the first factor is a map of the form $f_a(x, y) = (x, a(x) \cdot y)$, where the homotopy class $\{a\} \in \pi_n(SO_{n+1})$ and a is a smooth map of S^n to SO_{n+1} .

A bundle map f_a is a diffeomorphism. If $\{a\}, \{b\} \in \pi_n(SO_{n+1})$ are homotopic, then f_a and f_b are concordant. Conversely, if f_a and f_b are concordant, then it was shown in [19, 5.2] that a and b are homotopic. Actually, we have

LEMMA 2.2. [19]. *The concordance classes of orientation-preserving diffeomorphisms of $S^n \times S^n$, which can be extended to orientation-preserving diffeomorphisms of $S^n \times D^{n+1}$, are in 1-1 correspondence with $\pi_n(SO_{n+1})$.*

We write $S^n = \{x = (x_0, \dots, x_n) \in R^{n+1} \mid \sum_{i=0}^n x_i^2 = 1\}$. A acts on S^n as an $(n+1)$ -square matrix with -1 on its diagonal and 0 elsewhere. For $(x, y) \in S^n \times S^n$, $A(x, y) = (-x, -y)$.

LEMMA 2.3. *Let $b \in \pi_n(SO_{n+1})$. Then the bundle map f_b defined above is an A -equivariant bundle map if and only if b factors through P^n by the double covering map $\pi: S^n \rightarrow P^n$.*

PROOF. $f_b A = A f_b \iff f_b A(x, y) = A f_b(x, y) \iff (-x, b(-x) \cdot (-y)) = (-x, -b(x) \cdot y) \iff b(-x) \cdot A \cdot y = A \cdot b(x) \cdot y$. Since A lies in the center of SO_{n+1} , we have $b(x) \cdot y = b(-x) \cdot y$ for all $y \in S^n$. Hence $f_b A = A f_b$ iff $b(x) = b(-x)$ for all $x \in S^n$ iff b factors through $\pi: S^n \rightarrow P^n$. Q.E.D.

LEMMA 2.4. *Every A -diffeomorphism of $S^n \times S^n$, $n \geq 3$, which can be extended to an orientation-preserving A -diffeomorphism of $S^n \times D^{n+1}$ is A -concordant to an A -equivariant bundle map over the first factor.*

PROOF. Let f be such an A -diffeomorphism, and h be its A -equivariant extension to $S^n \times D^{n+1}$. $f'(h')$ denotes the map induced by $f(h)$ on the orbit space $S^n \times S^n/A(S^n \times D^{n+1}/A)$. Let $i': S^n \times 0/A \rightarrow S^n \times D^{n+1}/A$ be the

inclusion. $h'|S^n \times 0/A$ and i' are homotopic by [24]. Hence, they are isotopic by a theorem of Haefliger, [12]. By the equivariant isotopy extension theorem, [28], there exists an A -equivariant diffeomorphism H of $S^n \times D^{n+1}$ such that H is equivariantly isotopic to identity, $H|S^n \times S^n = \text{identity}$, and $H \circ h|S^n \times 0/A = i$, the inclusion. Let B^{n+1} be a small disk in D^{n+1} , with B^{n+1} and D^{n+1} concentric. Both $S^n \times B^{n+1}$ and $H \circ h(S^n \times B^{n+1})$ are equivariant tubular neighborhoods of $S^n \times 0$ in $S^n \times D^{n+1}$. Then by the uniqueness of the equivariant tubular neighborhoods, [4, p. 310], there exists an A -equivariant diffeomorphism G of $S^n \times D^{n+1}$ such that G is A -equivariantly isotopic to identity, $G|S^n \times S^n = \text{Id}$, and $G \circ H \circ h|S^n \times B^{n+1}$ is an A -equivariant bundle map covering the identity on S^n . The restriction of $G \circ H \circ h$ to $S^n \times D^{n+1}$ -interior $S^n \times B^{n+1}$ gives us an A -concordance between f and an A -equivariant bundle map. Q.E.D.

Similarly, every A -diffeomorphism of $S^n \times S^n$ which extends equivariantly to $D^{n+1} \times S^n$ is A -concordant to an A -bundle over the second factor.

PROPOSITION 2.5. *The A -concordance classes of orientation-preserving A -diffeomorphism H (or K) in (2.1) are in 1-1 correspondence with the Image π^* of $[P^n, SO_{n+1}]$ in $[S^n, SO_{n+1}]$.*

PROOF. Let h be an A -diffeomorphism of $S^n \times S^n$, which can be extended equivariantly to $S^n \times D^{n+1}$. h is A -concordant to an A -equivariant bundle map f_b by (2.4), where $b \in \pi_n(SO_{n+1})$. From (2.3), we know that b factors through $\pi: S^n \rightarrow P^n$, i.e. $b \in \text{Image } \pi^*$.

The above correspondence $H \rightarrow b$ is well-defined. If H' is A -concordant to H , and $b' \in \text{Image } \pi^*$ corresponds to H' , then f_b and $f_{b'}$ are A -concordant, hence concordant. b is homotopic to b' by (2.2). This correspondence is 1-1 and onto, since the mapping given by $b \rightarrow f_b$ for $b \in \text{Image } \pi^*$ is its inverse. Q.E.D.

3. The image of π^* : $[P^n, SO_{n+1}] \rightarrow [S^n, SO_{n+1}]$. In this section, we will compute $[P^n, SO_{n+1}]$ and its image under π^* : $[P^n, SO_{n+1}] \rightarrow [S^n, SO_{n+1}]$. Let us first recall some facts about $\pi_n(SO_{n+1})$, which, for instance, can be found in [16] or [18].

Let $s_{m*}: \pi_k(SO_m) \rightarrow \pi_k(SO_{m+1})$ denote the homomorphism induced by the natural embedding $s: SO_n \rightarrow SO_{n+1}$. Consider the following exact sequence,

$$\cdots \xrightarrow{\partial_m} \pi_k(SO_m) \xrightarrow{s_{m*}} \pi_k(SO_{m+1}) \xrightarrow{q_{m*}} \pi_k(S^m) \xrightarrow{\partial_m} \pi_{k-1}(SO_m) \rightarrow \cdots$$

Let ι_m denote the generator of $\pi_m(S^m)$. Write $\tau_m = \partial_{m+1} \iota_{m+1} \in \pi_m(SO_{m+1})$. Putting $m = n, n+1$ in the above exact sequence, we have the following proposition from [18]:

PROPOSITION 3.1. *For n odd, $\neq 1, 3, 7$, $\pi_n(SO_{n+1})$ is the direct sum of two cyclic subgroups image ∂_{n+1} and image s_{n*} . Moreover, $s_{n+1*}: \text{image } s_{n*} \subseteq$*

$\pi_n(SO_{n+1}) \rightarrow \pi_n(SO_{n+2})$ is an isomorphism. For n even, $\pi_n(SO_{n+1})$ is the direct sum of image ∂_{n+1} and a certain cyclic subgroup G such that $s_{n+1*}: G \subseteq \pi_n(SO_{n+1}) \rightarrow \pi_n(SO_{n+2})$ is an isomorphism.

From now on, we will write τ for $\tau_n = \partial_{n+1} \iota_{n+1}$, if no confusion will arise. Let σ denote the generator of the other cyclic summand of $\pi_n(SO_{n+1})$. Here we list the values of $\pi_n(SO_{n+1})$ and $\pi_n(SO_{n+2}) = \pi_n(SO)$ for $n > 3$ and $n \neq 7$, from [16].

$n(\bmod 8)$	0	1	2	3	4	5	6	7
$\pi_n(SO_{n+1})$	$Z_2 + Z_2$	$Z + Z_2$	Z_2	$Z + Z$	Z_2	Z	Z_2	$Z + Z$
	τ, σ	τ, σ	τ	τ, σ	τ	τ	τ	τ, σ
$\pi_n(SO)$	Z_2	Z_2	0	Z	0	0	0	Z

Consider the maps $\pi: S^n \rightarrow P^n$, $i: P^n \rightarrow P^{n+1}$, $j: P^n \rightarrow S^n$; which are the double covering, the inclusion, and the map pinching the complement of an open ball to a point. j generates $[P^n, S^n] = H^n(P^n) = Z$ or Z_2 , n odd or even; by Hopf's theorem, [23].

The element $\tau \in \pi_n(SO_{n+1})$ is also the characteristic map for the tangent bundle of S^{n+1} , [14] or [30]. We can choose a representative for τ such that $\tau(x) = \tau(-x)$ for $x \in S^n$, [14], which is defined as follows: let $\alpha: S^n \rightarrow SO_{n+1}$ be the map defined by the requirement that $\alpha(x)$ be a reflection through the hyperplane in R^{n+1} orthogonal to x , and let e denote the north pole of S^n , then we have $\tau(x) = \alpha(x)\alpha(e)$, [14, p. 89]. Hence this τ factors through $\pi: S^n \rightarrow P^n$, and $\tau_p \in [P^n, SO_{n+1}]$ is defined by $\tau = \tau_p \pi$.

Let $q: SO_{n+1} \rightarrow S^n$ denote the projection in the fibration $SO_n \rightarrow SO_{n+1} \rightarrow S^n$.

PROPOSITION 3.2. *The following diagram is homotopically commutative.*

$$\begin{array}{ccc}
 S^n & \xrightarrow{\pi} & P^n \\
 \tau \downarrow & \nearrow \tau_p & \downarrow j \\
 SO_{n+1} & \xrightarrow{q} & S^n
 \end{array}$$

PROOF. We have to show that $g\tau_p$ is homotopic to j . From the above description of τ , $\tau(x) = \alpha(x)\alpha(e)$; we see that $q\tau(x) = \alpha(x)\alpha(e) = \alpha(x)(-e)$, which is the point on S^n obtained by moving e toward x along the great circle passing through e and x by an angle twice the angle between e and x . We note that $q\tau$ maps the interior of the northern hemisphere D_+^n of S^n onto $S^n - \{e\}$ as a homeomorphism, and maps the equator S^{n-1} to $-e$. Since π maps interior D_+^n homeomorphically onto $P^n - P^{n-1}$, and $q\tau = q\tau_p\pi$, we see that $q\tau_p$ is just the

map pinching the complement of an open ball to a point, which is j . Q.E.D.

LEMMA 3.3. τj is null-homotopic in $[P^{2k}, SO_{2k+1}]$.

PROOF. We have $\tau j = \tau_p \pi j: P^{2k} \rightarrow S^{2k} \rightarrow SO_{2k+1}$. Consider $f = \pi j: P^{2k} \rightarrow P^{2k}$. Since f factors through S^{2k} , $f_*(\pi_1(P^{2k})) = 0$. Hence f is a non-orientation-true map in the language of [25]. By Theorem 1.2 of [25], (see also [25, 1.3(d)]), f is null-homotopic. Q.E.D.

Let $KO^{-k}(-)$ denote the reduced real K -theory. We have $KO^{-k}(X) = [\Sigma^k X, B_{SO}]$ for any finite CW complex X , [14]. $KO^{-1}(P^n) = [\Sigma P^n, B_{SO}] = [P^n, \Omega B_{SO}] = [P^n, SO]$, the latter one is equal to $[P^n, SO_{n+2}]$ because (SO, SO_{n+2}) is $(n+1)$ -connected.

For any fibration $F \rightarrow E \rightarrow B$, and any finite CW complex X , there is a fibre mapping sequence $\cdots \rightarrow [X, \Omega E] \rightarrow [X, \Omega B] \rightarrow [X, F] \rightarrow [X, E] \rightarrow [X, B]$, [23], which is exact.

LEMMA 3.4. For n odd, $Z \rightarrow [P^n, SO_{n+1}] \rightarrow [P^n, SO_{n+2}] \rightarrow 0$ is exact. For n even, $Z_2 \rightarrow [P^n, SO_{n+1}] \rightarrow [P^n, SO_{n+2}] \rightarrow 0$ is exact.

PROOF. Substitute $SO_{n+1} \rightarrow SO_{n+2} \rightarrow S^{n+1}$ and P^n into the above fibre mapping sequence. $[P^n, S^{n+1}] = 0$. $[P^n, \Omega S^{n+1}] = [\Sigma P^n, S^{n+1}] = H^{n+1}(\Sigma P^n) = H^n(P^n) = Z$, n odd; $= Z_2$, n even. Q.E.D.

We also need the following from [6] or [11].

PROPOSITION 3.5. $KO^{-1}(P^m) = Z_2$, for $m \not\equiv 3 \pmod{4}$; $KO^{-1}(P^m) = Z + Z_2$, for $m \equiv 3 \pmod{4}$, where the Z summand is the image of $KO^{-1}(S^m)$ under the degree 1 map $j: P^m \rightarrow S^m$. The inclusion $P^k \subseteq P^m$ induces $KO^{-1}(P^m) \rightarrow KO^{-1}(P^k)$, which is an isomorphism on Z_2 and annihilates the Z factor.

Replacing P^k by the mapping cylinder M_π of $\pi: S^k \rightarrow P^k$, we can change π into a cofibration $\pi': S^k \rightarrow M_\pi$, and we may consider $i: P^k \rightarrow P^{k+1}$ as the cofibre. For a simple space X , we have the following Puppe exact sequence, [27],

$$\begin{aligned} \cdots \longrightarrow [\Sigma P^k, X] &\xrightarrow{\Sigma \pi^*} \pi_{k+1}(X) \xrightarrow{j^*} [P^{k+1}, X] \\ &\xrightarrow{i^*} [P^k, X] \xrightarrow{\pi^*} \pi_k(X). \end{aligned}$$

Putting $k = n - 1$ and $X = SO_{n+1}$ in the above Puppe sequence, we have the following exact sequence:

$$\begin{aligned} (3.6) \quad \cdots \longrightarrow \pi_n(SO_{n+1}) &\xrightarrow{j^*} [P^n, SO_{n+1}] \\ &\xrightarrow{i^*} [P^{n-1}, SO_{n+1}] \xrightarrow{\pi^*} \pi_{n-1}(SO_{n+1}). \end{aligned}$$

For n odd, $j\pi: S^n \rightarrow P^n \rightarrow S^n$ is of degree 2. Hence $\pi^*j^*: \pi_n(SO_{n+1}) \rightarrow [P^n, SO_{n+1}] \rightarrow \pi_n(SO_{n+1})$ is just the multiplication by 2.

If $n \equiv 3 \pmod{4}$, then $\pi_n(SO_{n+1}) = Z + Z$, generated by τ and σ . π^*j^* is 1-1, hence j^* is 1-1. Thus $[P^n, SO_{n+1}]$ contains $Z + Z$. One of the generators is τ_p , since $\pi^*\tau_p = \tau_p\pi = \tau$. Therefore $j^*\tau = 2\tau_p$. For $m \equiv 2 \pmod{4}$, $\pi_m(SO_{m+2}) = 0$ and $[P^m, SO_{m+2}] = KO^{-1}(P^m) = Z_2$ by (3.5). Hence, the exact sequence (3.6) becomes $0 \rightarrow Z + Z \xrightarrow{j^*} [P^n, SO_{n+1}] \rightarrow Z_2 \rightarrow 0$. Since $j^*\tau = 2\tau_p$, we see that $[P^n, SO_{n+1}] = Z + Z$, generated by τ_p and $b = j^*\sigma$. $\pi^*b = \pi^*j^*\sigma = 2\sigma$.

If $n \equiv 5 \pmod{8}$, then $\pi_n(SO_{n+1}) = Z$, generated by τ . The argument in the preceding paragraph shows that $[P^n, SO_{n+1}] = Z$, generated by τ_p , and $j^*\tau = 2\tau_p$.

Now consider the following commutative diagram, where the rows are Puppe sequences and the columns are fibre mapping sequences.

$$(3.7) \quad \begin{array}{ccccc} \pi_{n+1}(S^{n+1}) & \xrightarrow{j_1} & [P^n, \Omega S^{n+1}] & \longrightarrow & [P^{n+1}, \Omega S^{n+1}] = 0 \\ \downarrow \partial & & \downarrow \partial_1 & & \downarrow \\ \pi_n(SO_{n+1}) & \xrightarrow{j^*} & [P^n, SO_{n+1}] & \xrightarrow{i^*} & [P^{n-1}, SO_{n+1}] \\ \downarrow s_* & & \downarrow s'_* & & \downarrow \\ \pi_n(SO_{n+2}) & \xrightarrow{j_2} & [P^n, SO_{n+2}] & \longrightarrow & [P^{n-1}, SO_{n+2}] \end{array}$$

If $n \equiv 1 \pmod{8}$, then $\pi_n(SO_{n+1}) = Z + Z_2$, generated by τ and σ respectively. Since j^* is 1-1 on the Z summand, and $\pi^*\tau_p = \tau$, $[P^n, SO_{n+1}]$ contains a Z subgroup which is generated by τ_p . Let ι_{n+1} denote the generator of $\pi_{n+1}(S^{n+1})$. In the diagram (3.7), $\partial\iota_{n+1} = \tau$ and $[P^n, \Omega S^{n+1}] = [\Sigma P^n, S^{n+1}] = H^n(P^n) = Z$ is generated by $j_1\iota_{n+1}$. $[P^n, SO_{n+2}] = KO^{-1}(P^n) = Z_2$ by (3.5). The middle column of (3.7) reads $Z \xrightarrow{\partial_1} [P^n, SO_{n+1}] \xrightarrow{s'_*} Z_2$. If $\partial_1(aj_1\iota_{n+1}) = \tau_p$ for some integer a , then $j^*(a\tau) = aj^*\partial\iota_{n+1} = \partial_1(aj_1\iota_{n+1}) = \tau_p$, a contradiction. Hence $\tau_p \notin \text{image } \partial_1$, $s'_*\tau_p \neq 0$. $[P^n, SO_{n+1}] = Z$, generated by τ_p . Also $j^*\tau = 2\tau_p$, $j^*\sigma = 0$.

We now consider the case n is even.

LEMMA 3.8. *For n even, $[P^n, SO_{n+1}] = Z_2$, which is generated by τ_p .*

PROOF. In the diagram (3.7), j_1 is onto. Let ι_{n+1} generate $\pi_{n+1}(S^{n+1})$. $j_1\iota_{n+1}$ generates $[P^n, \Omega S^{n+1}] = H^{n+1}(\Sigma P^n) = Z_2$. $\partial_1(j_1\iota_{n+1}) = j^*\partial\iota_{n+1} = j^*\tau = \tau_j = 0$ by (3.3). The map s'_* in the middle column in (3.7) is onto by (3.4). Hence $s'_*: [P^n, SO_{n+1}] \rightarrow [P^n, SO_{n+2}] = KO^{-1}(P^n) = Z_2$ is an isomorphism. Thus $[P^n, SO_{n+1}] = Z_2$, generated by τ_p . Q.E.D.

Summing up, we have the following:

THEOREM 3.9. Assume $n \neq 1, 2, 3, 7$. For n even, $[P^n, SO_{n+1}] = Z_2$, generated by τ_p , and $\pi^* \tau_p = \tau \in \pi_n(SO_{n+1})$. For $n \equiv 1 \pmod{4}$, $[P^n, SO_{n+1}] = Z$, generated by τ_p , and $\pi^* \tau_p = \tau$. For $n \equiv 3 \pmod{4}$, $[P^n, SO_{n+1}] = Z + Z$, generated by τ_p and b , where $b = j^* \sigma$, and $\pi^* \tau_p = \tau$, $\pi^* b = 2\sigma$.

Let $\text{Im } \pi^*$ denote the image of $[P^n, SO_{n+1}]$ in $\pi_n(SO_{n+1})$ under π^* .

$n(\pmod{8})$	0	1	2	3	4	5	6	7
$\pi_n(SO_{n+1})$	$Z_2 + Z_2$	$Z + Z_2$	$Z_2 + Z + Z$	Z_2	Z	Z_2	$Z + Z$	
	τ, σ	τ, σ	τ	τ, σ	τ	τ	τ	τ, σ
$\text{Im } \pi^*$	τ	τ	τ	$\tau, 2\sigma$	τ	τ	τ	$\tau, 2\sigma$

COROLLARY 3.10. $n > 3$ and $\neq 7$. Let $s'_*: [P^n, SO_{n+1}] \rightarrow [P^n, SO_{n+2}] = KO^{-1}(P^n)$ be induced by the inclusion. Then $s'_*(\tau_p)$ generates the Z_2 summand of $KO^{-1}(P^n)$; and for $n \equiv 3 \pmod{4}$, $s'_*(b)$ generates the Z summand of $KO^{-1}(P^n)$.

PROOF. We have proved the corollary for n even in (3.8). For n odd, $s'_*: [P^n, SO_{n+1}] \rightarrow [P^n, SO_{n+2}]$ is onto, (3.4). In (3.7), let ι_{n+1} generate $\pi_{n+1}(S^{n+1})$, $j_1 \iota_{n+1}$ generates $[P^n, \Omega S^{n+1}]$. $\partial_1(j_1 \iota_{n+1}) = j^* \partial \iota_{n+1} = j^* \tau = 2\tau_p$. Hence s'_* maps τ_p to the generator of the Z_2 summand of $KO^{-1}(P^n)$, and maps b to the generator of the Z summand for $n \equiv 3 \pmod{4}$ by the exactness of the middle column in (3.7). Q.E.D.

4. Proof of Proposition 1.2. We are going to prove Proposition 1.2 in this section. Let $D(S^n \times S^n)(D(S^n \times S^n, A))$ denote the group of concordance (A -concordance) classes of diffeomorphisms (A -diffeomorphisms) of $S^n \times S^n$. $\emptyset: D(S^n \times S^n, A) \rightarrow D(S^n \times S^n)$ is the homomorphism forgetting the action. Define $D_0(S^n \times S^n) =$ the subgroup of $D(S^n \times S^n)$ consisting of those elements which are homotopic to Id . Let $J_{2n+1} = \emptyset^{-1}(D_0(S^n \times S^n))$. Given a diffeomorphism (an A -diffeomorphism) f of $S^n \times S^n$, we will write $\{f\}$ ($\{f\}_A$) for its concordance class in $D(S^n \times S^n)(D(S^n \times S^n, A))$.

If f is a diffeomorphism of $S^n \times S^n$, then f_* induces an automorphism of $H_n(S^n \times S^n)$. We can associate to f_* its matrix representative M_f with respect to the natural basis $\{S^n \times 0, 0 \times S^n\}$ of $H_n(S^n \times S^n)$. M_f is an element of $GL(2, Z)$ the group of 2×2 -unimodular matrices. Let $\psi: D(S^n \times S^n) \rightarrow GL(2, Z)$ be the homomorphism defined by $f \rightarrow M_f$. We have $fg \rightarrow M_g \cdot M_f$.

From [19], we have the following:

PROPOSITION 4.1 [19]. If n is even, then image ψ consists of eight matrices:

$\begin{pmatrix} \pm 1 & 0 \\ 0 & \pm 1 \end{pmatrix}$ and $\begin{pmatrix} 0 & \pm 1 \\ \pm 1 & 0 \end{pmatrix}$. If $n = 1, 3, 7$, then image $\psi = GL(2, Z)$. If n is odd, but $n \neq 1, 3, 7$, then image ψ is the subgroup of $GL(2, Z)$ consisting of matrices $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ where $ab \equiv cd \equiv 0 \pmod{2}$.

LEMMA 4.2. If f is a diffeomorphism of $S^n \times S^n$ such that $S^n \times D^{n+1} \cup_f D^{n+1} \times S^n$ is a homotopy sphere, then $M_f = \begin{pmatrix} a & b \\ c & \pm 1 \end{pmatrix}$.

PROOF. Write $V_1 = S^n \times D^{n+1}$, $V_2 = D^{n+1} \times S^n$, $\partial V_1 = \partial V_2 = S^n \times S^n$. Let $i_k: \partial V_k \rightarrow V_k$, $k = 1$ or 2 , be the inclusion. Let $\{x, y\}$ denote the natural basis $\{S^n \times 0, 0 \times S^n\}$ of $H_n(S^n \times S^n)$, and u_1, u_2 the generator of $H_n(S^n \times D^{n+1})$, $H_n(D^{n+1} \times S^n)$ respectively. From the Mayer-Vietoris sequence, we have

$$0 \longrightarrow H_n(\partial V_1) \xrightarrow{(j_1, j_2)} H_n(V_1) \oplus H_n(V_2) \longrightarrow 0$$

where $j_1 = i_{1*}$ and $j_2 = i_{2*}f_*$. $i_{1*}x = u_1$, $i_{1*}y = 0$, $i_{2*}x = 0$, $i_{2*}y = u_2$. Let $M_f = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. $f_*x = ax + by$ and $f_*y = cx + dy$. Hence $j_1x = u_1$, $j_1y = 0$, $j_2x = i_{2*}f_*x = bu_2$, $j_2y = i_{2*}f_*y = du_2$. Thus the matrix for (j_1, j_2) with respect to the basis $\{x, y\}$ and $\{u_1, u_2\}$ is $\begin{pmatrix} 1 & b \\ 0 & d \end{pmatrix}$, which is unimodular, $d = \pm 1$. Q.E.D.

LEMMA 4.3. $n > 3$ and $n \neq 7$. If $(\Sigma^{2n+1}, T) = (\Sigma_f, T_f)$ for some A -diffeomorphism f of $S^n \times S^n$, then there exists another A -diffeomorphism g such that M_g is the identity matrix, and $(\Sigma^{2n+1}, T) = (\Sigma_g, T_g)$.

PROOF. If n is even, then it follows from (4.1) and (4.2) that $M_f = \begin{pmatrix} \pm 1 & 0 \\ 0 & \pm 1 \end{pmatrix}$. Consider the A -diffeomorphisms h_1 and h_2 of $S^n \times S^n$ defined by $h_1(x, y) = (x, -y)$ and $h_2(x, y) = (-x, y)$. h_1 (or h_2) extends equivariantly to $(S^n \times D^{n+1}, A)$ (or $(D^{n+1} \times S^n, A)$). One of the A -diffeomorphisms f, fh_1, h_2fh_1 , or h_2f has the corresponding matrix = identity matrix. Take g to be this map. Also, $(\Sigma^{2n+1}, T) = (\Sigma_g, T_g)$ by (2.1).

If n is odd and $n \neq 1, 3, 7$, then $M_f = \begin{pmatrix} a & b \\ c & \pm 1 \end{pmatrix}$ by (4.2). We can compose f with h_1 or h_2 if necessary, to make $M_f = \begin{pmatrix} a & b \\ c & 1 \end{pmatrix}$ and $\det M_f = +1$. From (4.1), $b \equiv c \equiv 0 \pmod{2}$. Since $a - bc = 1$, we see that $\begin{pmatrix} a & b \\ c & 1 \end{pmatrix} = \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ c & 1 \end{pmatrix}$.

We choose a representative for τ as in §3, satisfying $\tau(x) = \tau(-x)$. Consider the maps f_τ and g_τ , defined by $f_\tau(x, y) = (x, \tau(x) \cdot y)$ and $g_\tau(x, y) = (\tau(y) \cdot x, y)$. f_τ and g_τ are A -diffeomorphisms. Theorem 7.10.1 of [14] showed that the map $q \circ \tau$ in (3.2) is of degree 2. Hence $M_{f_\tau} = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$ and $M_{g_\tau} = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}$. Define $g = g_\tau^{(-c/2)} f f_\tau^{(-b/2)}$. Since $M_{fg} = M_g M_f$, we see that M_g is the identity matrix. $(\Sigma_f, T_f) = (\Sigma_g, T_g)$ by (2.1). Q.E.D.

LEMMA 4.4. A diffeomorphism f of $S^n \times S^n$ is homotopic to Id if and only if M_f is the identity matrix.

PROOF. If f and g are diffeomorphisms of $S^n \times S^n$ such that $\{f\} = \{g\}$ in $D(S^n \times S^n)$, then $M_f = M_g$. Let $G = \{\{f\} \in D(S^n \times S^n) | M_f \text{ is the identity matrix}\}$. Theorem II of [28] showed that $0 \rightarrow H_1 + \Gamma^{2n+1} \rightarrow G \rightarrow H_2 \rightarrow 0$ is exact, where H_i , $i = 1$ or 2 , is isomorphic to $\text{image}\{s_*: \pi_n(SO_n) \rightarrow \pi_n(SO_{n+1})\}$. Let a be a smooth map representing $\{a\} \in \text{image } s_* \subseteq \pi_n(SO_{n+1})$. We define two diffeomorphisms f_a and g_a of $S^n \times S^n$ by $f_a(x, y) = (x, a(x) \cdot y)$ and $g_a(x, y) = (a(y) \cdot x, y)$. The maps $\{a\} \rightarrow \{f_a\}$ and $\{a\} \rightarrow \{g_a\}$ are isomorphisms of $\text{image } s_*$ to H_1 and H_2 respectively, (compare 2.2). Γ^{2n+1} is the Kervaire-Milnor group [17] and acts by leaving the complement of a $2n$ -disk in $S^n \times S^n$ fixed.

Let a be a smooth map representing a homotopy class $\{a\} \in \pi_n(SO_{n+1})$ such that $\{a\} = s_*\beta$ for some $\beta \in \pi_n(SO_n)$. We can take $a|D_-^n = \text{Id}$, where D_-^n (D_+^n) denotes the lower (upper) hemisphere of S^n . Hence $f_a|D_-^n \times S^n = \text{Id}$. Let e denote the north pole of S^n . Since $\{a\} = s_*\beta$, we can take $f_a|S^n \times \{e\} = \text{Id}$. By the homotopy extension theorem, f_a is homotopic to a map h such that h is the identity on a neighborhood N of $S^n \times \{e\}$ and $f_a = h$ on $D_-^n \times S^n$. h keeps the complement of a disk $\overline{S^n \times S^n} | M_f \text{ is the identity matrix}\}$.

Since every element $\{h\}$ of G has a representative h such that h leaves the complement of a $2n$ -disk fixed, we can apply the Alexander trick to see that h is homotopic to Id . Thus $G = D_0(S^n \times S^n)$. Q.E.D.

Combining (4.3) and (4.4) together, we have proved Proposition (1.2). From now on, when $n > 3$ and $\neq 7$, we will assume the A -diffeomorphism f in the decomposition $(\Sigma^{2n+1}, T) = (\Sigma_f, T_f)$, be homotopic to Id .

5. Group structure on $hS(P^{2n+1})$. Given an involution (Σ^{2n+1}, T) , $n > 3$ and $\neq 7$, there exists an A -diffeomorphism f of $S^n \times S^n$ such that $(\Sigma^{2n+1}, T) = (\Sigma_f, T_f)$, and $(\Sigma_f, T_f) = (\Sigma_g, T_g)$ if $g \in \{f\}_A \in J_{2n+1} = \emptyset^{-1}(D_0(S^n \times S^n))$. But $\{f\}_A \in J_{2n+1}$ is not uniquely determined by (Σ^{2n+1}, T) . Suppose we can find a subgroup G_{2n+1} of J_{2n+1} such that G_{2n+1} is in 1-1 correspondence with $hS(P^{2n+1})$ under the mapping $\{f\}_A \rightarrow (\Sigma_f, T_f)$. Then $(hS(P^{2n+1}), *)$ forms a group by carrying over the composition law in G_{2n+1} : $(\Sigma_f, T_f)^*(\Sigma_g, T_g) = (\Sigma_{fg}, T_{fg})$. In this section, we will show that such a subgroup G_{2n+1} exists for $n \equiv 0, 1, 2 \pmod{4}$.

THEOREM 5.1. *For n even, and > 2 , such a subgroup G_{2n+1} of J_{2n+1} exists, hence $(hS(P^{2n+1}), *)$ is a subgroup.*

PROOF. From [19] or (4.4) above, we know that $D_0(S^n \times S^n)$ is the semi-direct product of $H_1 + \Gamma^{2n+1}$ and H_2 , where H_1 and H_2 are isomorphic to $\text{image}\{s_*: \pi_n(SO_n) \rightarrow \pi_n(SO_{n+1})\}$, which is equal to $\pi_n(SO_{n+1})$, [18]. $\pi_n(SO_{n+1}) = Z_2$, generated by τ for $n \not\equiv 0 \pmod{8}$; $\pi_n(SO_{n+1}) = Z_2 + Z_2$, generated by τ and σ for $n \equiv 0 \pmod{8}$. Every element of $D_0(S^n \times S^n)$ can be

uniquely expressed in the form $h_2 y h_1$, where $h_i \in H_i$ and $y \in \Gamma^{2n+1}$, [17]. We define a subgroup F_{2n+1} of $D_0(S^n \times S^n)$ as follows: $F_{2n+1} = \Gamma^{2n+1}$ for $n \not\equiv 0 \pmod{8}$; if $n \equiv 0 \pmod{8}$, then F_{2n+1} is the semidirect product of $(\sigma_1) + \Gamma^{2n+1}$ and (σ_2) , where (σ_i) denotes the cyclic group of order 2 generated by σ_i : $\sigma_1(x, y) = (x, \sigma(x) \cdot y)$, $\sigma_2(x, y) = (\sigma(y) \cdot x, y)$. Let $G_{2n+1} = \partial^{-1}(F_{2n+1})$.

We choose a smooth representative for $\tau \in \pi_n(SO_{n+1})$ such that $\tau(x) = \tau(-x)$. Define τ_1 and τ_2 by $\tau_1(x, y) = (x, \tau(x) \cdot y)$ and $\tau_2(x, y) = (\tau(y) \cdot x, y)$. τ_1 and τ_2 are A -diffeomorphisms of $S^n \times S^n$. Since τ commutes with σ in $\pi_n(SO_{n+1})$, [19], we see that any element of $D_0(S^n \times S^n)$ can be uniquely expressed in the form $b y a$, where $b \in (\tau_2)$, $a \in (\tau_1)$, and $y \in F_{2n+1}$.

If $h \in \{h\}_A \in J_{2n+1}$, then $\partial(\{h\}_A) = \{\tau_2^d\}\{f\}\{\tau_1^c\}$, where $\{f\} \in F_{2n+1}$, and $c, d = 0$ or 1 uniquely determined by $\{h\}_A$. We define $g = \tau_2^d h \tau_1^c$, which is an A -diffeomorphism of $S^n \times S^n$. Since $\{\tau_i\}$ is of order two in $D_0(S^n \times S^n)$, $\partial(\{g\}_A) \in F_{2n+1}$. $(\Sigma_g, T_g) = (\Sigma_h, T_h)$ by (2.1). On the other hand, if $\{f\}_A$, $\{g\}_A \in G_{2n+1}$ and $(\Sigma_f, T_f) = (\Sigma_g, T_g)$, then g is A -concordant to $\tau_2^d f \tau_1^c$, where $c, d = 0$ or 1 , by (2.1) and (3.9). But $\partial(\{f\}_A), \partial(\{g\}_A) \in F_{2n+1}$. Hence $c = d = 0$, $\{f\}_A = \{g\}_A$. Q.E.D.

THEOREM 5.2. *For $n \equiv 1 \pmod{4}$, $hS(P^{2n+1})$ is in 1-1 correspondence with J_{2n+1} ; hence $(hS(P^{2n+1}), *)$ forms a group.*

PROOF. As in (5.1), we know that $D_0(S^n \times S^n)$ is the semidirect product of $H_1 + \Gamma^{2n+1}$ and H_2 , where H_1 and H_2 are isomorphic to $\text{image}\{s_*: \pi_n(SO_n) \rightarrow \pi_n(SO_{n+1})\}$, which is 0 for $n \equiv 5 \pmod{8}$; and Z_2 , generated by σ for $n \equiv 1 \pmod{8}$. $\sigma \notin \text{image}\{\pi^*: [P^n, SO_{n+1}] \rightarrow [S^n, SO_{n+1}]\}$ by (3.9). Hence no element in $D_0(S^n \times S^n)$ is concordant to an A -bundle map by (2.5). Thus $hS(P^{2n+1})$ is in 1-1 correspondence with $J_{2n+1} = \partial^{-1}(D_0(S^n \times S^n))$ by (1.2) and (2.1). Q.E.D.

REMARK 5.3. For $n > 7$ and $n \equiv 3 \pmod{4}$, we have the exact sequence: $0 \rightarrow Z + \Gamma^{2n+1} \rightarrow D_0(S^n \times S^n) \rightarrow Z \rightarrow 0$, where each Z is isomorphic to $\text{image}\{s_*: \pi_n(SO_n) \rightarrow \pi_n(SO_{n+1})\}$, generated by σ . We know that $2m\sigma \in \text{image } \pi^*$ but $(2m+1)\sigma \notin \text{image } \pi^*$ by (3.9). Let σ_1 and σ_2 be defined by $\sigma_1(x, y) = (x, \sigma(x) \cdot y)$, $\sigma_2(x, y) = (\sigma(y) \cdot x, y)$. If none of the four diffeomorphisms $\sigma_1, \sigma_2, \sigma_2\sigma_1$, and $\sigma_1\sigma_2$ is concordant to an A -diffeomorphism, i.e. $\{\sigma_1\}, \{\sigma_2\}, \{\sigma_2\sigma_1\}, \{\sigma_1\sigma_2\} \notin \text{image } \partial$, then we can take $G_{2n+1} = \partial^{-1}(\Gamma^{2n+1})$ as in (5.1) and (5.2).

REMARK 5.4. *The case $n = 3$, $hS(P^7)$.* Viewing S^3 as the unit sphere in the quaternionic space, we define $\{r\}, \{t\} \in \pi_3(SO_4)$ by $r(x) \cdot y = xyx^{-1}$, $t(x) \cdot y = xy$. $\pi_3(SO_4) = Z + Z$ is generated by $\{r\}$ and $\{t\}$, [14, p. 94]. As in (3.9), we can show that $\text{image}\{\pi^*: [P^3, SO_4] \rightarrow \pi_3(SO_4)\}$ is generated by $\{r\}$ and $2\{t\}$.

Let t_1 and t_2 be defined in the same way as σ_1 and σ_2 in (5.3). If $\{t_1\}, \{t_2\}, \{t_1 t_2\}, \{t_2 t_1\} \notin \text{image } \varnothing$, then (1.2) is also true in this case and $(hS(P^7), *)$ forms a group.

6. Additivity of Browder-Livesay index invariant. In [7], [8], Browder and Livesay defined a desuspension invariant σ for any free involution (Σ^{2n+1}, T) as follows: construct an $(n-1)$ -connected characteristic submanifold N^{2n} for (Σ^{2n+1}, T) , i.e. $\Sigma^{2n+1} = A \cup B$, $A \cap B = N$, $TA = B$, $TN = N$, and A, B are $(2n+1)$ -submanifolds of Σ^{2n+1} . Let $K_n = \text{Ker}(H_n(N) \rightarrow H_n(A))$. If n is odd, they define a unimodular even symmetric bilinear form B on K_n (modulo torsion) by $B(x, y) = x \cdot T_* y$. Let $\sigma(\Sigma^{2n+1}, T) = (1/8) \text{index } B \in \mathbb{Z}$. If n is even, they use \mathbb{Z}_2 as coefficients and define the unimodular bilinear form B_2 on K_n (with \mathbb{Z}_2 coefficients) by $B_2(x, y) = x \cdot T_* y$. They also defined a quadratic form $\psi: K_n \rightarrow \mathbb{Z}_2$ associated to B_2 , $B_2(x, y) = \psi(x+y) + \psi(x) + \psi(y)$, such that, if $x \in K_n$ is represented by an immersed sphere d , then $\psi(x) = 1$ iff $d \cap Td$ in general position consists of an odd number of pairs of points. Write $\sigma(\Sigma^{2n+1}, T) = c(\psi) \in \mathbb{Z}_2$, the Art invariant of ψ . They also showed that for $n \geq 3$, $\sigma(\Sigma^{2n+1}, T) = 0$ iff (Σ^{2n+1}, T) admits a codimension 1 invariant sphere (for details, see [8]).

From now on, we assume $n \geq 3$. Suppose $(\Sigma^{2n+1}, T) = (\Sigma_f, T_f)$ for an A -diffeomorphism f of $S^n \times S^n$, let $j: S^n \times S^n/A \rightarrow S^n \times S^{m-1}/A \rightarrow P^{n+m} = (S^n \times D^m/A) \cup_{\text{Id}} (D^{n+1} \times S^{m-1}/A)$, with m large, be the natural inclusion. Both $j \circ (f/A)$ and j are classifying maps for the same \mathbb{Z}_2 -bundle. Hence they are homotopic, by a map $F: (S^n \times S^n \times I)/A \times \text{Id} \rightarrow P^{n+m}$. We may suppose that F is smooth and transverse regular on $P^{n+m-1} = (S^n \times D^{m-1}/A) \cup_{\text{Id}} (D^{n+1} \times S^{m-2}/A)$, relative boundary. The double cover M of $F^{-1}(P^{n+m-1})$ is a characteristic submanifold of $(S^n \times S^n \times I, A \times \text{Id})$, with $\partial M = M_1 - M_0$, where $M_1 = f^{-1}(S^n \times S^{n-1}) \times 1$, $M_0 = (S^n \times S^{n-1}) \times 0$. We recall that a codim 1 characteristic submanifold M of a free involution (W, T) is a codim 1 submanifold of W such that $W = A \cup B$, where A and B are codim 0 submanifolds of W , $A \cap B = M$, and $TA = B$, [8].

In the rest of this section, we will write $(W, T) = (S^n \times S^n \times I, A \times \text{Id})$, $W_1 = S^n \times S^n \times 1$, $W_0 = S^n \times S^n \times 0$; also, let $W = V \cup TV$, $V \cap TV = M$, $V_i = V \cap W_i$ for $i = 0, 1$, where $V_0 = S^n \times D^n$, $V_1 = f^{-1}(S^n \times D^n)$.

Since $S^n \times S^n/A$ is the total space of a spherical fibre bundle $S^n \rightarrow S^n \times S^n/A \rightarrow S^n/A$, (§1). By Gysin sequence, $H_k(S^n \times S^n/A; \mathbb{Z}_2) = \mathbb{Z}_2$ for $k \neq n$, and $= \mathbb{Z}_2 + \mathbb{Z}_2$ for $k = n$. Hence $H_k(S^n \times S^n/A; \mathbb{Z}_2) = H_k(W/T; \mathbb{Z}_2) = H^{2n+1-k}(W/T, \partial W/T; \mathbb{Z}_2)$.

We want to make a characteristic submanifold M of (W, T) as highly connected as possible.

LEMMA 6.1. *There exists a connected characteristic submanifold M for (W, T) with $\partial M = M_1 - M_0$.*

PROOF. Let M be a characteristic submanifold constructed above with $\partial M = M_1 - M_0$. Then $(M, \partial M)/T$ carries the unique nonzero element of $H_{2n}(W/T, \partial W/T; Z_2) = Z_2$, dual to the 1-dimensional cohomology class F^*x , where x generates $H^1(P^{n+m}; Z_2)$ and F is the classifying map constructed above, [32]. Hence a component of M/T carries this element. Let M' be the double cover of this component. If $M' \cap M_0 = \emptyset$, then we can take $y \in H^1(W_0/T; Z_2) = H^1(W/T; Z_2) = Z_2$ representing F^*x , hence $H_{2n}(M'/T, \partial M'/T; Z_2) \rightarrow H_{2n}(W/T, \partial W/T; Z_2)$ is trivial, a contradiction. Hence $M_0 \cap M' \neq \emptyset$. Since M_0 is a closed connected manifold, $M_0 \subseteq M'$. Similarly $M_1 \subseteq M'$. It is clear that T interchanges the two components of $W - M'$, so that M' is a characteristic submanifold. We will write M for M' . Q.E.D.

LEMMA 6.2. *There exists a simply connected characteristic submanifold M for (W, T) with $\partial M = M_1 - M_0$.*

PROOF. Since $\dim W = 2n + 1 \geq 7$ and $\pi_j(W) = 0$ for $j \leq 2$, the proof is exactly the same as in Lemma 2.2 of [8] by applying [4]: We apply equivalent handle exchanges in the interior of W to make M 1-connected. Q.E.D.

From now on, we assume the characteristic submanifold M is 1-connected, with $\partial M = M_1 - M_0$. Consider the following diagram of exact sequences:

$$\begin{array}{ccccccc}
 & H_k(M_1) & \longrightarrow & H_k(V_1) \oplus H_k(TV_1) & \longrightarrow & H_k(W_1) & \longrightarrow H_{k-1}(M_1) \\
 & \downarrow & & \downarrow & & \downarrow & \\
 H_{k+1}(W) & \rightarrow & H_k(M) & \longrightarrow & H_k(V) \oplus H_k(TV) & \longrightarrow & H_k(W) \longrightarrow H_{k-1}(M) \\
 & \downarrow & & \downarrow & & \downarrow & \\
 0 & \rightarrow & H_k(M, M_1) & \longrightarrow & H_k(V, V_1) \oplus H_k(TV, TV_1) & & \\
 (6.3) & & \downarrow & & \downarrow & & \\
 & H_{k-1}(M_1) & \longrightarrow & H_{k-1}(V_1) \oplus H_{k-1}(TV_1) & & & \\
 & \downarrow & & \downarrow & & & \\
 & H_{k-1}(M) & & & & &
 \end{array}$$

We can replace M_1, V_1 by M_0, V_0 in (6.3).

LEMMA 6.4. *By performing equivariant surgery (equivariant handle exchanges) in the interior of W , we can transform M into an $(n - 2)$ -connected $2n$ -characteristic submanifold. We also have $\pi_k(M, M_1) = \pi_k(M, M_0) = \pi_k(V, V_1) = \pi_k(V, V_0) = 0$ for $k \leq n - 2$.*

PROOF. Suppose M is already $(k - 1)$ -connected, $k - 1 < n - 2$. From (6.3), we have

$$\begin{array}{ccccccc}
0 & \longrightarrow & H_k(M) & \longrightarrow & H_k(V) \oplus H_k(TV) & \longrightarrow & 0 \\
& & \downarrow & & \downarrow \approx & & \\
0 & \longrightarrow & H_k(M, M_1) & \longrightarrow & H_k(V, V_1) \oplus H_k(TV, TV_1) & \longrightarrow & 0
\end{array}$$

Using the first exact sequence, we can perform equivariant handle exchanges in the interior of W to kill $H_k(M)$ as in [8], (see [8, 2.3] for details). The other part of the lemma follows from the above diagram, the induction hypothesis, and the Hurewicz theorem. Q.E.D.

From now on, we will assume M to be $(n-2)$ -connected. Letting $k = n-1$ in (6.3), we have

$$\begin{array}{ccccccc}
Z + Z & \twoheadrightarrow & H_{n-1}(M_1) & & & & \\
\downarrow & & \downarrow & & & & \\
H_n(W) & \longrightarrow & H_{n-1}(M) & \xrightarrow{(\alpha, \alpha')} & H_{n-1}(V) \oplus H_{n-1}(TV) & & \\
& & \downarrow & & \downarrow & & \\
& & H_{n-1}(M, M_1) & \xrightarrow{(\beta, \beta')} & H_{n-1}(V, V_1) \oplus H_{n-1}(TV, TV_1) & &
\end{array}$$

where “ \twoheadrightarrow ” means 1-1, “ \longrightarrow ” onto, and “ $\xrightarrow{\sim}$ ” isomorphic; $\alpha, \alpha', \beta, \beta'$ are maps induced by inclusions.

As in [8, 2.3], we see that $H_{n-1}(M, M_1) = \text{Ker } \beta \oplus \text{Ker } \beta'$, $T_*(\text{Ker } \beta) = \text{Ker } \beta'$, both $\beta: \text{Ker } \beta' \rightarrow H_{n-1}(V, V_1)$ and $\beta': \text{Ker } \beta \rightarrow H_{n-1}(TV, TV_1)$ are isomorphisms. Similarly, $H_{n-1}(M) = \{a + b | a \in \text{Ker } \alpha, b \in \text{Ker } \alpha'\}$ with

$$\begin{aligned}
\text{Ker } \alpha \cap \text{Ker } \alpha' &= \text{Image}(H_n(W) \rightarrow H_{n-1}(M)) \\
&= \text{Image}(H_{n-1}(M_1) \rightarrow H_{n-1}(M)) = \{a\},
\end{aligned}$$

a cyclic group. Suppose $a = px$, for some integer p , where $\{x\}$ is a direct summand of $H_{n-1}(M)$. Since both $\alpha = \text{Ker } \alpha' \rightarrow H_{n-1}(V)$ and $\alpha': \text{Ker } \alpha \rightarrow H_{n-1}(TV)$ are onto with kernel $= \{a\}$, if $p \neq 0$ or 1, then x contributes two copies of Z_p in $H_{n-1}(V) \oplus H_{n-1}(TV)$, which is impossible by a simple counting argument. Hence $\text{Ker } \alpha \cap \text{Ker } \alpha' = \{a\}$ is a direct summand of $H_{n-1}(M)$. $H_{n-1}(M) = \{a\} + H$, and $H \twoheadrightarrow H_{n-1}(V) \oplus H_{n-1}(TV)$. We can perform equivariant handle exchanges in the interior of W to kill H as in [8, 2.3]. Thus we have the following:

LEMMA 6.6. *By equivariant handle exchanges in the interior of W , we can make M $(n-2)$ -connected, with $H_{n-1}(M)$ a cyclic group; and $\pi_k(M, M_1) = \pi_k(M, M_0) = 0$ for $k < n$.*

In the rest of this section, all homology will be taken with rational coefficient \mathbb{Q} , except where explicitly stated.

Letting $k = n$ in (6.3), we have

$$\begin{array}{ccccccc}
 Q & & Q + Q & & Q + Q & & Q \\
 \parallel & & \parallel & & \parallel & & \parallel \\
 H_n(M_1) & \xrightarrow{(\delta, \delta')} & H_n(V_1) \oplus H_n(TV_1) & \longrightarrow & H_n(W_1) & \longrightarrow & H_{n-1}(M_1) \\
 \downarrow i_1 & & \downarrow j_1 & & \downarrow & & \downarrow \\
 H_n(M) & \xrightarrow{(\alpha, \alpha')} & H_n(V) \oplus H_n(TV) & \longrightarrow & H_n(W) & \longrightarrow & H_{n-1}(M) \\
 (6.7) \quad \downarrow i_2 & & \downarrow j_2 & & & & \\
 H_n(M, M_1) & \xrightarrow{(\beta, \beta')} & H_n(V, V_1) \oplus H_n(TV, TV_1) & & & & \\
 \downarrow i_3 & & & & & & \\
 Q = H_{n-1}(M_1) & & & & & & \\
 \downarrow & & & & & & \\
 H_{n-1}(M) & & & & & &
 \end{array}$$

where M is the characteristic submanifold for (W, T) as in (6.6).

LEMMA 6.8. *In the diagram (6.7), i_2 maps $\text{Ker } \alpha$ into $\text{Ker } \beta$ injectively.*

PROOF. i_2 maps $\text{Ker } \alpha$ into $\text{Ker } \beta$ by the commutativity of (6.7). $H_n(M_1) \rightarrow H_n(V_1)$ is an isomorphism. Since $H_n(V_1) \rightarrow H_n(TV)$ is trivial, $H_n(V_1) \rightarrow H_n(V)$ is 1-1. But $\alpha i_1 = j_1 \delta$ from (6.7). Hence $\text{Ker } \alpha \cap \text{Im } i_1 = 0$. Q.E.D.

LEMMA 6.9. *In the diagram (6.7), if $x \in \text{Ker } \beta \cap \text{Ker } i_3$, then there exists $y \in \text{Ker } \alpha$ such that $i_2(y) = x$.*

PROOF. By exactness, there exists $w \in H_n(M)$ such that $i_2(w) = x$. Let $\alpha(w) = t$, $j_2(t) = j_2 \alpha(w) = \beta i_2(w) = 0$. Hence there is a $z \in H_n(V_1)$ such that $j_1(z) = t$. But $\delta: H_n(M_1) \rightarrow H_n(V_1)$ is an isomorphism. Define $y = w - i_1 \delta^{-1}(z)$. We have $\alpha(y) = 0$ and $i_2(y) = x$. Q.E.D.

In the diagram (6.7), $H_{n-1}(M) = Q$ or 0 by (6.6).

LEMMA 6.10. *Let $m = \text{rank } H_n(V, V_1)$ in the diagram (6.7)*

(a) *If $H_{n-1}(M) = Q$, then $\text{rank } H_n(M) = 2m + 1$, $\text{rank Ker } \alpha = m$.*

(b) *If $H_{n-1}(M) = 0$, then $\text{rank } H_n(M) = 2m$, $\text{rank Ker } \alpha = m - 1$.*

PROOF. (a) If $H_{n-1}(M) = Q$, then $H_{n-1}(M_1) \rightarrow H_{n-1}(M)$ is an isomorphism. Since $\text{rank } H_n(M, M_1) = 2 \text{ rank } \beta = 2m$, $\text{rank } H_n(M) = 2m + 1$. Hence $\text{rank Ker } \alpha = \text{rank Ker } \alpha' \leq m$. But $i_3 = 0$. $\text{rank Ker } \alpha \geq \text{rank Ker } \beta = m$ by (6.8) and (6.9). Thus $\text{rank Ker } \alpha = m$.

(b) If $H_{n-1}(M) = 0$, then i_3 is onto. Since $\text{rank } H_n(M, M_1) = 2m$ as in (a), $\text{rank } H_n(M) = 2m$. Hence $\text{rank Ker } \alpha \leq m - 1$ from (6.8). But $\text{rank Ker } \alpha \geq m - 1$ by (6.8) and (6.9). Thus $\text{rank Ker } \alpha = m - 1$. Q.E.D.

Given such a characteristic submanifold M for (W, T) as in (6.6), we can define a bilinear form C_M^f on $H_n(M)$ by $C_M^f(x, y) = x \cdot T_* y$. Let $B_M^f = C_M^f|_{\text{Ker } \alpha}$, where $(\alpha, \alpha'): H_n(M) \rightarrow H_n(V) \oplus H_n(TV)$ is the map induced by inclusion as in (6.7).

Now, we assume n odd in the rest of this section, hence C_M^f is symmetric.

LEMMA 6.11. $\text{index } C_M^f = 2 \text{ index } B_M^f$.

PROOF. There are two cases: (a) $H_{n-1}(M) = Q$. $H_n(M) = \text{Ker } \alpha \oplus \text{Ker } \alpha' \oplus \text{Im } i_1$ by (6.8) and (6.10). For $x \in H_n(M)$, let $\bar{x} \in H^n(M, \partial M)$ denote its Poincaré dual. If $x, y \in \text{Ker } \alpha$, then $\bar{x}, \bar{y} \in \text{Image}\{\alpha^*: H^n(V, \partial V) \rightarrow H^n(M, \partial M)\}$. Let $\bar{x} = \alpha^* u$, $\bar{y} = \alpha^* v$. $x \cdot y = \langle \alpha^* u \cup \alpha^* v, [M] \rangle = \langle u \cup v, \alpha_* [M] \rangle = \langle u \cup v, 0 \rangle = 0$. Similarly, $x \cdot y = 0$ for $x, y \in \text{Ker } \alpha'$. Since $\text{Im } i_1 \subseteq \text{Image}\{i: H_n(\partial M) \rightarrow H_n(M)\}$, we see that $x \cdot z = 0$ for $x \in H_n(M)$ and $z \in \text{Im } i_1$. We have $T_* \text{Ker } \alpha = \text{Ker } \alpha'$, $T_* \text{Ker } \alpha' = \text{Ker } \alpha$, and $T_* \text{Im } i_1 = \text{Im } i_1$. Hence $C_M^f(x, y) = 0$ for $x \in \text{Ker } \alpha$, $y \in \text{Ker } \alpha'$, or $x \in \text{Im } i_1$; $y \in H_n(M)$. Thus $C_M^f = C_M^f|_{\text{Ker } \alpha} + C_M^f|_{\text{Ker } \alpha'} + a$ 1-dim trivial form. But $C_M^f|_{\text{Ker } \alpha} = C_M^f|_{\text{Ker } \alpha'} = B_M^f$. Hence $\text{index } C_M^f = 2 \text{ index } B_M^f$.

(b) $H_{n-1}(M) = 0$. Let $i: H_n(\partial M) \rightarrow H_n(M)$ be induced by the inclusion. i is injective, because $H_{n+1}(M, \partial M) \cong H^{n-1}(M) \cong H_{n-1}(M) = 0$. $H_n(\partial M) = H_n(M_0) \oplus H_n(M_1) = Q + Q$. The proof of (6.8) shows that

$$\text{Im } i \cap (\text{Ker } \alpha \oplus \text{Ker } \alpha') = 0.$$

Hence $H_n(M) = \text{Ker } \alpha \oplus \text{Ker } \alpha' \oplus \text{Im } i$ by (6.10). As in the case (a), we can show that $C_M^f = C_M^f|_{\text{Ker } \alpha} + C_M^f|_{\text{Ker } \alpha'} + a$ 2-dim trivial form. Hence $\text{index } C_M^f = 2 \text{ index } B_M^f$. Q.E.D.

Let S be an involution on a manifold X^{2p} . Let B' denote the bilinear form defined on $H_p(X^{2p})$ by $B'(x, y) = x \cdot S_* y$. If B' is symmetric, define $\sigma_s(X) = \text{index } B'$. We need the following theorem from [31].

THEOREM 6.12. [31, II. 4]. Let S_1 and S_2 be involutions on X_1 and X_2 , with $\partial X_1 = \text{disjoint union } Y_1 \cup X_0$ and $\partial X_2 = \text{disjoint union } Y_2 \cup X_0$, and $S_1|_{X_0} = S_2|_{X_0}$. Let (X, S) denote $(X_1 \cup_{X_0} X_2, S_1 \cup S_2)$. Then $\sigma_s(x) = \sigma_{s_1}(X_1) + \sigma_{s_2}(X_2)$.

Now, we are ready to prove the following:

THEOREM 6.13. Let (Σ^{2n+1}, T) be a free involution on homotopy sphere Σ^{2n+1} , where n is odd and ≥ 3 , and $(\Sigma^{2n+1}, T) = (\Sigma_f, T_f)$ for an A -diffeomorphism f of $S^n \times S^n$. Then $\text{index } B_M^f = 8\sigma(\Sigma^{2n+1}, T)$.

PROOF. From the Mayer-Vietoris sequences, we see that $N = S^n \times D^n \cup_{\text{Id}} M \cup_f D^{n+1} \times S^{n-1}$ is an $(n-1)$ -connected characteristic submanifold of

$$(\Sigma_f, T_f) = (S^n \times D^{n+1}, A) \cup_{\text{Id}} (S^n \times S^n \times I, A \times \text{Id}) \cup_f (D^{n+1} \times S^n, A).$$

$$\Sigma_f = E \cup T_f E, E \cap T_f E = N, \text{ and}$$

$$H_n(N) = \text{Ker}(H_n(N) \rightarrow H_n(E)) \oplus \text{Ker}(H_n(N) \rightarrow H_n(T_f E)),$$

[8]. Let $K_n = \text{Ker}(H_n(N) \rightarrow H_n(E))$, and B = the symmetric unimodular bilinear form defined on K_n by $B(x \cdot y) = x \cdot T_{f*} y$. $\sigma(\Sigma_f, T_f)$ is defined to be $(1/8)$ index B .

Consider the symmetric bilinear forms C, C_1, C_2 , defined on

$$H_n(N), H_n(S^n \times D^n), H_n(D^{n+1} \times S^{n-1})$$

respectively by $C(x, y) = x \cdot T_{f*} y$, $C_1(x, y) = x \cdot A_* y$, $C_2(x, y) = x \cdot A_* y$. Since $H_n(D^{n+1} \times S^{n-1}) = 0$ and $H_n(S^n \times S^{n-1}) \rightarrow H_n(S^n \times D^n)$ is onto, index $C_2 = \text{index } C_1 = 0$. Hence index $C = \text{index } C_M^f$ by (6.12). But index $C_M^f = 2 \text{ index } B_M^f$ by (6.11). Similarly index $C = 2 \text{ index } B$ [8, p. 75]. Hence index $B_M^f = \text{index } B = 8\sigma(\Sigma_f, T_f) = 8\sigma(\Sigma^{2n+1}, T)$. Q.E.D.

REMARK 6.14. If the characteristic submanifold M in (6.13) satisfies (6.8)(a), i.e. $H_{n-1}(M) = Q$, then the symmetric bilinear form B_M^f defined on $\text{Ker } \alpha$ in (6.7) is actually isomorphic to the unimodular symmetric bilinear form B defined on K_n in (6.14). By the Mayer-Vietoris sequence we can show that if $j: \text{Ker } \alpha \rightarrow K_n$ is an isomorphism under the map induced by the inclusion, then we show that for $x, y \in \text{Ker } \alpha$, $x \cdot T_* y = jx \cdot jT_* y$, which follows from the fact that some multiples of x and $T_* y$ can be represented by the immersions h_1 and h_2 of manifolds $M^P, N^P \rightarrow X^{2P}$, [32], and Theorem V.1.3 of [5]: The geometric intersection number of M and N = the intersection number of the homology classes $h_{1*}[M] \cdot h_{2*}[N]$. Hence B_M^f is isomorphic to B . But given an involution (Σ_f, T_f) , we do not know whether we can always find such an M or not.

Given a free involution on a homotopy sphere (Σ^{2n+1}, T) , $n \neq 3, 7$, we can always find an A -diffeomorphism f of $S^n \times S^n$, which is homotopic to identity, such that $(\Sigma^{2n+1}, T) = (\Sigma_f, T_f)$ by (1.2). The next theorem tells us that the Browder-Livesay index invariant is additive: Given two involutions (Σ_f, T_f) and (Σ_g, T_g) with f and g homotopic to identity, we have $\sigma(\Sigma_f, T_f) + \sigma(\Sigma_g, T_g) = \sigma(\Sigma_{gf}, T_{gf})$.

THEOREM 6.15. If $f, g \in J_{2n+1} = \varnothing^{-1}(D_0(S^n \times S^n))$, n odd, then $\sigma(\Sigma_f, T_f) + \sigma(\Sigma_g, T_g) = \sigma(\Sigma_{gf}, T_{gf})$.

PROOF. Let M, N' be characteristic submanifolds for $(W, T) = (S^n \times S^n \times I, A \times \text{Id})$ associated to f, g respectively as in (6.6). M and N' are $(n-2)$ -connected, $H_{n-1}(M)$ and $H_{n-1}(N')$ are cyclic, $\partial M = f^{-1}(S^n \times S^{n-1}) \times 1 - S^n \times S^{n-1} \times 0$ and $\partial N' = g^{-1}(S^n \times S^{n-1}) \times 1 - S^n \times S^{n-1} \times 0$. Since $f^{-1} \times \text{Id}$ is an equivariant diffeomorphism for (W, T) , $N = (f^{-1} \times \text{Id})(N')$ is a characteristic submanifold of (W, T) and $\partial N = f^{-1}g^{-1}(S^n \times S^{n-1}) \times 1 - f^{-1}(S^n \times S^{n-1})$

$\times 0$. Furthermore, the bilinear form $C_{N'}$, defined on $H_n(N')$ by $C_{N'}(x, y) = x \cdot T_*y$ is isomorphic to C_N^g , hence, $\text{index } C_{N'} = \text{index } C_N^g$.

Now, we glue two copies W' and W'' of W together along W_1 of W' and W_0 of W'' by the identity. Consider $M \subseteq W'$ and $N \subseteq W''$. $P = M \cup N$ is a characteristic submanifold for $(W' \cup W'', T) = (W, T)$ with $\partial P = f^{-1}g^{-1}(S^n \times S^{n-1}) \times 1 - S^n \times S^{n-1} \times 0$, and P satisfies (6.6). P is $(n-2)$ -connected and $H_{n-1}(P)$ is cyclic by the Mayer-Vietoris sequences. Hence C_P^{gf} and B_P^{gf} are defined.

By (6.12) again, we have $\text{index } C_P^{gf} = \text{index } C_{N'} + \text{index } C_M^f$. Hence $\text{index } C_P^{gf} = \text{index } C_N^g + \text{index } C_M^f$, which implies $\text{index } B_P^{gf} = \text{index } B_N^g + \text{index } B_M^f$ by (6.11). Thus $\sigma(\Sigma_{gf}, T_{gf}) = \sigma(\Sigma_g, T_g) + \sigma(\Sigma_f, T_f)$ by (6.13). Q.E.D.

REMARK 6.16. Actually, we have showed that given an A -diffeomorphism f of $S^n \times S^n$, n odd, we can associate an index $\beta(f)$ to f which is defined to be the index of the form B_M^f above. By the standard argument as in [8, 3.2], we see that B_M^f is independent of the choice of the characteristic submanifold M . The proof of Theorem 6.15 shows that the induced map $\bar{\beta}: D(S^n \times S^n, A) \rightarrow Z$ is a homomorphism for $n \geq 3$.

7. The Arf invariant case. Theorem (6.15) is no longer valid in the case n is even, as shown by the following example.

PROPOSITION 7.1. *If n is even and > 2 , then there exist two A -diffeomorphisms f, g of $S^n \times S^n$ such that f and g are homotopic to Id , and $\sigma(\Sigma_f, T_f) = \sigma(\Sigma_g, T_g) = \sigma(\Sigma_{fg}, T_{fg}) = 0$ but $\sigma(\Sigma_{fg}, T_{fg}) = 1$.*

PROOF. Let τ be one of the generators of $\pi_n(SO_{n+1})$. We choose a representative for τ such that $\tau(x) = \tau(-x)$, and define two A -diffeomorphisms f and g by $f(x, y) = (x, \tau(x) \cdot y)$, $g(x, y) = (\tau(y) \cdot x, y)$. f and g are homotopic to Id , (4.4). It follows from (2.1) that $(\Sigma_f, T_f) = (\Sigma_g, T_g) = (\Sigma_{fg}, T_{fg}) = (S^{2n+1}, A)$. Hence their Browder-Livesay invariant is 0.

P. Orlik showed that if (Σ^{4k+1}, T) extends to an involution with fixed point on a π -manifold W^{4k+2} whose boundary is Σ^{4k+1} , then $\sigma(\Sigma^{4k+1}, T) = C(W^{4k+2})$, the Arf invariant of W^{4k+2} , [26], [21, p. 69]. We will construct such a W to show that $\sigma(\Sigma_{fg}, T_{fg}) = 1$.

Following [22], we define W to be

$$(D^{n+1} \times D^{n+1})_1 \cup_g (D^{n+1} \times D^{n+1})_2 \cup_f (D^{n+1} \times D^{n+1})_3,$$

where g is the diffeomorphism gluing $(D^{n+1} \times S^n)_1$ and $(D^{n+1} \times S^n)_2$ together, f is the gluing map from $(S^n \times D^{n+1})_2$ to $(S^n \times D^{n+1})_3$. Since f and g are A -equivariant, we define an involution T' on W by gluing the antipodal map A on each summand. The restriction of T' to $\partial W = \Sigma_{fg}$ is T . $C(W^{4k+2}) = 1$ follows from [6, V]. Q.E.D.

As in the index case, given an involution $(\Sigma^{2n+1}, T) = (\Sigma_f, T_f)$ we can find an

$(n-2)$ -connected characteristic submanifold M of $(W, T) = (S^n \times S^n \times I, A \times \text{Id})$ with $\partial M = S^n \times S^{n-1} \times 0 \cup f^{-1}(S^n \times S^{n-1}) \times 1$ and $H_{n-1}(M)$ is cyclic.

In the rest of this section, all homology will be taken with Z_2 coefficients, unless stated otherwise explicitly.

As in (6.10), we have two cases: either (a) $H_{n-1}(M) = Z_2$, or (b) $H_{n-1}(M) = 0$. In case (a), since $Z = H_{n-1}(M_i; Z) \rightarrow H_{n-1}(M; Z)$ is onto (6.7), we see that $H_{n-1}(M_1) \rightarrow H_{n-1}(M)$ is an isomorphism.

Suppose $H_{n-1}(M) = Z_2$. We see that the map i_3 in (6.7) is trivial (taken with Z_2 coefficient). We define a bilinear form B_M^f on $\text{Ker } \alpha$ in (6.7) by $B_M^f = x \cdot T_* y$. By using the Mayer-Vietoris sequences, and applying [32] and [5, V. 1.3] as in (6.14), we can show that B_M^f is isomorphic to the bilinear form B_2 defined by Browder and Livesay in [8]. In particular, B_M^f is unimodular.

Following [8], we can define a cohomology operation ψ_M^f on $\text{Ker } \alpha$, (for details, see [8, §4]). For $x, y \in \text{Ker } \alpha$, we have $B_M^f(x, y) = \psi_M^f(x + y) + \psi_M^f(x) + \psi_M^f(y)$, [8, 4.5]. Since B_M^f is unimodular, the Arf invariant for ψ_M^f is well-defined, [5], as follows. Choose a symplectic basis $x_1, \dots, x_n, y_1, \dots, y_n$ for $\text{Ker } \alpha$ such that $B_M^f(x_i, y_j) = \delta_{ij}$, $B_M^f(x_i, x_j) = B_M^f(y_i, y_j) = 0$, the Arf invariant $c_1(f, M) = \sum_{i=1}^n \psi_M^f(x_i) \psi_M^f(y_i)$.

LEMMA 7.2. *Suppose $H_{n-1}(M) = Z_2$ in (6.7), and ψ_M^f is defined as in [8]. Then any $x \in \text{Ker } \alpha$ can be represented by an immersed manifold X^n , and $\psi_M^f(x) = 1$ iff $X \cap TX$ in general position consists of an odd number of pairs of points.*

PROOF. The representability of $x \in \text{Ker } \alpha$ by an immersed manifold follows from [32], since the coefficient group is Z_2 . The other assertion can be proved by exactly the same argument in [8, 4.6]. Q.E.D.

PROPOSITION 7.3. *For n even, if the manifold M associated to (Σ_f, T_f) as in (6.7) with Z_2 -coefficient satisfies $H_{n-1}(M) = Z_2$, then $c_1(f, M) = \sigma(\Sigma_f, T_f)$.*

PROOF. We noted before that B_M^f is isomorphic to B_2 in this case. Let ψ be the quadratic form associated to B_2 defined in [8]. From (7.2), [8, 4.6], and [5, V. 1.3] again, we see that ψ_M^f and ψ are isomorphic. Hence their Arf invariants are equal. Q.E.D.

PROPOSITION 7.4. *For n even, and $f, g \in J_{2n+1}$. If there exist M, N associated to (Σ_f, T_f) , (Σ_g, T_g) as in (6.6) such that $H_{n-1}(M) = H_{n-1}(N) = Z_2$, then $\sigma(\Sigma_f, T_f) + \sigma(\Sigma_g, T_g) = \sigma(\Sigma_{gf}, T_{gf})$.*

PROOF. As in (6.15), let P denote $M \cup N$, the characteristic submanifold associated to (Σ_{gf}, T_{gf}) . We denote the domain on which B_M^f, B_N^g, B_P^{gf} is defined by $\text{Ker } \alpha$, $\text{Ker } \beta$, $\text{Ker } \gamma$ respectively. By the Mayer-Vietoris sequence, we see that $H_{n-1}(P) = Z_2$, and $\text{Ker } \gamma = \text{Ker } \alpha \oplus \text{Ker } \beta$ under the inclusion. Using (7.2) and

[5, V. 1.3] as in (7.3), we see that the quadratic form ψ_M^f is the direct sum of ψ_M^f and ψ_N^f . Hence $\sigma(\Sigma_f, T_f) + \sigma(\Sigma_{gf}, T_{gf})$ by (7.3). Q.E.D.

COROLLARY 7.5. *Let f be the A -diffeomorphism defined in (7.1). M is an $(n-2)$ -connected characteristic submanifold for $(S^n \times S^n \times I, A \times \text{Id})$ such that $\partial M = f^{-1}(S^n \times S^{n-1}) \times 1 - S^n \times S^{n-1} \times 0$. If $H_{n-1}(M)$ is cyclic, then $H_{n-1}(M; Z_2) = 0$.*

PROOF. Let g be the A -diffeomorphism defined in (7.1) by $g(x, y) = (\tau(y) \cdot x, y)$. Since $g(S^n \times S^{n-1}) = S^n \times S^{n-1}$, we can take $N = S^n \times S^{n-1} \times I$ to be a characteristic manifold associated to (Σ_g, T_g) as in (6.6). $H_{n-1}(N; Z_2) = Z_2$. Assume $H_{n-1}(M; Z_2) = Z_2$. From (7.4), we would have $\sigma(\Sigma_{fg}, T_{fg}) = \sigma(\Sigma_f, T_f) + \sigma(\Sigma_g, T_g) = 0$. This contradicts (7.1). Hence $H_{n-1}(M; Z_2) = 0$. Q.E.D.

8. Curious involutions. Let Σ_0^{4k-1} denote the generator of bP^{4k} , a cyclic subgroup of Γ^{4k-1} , consisting of those homotopy spheres which bound parallelizable manifolds, [17]. Let (Σ^{4k-1}, T) be a fixed point free involution such that $\Sigma^{4k-1} \in bP_{4k}$, we can write $\Sigma^{4k-1} = m \Sigma_0^{4k-1}$ for some integer m , which is well-defined mod 2. Following [13], we will call an involution (Σ^{4k-1}, T) curious if $m + \sigma(\Sigma^{4k-1}, T) \bmod 2$ is equal to 1.

LEMMA 8.1. *The number of curious involutions (Σ_0^{4k-1}, T) with $\sigma(\Sigma_0^{4k-1}, T) = 0$ is finite.*

PROOF. The number of the normal cobordism classes $[P^{4k-1}, G/O]$ is finite, [21]. In each normal cobordism class, there is exactly one p.l. involution with the zero Browder-Livesay's index invariant, [21] or [33]. Since $\pi_j(PL/O)$ is finite, the number of differentiable involutions with zero index invariant in each normal cobordism class is finite by smoothing theory. Q.E.D.

Let S^n denote the standard sphere. As an application of our previous theorems, we have the following:

THEOREM 8.2. *Let $n = 8k + 3$; the number of curious involutions (Σ_0^n, T) with $\sigma(\Sigma_0^n, T) = 0$ is either 0 or equal to the number of involutions (S^n, T) with $\sigma(S^n, T) = 0$.*

PROOF. From (5.2), we know that $hS(P^{8k+3})$ is in 1-1 correspondence with $J_{8k+3} = \varnothing^{-1}(D_0(S^{4k+1} \times S^{4k+1}))$, a subgroup of the group

$$D_0(S^{4k+1} \times S^{4k+1}, A)$$

of concordance classes of A -diffeomorphisms of $S^{4k+1} \times S^{4k+1}$, consisting of those A -diffeomorphisms which are homotopic to identity.

Let $C = \{f \in J_{8k+3} \mid \Sigma_f = \Sigma_0^{8k+3}, \sigma(\Sigma_f, T_f) = 0\}$, and $C' = \{f \in J_{8k+3} \mid \Sigma_f = S^{8k+3}, \sigma(\Sigma_f, T_f) = 0\}$. There are two cases:

(a) k is odd. $D_0(S^{4k+1} \times S^{4k+1}) = \Gamma^{8k+3}$ by (5.2). Let $\gamma \in \Gamma^{8k+3}$ be the element which corresponds to Σ_0^{8k+3} . If C is not empty, then $\varnothing(f) = \gamma$ for all $f \in C$, where $\varnothing: D_0(S^{4k+1} \times S^{4k+1}, A) \rightarrow D_0(S^{4k+1} \times S^{4k+1})$ is the forgetting map in §4. Take $g \in C$, $\varnothing(g^{-1}) = \gamma^{-1}$, hence $\Sigma_{g^{-1}} = -\Sigma_0^{4k+3}$; and $\sigma(\Sigma_{g^{-1}}, T_{g^{-1}}) = -\sigma(\Sigma_g, T_g) = 0$ by (6.15). Using (6.15) again, we see that the mapping $f \rightarrow f \circ g^{-1}$ for $f \in C$ maps C into C' because $\Sigma_{fg^{-1}} = S^{4k+3}$. This correspondence is 1-1 and onto, since the inverse is given by $h \rightarrow h \circ g$ for $h \in C'$.

(b) k is even. From (5.2), we know that $D_0(S^{4k+1} \times S^{4k+1})$ is the semi-direct product of $(\sigma_1) + \Gamma^{8k+3}$ and (σ_2) , where σ_1 and σ_2 are defined by $\sigma_1(x, y) = (x, \sigma(x) \cdot y)$ and $\sigma_2(x, y) = (\sigma(y) \cdot x, y)$, $\sigma_1^2 = \sigma_2^2 = \text{Id}$ in $D_0(S^{4k+1} \times S^{4k+1})$, [19]. If C is not empty, then for $f \in C$, $\varnothing(f) = \gamma, \sigma_2\gamma\sigma_1, \gamma\sigma_1$, or $\sigma_2\gamma$, [19], where γ is the element of Γ^{8k+3} corresponding to Σ_0^{8k+3} .

(i) If there exists a $g \in C$ such that $\varnothing(g) = \gamma$. Since γ^{-1} lies in the center of $D_0(S^{4k+1} \times S^{4k+1})$, [19], and $\varnothing(g^{-1}) = \gamma^{-1}$, we see that $\varnothing(gf^{-1}) = \text{Id}, \sigma_2\sigma_1, \sigma_1$, or σ_2 . Hence $\Sigma_{fg^{-1}} = S^{8k+3}$. By applying (6.15) as in (a), we see that the mapping $f \rightarrow f \circ g^{-1}$ for $f \in C$ gives a 1-1 correspondence between C and C' .

(ii) If $\varnothing(f) \neq \gamma$ for every $f \in C$, but there exists $g \in C$ such that $\varnothing(g) = \gamma\sigma_1$. Then $\varnothing(g^{-1}) = \sigma_1^{-1}\gamma^{-1} = \sigma_1\gamma^{-1}$, and $\sigma(\Sigma_{g^{-1}}, T_{g^{-1}}) = -\sigma(\Sigma_g, T_g) = 0$ by (6.15). In this case, $\varnothing(f) = \gamma\sigma_1, \sigma_2\gamma$, or $\sigma_2\gamma\sigma_1$ for $f \in C$. As in (i), we have $\varnothing(fg^{-1}) = \text{Id}, \sigma_2\sigma_1$, or σ_2 . Hence $\Sigma_{fg^{-1}} = S^{8k+3}$. By (6.15) again, the mapping $f \rightarrow f \circ g^{-1}$ for $f \in C$ gives a 1-1 correspondence between C and C' .

(iii) If $\varnothing(f) \neq \gamma, \gamma\sigma_1$, for every $f \in C$, but there exists $g \in C$ such that $\varnothing(g) = \sigma_2\gamma$. Then the mapping $f \rightarrow g^{-1}f$ gives a 1-1 correspondence between C and C' as in (i).

(iv) If $\varnothing(f) = \sigma_2\gamma\sigma_1$ for all $f \in C$. Take $g \in C$, $\varnothing(g^{-1}) = \sigma_1^{-1}\gamma^{-1}\sigma_2^{-1}$, and $\sigma(\Sigma_{g^{-1}}, T_{g^{-1}}) = 0$ by (6.15) as before. For $f \in C$, $f \rightarrow f \circ g^{-1}$ gives 1-1 correspondence between C and C' by (6.15) as before. Q.E.D.

9. **Decomposition of (Σ^{2n}, T) .** In this section, we will prove an analogue of (1.1) for free involutions on even dimensional homotopy spheres.

PROPOSITION 9.1. *For $n > 3$, $(\Sigma^{2n}, T) = (S^n \times D^n, A) \cup_g (D^{n+1} \times S^{n-1}, A)$ for some A -diffeomorphism g of $(S^n \times S^{n-1}, A)$.*

PROOF. Let P^m denote the real projective space. There is a homotopy equivalence $f: P^{2n} \rightarrow Q^{2n} = \Sigma^{2n}/T$. Let $i: P^n \rightarrow P^{2n}$ be the inclusion. For dimensional reasons, $f|P^n$ is homotopic to an embedding by [12]. By the homotopy extension theorem, we see that f homotopic to a smooth map g such that $g|P^n$ is an embedding.

Let ν_1 denote the normal bundle of P^n in P^{2n} , and ν_2 the normal bundle of gP^n in Q^{2n} . Let τ_1 and τ_2 denote the tangent bundles of P^{2n} and Q^{2n} . By

Theorem 3.6 in [1], $g^*\tau_2$ and τ_1 are J -equivalent. Since the projection $\widetilde{KO}(P^{2n}) \rightarrow J(P^{2n})$ is an isomorphism, $g^*\tau_2$ and τ_1 are stably equivalent. Let τ_p denote the tangent bundle of P^n . $g^*\tau_2|P^n$ is stably equivalent to $\tau_1|P^n$. Since $g|P^n$ is an embedding, $(g|P^n)^*(\tau_1|gP^n)$ is stably equivalent to $\tau_1|P^n$. The induced map commutes with the Whitney sum; hence $\tau_p \oplus \nu_1$ is stably equivalent to $\tau_p \oplus g^*\nu_2$. By adding a stable inverse for τ_p , we see that ν_1 is stably equivalent to $g^*\nu_2$. But ν_1 , the normal bundle of P^n in P^{2n} , is equivalent to $n\eta = \eta \oplus \eta \oplus \cdots \oplus \eta$, n times, where η is the canonical line bundle over P^n . Hence by Corollary 1.10 in [5], $g^*\nu_2$ and ν_1 are actually equivalent. By lifting this equivalence of normal bundles to the double cover, we see that there is an equivariant embedding $h = (S^n \times D^n, A) \rightarrow (\Sigma^{2n}, T)$. The image solid torus is unknotted by [12], the complement is diffeomorphic to $D^{n+1} \times S^{n-1}$ by the h -cobordism theorem.

Consider $(S^n \times D^n, A) \xrightarrow{h} (\Sigma^{2n}, T) \xleftarrow{k'} (D^{n+1} \times S^{n-1}, U)$, where we define an involution on the right-hand torus by $U = k'^{-1}Tk'$. Both h and k' are equivariant embeddings. The A -invariant diagonal sphere in $S^{n-1} \times S^{n-1} \subseteq S^n \times S^{n-1}$ on the left-hand side is mapped by $k'h|S^n \times S^n$ onto a U -invariant sphere S_{Δ}^{n-1} on the right. On the boundary of $D^{n+1} \times S^{n-1}$, U is equivalent to A . We equivariantly collar $S^n \times S^{n-1}$ in $(D^{n+1} \times S^{n-1}, U)$ by [9, 21.2], and push S_{Δ}^{n-1} a little way inside the boundary. U is equivalent to A on a tubular neighborhood N of this interior copy of S_{Δ}^{n-1} , which can be proved by applying Lemma 2 of [20] to show that the normal bundle of $P^{n-1} = S_{\Delta}^{n-1}/A$ in N is equivalent to the normal bundle of P^{n-1} in P^{2n} . The orbit space $(D^{n-1} \times S^{n-1} - \dot{N})/U$ is an h -cobordism between two copies of a manifold diffeomorphic to $S^n \times S^{n-1}/A$. Since the Whitehead group $Wh(Z_2) = 0$, this h -cobordism is diffeomorphic to $(S^n \times S^{n-1}/A) \times I$ by the s -cobordism theorem. Therefore $(D^{n+1} \times S^{n-1}, U)$ is equivalent to $(D^{n+1} \times S^{n-1}, A)$. Q.E.D.

Similar to (2.1), we have the following:

PROPOSITION 9.2. $n > 3$, $(\Sigma^{2n}, T) = (\Sigma_f, T_f) = (\Sigma_g, T_g)$ for some A -diffeomorphisms f and g of $S^n \times S^{n-1}$ iff there exists A -diffeomorphisms $H: S^n \times D^n \rightarrow S^n \times D^n$ and $K: D^{n+1} \times S^{n-1} \rightarrow D^{n+1} \times S^{n-1}$ such that $f = KgH$ on $S^n \times S^{n-1}$.

PROOF. Exactly the same as in [20]. Suppose we have two distinct decompositions, $(S^n \times D^n, A) \xrightarrow{h_i} (\Sigma^{2n}, T) \xleftarrow{k_i} (D^{n+1} \times S^{n-1}, A)$, $i = 1$ or 2 . On $S^n \times S^{n-1}$, $f = k_1^{-1}h_1$ and $g = k_2^{-1}h_2$. Note that $h_1, h_2 = S^n \times 0 \rightarrow \Sigma^{2n+1}$ are equivariantly homotopic embeddings, since both are lifted classifying maps for P^n in Q^{2n} . Hence $h_1(S^n \times 0)$ and $h_2(S^n \times 0)$ are equivariantly isotopic by a global isotopy by [12] as in the proof of (2.4). By the equivariant tubular neighborhood theorem, [3], there is an equivariant isotopy of Σ^{2n} such that, after composing with the first isotopy, there is an equivariant diffeomorphism $r: \Sigma^{2n} \rightarrow$

Σ^{2n} with $H = h_2^{-1} r h_1$, an equivariant bundle map. Let $K = k_1^{-1} r^{-1} k_2: D^{n+1} \times S^{n-1}$; we note that $k_1 = r^{-1} k_2 K^{-1}$. Hence $f = k_1^{-1} h_1 = K k_2^{-1} r h_1 = K k_2^{-1} h_2 H = K g H$. H extends equivariantly to all of $S^n \times D^n$, and H extends equivariantly to all of $D^{n+1} \times S^{n-1}$. Q.E.D.

10. Equivariant Milnor's pairing. Milnor defined in [22] a certain pairing Σ on $s_* \pi_n(SO_n) \otimes \pi_n(SO_{n+1})$ to Γ^{2n+1} , the group of homotopy spheres, where $s_* \pi_n(SO_n)$ denotes the image of $s_*: \pi_n(SO_n) \rightarrow \pi_n(SO_{n+1})$. The pairing is defined as follows. Let $a \in \pi_n(SO_n)$, $a' \in \pi_n(SO_{n+1})$; define two diffeomorphisms f_1 and f_2 on $S^n \times S^n$ by $f_1(x, y) = (x, s_* a(x) \cdot y)$ and $f_2(x, y) = (a'(y) \cdot x, y)$. Let $h = f_1 \circ f_2$, $\Sigma(s_* a, a') = S^n \times D^{n+1} \cup_h D^{n-1} \times S^n$.

If the above $s_* a$ and a' lie in the image of $\pi^*: [P^n, SO_{n+1}] \rightarrow \pi_n(SO_{n+1})$, then we can take representatives of $s_* a$ and a' such that $s_* a(x) = s_* a(-x)$ and $a'(x) = a'(-x)$. Hence $h = f_1 \circ f_2$ is an A -diffeomorphism, and $\Sigma(s_* a, a')$ admits a free involution, which is (Σ_h, T_h) . In §7, we used $a' = s_* a = \tau$ in $\pi_n(SO_{n+1})$, n even, to construct an involution on the Kervaire sphere with non-zero Arf invariant. For n odd, we will see that all the involutions obtained in this way have zero index invariant.

From (3.1) and (3.9), we know that in $\pi_n(SO_{n+1})$, $s_* \pi_n(SO_n) \cap \pi^*[P^n, SO_{n+1}] = \{2\sigma\}$ for $n > 7$, $n \equiv 3 \pmod{4}$; and $= 0$ for $n \equiv 1 \pmod{4}$. Hence we only have to consider the case where $n \equiv 3 \pmod{4}$ and $n \neq 3, 7$.

LEMMA 10.1. *If f is an A -diffeomorphism of $S^n \times S^n$ such that f leaves $S^n \times S^{n-1}$ or $S^{n-1} \times S^n$ invariant, then $\sigma(\Sigma_f, T_f) = 0$.*

PROOF. $S^n \times D^n \cup_f D^{n+1} \times S^{n-1}$ or $S^{n-1} \times D^{n+1} \cup_f D^n \times S^n$ is a codim 1 invariant sphere of (Σ_f, T_f) , because $S^n \times S^{n-1}$, $S^n \times D^n$, $D^{n+1} \times S^{n-1}$, etc. are all invariant under A . Q.E.D.

In $\pi_n(SO_{n+1})$ for $n \equiv 3 \pmod{4}$ and $n \neq 3, 7$, $\pi^*[P^n, SO_{n+1}] = Z + Z$, generated by τ and 2σ ; $\pi^*[P^n, SO_{n+1}] \cap s_* \pi_n(SO_n) = Z$, generated by 2σ , (3.9).

We first consider $\Sigma(2\sigma, 2\sigma)$. Let $\sigma = s_* a$, where $a \in \pi_n(SO_n)$, [18]. We can choose a representative for $2a$ such that $2a(x) = 2a(-x)$. Let f represent $a \in \pi_n(SO_n)$ such that $f|_{\text{the southern hemisphere}} = \text{identity}$. Since n is odd, Af is homotopic to f . The map $g: S^n \rightarrow SO_n$ defined by $g = f$ on the northern hemisphere and $= Af$ on the southern hemisphere represents $f + Af$, hence $2a$, and $g(x) = g(-x)$. We have $f_1(x, y) = (x, s_* 2a(x) \cdot y)$, $f_2(x, y) = (s_* 2a(y) \cdot x, y)$, and $h = f_1 \circ f_2$. Since $2a \in \pi_n(SO_n)$, $s_* 2a(S^n) \cdot S^{n-1} \subseteq S^{n-1}$, $f_1(S^n \times S^{n-1}) \subseteq S^n \times S^{n-1}$. But $2a(x) \in SO_n$, which has a matrix representation $[2a(x)]$. We define $d \in \pi_n(SO_n)$ by $d(x) = [2a(x)]^{-1}$, the inverse matrix for $[2a(x)]$. The diffeomorphism f_3 defined by $f_3(x, y) = (x, s_* d(x) \cdot y)$ is the inverse for f_1 , and $f_3(S^n \times S^{n-1}) \subseteq S^n \times S^{n-1}$. Hence $f_1(S^n \times S^{n-1}) = S^n \times S^{n-1}$. Similarly,

$g_2(S^n \times S^{n-1}) = S^n \times S^{n-1}$. Thus $h = f_1 \circ f_2$ leaves $S^n \times S^{n-1}$ invariant. $\sigma(\Sigma_h, T_h) = 0$ by (10.1).

Now, we consider the A -diffeomorphisms g_1 and g_2 of $S^n \times S^n$ defined by $g_2 = f_2$ in the preceding paragraph and $g_1(x, y) = (\tau(y) \cdot x, y)$, where $\tau(y) = \alpha(y)\alpha(e)$ as in §3. e is the north pole of S^n , $\alpha(y)$ = the reflection through the hyperplane orthogonal to y . This representative of τ satisfies $\tau(x) = \tau(-x)$. $\tau(S^{n-1}) \cdot S^n = \alpha(S^{n-1})\alpha(e)S^n = \alpha(S^{n-1})S^n = S^n$. Hence $g_1(S^n \times S^{n-1}) \subseteq S^n \times S^{n-1}$. But $\alpha(y)\alpha(y) = \text{identity}$. Thus $g_1(S^n \times S^{n-1}) = S^n \times S^{n-1}$. $h = g_2 \circ g_1$ leaves $S^n \times S^{n-1}$ invariant. $\sigma(\Sigma_h, T_h) = 0$ by (10.1).

Suppose β_1 and β_2 are two representatives for $\beta \in \pi_n(SO_{n+1})$ such that $\beta_i(x) = \beta_i(-x)$; we define two A -diffeomorphisms h_1 and h_2 by $h_i(x, y) = (x, \beta_i(x) \cdot y)$. Since $\pi^*[P^n, SO_{n+1}] \rightarrow \pi_n(SO_{n+1})$ is 1-1 for $n \equiv 3 \pmod 4$ and $n > 7$, (3.9), we see that h_1 and h_2 are A -concordant by (2.4) and (2.5). Hence the construction of the (Σ_h, T_h) is independent of the choice of representatives for $\beta \in \text{Image } \pi^*$. Thus we have

PROPOSITION 10.2. *Every involution (Σ^{8n+7}, T) constructed above by using Milnor's pairing: $\Sigma = \Sigma(2m\sigma, 2no)$ or $\Sigma(2m\sigma, n\tau)$, has zero Browder-Livesay index invariant.*


Added in proof. Lemma 6.11 is not true. Since T changes the orientation of the characteristic submanifold M , we have

$$\text{index } C_M^f | \text{Ker } \alpha = -\text{index } C_M^f | \text{Ker } \alpha'.$$

Thus $\text{index } C_M^f = 0$.

Here we will adapt the proof of [31, Theorem II.4] (instead of applying the Theorem itself, which was stated as Theorem 6.12 above) to verify Theorems 6.13 and 6.15.

Let M, N , and P be the characteristic submanifolds for $(W, T) = (S^n \times S^n \times I, A \times \text{id})$ associated with f, g , and gf respectively as in (6.15)



$$M_0 = S^n \times S^{n-1}, M_1 = f^{-1}(S^n \times S^{n-1}) = N_0, N_1 = f^{-1}g^{-1}(S^n \times S^{n-1}), \\ V_1 = f^{-1}(S^n \times D^n), P = M \cup_{M_1} N, V = V_M \cup_{V_1} V_N, \text{ etc.}$$

Let

$$\begin{aligned}
 (\alpha_1, \alpha'_1): H_n M &\rightarrow H_n V_M \oplus H_n TV_M, \\
 (\alpha_2, \alpha'_2): H_n N &\rightarrow H_n V_N \oplus H_n TV_N, \quad \text{and} \\
 (\alpha, \alpha'): H_n P &\rightarrow H_n V \oplus H_n TV
 \end{aligned}$$

be the maps induced by inclusion as in (6.7).

THEOREM.

$$\text{index } B_P^{g,f} = \text{index } B_N^g + \text{index } B_M^f.$$

PROOF. (6.10) and (6.11) stated that

$$H_n M = \text{Ker } \alpha_1 \oplus \text{Ker } \alpha'_1 \oplus \text{Im } i_1,$$

where $i_1: H_n \partial M \rightarrow H_n M$ is the inclusion. We take Q as coefficient from now on. Let $m = \text{rank } H_n(V, V_1)$. We can classify M into two types:

(a) $H_{n-1}M = Q$, $\text{rank } H_n M = 2m + 1$, $\text{rank Ker } \alpha_1 = m$, and $Q = H_{n-1}M_1 \rightarrow H_{n-1}M$ is onto.

(b) $H_{n-1}M = 0$, $\text{rank } H_n M = 2m$, $\text{rank Ker } \alpha_1 = m - 1$.

Consider the following exact sequence.

$$\begin{array}{ccccccc}
 H_n(M_1) & \longrightarrow & H_n M \oplus H_n N & \longrightarrow & H_n P & \longrightarrow & H_{n-1}M_1 \longrightarrow \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 Q + Q & \longrightarrow & H_n V_M \oplus H_n TV_M \oplus H_n V_N \oplus H_n TV_N & \longrightarrow & H_n V \oplus H_n TV & \longrightarrow & 0
 \end{array}$$

If one of M, N is of type (a), then we can show that $\text{Ker } \alpha_1 + \text{Ker } \alpha_2 \rightarrow \text{Ker } \alpha$ is an isomorphism by a simple counting argument, and $\text{index } B_P^{g,f} = \text{index } B_N^g + \text{index } B_M^f$ follows from the statement in (6.14).

Now we assume that both M and N are of type (b). Let $(\alpha^*, \alpha'^*): H^n V \oplus H^n TV \rightarrow H^n P$ be the map induced by inclusions. We have the following exact sequence.

$$\begin{array}{ccccc}
 & & k & & \\
 & \swarrow & & \searrow & \\
 H_n M_1 & \longrightarrow & \text{Ker } \alpha_1 \oplus \text{Ker } \alpha_2 & \xrightarrow{j_1 + j_2} & \text{Ker } \alpha \longrightarrow H_{n-1}M_1 \\
 & & \uparrow \cong & & \uparrow \cong \\
 & & \text{Im } \alpha'^* & \xrightarrow{k^*} & H^n M_1
 \end{array}$$

Let $A_B \text{Im } k$ denote the annihilator of $\text{Im } k$ under $B_P^{g,f}$. Then we may use the argument in [31, II.4] to show that $A_B \text{Im } k = \text{Im } j_1 + \text{Im } j_2$. But we also have

LEMMA [31, II.3]. *If B is a symmetric bilinear form on a vector space V , and if there is a subspace $C \subseteq V$ with $C \subseteq AC$, then $\text{index } B = \text{index } B|_C$.*

Thus our theorem follows from this Lemma and (6.14) as above. Q.E.D.
The proof of Theorem 6.13 is similar.

BIBLIOGRAPHY

1. J. F. Adams, *On the groups $J(X)$* . II, *Topology* 3 (1965), 137–171. MR 33 #6626.
2. I. Bernstein, *Involutions with nonzero Arf invariant*, *Bull. Amer. Math. Soc.* 74 (1968), 678–682. MR 38 #5225.
3. G. E. Bredon, *Introduction to compact transformation groups*, Academic Press, New York, 1972.
4. W. Browder, *Structures on $M \times R$* , *Proc. Cambridge Philos. Soc.* 61 (1965), 337–345. MR 30 #5321.
5. ———, *Surgery on simply-connected manifolds*, *Ergebnisse der Mathematik und ihrer Grenzgebiete*, Band 65, Springer-Verlag, Berlin and New York, 1972.
6. ———, *Cobordism invariants, the Kervaire invariant and fixed point free involutions*, *Trans. Amer. Math. Soc.* 178 (1973), 193–225. MR 48 #3067.
7. W. Browder and G. R. Livesay, *Fixed point free involutions on homotopy spheres*, *Bull. Amer. Math. Soc.* 73 (1967), 242–245. MR 34 #6781.
8. ———, *Fixed point free involutions on homotopy spheres*, *Tôhoku Math. J.* (2) 25 (1973), 69–87. MR 47 #9610.
9. P. E. Conner and E. E. Floyd, *Differentiable periodic maps*, *Ergebnisse der Mathematik und ihrer Grenzgebiete*, N. F., Band 33, Springer-Verlag, Berlin; Academic Press, New York, 1964. MR 31 #750.
10. A. Dold, *Über faserweise homotopieäquivalenz von Faserräumen*, *Math. Z.* 62 (1955), 111–136. MR 17, 519.
11. M. Fujii, *KO-groups of projective spaces*, *Osaka J. Math.* 4 (1967), 141–149. MR 36 #2143.
12. A. Haefliger, *Plongements différentiable de variétés dans variétés*, *Comment. Math. Helv.* 36 (1961), 47–82. MR 26 #3069.
13. M. W. Hirsch and J. Milnor, *Some curious involutions of spheres*, *Bull. Amer. Math. Soc.* 70 (1964), 372–377. MR 31 #751.
14. D. Husemoller, *Fibre bundles*, McGraw-Hill, New York, 1966. MR 37 #4821.
15. I. James and E. Thomas, *An approach to the enumeration problem for non-stable vector bundles*, *J. Math. Mech.* 14 (1965), 485–506. MR 30 #5319.
16. M. A. Kervaire, *Some nonstable homotopy groups of Lie groups*, *Illinois J. Math.* 4 (1960), 161–169. MR 22 #4075.
17. M. A. Kervaire and J. W. Milnor, *Groups of homotopy spheres*. I, *Ann. of Math.* (2) 77 (1963), 504–537. MR 26 #5584.
18. A. Kosinski, *On the inertia group of π -manifolds*, *Amer. J. Math.* 89 (1967), 227–248. MR 35 #4936.
19. J. Levine, *Self-equivalences of $S^n \times S^k$* , *Trans. Amer. Math. Soc.* 143 (1969), 523–543. MR 40 #2098.
20. G. R. Livesay and C. B. Thomas, *Free Z_2 and Z_3 actions on homotopy spheres*, *Topology* 7 (1968), 11–14. MR 36 #3343.
21. S. López de Medrano, *Involutions on manifolds*, *Ergebnisse der Mathematik und ihrer Grenzgebiete*, Band 59, Springer-Verlag, New York, 1971. MR 45 #7747.
22. J. W. Milnor, *Differentiable structures on spheres*, *Amer. J. Math.* 81 (1959), 962–972. MR 22 #990.
23. R. E. Mosher and M. C. Tangora, *Cohomology operations and application in homotopy theory*, Harper & Row, New York and London, 1968. MR 37 #2223.
24. P. Olum, *Mappings of manifolds and the notion of degree*, *Ann. of Math.* (2) 58 (1953), 458–480. MR 15, 338.
25. ———, *Cocycle formulas for homotopy classification; maps into projective and lens space*, *Trans. Amer. Math. Soc.* 103 (1962), 30–44. MR 25 #576.

- 26. P. Orlik, *On the Arf invariant of an involution*, Canad. J. Math. 22 (1970), 519–524. MR 41 #7718.
- 27. D. Puppe, *Homotopiemengen und ihre induzierten Abbildungen. I*, Math. Z. 69 (1958), 299–344. MR 20 #6698.
- 28. M. Rothenberg and J. Sondow, *Non-linear smooth representatives of compact lie groups* (preprint).
- 29. H. Sato, *Diffeomorphism groups of $S^p \times S^q$ and exotic spheres*, Quart. J. Math. Oxford Ser. (2) 20 (1969), 255–276. MR 40 #6584.
- 30. N. E. Steenrod, *The topology of fibre bundles*, Princeton Math. Ser., vol. 14, Princeton Univ. Press, Princeton, N. J., 1951. MR 12, 522.
- 31. E. Stone, Ph. D. Thesis, Cornell University, 1972 (to appear).
- 32. R. Thom, *Quelques propriétés globales des variétés différentiables*, Comment. Math. Helv. 28 (1954), 17–86. MR 15, 890.
- 33. C. T. C. Wall, *Free piecewise linear involutions on spheres*, Bull. Amer. Math. Soc. 74 (1968), 554–558. MR 36 #5955.
- 34. ———, *Surgery on compact manifolds*, Academic Press, New York, 1972.

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA, NEW YORK
14853

Current address: Department of Mathematics, University of Kansas, Lawrence, Kansas
66045