

ASYMMETRIC MAXIMAL IDEALS IN $M(G)$

BY

SADAHIRO SAEKI

ABSTRACT. Let G be a nondiscrete LCA group, $M(G)$ the measure algebra of G , and $M_0(G)$ the closed ideal of those measures in $M(G)$ whose Fourier transforms vanish at infinity. Let Δ_G , Σ_G and Δ_0 be the spectrum of $M(G)$, the set of all symmetric elements of Δ_G , and the spectrum of $M_0(G)$, respectively. In this paper this is shown: Let Φ be a separable subset of $M(G)$. Then there exist a probability measure τ in $M_0(G)$ and a compact subset X of $\Delta_0 \setminus \Sigma_G$ such that for each $|c| \leq 1$ and each

$$\nu \in \Phi \text{ Card } \{f \in X: \hat{\tau}(f) = c \text{ and } |\hat{\nu}(f)| = r(\nu)\} \geq 2^c.$$

Here $r(\nu) = \sup \{|\hat{\nu}(f)|: f \in \Delta_G \setminus \hat{G}\}$. As immediate consequences of this result, we have (a) every boundary for $M_0(G)$ is a boundary for $M(G)$ (a result due to Brown and Moran), (b) $\Delta_G \setminus \Sigma_G$ is dense in $\Delta_G \setminus \hat{G}$, (c) the set of all peak points for $M(G)$ is \hat{G} if G is σ -compact and is empty otherwise, and (d) for each $\mu \in M(G)$ the set $\hat{\mu}(\Delta_0 \setminus \Sigma_G)$ contains the topological boundary of $\hat{\mu}(\Delta_G \setminus \hat{G})$ in the complex plane.

Throughout the paper, let G be a nondiscrete locally compact abelian group with dual \hat{G} , $M(G)$ the convolution measure algebra of G , and $M_0(G)$ the ideal in $M(G)$ which consists of all measures with Fourier transforms vanishing at infinity. As is well known, we then have $L^1(G) = M_a(G) \subset M_0(G) \subset M_c(G)$. Let Δ_G denote the spectrum of $M(G)$, i.e., the space of all nonzero complex homomorphisms of $M(G)$, and let $\hat{\mu}$ denote the Gelfand transform of $\mu \in M(G)$. We define

$$\Delta_0 = \{f \in \Delta_G: \hat{\sigma}(f) \neq 0 \text{ for some } \sigma \in M_0(G)\},$$

$$\Sigma_G = \{f \in \Delta_G: f(\sigma^*) = \overline{f(\sigma)} \text{ for all } \sigma \in M(G)\},$$

where $\sigma^*(E) = \sigma(-E)$ for all Borel sets E in G and $f(\sigma) = \hat{\sigma}(f)$. Then Δ_0 is open (in Δ_G), Σ_G is closed, and $\hat{G} \subset \Delta_0 \cap \Sigma_G$. Moreover, Δ_0 may be identified with the spectrum of $M_0(G)$, since $M_0(G)$ is an ideal in $M(G)$.

It is shown in [11] that given $\mu \in M_c(G)$, there exist fairly many elements $f \in \Delta_G$ such that $M_a(G) + L^1(\mu) \subset \text{Ker}(f)$ but $M_0(G) \not\subset \text{Ker}(f)$. In fact, it is not difficult to improve Theorem 2 of [11] as follows.

THEOREM A. *Let $0 \neq \lambda \in M_0(G)$, $\mu \in M_c(G)$, and H a subgroup of G which is a G_δ -set. Then there exists a probability measure $\sigma = \tau * \tau^*$, with $\tau \in$*

Received by the editors July 11, 1975.

AMS (MOS) subject classifications (1970). Primary 43A10.

Key words and phrases. LCA group, measure algebra, asymmetric maximal ideal, M_0 -boundary.

$M_0^+(\text{supp } \lambda)$, having the following properties:

(i) Given $0 \leq r \leq 1$, the set of all $f \in \Sigma_G$ such that

$$\hat{o}(f) = r, \quad \text{Ker}(f) \supset L^1(\mu), \quad \text{and} \quad \hat{v}(f) = \hat{v}(1) \quad \forall v \in M_d(G)$$

has cardinality $\geq 2^c$. Here c denotes the cardinal number of the continuum.

(ii) Given a complex number c of modulus ≤ 1 and $g \in \Delta_G$ with $g(\delta_x) = 1$ for all $x \in H$, the set of all $f \in \Delta_G \setminus \Sigma_G$ such that

$$\hat{o}(f) = c, \quad \text{Ker}(f) \supset L^1(\mu), \quad \text{and} \quad \hat{v}(f) = \hat{v}(g) \quad \forall v \in M_d(G)$$

has cardinality $\geq 2^c$.

For some related results, we refer the reader to Izuchi and Shimizu [8], Saka [12], Shimizu [13], and Williamson [15]. Now let $\mu \in M(G)$ be given, and define

$$r(\mu) = \sup \{ |\hat{\mu}(f)| : f \in \Delta_G \setminus \hat{G} \}.$$

Since $\Delta_G \setminus \hat{G}$ is compact, there exists at least one f in this set such that $|\hat{\mu}(f)| = r(\mu)$. It seems to be a natural problem to ask how many f as above there exist. Our answer is as follows.

THEOREM B. Let $\mu \in M(G)$, and Φ a separable subset of $L^1(\mu)$. Then there exist a probability measure τ in $M_0(G)$ and a compact set X in $\Delta_0 \setminus \Sigma_G$ such that

$$\text{Card} \{ f \in X : \hat{\tau}(f) = c \text{ and } |\hat{v}(f)| = r(v) \} \geq 2^c$$

for every complex number c of modulus ≤ 1 and every measure v in $[L^1(\mu) \cap M^+(G)] \cup \Phi$.

Notice that we can set $\Phi = L^1(\mu)$ if G is metrizable, since then $L^1(\mu)$ is separable. As easy consequences of the last theorem, we have the following results.

COROLLARY 1.

- (a) Every boundary of $M_0(G)$ is a boundary of $M(G)$.
- (b) The set $\Sigma_G \setminus \hat{G}$ is the topological boundary of $\Delta_G \setminus \Sigma_G$ in Δ_G . In other words, $\Delta_G \setminus \Sigma_G$ is dense in $\Delta_G \setminus \hat{G}$.
- (c) If G is σ -compact, then the set P_G of all peak points for $M(G)$ is precisely \hat{G} . If not, then $P_G = \emptyset$.

COROLLARY 2. For each $\mu \in M(G)$, the set $\hat{\mu}(\Delta_0 \setminus \Sigma_G)$ contains the topological boundary of $\hat{\mu}(\Delta_G \setminus \hat{G})$ in the complex plane \mathbb{C} . In particular, we have

- (a) If $\text{Card} [\hat{\mu}(\Delta_0 \setminus \Sigma_G)] < c$, then $\hat{\mu}(\Delta_G \setminus \hat{G})$ is (at most) countable and coincides with $\hat{\mu}(\Delta_0 \setminus \Sigma_G)$.
- (b) If $\hat{\mu}$ is real on $\Delta_0 \setminus \Sigma_G$, then $\hat{\mu}(\Delta_G \setminus \hat{G}) = \hat{\mu}(\Delta_0 \setminus \Sigma_G)$.

Notice that Theorem B implies the result of Brown and Moran [2] and Graham [5]: If $\mu \in M(G)$ and $\hat{\mu} = 0$ off Σ_G , then $r(\mu) = 0$. Part (a) of Corollary 1 is due to Brown and Moran [3]. We also refer to Brown's result in [1]: $\Delta_0 \cap \Sigma_G$ is not entirely contained in the Shilov boundary of $M_0(G)$. It may be an interesting problem to ask whether or not we have $\hat{\mu}(\Delta_0 \setminus \Sigma_G) = \hat{\mu}(\Delta_G \setminus \hat{G})$ for all $\mu \in M(G)$.

To prove Theorem B, we shall first construct a measure of a certain type (assuming that G is metrizable). The construction of such a measure is almost the same as the corresponding one in [11], and Körner's method [9] plays an important role in our construction.

We now introduce some notation. Let m_G denote the Haar measure of G , and \mathbf{Z} the group of all integers. For a set K in G and $p \in \mathbf{Z}^+$, we define

$$pK = \{x_1 + \cdots + x_p : x_j \in K \text{ for all } 1 \leq j \leq p\}$$

if $p \geq 1$, $pK = \{0\}$ if $p = 0$, and $(-p)K = -(pK)$. The subgroup of G which the set K generates is denoted by $Gp(K)$. We say that a Borel set K is of type M_0 if $M_0(K) = M_0(G) \cap M(K)$ is nonzero. Let $q(G)$ denote the supremum of all natural numbers p such that every neighborhood of the identity 0 of G contains an element of order $\geq p$. Then it is easy to see that if $q(G) = \infty$, then G is an I -group, and that if $q(G)$ is finite, then G contains an open-and-compact subgroup H such that $\text{ord}(x) \leq q(G)$ for all x in H . A set K in G is called *strongly independent* if it is independent in the usual sense [10, p. 97] and if all of its elements have order $q(G)$. Finally, we denote by $Gp'(K)$ the set of all points x of the form $x = k_1x_1 + \cdots + k_ux_u$, where $u = u_x$ is a natural number, x_1, \dots, x_u are distinct elements of K , $k_j \in \mathbf{Z}$ for all $1 \leq j \leq u$, and $|k_j| = 1$ for at least one index j .

LEMMA 1. Let $\mu_0 \in M^+(G)$, D a compact subset of G with Haar measure zero, and N a natural number. Let also V_1, V_2, \dots, V_u be nonempty open sets in G . Then we can find nonempty open sets $U_j \subset V_j$ ($1 \leq j \leq u$) subject to the following conditions:

(i) If $p_j \in \mathbf{Z}$, $|p_j| < q(G)$, and $1 \leq \sum_{j=1}^u |p_j| \leq N$, then the set $\sum_{j=1}^u p_j U_j$ does not contain 0 in G , and

$$m_G \left[D + \sum_{j=1}^u p_j U_j \right] < 1/N.$$

(ii) If $q_j \in \mathbf{Z}$, $\sum_{j=1}^u |q_j| \leq N$, and $|q_j| = 1$ for at least one index j , then

$$\mu_0 \left[D + \sum_{j=1}^u q_j U_j \right] < 1/N.$$

PROOF. Let P be the set of all $p = (p_1, \dots, p_u) \in Z^u$ as in (i). Similarly, let Q be the set of all $q = (q_1, \dots, q_u) \in Z^u$ as in (ii).

The standard Baire category argument [10, 5.2.3] shows that there are points $x_j \in V_j$ ($1 \leq j \leq u$) of order $\geq q(G)$ such that $\{x_j: 1 \leq j \leq u\}$ is independent. Since P is finite and D is a compact set with Haar measure zero, we can find a neighborhood W of $0 \in G$ so that

$$(1) \quad 0 \notin \sum_1^u p_j(x_j + W) \quad \text{and} \quad m_G \left[D + \sum_1^u p_j(x_j + W) \right] < 1/N$$

for all $p \in P$. We may assume that $x_j + W \subset V_j$ ($1 \leq j \leq u$).

Put $E = \{x_j: 1 \leq j \leq u\}$, and take a compact neighborhood X of $0 \in G$ such that $X + X \subset W$. Since $M_a(G)$ is an ideal of $M(G)$, it follows from the Fubini theorem and the definition of Q that

$$(2) \quad \int_{X^u} \sum_{q \in Q} \mu_0 \left[D + Gp(E) + \sum_1^u q_j t_j \right] dt_1 \cdots dt_u = 0.$$

Therefore there are u points t_1, t_2, \dots, t_u in X for which the integrand in (2) is zero. Hence, in particular, we have

$$(3) \quad \mu_0 \left[D + \sum_1^u q_j y_j \right] = 0 \quad (q \in Q),$$

where $y_j = x_j + t_j$. Upon comparing (1) and (3), we see that if $U \subset X$ is a sufficiently small neighborhood of $0 \in G$, then the sets $U_j = y_j + U$ have the required properties.

LEMMA 2. Suppose that G is metrizable. Let $\mu_0 \in M^+(G)$, and let C_0 be a σ -compact subset of G with Haar measure zero. Then there exists a strongly independent compact set K in G of type M_0 such that

$$m_G[C_0 + Gp(K)] = \mu_0[C_0 + Gp'(K)] = 0.$$

PROOF. If $q(G)$ is finite, we fix an open-and-compact subgroup H of G such that $\text{ord}(x) \leq q(G)$ for all x in H . In the other case, we set $H = G$.

Let $\{D_n\}_1^\infty$ be an increasing sequence of compact subsets of G with $C_0 = \bigcup_1^\infty D_n$, and $\{\hat{E}_n\}_1^\infty$ a sequence of compact subsets of \hat{G} with $\hat{G} = \bigcup_1^\infty \hat{E}_n$. We shall construct a sequence $\{\sigma_n\}_1^\infty$ of probability measures in $M_0(H)$, a sequence $\{I_n\}_1^\infty$ of finite collections of disjoint compact sets in H , a sequence $\{\hat{F}_n\}_1^\infty$ of compact sets in \hat{G} , and also a sequence $\{n_p\}_1^\infty$ of natural numbers. They will satisfy the following conditions (and some other conditions):

$$(1) \quad \text{supp } \sigma_n \subset \bigcup \{\text{int}(I): I \in I_n\}.$$

$$(2) \quad \sup \{|\widehat{\sigma_n}|_I(\chi)|: \chi \in \hat{G} \setminus \hat{F}_n\} < 2^{-n} \sigma_n(I) \quad \forall I \in \mathcal{I}_n.$$

It is also assumed that each set in \mathcal{I}_{n+1} is a subset of some set in \mathcal{I}_n .

We first take any probability measure $\sigma_1 \in M_0(H)$ with compact support of diameter $< 1/2$. Let I be any compact neighborhood of $\text{supp } \sigma_1$ such that $\text{diam } I < 1$, $\mathcal{I}_1 = \{I\}$, and $n_1 = 1$. Since $\sigma_1 \in M_0(G)$, we can take a compact set \hat{F}_1 in \hat{G} subject to (2) with $n = 1$.

Suppose that p is a natural number, and that n_j ($1 \leq j \leq p$), σ_n , \mathcal{I}_n , \hat{F}_n ($1 \leq n \leq m = n_p$) have been defined. Let M_p be the largest natural number such that

$$(3) \quad \max \{\sigma_m(I): I \in \mathcal{I}_m\} \leq M_p^{-2},$$

and write

$$(4) \quad \{A \subset \mathcal{I}_m: 1 \leq \text{Card } A \leq M_p\} = \{A_r: 1 \leq r \leq s_p\}.$$

Setting $n_{p+1} = n_p + s_p$, we shall construct σ_n , \mathcal{I}_n , and \hat{F}_n for all $m < n \leq n_{p+1}$ as follows.

Suppose that these objects have been defined for some $n = m + r - 1$ with $1 \leq r \leq s_p$, and put

$$(5) \quad K_n = \{I \in \mathcal{I}_n: I \subset J \text{ for some } J \in A_r\}.$$

Then, for each set K in K_n , there are a finite collection $\{L_j^K\}_j$ of disjoint compact sets in K and a collection $\{\nu_j^K\}_j$ of measures in $M_0^+(K)$, with $\text{supp } \nu_j^K \subset \text{int}(L_j^K)$, such that

$$(6) \quad 0 < \|\nu_j^K\| < n^{-1} \sigma_n(K);$$

$$(7) \quad \sum_j \|\nu_j^K\| = \sigma_n(K);$$

$$(8) \quad \left| \sum_j (\nu_j^K)^\wedge(\chi) - (\sigma_n|_K)^\wedge(\chi) \right| < 2^{-n} \sigma_n(K) \quad \forall \chi \in \hat{F}_n.$$

To see this, it suffices to apply Lemma 3 of [11] and its obvious modification. By virtue of Lemma 1, we can demand that the sets L_j^K satisfy the following additional conditions:

$$(9) \quad \text{diam } L_j^K < 1/n;$$

$$(10) \quad 0 \notin \sum_{K \in K_n} \sum_j p_j^K L_j^K \quad \forall (p_j^K) \in P_n;$$

$$(11) \quad m_G \left[D_n + \sum_{K \in K_n} \sum_j p_j^K L_j^K \right] < 2^{-n} / \text{Card } P_n \quad \forall (p_j^K) \in P_n;$$

$$(12) \quad \mu_0 \left[D_n + \sum_{K \in K_n} \sum_j q_j^K L_j^K \right] < 2^{-n} / \text{Card } Q_n \quad \forall (q_j^K) \in Q_n.$$

Here P_n is the set of all tuples (p_j^K) of integers such that $|p_j^K| < q(G)$ for all j and K and $1 \leq \sum_{K,j} |p_j^K| \leq n$. Similarly Q_n is the set of all tuples (q_j^K) of integers such that $|q_j^K| = 1$ for some (K, j) and $\sum_{K,j} |q_j^K| \leq n$. Define

$$(13) \quad \sigma_{n+1} = \sum_{I \in K_n} \sigma_n |I| + \sum_{K \in K_n} \sum_j v_j^K,$$

$$(14) \quad I_{n+1} = (I_n \setminus K_n) \cup \bigcup_{K \in K_n} \{L_j^K\}_j.$$

Then (1), with n replaced by $n + 1$, is satisfied. Finally we choose a compact set \hat{F}_{n+1} in \hat{G} , with $\hat{F}_{n+1} \supset \hat{F}_n \cup \hat{E}_n$, so that (2) holds for $n + 1$.

This completes our induction. It is a routine matter to prove that the sequence $\{\sigma_n\}_1^\infty$ converges to some probability measure $\sigma \in M_0(H)$ in the weak-* topology of $M(G)$, that

$$(15) \quad K = \text{supp } \sigma \subset \bigcap_{n=1}^\infty \bigcup \{I : I \in I_n\},$$

and that K is strongly independent. (See the proof of Lemma 4 of [11], and notice that every element of H has order $\leq q(G)$.)

Now we want to confirm

$$m_G [C_0 + Gp(K)] = \mu_0 [C_0 + Gp'(K)] = 0.$$

Let $0 \neq x \in Gp(K)$ be given. We have $x = \sum_1^u k_i x_i$ for some $(k_1, \dots, k_u) \in \mathbb{Z}^u$ and some distinct elements x_1, \dots, x_u of K . By (9), (14), and (15), there exists a natural number N_x such that the points x_i belong to distinct sets in I_n whenever $n > N_x$. Choose any natural number p so that

$$n_p > N_x + \sum_1^u |k_i| \quad \text{and} \quad M_p > u,$$

and let A be the collection of all I in I_{n_p} which contain some x_i ($1 \leq i \leq u$). Then $1 \leq \text{Card } A = u < M_p$, and so $A = A_r$ for some $1 \leq r \leq s_p$ by (4). Setting $n = n_p + r - 1$, we therefore infer from (5), (14) and (15) that x belongs to the set

$$\bigcup_{P_n} \left(\sum_{K \in K_n} \sum_j p_j^K L_j^K \right).$$

Since p can be chosen as large as one pleases, we conclude that

$$(16) \quad Gp(K) \setminus \{0\} \subset \bigcup_{n=N}^{\infty} \bigcup_{P_n} \sum_{K_n} \sum_j p_j^K L_j^K \quad (N = 1, 2, \dots).$$

Similarly we have

$$(17) \quad Gp'(K) \subset \bigcup_{n=N}^{\infty} \bigcup_{Q_n} \sum_{K_n} \sum_j q_j^K L_j^K \quad (N = 1, 2, \dots).$$

It follows from (11) and (16) that

$$(18) \quad \begin{aligned} m_G[D_N + Gp(K)] &\leq \sum_{n=N}^{\infty} \sum_{P_n} m_G \left[D_N + \sum_{K_n} \sum_j p_j^K L_j^K \right] \\ &\leq \sum_{n=N}^{\infty} \sum_{P_n} m_G \left[D_n + \sum_{K_n} \sum_j p_j^K L_j^K \right] < 2^{-N+1} \end{aligned}$$

for all $N \geq 1$. (Notice that $m_G(D_N) = 0$.) Letting $N \rightarrow \infty$ in (18), we have $m_G[C_0 + Gp(K)] = 0$. Similarly we have $\mu_0[C_0 + Gp'(K)] = 0$ by (12) and (17). This completes the proof.

LEMMA 3. Let $\mu_0 \in M^+(G)$, C_0 a σ -compact subset of G which carries μ_0 , and K a compact subset of G such that

$$(*) \quad \mu_0[C_0 + Gp'(K)] = 0.$$

Suppose that K_1, K_2, \dots, K_p are disjoint compact subsets of K and that $\sigma_j \in M_c(K_j \cup (-K_j))$ for all $1 \leq j \leq p$.

(a) If $m = (m_j)_1^p$ and $n = (n_j)_1^p$ are different tuples of nonnegative integers, then

$$\mu_0 * \sigma_1^{m_1} * \dots * \sigma_p^{m_p} \perp \mu_0 * \sigma_1^{n_1} * \dots * \sigma_p^{n_p}.$$

(b) If $\sigma_j \in M_c(K_j)$ for all $1 \leq j \leq p$ and $\nu \in L^1(\mu_0)$, then

$$\|\nu * \sigma_1^{n_1} * \dots * \sigma_p^{n_p}\| = \|\nu\| \cdot \|\sigma_1\|^{n_1} \dots \|\sigma_p\|^{n_p}.$$

PROOF. To prove (a), we use the well-known method of Hewitt and Kakutani [6] (see also [10, 5.4.2]). Without loss of generality, assume that $\sigma_j \geq 0$ for all $1 \leq j \leq p$ and that $m_1 < n_1$. Write $\tau_m = \sigma_1^{m_1} * \dots * \sigma_p^{m_p}$, and similarly for τ_n . Putting $E_j = K_j \cup (-K_j)$ for $1 \leq j \leq p$, we then see that $\mu_0 * \tau_m$ is carried by the set $A_m = C_0 + m_1 E_1 + \dots + m_p E_p$. Therefore it suffices to show $(\mu_0 * \tau_n)(A_m) = 0$. Let $\lambda_j \in M(E_j^{n_j})$ be the n_j -fold product of σ_j , and let B_j be the set of all points $x_j = (x_{j1}, \dots, x_{jn_j})$ of $E_j^{n_j}$ such that $x_{ji} \neq \pm x_{jk}$ whenever $1 \leq i < k \leq n_j$. Since σ_j is a continuous measure, we then have

$\lambda_j(G^n \setminus B_j) = 0$ by the Fubini theorem. On the other hand, $(x_{ji}) \in B_1 \times \cdots \times B_p$ implies

$$(1) \quad \mu_0 \left[A_m - \sum_{j=1}^p \sum_{i=1}^{n_i} x_{ji} \right] \leq \mu_0 \left[C_0 + m_1 E_1 - \sum_{i=1}^{n_1} x_{1i} + \sum_{j=2}^p Gp(K_j) \right] \\ \leq \mu_0 [C_0 + Gp'(K)] = 0$$

by (*) and the definition of $Gp'(K)$. Evidently these two facts imply $(\mu_0 * \tau_n)(A_m) = 0$, as was required.

To prove (b), we need the following fact: Given $\mu \in M(G)$ and $\epsilon > 0$, there is a neighborhood V of $0 \in G$ such that

$$(2) \quad \sigma \in M^+(G), \quad \text{supp } \sigma - \text{supp } \sigma \subset V \Rightarrow \|\mu * \sigma\| \geq (\|\mu\| - \epsilon)\|\sigma\|.$$

Suppose by way of contradiction that this is false for some μ and ϵ . Then, to each neighborhood V of 0 there corresponds a probability measure $\sigma_V \in M(G)$ such that $\|\mu * \sigma_V\| < \|\mu\| - \epsilon$ and $\text{supp } \sigma_V \subset V - x_V$ for some $x_V \in G$. Upon replacing σ_V by $\sigma_V * \delta_{x_V}$, we may assume that $x_V = 0$. But then the net $\{\mu * \sigma_V\}$ converges to μ in the weak-* topology of $M(G)$. Hence

$$\|\mu\| \leq \liminf_V \|\mu * \sigma_V\| \leq \|\mu\| - \epsilon,$$

a contradiction.

We now prove (b) as follows. By the continuity of convolution, we can retain generality in assuming that each σ_j has the form $\sigma_j = \sum_{k=1}^q c_{jk} \tau_{jk}$, where the c_{jk} are complex numbers of absolute modulus one and the τ_{jk} are mutually singular measures in $M_c^+(K_j)$. Expanding $\sigma_j^n = (\sum_{k=1}^q c_{jk} \tau_{jk})^n$ for all $1 \leq j \leq p$ and applying part (a), we reduce (b) to the case where $\sigma_j \geq 0$ ($1 \leq j \leq p$), and hence to the case where $c_{jk} = 1$ for all j and k . Since we can demand that every τ_{jk} has support of sufficiently small diameter, part (b) follows from (2). This completes the proof.

PROOF OF THEOREM B. Let $\mu \in M(G)$, and Φ a separable subset of $L^1(\mu)$. Given $\sigma \in M(G)$, we let σ_s denote the singular part of σ with respect to m_G . Notice that

$$(*) \quad r(\sigma) = \lim_{n \rightarrow \infty} \|\sigma^n + M_a(G)\|^{1/n} = \lim_{n \rightarrow \infty} \|(\sigma^n)_s\|^{1/n},$$

since $M_a(G)$ is an ideal in $M(G)$ with spectrum \hat{G} . Now define μ_0 to be the singular part of $\exp(|\mu|)$, and choose a σ -compact subset C_0 of G so that $m_G(C_0) = \mu_0(G \setminus C_0) = 0$. Then $\nu \in L^1(\mu)$ implies $(\nu^n)_s \in L^1(\mu_0)$ for all $n \in \mathbb{Z}^+$.

We first assume that G is metrizable, and take a compact subset K of G as

where $H = H_\Gamma$ is the annihilator of Γ in G and m_H denotes the Haar measure of H of norm one. This can be proved by considering the Fourier transform of ν and by applying Theorem 1.9.1 of [10]. Since $\Phi \subset L^1(\mu)$ is separable, there is a σ -compact open subgroup Γ of G such that

$$(9) \quad \|(\nu^n)_s\| = \|(\nu^n)_s * m_H\| \quad \forall \nu \in \Phi \text{ and } \forall n \in \mathbb{Z}^+.$$

By Lemma 6 of [11], we may assume that $G_0 = G/H$ is metrizable and $m_G(C_0 + H) = 0$. Let $\pi: G \rightarrow G_0$ be the natural quotient map, and let

$$\nu \rightarrow \pi^*(\nu) = \nu \circ \pi^{-1}: M(G) \rightarrow M(G_0)$$

be the measure algebra homomorphism induced by π . Then it is easy to check that π^* maps $M_a^+(G)$ onto $M_a^+(G_0)$, $M_0^+(G)$ onto $M_0^+(G_0)$, and $L^1(\mu_0)$ onto $L^1(\pi^*(\mu_0))$ (cf. [14, 2.2.4]). Moreover, we have $\|\pi^*(\nu)\| = \|\nu * m_H\|$ for all $\nu \in M(G)$, as is easily seen. It follows from (9) that

$$(10) \quad \|\pi^*[(\nu^n)_s]\| = \|(\nu^n)_s\| \quad \forall n \in \mathbb{Z}^+$$

for all $\nu \in \Phi$. Obviously (10) is satisfied for every $\nu \in M^+(G)$ as well.

Since $m_{G_0}[\pi(C_0)] = m_G(C_0 + H) = 0$ and $\pi^*(\mu_0)$ is carried by the set $\pi(C_0)$, we have $L^1(\pi^*(\mu_0)) \subset M_s(G_0)$. In particular $\pi^*[(\nu^n)_s]$ is the singular part of $(\pi^*(\nu))^n = \pi^*(\nu^n)$ for every $\nu \in L^1(\mu)$ and every $n \in \mathbb{Z}^+$. Hence $r[\pi^*(\nu)] = r(\nu)$ for all $\nu \in [L^1(\mu) \cap M^+(G)] \cup \Phi$, by (10). To complete the proof, it therefore suffices to note that $\pi^*[M_0^+(G)] = M_0^+(G_0)$, that $\pi^*[M(G)] = M(G_0)$, and that the adjoint map of π^* sends $\Delta_{G_0} \setminus \Sigma_{G_0}$ into $\Delta_G \setminus \Sigma_G$ in a one-to-one way. This establishes Theorem B for all nondiscrete groups.

PROOF OF COROLLARY 1. Let $Y \subset \Delta_0$ be a boundary of $M_0(G)$, and $\mu \in M(G)$. Choose any $f \in \Delta_G$ such that $|\hat{\mu}(f)| = \|\hat{\mu}\|_{\Delta_G}$. If $f \in \hat{G}$, we take $\lambda \in M_a(G)$ so that $0 \leq \hat{\lambda} \leq 1$ on \hat{G} and $\hat{\lambda}(f) = 1$. Then we have $\lambda * \mu \in M_0(G)$ and $\|\widehat{\lambda * \mu}\|_{\Delta_G} = |\hat{\mu}(f)|$; hence $|\hat{\mu}(g)| = |\widehat{\lambda * \mu}(g)| = |\hat{\mu}(f)|$ for some $g \in Y$. If $f \notin \hat{G}$, then $r(\mu) = |\hat{\mu}(f)|$. By Theorem B, we can find a probability measure $\tau \in M_0(G)$ such that $r(\tau * \mu) = r(\mu)$. Then $|\hat{\mu}(g)| = |\widehat{\tau * \mu}(g)| = r(\mu) = |\hat{\mu}(f)|$ for some $g \in Y$, which establishes part (a).

To prove (b), first notice that $\overline{\Delta_G \setminus \Sigma_G} \subset \Delta_G \setminus \hat{G}$ since \hat{G} is open and is contained in Σ_G . If the above two sets were different, there would exist a nonempty open set U in Δ_G such that $U \cap \overline{\Delta_G \setminus \Sigma_G} = \emptyset \neq U \cap \hat{G}$. Since the space of all Gelfand transforms of measures is closed under the complex conjugation on Σ_G , it would follow from the Stone-Weierstrass theorem that there would exist a $\hat{\mu} \in M(G)$ such that $0 \leq \hat{\mu} \leq 1$ on Σ_G , $\hat{\mu}(f) = 1$ for some $f \in U \cap \hat{G}$, and $\hat{\mu} < 1/2$ on $\Sigma_G \setminus U$. Then the set $U \cap \hat{\mu}^{-1}(1)$ would be a local peak set for $M(G)$, and therefore would be a peak set for $M(G)$ by Rossi's theorem [4]. Consequently

in Lemma 2. Let $\sigma_1, \sigma_2, \dots, \sigma_p$ be mutually singular measures in $M_c(K)$, and let z_1, z_2, \dots, z_p be complex numbers satisfying $|z_j| \leq \|\sigma_j\|$ ($1 \leq j \leq p$). We then claim that given $\nu \in L^1(\mu)$ there exists an element $f \in \Delta_G \setminus \hat{G}$ such that

$$(1) \quad |f(\nu)| = r(\nu) \quad \text{and} \quad f(\sigma_j) = z_j \quad (1 \leq j \leq p).$$

There is no loss of generality in assuming $\|\sigma_j\| = 1$ for all j . Let τ_{2j-1} and τ_{2j} be mutually singular measures in $L^1(\sigma_j)$ such that $\sigma_j = (\tau_{2j-1} + \tau_{2j})/2$ and $\|\tau_{2j-1}\| = \|\tau_{2j}\| = 1$, and write $z_j = (w_{2j-1} + w_{2j})/2$ with $|w_{2j-1}| = |w_{2j}| = 1$. Since $m_G[C_0 + Gp(K)] = 0$, it follows from Lemma 3 that

$$\begin{aligned} (2) \quad & \left\| \left[\nu * \left(\delta_0 + \sum_{k=1}^{2p} \bar{w}_k \tau_k \right) \right]^n + M_a(G) \right\| \\ &= \left\| (\nu^n)_s * \left(\delta_0 + \sum_{k=1}^{2p} \bar{w}_k \tau_k \right)^n \right\| \\ &= \|(\nu^n)_s\| \left(1 + \sum_{k=1}^{2p} \|\tau_k\| \right)^n = \|(\nu^n)_s\| (1 + 2p)^n, \end{aligned}$$

which yields

$$(3) \quad r \left[\nu * \left(\delta_0 + \sum_{k=1}^{2p} \bar{w}_k \tau_k \right) \right] = r(\nu) \cdot (1 + 2p).$$

We can therefore find an element $f \in \Delta_G \setminus \hat{G}$ such that

$$(1)' \quad |f(\nu)| = r(\nu) \quad \text{and} \quad f(\tau_k) = w_k \quad (1 \leq k \leq 2p).$$

By the choices of τ_k and w_k , (1)' implies (1), which establishes our claim.

We next assert that, given $\nu \in L^1(\mu)$, every linear functional on $M_c(K)$, of norm ≤ 1 , extends to an element $f \in \Delta_G \setminus \hat{G}$ such that $|f(\nu)| = r(\nu)$. In fact, this is an easy consequence of (1) and the arguments of Hewitt and Kakutani in [6]. We leave the details to the reader.

Now choose three disjoint compact sets K_j in K ($j = 1, 2, 3$), each of type M_0 , and fix two probability measures $\lambda \in M_0(K_1)$ and $\tau \in M_0(K_2)$. We now prove that τ and the set

$$X = \{f \in \Delta_G : f(\lambda) = 1, |f(\lambda^*)| \leq 1/2\} \cup \{f \in \Delta_G : |1 - f(\lambda * \lambda^*)| \geq 3/2\}$$

have the required property. It is obvious that X is a compact subset of $\Delta_0 \setminus \Sigma_G$. Let c be a complex number of modulus ≤ 1 , and $\nu \in L^1(\mu)$. Let also φ be an arbitrary (linear) functional on $M_c(K_3)$, of norm ≤ 1 . By the Hahn-Banach theorem, φ extends to a functional ψ on $M_c(K)$, of norm one, such that $\psi(\lambda) = 1$ and $\psi(\tau) = c$. It follows from the result asserted in the last paragraph that there

is an f in $\Delta_G \setminus \hat{G}$ such that $|f(\nu)| = r(\nu)$, $f(\lambda) = 1$, $f(\tau) = c$ and $f = \varphi$ on $M_c(K_3)$. We want to show that such an f can be chosen from the set X . If $|f(\lambda^*)|$ is less than $1/2$, then there is nothing to prove; so assume $|f(\lambda^*)| \geq 1/2$. Setting $\tau_1 = \lambda * \lambda^*$, we then have

$$\|(\nu^m)_s * \tau_1^n\| \geq |f(\nu^m * \tau_1^n)| \geq r(\nu)^m (1/2)^n$$

for all m and $n \in \mathbb{Z}^+$, so that

$$(4) \quad r[\nu^m * (\delta_0 - \tau_1)] \geq r(\nu)^m (3/2) \quad (m \in \mathbb{Z}^+)$$

by (*) and Lemma 3. Putting $\mu_1 = \mu_0 * \exp(\tau_1)$, we also see that μ_1 is carried by the σ -compact set $C_1 = C_0 + Gp(K_1)$ and that

$$\begin{aligned} \mu_1[C_1 + Gp'(K_2 \cup K_3)] &= \int \mu_0[C_1 + Gp'(K_2 \cup K_3) - y] d\theta(y) \\ &\leq \mu_0[C_0 + Gp'(K)] \cdot e = 0, \end{aligned}$$

where $\theta = \exp(\tau_1)$. Therefore, if τ_2, \dots, τ_p are mutually singular probability measures in $M_c(K_2 \cup K_3)$ and if $m, n, n_2, \dots, n_p \in \mathbb{Z}^+$, then

$$(5) \quad \|[\nu^m * (\delta_0 - \tau_1)]^n * \tau_2^{n_2} * \dots * \tau_p^{n_p} + M_a(G)\| \geq r[\nu^m * (\delta_0 - \tau_1)]^n$$

by Lemma 3 (applied to μ_1 and C_1). Consequently, one more application of Lemma 3, combined with (5), yields

$$(6) \quad r\left[\nu^m * (\delta_0 - \tau_1) * \left(\delta_0 + \sum_{j=2}^p \bar{z}_j \tau_j\right)\right] = r[\nu^m * (\delta_0 - \tau_1)] \cdot p$$

for all complex numbers z_2, \dots, z_p of absolute modulus one. (Notice that the left-hand side of (6) cannot be larger than the right-hand one.) Therefore there is a $g_m \in \Delta_G \setminus \hat{G}$ such that

$$(7) \quad |g_m[\nu^m * (\delta_0 - \tau_1)]| = r[\nu^m * (\delta_0 - \tau_1)], \quad g_m(\tau_j) = z_j \quad (2 \leq j \leq p).$$

It follows from (4) and the first equality of (7) that $|1 - g_m(\tau_1)| \geq 3/2$, and so $g_m \in X$; moreover $|g_m(\nu)| = |g_m(\nu^m)|^{1/m} \geq r(\nu) (3/4)^{1/m}$ by (7) and (4). Recalling that X is compact and letting $m \rightarrow \infty$, we find an element $h \in X$ such that

$$(8) \quad |h(\nu)| = r(\nu) \quad \text{and} \quad h(\tau_j) = z_j \quad (2 \leq j \leq p).$$

We repeat almost the same argument as before to obtain an $f \in X$ with the required property. Since it is easy to prove that the conjugate space of $M_c(K_3)$ has cardinality equal to 2^c , this establishes Theorem B for metrizable groups.

The proof for the nonmetrizable case is now easy. We first note that given $\nu \in M(G)$ there is a σ -compact open subgroup Γ of \hat{G} such that $\|\nu * m_H\| = \|\nu\|$,

there would exist a $\nu \in M(G)$ such that $\hat{\nu}(f) = 1$, $|\hat{\nu}| \leq 1$ on Δ_G , and $|\hat{\nu}| < 1/2$ on $\Delta_G \setminus \Sigma_G$. But then $r(\nu) = 1$, which contradicts Theorem B. This establishes part (b).

By Theorem B, no element of $\Delta_G \setminus \hat{G}$ can be a peak point for $M(G)$; hence $P_G \subset \hat{G}$. Therefore part (c) is an easy consequence of the fact that G is σ -compact if and only if \hat{G} is metrizable [7]. This completes the proof.

PROOF OF COROLLARY 2. Let $\mu \in M(G)$ be given. Notice that $\Delta_G \setminus \hat{G}$ is the spectrum of the quotient algebra $M(G)/M_d(G)$. Choose a countable dense subset D of $\mathbb{C} \setminus \hat{\mu}(\Delta_G \setminus \hat{G})$. For each $c \in D$, there is a $\nu_c \in M(G)$ such that $\hat{\nu}_c = (c - \hat{\mu})^{-1}$ on $\Delta_G \setminus \hat{G}$. Setting $\Phi = \{\nu_c : c \in D\}$, we apply Theorem B to find a compact set X in $\Delta_0 \setminus \Sigma_G$ such that

$$\sup \{|c - \hat{\mu}(f)|^{-1} : f \in X\} = \sup \{|c - \hat{\mu}(g)|^{-1} : g \in \Delta_G \setminus \hat{G}\}$$

for all $c \in D$. Since $\hat{\mu}(X)$ is compact, this implies that $\hat{\mu}(X)$ contains all the boundary points of $\hat{\mu}(\Delta_G \setminus \hat{G})$ in \mathbb{C} .

If $\text{Card}[\hat{\mu}(\Delta_0 \setminus \Sigma_G)] < \aleph_1$, then $\hat{\mu}(\Delta_G \setminus \hat{G})$ has a countable boundary since it is compact. Therefore $\hat{\mu}(\Delta_G \setminus \hat{G})$ itself is countable, so that $\hat{\mu}(\Delta_G \setminus \hat{G}) = \hat{\mu}(\Delta_0 \setminus \Sigma_G)$ by the result already established. If $\hat{\mu}$ is real on $\Delta_0 \setminus \Sigma_G$, then $\hat{\mu}$ must be real on $\Delta_G \setminus \hat{G}$, hence $\hat{\mu}(\Delta_G \setminus \hat{G})$ has no interior point, and hence $\hat{\mu}(\Delta_G \setminus \hat{G}) = \hat{\mu}(\Delta_0 \setminus \Sigma_G)$. This establishes Corollary 2.

REMARKS. (a) Theorem A implies $\hat{\mu}_d(\Delta_G) \subset \hat{\mu}(\Delta_0 \setminus \Sigma_G)$ for all $\mu \in M(G)$. Moreover, we can prove that $\hat{\mu}_d(\Delta_G) \subset \hat{\mu}(\Delta_0 \cap \Sigma_G \setminus \hat{G})$ by applying the methods in [11].

(b) Notice that $\delta_0(C_1 + C_2) = 0$ if and only if $C_1 \cap (-C_2)$ is empty. If we only require that $C_0 \cap Gp'(K) = \emptyset$ in Lemma 2 instead of that $m_G[C_0 + Gp(K)] = \mu_0[C_0 + Gp'(K)] = 0$, then the assumption that C_0 is a σ -compact set with $m_G(C_0) = 0$ can be weakened to be that C_0 is a set of the first category in G (cf. [5, 2.1]).

(c) In some special cases, the proof of Theorem B can be somewhat simplified and we have a result slightly stronger than Theorem B.

Let H_0 be an open subgroup of G of the form $H_0 = \mathbb{R}^n \times H_1$, where n is a nonnegative integer and H_1 is a compact subgroup of G (cf. [10, 2.4.1]). Let P be the set of all $p \in \mathbb{Z}$ such that $1 \leq p < q(H_1)$ and $\text{Card}\{\chi \in \hat{H}_1 : \chi^p = 1\} < \infty$. Then the last condition in Lemma 2 can be strengthened to be that $m_G[C_0 + Gp(K)] = \mu_0[C_0 + K(P)] = 0$. Here $K(P)$ denotes the set of all points x of the form $x = \sum_1^u k_j x_j$, where $u = u_x$ is a natural number, x_1, x_2, \dots, x_u are distinct elements of K , and k_1, k_2, \dots, k_u are integers such that $|k_j| \in P$ for some $1 \leq j \leq u$. The case where $2 \in P$ is particularly interesting.

Suppose in Lemma 3 that μ_0 , C_0 and K are such that $\mu_0[C_0 + K(\{1, 2\})] = 0$. Then we can prove that

$$\|\nu * \sigma_1^{n_1} * \cdots * \sigma_p^{n_p}\| = \|\nu\| \cdot \|\sigma_1\|^{n_1} \cdots \|\sigma_p\|^{n_p}$$

for all $\nu \in L^1(M_0)$ and all $\sigma_j \in M_c(K_j \cup (-K_j))$. Therefore a moment's glance at the proof of Theorem B yields this result: If either $q(G) = 2$, or G contains an open subgroup H_0 as above with $2 \in P$, then the measure τ in Theorem B can be taken so that $\tau = \lambda * \lambda^*$ for some $\lambda \in M_0^+(G)$. We omit the details.

(d) If $\mu \in M^+(G)$, then the number $r(\mu)$ is in $\hat{\mu}(\Delta_0 \setminus \Sigma_G)$. To see this, choose a complex number z of absolute modulus one so that $zr(\mu) \in \hat{\mu}(\Delta_G \setminus \hat{G})$. Then we have

$$\begin{aligned} r(\delta_0 + \mu) &= \lim_{n \rightarrow \infty} \|[(\delta_0 + \mu)^n]_s\|^{1/n} \\ &\geq \lim_{n \rightarrow \infty} \|[(\delta_0 + \bar{z}\mu)^n]_s\|^{1/n} = 1 + r(\mu), \end{aligned}$$

and so $r(\delta_0 + \mu) = 1 + r(\mu)$. Thus our assertion follows from Theorem B with $\Phi = \{\delta_0 + \mu\}$.

(e) Let $M_0^\infty(G)$ denote the L -ideal in $M(G)$ generated by all measures μ of the form $\mu = \mu_1 * \mu_2 * \cdots$, where the μ_j are probability measures in $M_0(G)$ and the infinite convolution product is assumed to converge in the weak-* topology of $M(G)$. Let also Δ_0^∞ denote the spectrum of $M_0(G)$ identified with an open subset of Δ_G . Then it is not difficult to prove that $\text{Card}(\Delta_0 \setminus \Delta_0^\infty) \geq 2^c$. Moreover, in Theorem B, we can replace $M_0(G)$ and Δ_0 by $M_0^\infty(G)$ and Δ_0^∞ , respectively. Using this result, we can prove that if Y is a boundary of $M_0^\infty(G)$, then $(Y \setminus \Sigma_G) \cup (Y \cap \hat{G})$ is a boundary of $M(G)$, which of course improves part (a) of Corollary 1. Similarly the set Δ_0 in Corollary 2 can be replaced by Δ_0^∞ .

(f) Finally we list three problems which the author has been unable to solve.

- (i) Is it true that $\hat{\mu}(\Delta_G \setminus \hat{G}) = \hat{\mu}(\Delta_0 \setminus \Sigma_G)$ for all $\mu \in M(G)$?
- (ii) Does $\hat{\mu}(\Sigma_G \setminus \hat{G}) = \{0\}$ imply $r(\mu) = 0$?
- (iii) Does $\Sigma_G \setminus \hat{G}$ contain any strong boundary point for $M(G)$?

REFERENCES

1. G. Brown, $M_0(G)$ has a symmetric maximal ideal off the Šilov boundary, Proc. London Math. Soc. (3) 27 (1973), 484–504. MR 48 #2666.
2. G. Brown and W. Moran, $L^{1/2}(G)$ is the kernel of the asymmetric maximal ideals of $M(G)$, Bull. London Math. Soc. 5 (1973), 179–186. MR 49 #3450.
3. ———, $M_0(G)$ -boundaries are $M(G)$ -boundaries, J. Functional Analysis 18 (1975), 350–368. MR 50 #14060.
4. T. W. Gamelin, *Uniform algebras*, Prentice-Hall, New York, 1969.
5. C. C. Graham, Measures vanishing off the symmetric maximal ideals of $M(G)$, Proc. Cambridge Philos. Soc. 75 (1974), 51–61. MR 49 #9539.
6. E. Hewitt and S. Kakutani, A class of multiplicative linear functionals on the

measure algebra of a locally compact Abelian group, Illinois J. Math. 4 (1960), 553–574. MR 23 #A527.

7. E. Hewitt and K. R. Stromberg, *A remark on Fourier-Stieltjes transforms*, An. Acad. Brasil. Ci. 34 (1962), 175–180. MR 27 #533.

8. K. Izuchi and T. Shimizu, *Topologies on groups and a certain L -ideal of measure algebras*, Tôhoku Math. J. 25 (1973), 53–60.

9. L.-A. Lindahl and F. Poulsen, *Thin sets in harmonic analysis*, Dekker, New York, 1971.

10. W. Rudin, *Fourier analysis on groups*, Interscience Tracts in Pure and Appl. Math., no. 12, Interscience, New York and London, 1962. MR 27 #2808.

11. S. Saeki, *Symmetric maximal ideals in $M(G)$* , Pacific J. Math. 53 (1974), 229–243.

12. K. Saka, *A note on subalgebras of a measure algebra vanishing on non-symmetric homomorphisms*, Tôhoku Math. J. (2) 25 (1973), 333–338. MR 50 #10686.

13. T. Shimizu, *L -ideals of measure algebras*, Proc. Japan Acad. 48 (1972), 172–176. MR 47 #7326.

14. J. L. Taylor, *Measure algebras*, CBMS Regional Conf. Ser. in Math., no. 16, Amer. Math. Soc., Providence, R. I., 1973.

15. J. H. Williamson, *Banach algebra elements with independent powers and theorems of Wiener-Pitt type*, Function Algebras (Proc. Internat. Sympos. on Function Algebras, Tulane Univ., 1965), Scott, Foresman, Chicago, Ill., 1966, pp. 186–197. MR 33 #6302.

DEPARTMENT OF MATHEMATICS, TOKYO METROPOLITAN UNIVERSITY,
SETAGAYA, TOKYO, JAPAN