

A NOTE ON LIMITS OF UNITARILY EQUIVALENT OPERATORS

BY

LAWRENCE A. FIALKOW

In loving memory of my mother

ABSTRACT. Let $\mathcal{U}(\mathcal{H})$ denote the set of all unitary operators on a separable complex Hilbert space \mathcal{H} . If T is a bounded linear operator on \mathcal{H} , let π_T denote the mapping of $\mathcal{U}(\mathcal{H})$ onto $\mathcal{U}(T)$ given by conjugation. It is proved that if T is normal or isometric, then there exists a locally defined continuous cross-section for π_T if and only if the spectrum of T is finite. Examples of nonnormal operators with local cross-sections are given.

1. Introduction. Let \mathcal{H} denote a separable complex Hilbert space and let $\mathcal{L}(\mathcal{H})$ denote the algebra of all bounded linear operators on \mathcal{H} . Let $\mathcal{U}(\mathcal{H})$ denote the set of all unitary operators on \mathcal{H} , and for T in $\mathcal{L}(\mathcal{H})$, let $\mathcal{U}(T)$ denote the unitary orbit of T , i.e., $\mathcal{U}(T) = \{U^*TU \mid U \text{ in } \mathcal{U}(\mathcal{H})\}$. We denote by π_T the norm continuous mapping of $\mathcal{U}(\mathcal{H})$ onto $\mathcal{U}(T)$ defined by $\pi_T(U) = U^*TU$ (U in $\mathcal{U}(\mathcal{H})$). A *local cross-section* for π_T is a pair $(\varphi_T, \mathfrak{B})$ such that \mathfrak{B} is a relatively open subset of $\mathcal{U}(T)$ that contains T and $\varphi_T: \mathfrak{B} \rightarrow \mathcal{U}(\mathcal{H})$ is a norm continuous function such that $\varphi_T(T) = 1$ and $\pi_T(\varphi_T(S)) = S$ for each S in \mathfrak{B} . If π_T has a local cross-section, we say that T has a (*local unitary*) *cross-section*, and in this case T clearly satisfies the following *sequential unitary lifting property*:

(P) If $\{U_n\} \subset \mathcal{U}(\mathcal{H})$ and $\lim \|U_n^*TU_n - T\| = 0$, then there exists a sequence $\{W_n\} \subset \mathcal{U}(\mathcal{H})$ such that $\lim \|W_n - 1\| = 0$ and such that $W_n^*TW_n = U_n^*TU_n$ for each n . The problem to which this note is addressed is that of obtaining necessary and sufficient conditions for an operator on \mathcal{H} to have a local cross-section or to satisfy (P).

The fact that an orthogonal projection satisfies (P) has appeared in the literature in diverse contexts. In [4, Lemma 1], P. R. Halmos proved that if P and Q are finite rank projections and $\|P - Q\| = \varepsilon < 1$, then there exists a unitary operator W such that $W^*PW = Q$ and $\|W - 1\| < 2\varepsilon^{1/2}$. This result was used in proving a principal characterization of quasi-triangular operators [4, Theorem 2]. Earlier results related to this lemma appeared in [1]. As we remark below, this lemma implies that a finite rank projection satisfies (P).

Received by the editors April 24, 1975 and, in revised form, February 13, 1976.

AMS (MOS) subject classifications (1970). Primary 47A10.

© American Mathematical Society 1977

An extension of the lemma to arbitrary projections in $\mathcal{L}(\mathcal{H})$ was given in [2] as a tool in studying the closure of the unitary orbit of a partial isometry.

In §3 we prove that if an operator T is normal or isometric, then it satisfies (P) if and only if its spectrum is finite; if the spectrum of T is finite, then there exists a local cross-section $(\varphi_T, \mathfrak{B})$ such that for each operator S in \mathfrak{B} , $\varphi_T(S)$ is in the norm closed subalgebra of $\mathcal{L}(\mathcal{H})$ generated by S , T , and 1. Furthermore, if a normal operator with infinite spectrum is a direct summand of an operator S , then S does not satisfy (P). We prove in Theorem 3.5 that if T is an irreducible operator and T has a reducing essential eigenvalue, then T does not satisfy (P); if such an operator is a direct summand of an operator S , then S does not satisfy (P). This result is used to show that there exist compact, quasi-nilpotent operators that do not satisfy (P); these include the Volterra integral operator and every compact injective bilateral weighted shift.

In §§2 and 4 we show that there do exist nonnormal operators that satisfy (P) or have local unitary cross-sections. In Corollary 2.4 we prove that each operator on a finite dimensional Hilbert space satisfies (P). Suppose that T has a cross-section and that S is an operator such that $T^*T + S^*S$ is invertible. In Theorems 4.2 and 4.3 we prove that the operator on $\mathcal{H} \oplus \mathcal{H}$ whose operator matrix is $\begin{pmatrix} T & 0 \\ S & 0 \end{pmatrix}$ has a cross-section if either (1) S is a right invertible operator in the C^* -algebra generated by T , or (2) S is an isometry. Additional examples of operators with cross-sections may be constructed by application of Theorem 2.8, which states that a finite direct sum of operators (having mutually disjoint spectra) has a cross-section if and only if each summand has a cross-section. Analogues of these results for the sequential unitary lifting property are also given. In [2] it was proven that if T has a local cross-section, then each path in $\mathcal{U}(T)$ can be lifted to a path in $\mathcal{U}(\mathcal{H})$; we prove in Proposition 2.2 that if T satisfies (P), then $\mathcal{U}(T)^-$ (the norm closure of $\mathcal{U}(T)$ in $\mathcal{L}(\mathcal{H})$) is path connected (cf. [3]). At the end of §4 we list some questions related to characterizing the operators having local cross-sections.

I wish to thank E. Azoff, L. Brown, D. Hadwin, C. Pearcy, and the referee for their helpful comments and suggestions concerning this paper.

2. On the sequential unitary lifting property. In this section we give some general properties of operators satisfying (P). We denote the spectrum of an operator T by $\sigma(T)$. For $\epsilon > 0$, we denote the set $\{X \text{ in } \mathcal{U}(T) \mid \|X - T\| < \epsilon\}$ by $\mathfrak{B}(T, \epsilon)$.

LEMMA 2.1. *If U is a unitary operator in $\mathcal{L}(\mathcal{H})$ and $\|U - 1\| < 2$, then there exists a path $\{U_t\}_{0 \leq t < 1} \subset \mathcal{U}(\mathcal{H})$ such that $U_0 = U$, $U_1 = 1$, and such that, for each t , $\|U_t - 1\| < \|U - 1\|$.*

PROOF. Let A denote a selfadjoint operator such that $U = e^{iA}$ (see [5, p. 66]). Since $2 > \|U - 1\| = \|e^{iA} - 1\| = \sup_{s \in \sigma(A)} |e^{is} - 1|$, it is clear that $\sigma(A)$ contains no number of the form $\pi + 2\pi n$, where n is an integer. For each n , let $I_n = [2\pi n, 2\pi n + \pi]$ and $J_n = [2\pi n + \pi, 2\pi(n + 1)]$. For $0 < t < 1$ and s in $I_n \cap \sigma(A)$, we define $f_t(s) = (1 - t)s + 2\pi nt$; for s in $J_n \cap \sigma(A)$, we define $f_t(s) = (1 - t)s + 2\pi(n + 1)t$. For $0 < t < 1$ and s in $\sigma(A)$, we set $g_t(s) = e^{if_t(s)}$. The hypothesis implies that g_t is well defined and continuous on $\sigma(A)$, and it follows readily from the functional calculus that $U_t = g_t(A)$ ($0 < t < 1$) defines a path satisfying the desired properties.

If T is in $\mathcal{L}(\mathcal{H})$, let $\mathfrak{S}(T)$ denote the set of all sequences $\{U_n\} \subset \mathcal{U}(\mathcal{H})$ such that $\lim \|U_n^* T U_n - T\| = 0$. If $U = \{U_n\}$ is in $\mathfrak{S}(T)$ and $V = \{V_n\} \subset \mathcal{U}(\mathcal{H})$, we say that V re-implements U in case $\lim \|1 - V_n\| = 0$ and $U_n^* T U_n = V_n^* T V_n$ for each n .

PROPOSITION 2.2. *If an operator T in $\mathcal{L}(\mathcal{H})$ satisfies (P), then $\mathcal{U}(T)^-$ is path connected.*

PROOF. Let S be in $\mathcal{U}(T)^-$ and let $\{U_n\} \subset \mathcal{U}(\mathcal{H})$ be a sequence such that $\lim \|U_n T U_n^* - S\| = 0$. Then $U = \{U_n^* U_{n+1}\}$ is in $\mathfrak{S}(T)$ and the hypothesis implies that there is a sequence $\{W_n\} \subset \mathcal{U}(\mathcal{H})$ such that $\{W_n\}$ re-implements U . If $T_n = U_n T U_n^*$ and $V_n = U_{n+1} W_n^* U_{n+1}^*$, then $T_{n+1} = V_n T_n V_n^*$ and $\lim \|1 - V_n\| = \lim \|1 - W_n\| = 0$. We may thus assume that $\|1 - V_n\| < 2$, and the preceding lemma implies that, for fixed n , there exists a path $\{V_{t,n}\}$ ($0 < t < 1$) from V_n to 1 such that $\|V_{t,n} - 1\| < \|V_n - 1\|$ for each t . It follows that $K_n = \{V_{t,n} T_n V_{t,n}^*\}$ ($0 < t < 1$) is a path from T_{n+1} to T_n such that for each t we have

$$\begin{aligned} \|V_{t,n} T_n V_{t,n}^* - T_n\| &\leq \|V_{t,n} T_n V_{t,n}^* - V_{t,n} T_n\| + \|V_{t,n} T_n - T_n\| \\ &\leq \|V_{t,n} T_n\| \|V_{t,n}^* - 1\| + \|V_{t,n} - 1\| \|T_n\| \\ &\leq 2\|T\| \|V_{t,n} - 1\| \leq 2\|T\| \|V_n - 1\|. \end{aligned}$$

Thus $\|V_{t,n} T_n V_{t,n}^* - S\| \leq 2\|T\| \|V_n - 1\| + \|T_n - S\|$, and it follows that the paths $\{K_n\}$ may be combined and reparametrized to form a single path from T to S , each member of which, except perhaps S itself, is in $\mathcal{U}(T)$.

LEMMA 2.3. *For an operator T in $\mathcal{L}(\mathcal{H})$ the following properties are equivalent:*

(P1) *Given $\epsilon > 0$, there exists $\delta > 0$ such that if S is in $\mathcal{U}(T)$ and $\|S - T\| < \delta$, then there exists a unitary operator W such that $\|W - 1\| < \epsilon$ and $W^* T W = S$.*

(P2) *T satisfies property (P).*

(P3) *If $\{U_n\}$ is in $\mathfrak{S}(T)$, then there exists a subsequence $V = \{U_{n_k}\}$ and a sequence $W = \{W_k\}$ in $\mathcal{U}(\mathcal{H})$ such that W re-implements V .*

PROOF. To prove that (P1) implies (P2), suppose that $U = \{U_n\}$ is in $\mathfrak{S}(T)$. For each integer $k > 0$, corresponding to $\epsilon = 1/k$, there exists $\delta_k > 0$ satisfying the conditions of (P1). Let N_k be a positive integer such that if $n > N_k$, then $\|U_n^*TU_n - T\| < \delta_k$; we may also assume that $N_k > N_{k-1}$ for each k . Thus, for $n > N_k$, there exists a unitary operator $W_{n,k}$ such that $\|W_{n,k} - 1\| < 1/k$ and $W_{n,k}^*TW_{n,k} = U_n^*TU_n$. For each integer $n > N_1$, let $k(n)$ denote the unique integer such that $N_{k(n)+1} > n > N_{k(n)}$, and define $W_n = W_{n,k(n)}$; for each $n < N_1$, let $W_n = U_n$. It now follows readily that $\{W_n\}$ re-implements U , and therefore T satisfies (P).

It is clear that (P2) implies (P3). That (P3) implies (P1) may be proven by a straightforward proof by contradiction; we omit the details.

REMARK. To prove that an operator T satisfies (P1), it clearly suffices to assume that $0 < \epsilon < 2$. If T is a finite rank projection and $0 < \epsilon < 2$, the above-mentioned result of P. R. Halmos shows that T satisfies (P1) with $\delta = \epsilon^2/4$.

COROLLARY 2.4. *If \mathfrak{H} is a finite dimensional Hilbert space and T is in $\mathcal{L}(\mathfrak{H})$, then T satisfies (P).*

PROOF. Let $U = \{U_n\}$ be in $\mathfrak{S}(T)$. Since $\mathcal{U}(\mathfrak{H})$ is compact, there exists a norm-convergent subsequence $U_{n_k} \rightarrow V$. Now V is a unitary operator that commutes with T , and it follows that $\{V^*U_{n_k}\}$ re-implements $\{U_{n_k}\}$. Thus T satisfies (P3), and Lemma 2.3 implies that T satisfies (P).

Let \mathfrak{H}_1 and \mathfrak{H}_2 denote separable complex Hilbert spaces and let $\mathcal{L}(\mathfrak{H}_2, \mathfrak{H}_1)$ denote the Banach space of all bounded linear operators from \mathfrak{H}_2 into \mathfrak{H}_1 . For operators A in $\mathcal{L}(\mathfrak{H}_1)$ and B in $\mathcal{L}(\mathfrak{H}_2)$ we define the operator \mathfrak{T} on $\mathcal{L}(\mathfrak{H}_2, \mathfrak{H}_1)$ by $\mathfrak{T}(X) = AX - XB$ (X in $\mathcal{L}(\mathfrak{H}_2, \mathfrak{H}_1)$).

LEMMA 2.5. *If \mathfrak{T} is bounded below on $\mathcal{L}(\mathfrak{H}_2, \mathfrak{H}_1)$, then $A \oplus B$ (in $\mathcal{L}(\mathfrak{H}_1 \oplus \mathfrak{H}_2)$) satisfies (P) if and only if both A and B satisfy (P).*

PROOF. We assume first that A and B satisfy (P). If $\{U_n\}$ is in $\mathfrak{S}(A \oplus B)$, we denote the operator matrix of U_n by

$$\begin{pmatrix} X_n & Y_n \\ Z_n & V_n \end{pmatrix}.$$

Since $\lim \|U_n^*(A \oplus B)U_n - A \oplus B\| = 0$, a calculation yields the relations

- (1) $\lim \|X_n^*AX_n + Z_n^*BZ_n - A\| = 0$,
- (2) $\lim \|X_n^*AY_n + Z_n^*BV_n\| = 0$, and
- (3) $\lim \|Y_n^*AY_n + V_n^*BV_n - B\| = 0$.

If we multiply (2) on the left by X_n , multiply (3) on the left by Y_n , and add the new limits, we obtain $\lim \|AY_n - Y_nB\| = 0$; since \mathfrak{T} is bounded below, it follows that $\lim \|Y_n\| = 0$. Similarly, if we multiply (1) on the right by Z_n^* ,

multiply (2) on the right by V_n^* , and add the new limits, we obtain $\lim \|AZ_n^* - Z_n^*B\| = 0$, whence $\lim \|Z_n^*\| = 0$. Now

$$\lim \|1 - X_n X_n^*\| = \lim \|Y_n Y_n^*\| = 0$$

and

$$\lim \|1 - X_n^* X_n\| = \lim \|Z_n^* Z_n\| = 0,$$

and it follows that X_n is invertible for all but finitely many values of n . A similar argument with V_n allows us to assume that each X_n and each V_n is invertible.

If $X_n = R_n(X_n^* X_n)^{1/2}$ and $V_n = S_n(V_n^* V_n)^{1/2}$ denote, respectively, the polar decompositions of X_n and V_n , then R_n and S_n are unitary; we will show that $\{R_n\}$ is in $\mathfrak{S}(A)$ and $\{S_n\}$ is in $\mathfrak{S}(B)$. Since $\lim \|1 - X_n^* X_n\| = 0$, the functional calculus implies that

$$(4) \lim \|1 - (X_n^* X_n)^{1/2}\| = 0.$$

Now

$$\begin{aligned} R_n^* A R_n - A &= R_n^* A R_n (1 - (X_n^* X_n)^{1/2}) \\ &\quad + (1 - (X_n^* X_n)^{1/2}) R_n^* A R_n (X_n^* X_n)^{1/2} + X_n^* A X_n - A. \end{aligned}$$

Since $\lim \|Z_n^*\| = 0$, the last equation, together with (1) and (4), imply that $\{R_n\}$ is in $\mathfrak{S}(A)$. Similarly, using (3), the fact that $\lim \|1 - (V_n^* V_n)^{1/2}\| = 0$, and $\lim \|Y_n\| = 0$, it follows that $\{S_n\}$ is in $\mathfrak{S}(B)$.

Since A and B satisfy (P), there exist sequences $\{P_n\} \subset \mathfrak{U}(\mathcal{H}_1)$ and $\{Q_n\} \subset \mathfrak{U}(\mathcal{H}_2)$ that re-implement, respectively, $\{R_n\}$ (for A) and $\{S_n\}$ (for B). Since $P_n R_n^*$ commutes with A and $Q_n S_n^*$ commutes with B , the unitary operator $U'_n = (P_n R_n^* \oplus Q_n S_n^*) U_n$ satisfies $U_n^* (A \oplus B) U'_n = U_n^* (A \oplus B) U_n$. Since the matrix of U'_n is

$$\begin{bmatrix} P_n (X_n^* X_n)^{1/2} & P_n R_n^* Y_n \\ Q_n S_n^* Z_n & Q_n (V_n^* V_n)^{1/2} \end{bmatrix},$$

it follows that $\lim \|1 - U'_n\| = 0$, and therefore $A \oplus B$ satisfies (P).

For the converse, we assume that $A \oplus B$ satisfies (P) but that A does not. If $\{W_n\}$ is an element of $\mathfrak{S}(A)$ that cannot be re-implemented, then $\{W_n \oplus 1\}$ is in $\mathfrak{S}(A \oplus B)$, and thus may be re-implemented by a sequence $\{U_n\}$. We use the notation for the matrix of U_n of the first part of the proof. A calculation yields the relations (1') $X_n^* A X_n + Z_n^* B Z_n - W_n^* A W_n = 0$, (2') $X_n^* A Y_n + Z_n^* B V_n = 0$, and (3') $Y_n^* A Y_n + V_n^* B V_n - B = 0$. Using (1')–(3'), calculations analogous to those above (using (1)–(3)) now yield $Z_n = 0$ and $Y_n = 0$. Thus, from (1'), it follows that $\{X_n\}$ re-implements $\{W_n\}$, which is a contradiction. A similar argument for B completes the proof; we omit these details.

THEOREM 2.6 *If \mathfrak{T} is bounded below on $\mathcal{L}(\mathcal{K}_2, \mathcal{K}_1)$, then $A \oplus B$ (in $\mathcal{L}(\mathcal{K}_1 \oplus \mathcal{K}_2)$) has a local cross-section if and only if A and B have local cross-sections.*

PROOF. Let $\delta > 0$ be such that $\|AX - XB\| \geq \delta\|X\|$ for each X in $\mathcal{L}(\mathcal{K}_2, \mathcal{K}_1)$. Let $\varepsilon > 0$ be chosen such that A and B have local cross-sections of the form $(\varphi_A, \mathfrak{B}(A, \varepsilon))$ and $(\varphi_B, \mathfrak{B}(B, \varepsilon))$ respectively. Let U be a unitary operator on $\mathcal{K}_1 \oplus \mathcal{K}_2$ with operator matrix $U = \begin{pmatrix} X & Y \\ Z & V \end{pmatrix}$, and let $X = R(X^*X)^{1/2}$ and $V = S(V^*V)^{1/2}$ denote the respective polar decompositions of X and V . We will show first that there exists $\varepsilon_0 > 0$ such that if

$$(*) \quad \|U^*(A \oplus B)U - (A \oplus B)\| < \varepsilon_0,$$

then S and R are unitary, $\|R^*AR - A\| < \varepsilon$, and $\|S^*BS - B\| < \varepsilon$.

If $(*)$ is satisfied for some $\varepsilon_0 > 0$, then we have (1) $\|X^*AX + Z^*BZ - A\| < \varepsilon_0$, (2) $\|X^*AY + Z^*BV\| < \varepsilon_0$, and (3) $\|Y^*AY + V^*BV - B\| < \varepsilon_0$. Now

$$\begin{aligned} \|Y\| &\leq (1/\delta)\|AY - YB\| \\ &= (1/\delta)\|(XX^* + YY^*)AY + (XZ^* + YV^*)BV - YB\| \\ &= (1/\delta)\|X(X^*AY + Z^*BV) + Y(Y^*AY + V^*BV - B)\| < 2\varepsilon_0/\delta. \end{aligned}$$

Similarly, we have $\|Z\| < 2\varepsilon_0/\delta$. If we set $K = 4\varepsilon_0^2/\delta^2$, then $\|1 - XX^*\| = \|YY^*\| = \|Y\|^2 < K$; likewise $\|1 - X^*X\|$, $\|1 - VV^*\|$, and $\|1 - V^*V\|$ are less than K . Thus if $K < 1$ (i.e., if $\varepsilon_0 < \delta/2$), then V and X are invertible and therefore R and S are unitary. If we assume also that $\varepsilon_0 < 1$, then

$$\begin{aligned} \|R^*AR - A\| &\leq \|R^*AR(1 - (X^*X)^{1/2})\| \\ &\quad + \|(1 - (X^*X)^{1/2})R^*AR(X^*X)^{1/2}\| \\ &\quad + \|Z^*BZ\| + \|X^*AX + Z^*BZ - A\| \\ &< 2\|A\|(1 - (1 - K)^{1/2}) + \|B\|\|Z\|^2 + \varepsilon_0 \\ &< K(2\|A\| + \|B\|) + \varepsilon_0 < \varepsilon_0((4/\delta^2)(2\|A\| + \|B\|) + 1) \\ &= \varepsilon_0(8\|A\| + 4\|B\| + \delta^2)/\delta^2. \end{aligned}$$

We may estimate $\|S^*BS - B\|$ similarly, and it follows that if $(*)$ is satisfied for

$$0 < \varepsilon_0 < \min \left\{ 1, \frac{\delta}{2}, \frac{\varepsilon\delta^2}{8\|A\| + 4\|B\| + \delta^2}, \frac{\varepsilon\delta^2}{8\|B\| + 4\|A\| + \delta^2} \right\},$$

then R and S are unitary, $\|R^*AR - A\| < \varepsilon$, and $\|S^*BS - B\| < \varepsilon$.

For $U^*(A \oplus B)U$ in $\mathfrak{B}(A \oplus B, \varepsilon_0)$, we now define $\varphi(U^*(A \oplus B)U) = (\varphi_A(R^*AR)R^* \oplus \varphi_B(S^*BS)S^*)U$, and we will prove that $(\varphi, \mathfrak{B}(A \oplus B, \varepsilon_0))$ is a local cross-section. To prove that φ is well defined, suppose that $U^*(A \oplus B)U = U_1^*(A \oplus B)U_1$ for U_1 in $\mathcal{U}(\mathcal{K}_1 \oplus \mathcal{K}_2)$. Since \mathfrak{T} is bounded

below and U_1U^* commutes with $A \oplus B$, it follows that $U_1U^* = L \oplus M$, where L and M are unitary operators that commute with A and B respectively. Thus the operator matrix of $U_1 = (L \oplus M)U$ is $\begin{pmatrix} LX & LY \\ MZ & MV \end{pmatrix}$. Now the respective polar decompositions of LX and MV are $LX = LR(X^*X)^{1/2}$ and $MV = MS(V^*V)^{1/2}$, and we have

$$\begin{aligned} & (\varphi_A((LR)^*ALR)R^*L^* \oplus \varphi_B((MS)^*BMS)S^*M^*)U_1 \\ &= (\varphi_A(R^*AR)R^* \oplus \varphi_B(S^*BS)S^*)(L^* \oplus M^*) \begin{pmatrix} LX & LY \\ MZ & MV \end{pmatrix} \\ &= \varphi(U^*(A \oplus B)U), \end{aligned}$$

which completes the proof that φ is well defined. Since $\varphi_A(R^*AR)R^*$ commutes with A and $\varphi_B(S^*BS)S^*$ commutes with B , it is clear that φ is a cross-section, and it is also clear that $\varphi(A \oplus B) = 1$. To complete the proof it suffices to prove that φ is continuous. Suppose that $\{U_n\}_{n=0}^\infty \subset \mathcal{U}(\mathcal{H}_1 \oplus \mathcal{H}_2)$ is a sequence such that $U_n^*(A \oplus B)U_n \rightarrow U_0^*(A \oplus B)U_0$ in $\mathfrak{B}(A \oplus B, \epsilon_0)$. Since A and B satisfy (P), Lemma 2.5 implies that there is a sequence $\{W_n\}_{n=1}^\infty \subset \mathcal{U}(\mathcal{H}_1 \oplus \mathcal{H}_2)$ such that $W_n \rightarrow 1$ and such that $W_n^*(A \oplus B)W_n = U_0U_n^*(A \oplus B)U_nU_0^*$ for each n . Thus

$$U_n^*(A \oplus B)U_n = U_0^*W_n^*(A \oplus B)W_nU_0,$$

and since φ is well defined, we may verify the continuity of φ using the sequence $\{W_nU_0\}$; moreover, the limit

$$\varphi(U_0^*W_n^*(A \oplus B)W_nU_0) \rightarrow \varphi(U_0^*(A \oplus B)U_0)$$

is now a simple consequence of the definition of φ and the fact that $W_nU_0 \rightarrow U_0$.

For the converse, suppose that $A \oplus B$ has a cross-section of the form $(\varphi, \mathfrak{B}((A \oplus B), \epsilon))$ for some $\epsilon > 0$. If X is in $\mathcal{U}(\mathcal{H}_1)$ and $\|X^*AX - A\| < \epsilon$, then $\varphi(X^*AX \oplus B)(X^* \oplus 1)$ commutes with $A \oplus B$ and therefore is of the form $Z \oplus W$, where Z commutes with A . We now define $\varphi_A(X^*AX) = ZX$ and it follows easily that φ_A is well defined, that φ_A is a cross-section, and that $\varphi_A(A) = 1$. To complete the proof we will prove that φ_A is continuous on $\mathfrak{B}(A, \epsilon)$. Suppose $\{X_n\}_{n=0}^\infty \subset \mathcal{U}(\mathcal{H}_1)$ is a sequence such that $X_n^*AX_n \rightarrow X_0^*AX_0$ in $\mathfrak{B}(A, \epsilon)$. Since $A \oplus B$ satisfies (P), Lemma 2.5 implies that there exists a sequence $\{W_n\} \subset \mathcal{U}(\mathcal{H}_1)$ such that $W_n \rightarrow 1$ and such that $W_n^*AW_n = X_0X_n^*AX_nX_0^*$ for each n . Since $X_n^*AX_n = X_0^*W_n^*AW_nX_0$ and since φ_A is well defined, we may use the sequence $\{W_nX_0\}$ in calculating $\varphi_A(X_n^*AX_n)$. It now follows readily from the definition of φ_A and from the fact that $W_nX_0 \rightarrow X_0$ that $\varphi_A(X_n^*AX_n) = \varphi_A(X_0^*W_n^*AW_nX_0) \rightarrow \varphi_A(X_0^*AX_0)$, and the proof is complete.

Let n be an integer greater than or equal to 2. Let A_i be in $\mathcal{L}(\mathcal{H}_i)$,

$i = 1, \dots, n$, and suppose that $\sigma(A_i) \cap \sigma(A_j) = \emptyset$ for $i \neq j$. Let $B = A_1 \oplus \dots \oplus A_n$ in $\mathcal{L}(\mathcal{H}_1 \oplus \dots \oplus \mathcal{H}_n)$.

THEOREM 2.7. *B satisfies (P) if and only if A_i satisfies (P), $i = 1, \dots, n$.*

PROOF. Let $n = 2$ and define $\mathfrak{T}(X) = A_1X - XA_2$ for X in $\mathcal{L}(\mathcal{H}_2, \mathcal{H}_1)$. Rosenblum's theorem [7, p. 8] implies that \mathfrak{T} is invertible, and the result follows from Lemma 2.5. The proof may now be completed by a straightforward induction on n .

By using Theorem 2.6 and an induction proof like that in Theorem 2.7, we obtain the following result.

THEOREM 2.8. *B has a local cross-section if and only if A_i has a local cross-section, $i = 1, \dots, n$.*

3. Normal operators and isometries. In this section we prove that if an operator is normal or isometric, then it satisfies (P) if and only if its spectrum is finite, in which case it also has a local cross-section.

THEOREM 3.1. *A normal operator N in $\mathcal{L}(\mathcal{H})$ satisfies (P) if and only if its spectrum is finite.*

PROOF. If $\sigma(N)$ is finite, then N is unitarily equivalent to a finite direct sum of distinct complex scalars, so the result follows from Theorem 2.6.

If $\sigma(N)$ is infinite, there exists a convergent sequence $\{\lambda_n\} \subset \sigma(N)$ and a sequence $\{D_n\}$ of open disks such that (1) D_n is centered at λ_n with radius r_n , (2) $\lim r_n = 0$, and (3) $D_n^- \cap D_m^- = \emptyset$ for $n \neq m$. Let E denote the spectral measure of N . Since D_n is open and $D_n \cap \sigma(N) \neq \emptyset$, then $E_n = E(D_n) \neq 0$ (see [7, Theorem 1.13, p. 18]), and we let e_n denote a unit vector in the range of E_n . Condition (3) implies that we may extend $\{e_n\}$ to an orthonormal basis $\{e_n\} \cup \{f_m\}$ for \mathcal{H} (where the f_m 's will be absent in some cases). We define a sequence $\{U_n\} \subset \mathcal{U}(\mathcal{H})$ as follows: $U_n(e_j) = e_j$ ($j \neq n, n + 1$), $U_n(e_n) = e_{n+1}$, $U_n(e_{n+1}) = e_n$, and $U_n(f_m) = f_m$ (all m). We have

$$\begin{aligned} & \|NU_n^*e_n - U_n^*Ne_n\| \\ & \leq \|(N - \lambda_{n+1})U_n^*e_n\| + \|(\lambda_{n+1} - \lambda_n)U_n^*e_n\| + \|\lambda_n U_n^*e_n - U_n^*Ne_n\| \\ & \leq \|(N - \lambda_{n+1})E_{n+1}e_{n+1}\| + |\lambda_{n+1} - \lambda_n| + \|(N - \lambda_n)E_n e_n\| \\ & \leq r_{n+1} + |\lambda_{n+1} - \lambda_n| + r_n. \end{aligned}$$

If x is in $E_n \mathcal{H} \ominus \langle e_n \rangle$, then

$$\begin{aligned} \|NU_n^*x - U_n^*Nx\| & \leq \|NU_n^*x - \lambda_n U_n^*x\| + \|\lambda_n U_n^*x - U_n^*Nx\| \\ & \leq 2\|(N - \lambda_n)E_n x\| \leq 2r_n \|x\|. \end{aligned}$$

Similarly, we have

$$\begin{aligned} \|(NU_n^* - U_n^*N)e_{n+1}\| &< r_n + |\lambda_n - \lambda_{n+1}| + r_{n+1} \quad \text{and} \\ \|NU_n^*x - U_n^*Nx\| &< 2r_{n+1}\|x\| \end{aligned}$$

for x in $E_{n+1}\mathcal{H} \ominus \langle e_{n+1} \rangle$. Since $NU_n^* - U_n^*N = 0$ on $(E_n\mathcal{H} \oplus E_{n+1}\mathcal{H})^\perp$, these relations imply that

$$\|U_n^*NU_n - N\| = \|U_n^*N - NU_n^*\| < 2(2r_n + 2r_{n+1} + |\lambda_{n+1} - \lambda_n|),$$

and therefore $\{U_n\}$ is in $\mathcal{S}(N)$. If W is a unitary operator such that $W^*NW = U_n^*NU_n$, then WU_n^* commutes with N and thus also with E_{n+1} (see [7, Fuglede's Theorem, p. 19]). We have $We_n = WU_n^*E_{n+1}e_{n+1} = E_{n+1}WU_n^*e_{n+1}$, and (3) now implies that $(We_n, e_n) = 0$. It follows that if $|\lambda| = 1$, then $\|W - \lambda\| \geq 2^{1/2}$, which completes the proof.

If M and N are in $\mathcal{L}(\mathcal{H})$, we denote by $\mathcal{Q}(N, M)$ the norm closed subalgebra of $\mathcal{L}(\mathcal{H})$ generated by N, M , and 1 .

THEOREM 3.2. *If N is a normal operator with finite spectrum, then π_N has a local cross-section $(\varphi_N, \mathfrak{B})$ such that for each M in \mathfrak{B} , $\varphi_N(M)$ is in $\mathcal{Q}(N, M)$.*

PROOF. We first give the proof for a projection P in $\mathcal{L}(\mathcal{H})$. Let $\mathfrak{B} = \{Q \text{ in } \mathcal{U}(P) \mid \|P - Q\| < 1\}$, and let $Q = \pi_P(U)$ be in \mathfrak{B} . With respect to the decomposition $\mathcal{H} = (\ker(P))^\perp \oplus \ker(P)$, we denote the matrix of U by $\begin{pmatrix} X & Y \\ Z & V \end{pmatrix}$; thus the matrix of Q is

$$\begin{pmatrix} X^*X & X^*Y \\ Y^*X & Y^*Y \end{pmatrix}.$$

Since $\|PU - UP\| = \|U^*PU - P\| = \|Q - P\| < 1$, a matrix calculation of $PU - UP$ shows that $\|Z\| < 1$ and $\|Y\| < 1$. Now $\|1 - V^*V\| = \|Y^*Y\| = \|Y\|^2 < 1$, and similarly $\|1 - XX^*\| = \|YY^*\| < 1$, $\|1 - VV^*\| = \|ZZ^*\| < 1$, and $\|1 - X^*X\| = \|Z^*Z\| < 1$; thus, the polar decomposition implies that X and V are invertible. In particular, X^*X and $1 - Y^*Y = V^*V$ are invertible, and we define $\varphi_P(Q)$ by the matrix

$$\begin{bmatrix} (X^*X)^{1/2} & (X^*X)^{-1/2}X^*Y \\ -(1 - Y^*Y)^{-1/2}Y^*X & (1 - Y^*Y)^{1/2} \end{bmatrix}.$$

Now $\varphi_P(Q)$ is independent of the choice of U since it is defined in terms of the matrix components of Q . Thus $\varphi_P(Q)$ is well defined and it follows from the identity

$$\varphi_P(Q) = ((X^*X)^{-1/2}X^* \oplus (V^*V)^{-1/2}V^*)U$$

that $\varphi_P(Q)$ is unitary, that $\varphi_P(P) = 1$ and that $\pi_P(\varphi_P(Q)) = Q$. It is clear from the matrix of Q that φ_P is continuous on \mathfrak{B} . That $\varphi_P(Q)$ is in $\mathcal{Q}(P, Q)$ follows from the identity

$$\begin{aligned} \varphi_P(Q) &= (PQP)^{1/2} + (1 - P + PQP)^{-1/2}PQ(1 - P) \\ &\quad + ((1 - P)(1 - Q(1 - P)))^{1/2} \\ &\quad - (P + (1 - P)(1 - Q(1 - P)))^{-1/2}(1 - P)QP. \end{aligned}$$

Indeed, the Weierstrass approximation theorem and the functional calculus for positive operators imply that each term in the preceding sum is in $\mathcal{Q}(P, Q)$.

If $\sigma(N) = \{\lambda_1, \dots, \lambda_n\}$, then N is diagonalizable, and thus there exist polynomials $p_1(z), \dots, p_n(z)$, such that $p_1(N), \dots, p_n(N)$ are pairwise orthogonal projections and such that $\sum_{i=1}^n p_i(N) = 1$ and $N = \sum_{i=1}^n \lambda_i p_i(N)$. Let $\delta > 0$ be a positive number such that if M is in $\mathcal{U}(N)$ and $\|M - N\| < \delta$, then $\|p_i(M) - p_i(N)\| < 1, i = 1, \dots, n$. Let $\mathfrak{B} = \{M \text{ in } \mathcal{U}(N) \mid \|M - N\| < \delta\}$; thus, if M is in \mathfrak{B} , then for each fixed $i, p_i(M)$ is in the domain of the local cross-section $(\varphi_i, \mathfrak{B}_i)$ for $p_i(N)$ that was defined above. We now define $\varphi_N(M) = \sum_{i=1}^n p_i(N)\varphi_i(p_i(M))p_i(M)$. It is clear that $\varphi_N(M)$ is in $\mathcal{Q}(N, M)$ and that φ_N is continuous on \mathfrak{B} . Since for each i we have $\varphi_i(p_i(M))^*p_i(N)\varphi_i(p_i(M)) = p_i(M)$, a calculation shows that $\varphi_N(M)$ is a unitary operator and that $\varphi_N(N) = 1$ and $\pi_N(\varphi_N(M)) = M$. Thus $(\varphi_N, \mathfrak{B})$ is a cross-section for N and the proof is complete.

Let \mathcal{Q} denote a C^* -subalgebra of $\mathcal{L}(\mathcal{H})$ and let $\mathcal{U}(\mathcal{Q}) = \mathcal{U}(\mathcal{H}) \cap \mathcal{Q}$. For T in \mathcal{Q} , let $\mathcal{U}_{\mathcal{Q}}(T)$ denote the $\mathcal{U}(\mathcal{Q})$ orbit of T in \mathcal{Q} and define $\pi(\mathcal{Q})_T: \mathcal{U}(\mathcal{Q}) \rightarrow \mathcal{U}_{\mathcal{Q}}(T)$ by $\pi(\mathcal{Q})_T(U) = U^*TU$.

COROLLARY 3.3. *If N is a normal operator in \mathcal{Q} with finite spectrum, then $\pi(\mathcal{Q})_N$ has a local cross-section.*

REMARK. In the case $\mathcal{Q} = \mathcal{L}(\mathcal{H})$, Corollary 3.3 is a direct consequence of Theorem 2.8. Note also that since any two projections differ in norm by at most 1, then the cross-section given in Theorem 3.2 for a projection P is densely defined in $\mathcal{U}(P)$; however, it is easy to prove that this cross-section cannot be extended to all of $\mathcal{U}(P)$. In [2] it was proven that if T has a cross-section, then each path in $\mathcal{U}(T)$ can be lifted to a path in $\mathcal{U}(\mathcal{H})$. By adapting the argument used in proving Theorem 3.1, it is not difficult to prove that there exists a diagonalizable normal operator (with infinite spectrum) which does not enjoy this path lifting property.

In the sequel \mathcal{H}_1 and \mathcal{H}_2 denote separable complex Hilbert spaces and $\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2$.

COROLLARY 3.4. *If N is a normal operator in $\mathcal{L}(\mathcal{H}_1)$ with infinite spectrum and if T is in $\mathcal{L}(\mathcal{H}_2)$, then $N \oplus T$ does not satisfy (P).*

PROOF. We use the notation of the proof of Theorem 3.1. Suppose that $N \oplus T$ does satisfy (P) and let $\{W_n\} \subset \mathcal{U}(\mathcal{H})$ denote a sequence that

re-implements $\{U_n \oplus 1\}$ for $N \oplus T$. Now $V_n = W_n(U_n^* \oplus 1)$ commutes with $N \oplus T$, and if we denote the matrix of V_n by

$$\begin{pmatrix} A_n & B_n \\ C_n & D_n \end{pmatrix},$$

then a calculation shows that A_n commutes with N . Fuglede's Theorem [7, Theorem 1.13] implies that A_n commutes with E_{n+1} and thus $A_n U_n e_n = A_n E_{n+1} e_{n+1} = E_{n+1} A_n e_{n+1}$; in particular $(A_n U_n e_n, e_n) = 0$. Now $\|A_n U_n - 1\| > |((A_n U_n - 1)e_n, e_n)| = 1$; however, since $V_n(U_n \oplus 1) = W_n \rightarrow 1$, a matrix calculation shows that $A_n U_n \rightarrow 1$, which is a contradiction.

If T is in $\mathcal{L}(\mathcal{H})$, let $R_\epsilon(T)$ denote the set of all reducing essential eigenvalues of T (see [8]).

LEMMA 3.5. *Let T be in $\mathcal{L}(\mathcal{H})$ and suppose $R_\epsilon(T)$ is nonempty. Then there exists a sequence $\{U_n\}$ in $\mathfrak{S}(T)$ such that for each n , $\inf_{|\lambda|=1} \|U_n - \lambda\| \geq 2^{1/2}$.*

PROOF. Let λ be in $R_\epsilon(T)$. Corollary 4.9 of [8] implies that there exists a sequence $\{W_n\}$ of unitary operators from \mathcal{H} onto $\mathcal{H} \oplus \mathcal{H}$ such that

$$W_n^*(T \oplus \lambda)W_n \rightarrow T.$$

Let V be a unitary operator on \mathcal{H} such that $\inf_{|\lambda|=1} \|V - \lambda\| \geq 2^{1/2}$ and let $U_n = W_n^*(1 \oplus V)W_n$. Then

$$\begin{aligned} \|U_n^* T U_n - T\| &\leq \|U_n^*(T - W_n^*(T \oplus \lambda)W_n)U_n\| \\ &\quad + \|U_n^* W_n^*(T \oplus \lambda)W_n U_n - T\| \\ &= 2\|T - W_n^*(T \oplus \lambda)W_n\| \rightarrow 0. \end{aligned}$$

Therefore, $\{U_n\}$ satisfies the required conditions, and the proof is complete.

LEMMA 3.6. *Let V be an irreducible operator in $\mathcal{L}(\mathcal{H}_1)$. Suppose there exist a sequence $\{U_n\}$ in $\mathfrak{S}(V)$ and a positive number ϵ such that for each n , $\inf_{|\lambda|=1} \|U_n - \lambda\| \geq \epsilon$. Then for each T in $\mathcal{L}(\mathcal{H}_2)$, $V \oplus T$ does not satisfy (P).*

PROOF. If $V \oplus T$ does satisfy (P), then there exists a sequence $\{W_n\} \subset \mathcal{U}(\mathcal{H})$ that re-implements $\{U_n \oplus 1\}$ for $V \oplus T$. If we denote the operator matrix of $(U_n \oplus 1)W_n^*$ by

$$\begin{pmatrix} A_n & B_n \\ C_n & D_n \end{pmatrix},$$

then since $W_n^* \rightarrow 1$, a calculation of $W_n^* = (U_n^* \oplus 1)(U_n \oplus 1)W_n^*$ shows that $U_n^* A_n \rightarrow 1$ and $C_n \rightarrow 0$. Since $(U_n \oplus 1)W_n^*$ commutes with $V \oplus T$ and $(V \oplus T)^*$, A_n commutes with V and V^* , and the irreducibility of V implies that $A_n = \lambda_n$. Now $\lim |\lambda_n|^2 - 1| = \lim \|C_n^* C_n\| = 0$, and thus $|\lambda_n| \rightarrow 1$. Since there exists a convergent subsequence $\lambda_{n_k} \rightarrow \lambda$ ($|\lambda| = 1$), it follows that

$\lim \|U_{n_k} - \lambda\| = 0$, which is a contradiction.

The preceding two lemmas now yield the following result.

THEOREM 3.5. *If V is an irreducible operator in $\mathcal{L}(\mathcal{H}_1)$ and $R_e(V)$ is nonempty, then for each operator T in $\mathcal{L}(\mathcal{H}_2)$, $V \oplus T$ does not satisfy (P).*

EXAMPLE 3.6. Using the preceding theorem we will show that there exist compact quasi-nilpotent operators that do not satisfy (P). Indeed, if K is compact, then 0 is in $R_e(K)$, and thus if K is irreducible, then K does not satisfy (P). One example of this type is the Volterra integral operator (see [7, Proposition 4.11, p. 82]). Let $\{e_n\}_{n=-\infty}^{+\infty}$ denote an orthonormal basis for \mathcal{H} , and let $\{\lambda_n\}_{n=-\infty}^{+\infty}$ denote a sequence of nonzero complex numbers. We define an injective bilateral shift W by $We_n = \lambda_n e_{n+1}$. If W is compact (or, equivalently, if $\lim_{n \rightarrow \infty} \lambda_n = \lim_{n \rightarrow -\infty} \lambda_n = 0$), then W is quasi-nilpotent (see [5, p. 48]), and since the sequence $\{\lambda_n\}$ is nonperiodic, it follows from [5, Problem 129] that W is irreducible. Therefore each injective bilateral weighted shift that is compact does not satisfy (P).

COROLLARY 3.6. *If V has a cross-section and $R_e(V)$ is nonempty, then V has a nontrivial reducing subspace.*

COROLLARY 3.7. *If V is an irreducible hyponormal operator in $\mathcal{L}(\mathcal{H}_1)$ ($\dim \mathcal{H}_1 > 1$), then for each operator T in $\mathcal{L}(\mathcal{H}_2)$, $V \oplus T$ does not satisfy (P).*

PROOF. If V is a hyponormal operator in $\mathcal{L}(\mathcal{H}_1)$ and \mathcal{H}_1 is finite dimensional, then V is normal, and thus if $\dim(\mathcal{H}_1) > 1$, then V is reducible. We may therefore assume that $\dim(\mathcal{H}_1) = \aleph_0$. Theorem 3.10 of [8] implies that if V is hyponormal, then $R_e(V)$ is nonempty. The result now follows from Theorem 3.5.

REMARK. The preceding result applies in particular if V is a unilateral shift of multiplicity one. If $\|T\| < 1$, the conclusion that $V \oplus T$ does not satisfy (P) may be obtained from Lemma 2.5. Indeed, for $\|T\| < 1$, it is not difficult to prove that the operator τ on $\mathcal{L}(\mathcal{H}_2, \mathcal{H}_1)$ defined by $\tau(X) = VX - XT$ (X in $\mathcal{L}(\mathcal{H}_2, \mathcal{H}_1)$) is left invertible (although not invertible) in $\mathcal{L}(\mathcal{L}(\mathcal{H}_2, \mathcal{H}_1))$ and is thus bounded below.

The preceding corollary implies that if a hyponormal operator satisfies (P), then it has no minimal infinite dimensional reducing subspace. We conjecture that a hyponormal operator satisfies (P) if and only if its spectrum is finite.

COROLLARY 3.8. *An isometry V in $\mathcal{L}(\mathcal{H})$ satisfies (P) if and only if its spectrum is finite, in which case V has a local cross-section.*

PROOF. If V is unitary, the result follows from Theorem 3.1 and Theorem 3.2. If V is nonunitary, the von Neumann decomposition implies that V has a unilateral shift of multiplicity one as a direct summand. Corollary 3.7 now

implies that V does not satisfy (P), and since the spectrum of each nonunitary isometry is infinite, the proof is complete.

4. Nonnormal operators with cross-sections. In this section we give examples of nonnormal operators with cross-sections. Let T be an operator in $\mathcal{L}(\mathcal{H})$ with closed range and let $C^*(T)$ denote the C^* -subalgebra of $\mathcal{L}(\mathcal{H})$ generated by T . We will make use of the following simple observations: If $T = UP$ denotes the polar decomposition of T , then P is in $C^*(T)$; if $\{U_n\}_{n=0}^\infty \subset \mathcal{U}(\mathcal{H})$ and $U_n^*TU_n \rightarrow U_0^*TU_0$, then $U_n^*SU_n \rightarrow U_0^*SU_0$ for each S in $C^*(T)$; if T commutes with U_0 , then S commutes with U_0 for each S in $C^*(T)$. Let $P = 0 \oplus 1$ in $\mathcal{L}(\mathcal{H} \oplus \mathcal{H})$; in the sequel $(\varphi, \mathfrak{B}(P, 1))$ will denote a local cross-section for P (cf. Theorem 3.2). If T and S are in $\mathcal{L}(\mathcal{H})$, $S \neq 0$, then $M \equiv M(T, S)$ denotes the nonnormal operator on $\mathcal{H} \oplus \mathcal{H}$ whose operator matrix is $\begin{pmatrix} T & 0 \\ S & 0 \end{pmatrix}$.

LEMMA 4.1. *Let T and S be in $\mathcal{L}(\mathcal{H})$ and suppose that $T^*T + S^*S$ is invertible. If $\varepsilon > 0$, then there exists $\delta > 0$ such that if X is in $\mathfrak{B}(M, \delta)$, then (i) $\|P_{\ker(X)} - P\| < 1$ and (ii) $\|VXV^* - M\| < \varepsilon$, where $V = \varphi(P_{\ker(X)})$.*

PROOF. Let f be a real continuous function on $[0, \|M\|]$. It follows from the functional calculus that the mapping $F: \mathcal{U}(M) \rightarrow \mathcal{L}(\mathcal{H})$ defined by $F(X) = f((X^*X)^{1/2})$ is norm continuous. Since M has closed range, 0 is an isolated point of $\sigma((M^*M)^{1/2})$, and thus the characteristic function of $\{0\}$ with respect to $\sigma((M^*M)^{1/2})$ can be extended to a function f that is continuous on $[0, \|M\|]$. Now $F(X) = P_{\ker(X)}$ for each X in $\mathcal{U}(M)$, and since F is continuous, there exists $\delta_1 > 0$ such that if $\|X - M\| < \delta_1$ (X in $\mathcal{U}(M)$), then $\|P_{\ker(X)} - P\| < 1$. Thus $V = \varphi(P_{\ker(X)})$ is defined and

$$\|VXV^* - M\| \leq 2\|V - 1\| \|M\| + \|X - M\|.$$

Since φ is continuous, there exists $\delta < \min\{\varepsilon/2, \delta_1\}$ such that if X is in $\mathfrak{B}(M, \delta)$, then $\|\varphi(P_{\ker(X)}) - 1\| < \varepsilon/(4\|M\|)$. With this δ , (i) and (ii) are satisfied.

THEOREM 4.2. *Suppose that T has a local cross-section and that S satisfies the following conditions: (i) $T^*T + S^*S$ is invertible, (ii) S is right invertible, and (iii) S is in $C^*(T)$. Then $M = M(T, S)$ has a local cross-section.*

PROOF. Let $(\psi, \mathfrak{B}(T, \varepsilon))$ denote a cross-section for T . We use the notation and results of Lemma 4.1. Thus, if X is in $\mathfrak{B}(M, \delta)$ and $V = \varphi(P_{\ker(X)})$, then $\|VXV^* - M\| < \varepsilon$. Now $V^*PV = P_{\ker(X)}$, so $P_{\ker(VXV^*)} = VP_{\ker(X)}V^* = P$, and thus $VXV^* = WMW^* = M(T_1, S_1)$ where W is unitary and $T_1^*T_1 + S_1^*S_1$ is invertible. It follows that $WP = PW$, and thus there exist unitary operators Z and Y such that $T_1 = Z^*TZ$ and $S_1 = Y^*SZ$. Since $\|T_1 - T\| < \|M(T_1, S_1) - M\| < \varepsilon$, we may now define

$$\rho(X) = (\psi(T_1) \oplus \psi(T_1)Z^*Y)\varphi(P_{\ker(X)}).$$

We will show that $(\rho, \mathfrak{B}(M, \delta))$ is a local cross-section for M .

To check that ρ is well defined, suppose there exist unitary operators Z_1 and Y_1 such that $Z_1^*TZ_1 = Z^*TZ$ and $Y_1^*SZ_1 = Y^*SZ$. Since T commutes with Z_1Z^* , so does S , and thus $Y_1^*Z_1Z^*SZ = Y_1^*SZ_1 = Y^*SZ = Y^*ZZ^*SZ$. Now (ii) implies that $Y_1^*Z_1 = Y^*Z$, and therefore ρ is well defined. Since $\psi(T_1)^*T\psi(T_1) = T_1 = Z^*TZ$, (iii) implies that $Z\psi(T_1)^*$ commutes with S . Now we have

$$\begin{aligned} \rho(X)^*M\rho(X) &= V^*M(\psi(T_1)^*T\psi(T_1), Y^*Z\psi(T_1)^*S\psi(T_1))V \\ &= V^*M(T_1, S_1)V = X. \end{aligned}$$

Since $\rho(M) = 1$, to complete the proof it suffices to verify that ρ is continuous. Suppose that $X_n \rightarrow X_0$ in $\mathfrak{B}(M, \delta)$. Lemma 4.1 implies that $\varphi(P_{\ker(X_n)}) \rightarrow \varphi(P_{\ker(X_0)})$; moreover, if we set $V_n = \varphi(P_{\ker(X_n)})$ for $n \geq 0$, then $V_nX_nV_n^* \rightarrow V_0X_0V_0^*$, and each $V_nX_nV_n^*$ is of the form $M_n = M(T_n, S_n)$, where $T_n^*T_n + S_n^*S_n$ is invertible. As above, there exist sequences $\{Z_n\}, \{Y_n\} \subset \mathcal{U}(\mathcal{H})$ such that $T_n = Z_n^*TZ_n$ and $S_n = Y_n^*SZ_n$ ($n \geq 0$). Since $T_n \rightarrow T_0$ in $\mathfrak{B}(T, \epsilon)$, then $\psi(T_n) \rightarrow \psi(T_0)$; it now suffices to prove that $Z_n^*Y_n \rightarrow Z_0^*Y_0$. From the remarks at the beginning of this section, (ii) and (iii) imply that there exists an operator R in $C^*(T)$ such that $SR = 1$, and since $Z_n^*TZ_n \rightarrow Z_0^*TZ_0$, it follows that $Z_n^*RZ_n \rightarrow Z_0^*RZ_0$. Since $Y_n^*Z_nZ_n^*SZ_n = S_n \rightarrow S_0 = Y_0^*Z_0Z_0^*SZ_0$, it now follows that $Y_n^*Z_n \rightarrow Y_0^*Z_0$, which completes the proof.

THEOREM 4.3. *If T has a local cross-section and S is an isometry in $\mathcal{L}(\mathcal{H})$, then $M = M(T, S)$ has a local cross-section.*

PROOF. Let $(\psi, \mathfrak{B}(T, \epsilon))$ denote a local cross-section for T , where $0 < \epsilon < \frac{1}{2}$. We use the notation and results of Lemma 4.1. If X is in $\mathfrak{B}(M, \delta)$ and $V = \varphi(P_{\ker(X)})$, then, as in the proof of the preceding result, $VXV^* = M(T_1, S_1)$, where $T_1^*T_1 + S_1^*S_1$ is invertible. It follows as before that there exist unitary operators Z and Y such that $T_1 = Z^*TZ$ and $S_1 = Y^*SZ$. Since $\|Z^*TZ - T\| < \epsilon$, $\psi(T_1)$ is defined. We next define a unitary operator W such that $W^*S = U \equiv Y^*SZ\psi(T_1)^*$. For each t in \mathcal{H} we set $W_1(S_t) = Ut$; W_1 is a linear mapping of $S\mathcal{H}$ isometrically onto $U\mathcal{H}$. Since $\|Y^*SZ - S\| < \epsilon$, then $\|P_{U\mathcal{H}} - P_{S\mathcal{H}}\| = \|UU^* - SS^*\| < 2\epsilon < 1$. Let $(\gamma, \mathfrak{B}(P_{S\mathcal{H}}, 1))$ denote a local cross-section for $P_{S\mathcal{H}}$. Now $\gamma(P_{U\mathcal{H}})^*P_{S\mathcal{H}} = P_{U\mathcal{H}}\gamma(P_{U\mathcal{H}})^*$, and we define $W_2: (S\mathcal{H})^\perp \rightarrow (U\mathcal{H})^\perp$ by $W_2 = \gamma(P_{U\mathcal{H}})^*(S\mathcal{H})^\perp$. Thus W_1 and W_2 together define a unitary operator W_3 on \mathcal{H} such that if we set $W = W_3^*$, then $W^*S\psi(T_1) = Y^*SZ$. Furthermore, W is independent of Y and Z since W is defined in terms of $S_1 = Y^*SZ$. We now define

$$\tau(X) = (\psi(T_1) \oplus W)\varphi(P_{\ker(X)}),$$

and we will show that $(\tau, \mathfrak{B}(M, \delta))$ is a local cross-section. It is clear that $\pi_M \tau = 1_{\mathfrak{B}(M, \delta)}$ and that $\tau(M) = 1$. Suppose that $X_n \rightarrow X_0$ in $\mathfrak{B}(M, \delta)$. If $V_n = \varphi(P_{\ker(X_n)})$ ($n \geq 0$), then $V_n \rightarrow V_0$, $V_n X_n V_n^* \rightarrow V_0 X_0 V_0^*$, and each $V_n X_n V_n^*$ is of the form $M_n = M(Z_n^* T Z_n, Y_n^* S Z_n)$, where Z_n and Y_n are unitary. Since $Z_n^* T Z_n \rightarrow Z_0^* T Z_0$ in $\mathfrak{B}(T, \epsilon)$, then $\psi(Z_n^* T Z_n) \rightarrow \psi(Z_0^* T Z_0)$. Thus it suffices to prove that $W_n \rightarrow W_0$, where W_n satisfies $W_n^* S = U_n \equiv Y_n^* S Z_n \psi(Z_n^* T Z_n)^*$ ($n > 0$), and $W_n^* |(S \mathfrak{J} \mathcal{C})^\perp = \gamma(P_{U_n \mathfrak{J} \mathcal{C}})^* |(S \mathfrak{J} \mathcal{C})^\perp$. Let $P_n = P_{U_n \mathfrak{J} \mathcal{C}}$; since $Y_n^* S Z_n \rightarrow Y_0^* S Z_0$, we have $\gamma(P_n) \rightarrow \gamma(P_0)$. For $\alpha > 0$, let N be a positive integer such that if $n > N$, then $\|U_n - U_0\| < \alpha$ and $\|\gamma(P_n) - \gamma(P_0)\| < \alpha$. If z is in $\mathfrak{J} \mathcal{C}$, $\|z\| = 1$, then $z = St + z'$, where z' is in $(S \mathfrak{J} \mathcal{C})^\perp$ and $\|t\|^2 + \|z'\|^2 = 1$. Now

$$\begin{aligned} \|(W_n - W_0)^* z\|^2 &= \|(U_n - U_0)t + (\gamma(P_n)^* - \gamma(P_0)^*)z'\|^2 \\ &< \alpha^2(\|t\| + \|z'\|)^2 \leq 2\alpha^2. \end{aligned}$$

Thus $\|W_n - W_0\| \leq 2^{1/2}\alpha$, and the proof is complete.

Straightforward modifications of the proofs of Theorem 4.2 and Theorem 4.3 yield the following results, whose proofs are omitted.

PROPOSITION 4.4. *If T satisfies (P) and S satisfies conditions (i)–(iii) of Theorem 4.2, then $M = M(T, S)$ satisfies (P).*

PROPOSITION 4.5. *If T satisfies (P) and S is an isometry in $\mathcal{L}(\mathfrak{J} \mathcal{C})$, then $M = M(T, S)$ satisfies (P).*

REMARK. The preceding results, together with those of §§2 and 3, provide examples of nonnormal operators that satisfy (P) or have local cross-sections. If we consider the classes of (1) algebraic operators, (2) operators with finite spectrum, (3) operators with a closed unitary orbit, (4) operators satisfying the sequential unitary lifting property, and (5) operators having a local cross-section, then our results, together with those of [6], show that for normal operators and isometries these five classes coincide. For arbitrary nonnormal operators, we have not been able to determine the relationships between all of these classes. In this regard, the following questions seem pertinent.

1. If an operator satisfies the sequential unitary lifting property, does it have a local cross-section; does it satisfy the unitary path lifting property?

2. If T satisfies (P), is $\mathcal{U}(T)$ closed? An affirmative answer to this question would of course render Proposition 2.2 vacuous; on the other hand, the answer is not obviously affirmative even for the operators given by the results of this section.

3. Does every operator that is unitarily equivalent to an $n \times n$ operator matrix, each of whose entries is a scalar multiple of $1_{\mathfrak{J} \mathcal{C}}$ have a local cross-section? Theorem 4.2 implies an affirmative answer when $n = 2$. We

note also that not every operator with a local cross-section is unitarily equivalent to such a scalar valued operator matrix. If U is a unilateral shift of multiplicity one in $\mathcal{L}(\mathcal{H})$ and $T = M(0, U)$, then Theorem 4.3 implies that T has a local cross-section; moreover, $\dim(\ker(T) \cap \ker(T^*)) = 1$. Now T is not unitarily equivalent to a scalar valued operator matrix; indeed, if S is unitarily equivalent to an $n \times n$ scalar valued operator matrix, then $\dim(\ker(S) \cap \ker(S^*)) = 0$ or $\dim(\ker(S) \cap \ker(S^*)) = \aleph_0$ (since S is unitarily equivalent to $A \oplus A \oplus A \oplus \cdots \oplus A \oplus \cdots$, where A is an operator on an n -dimensional Hilbert space).

4. Does each algebraic operator satisfy (P)? If Proposition 4.5 could be extended so that the isometry S may be replaced by an arbitrary bounded operator acting between two possibly different Hilbert spaces, then we could prove that each algebraic operator satisfies (P).

ADDED IN PROOF. In a forthcoming sequel to this paper, question 2 is answered affirmatively. As a consequence, it is proved that each operator satisfying (P) is unitarily equivalent to an operator of the form $A \oplus B \oplus \cdots \oplus B \oplus \cdots$, where A and B are operators on finite dimensional spaces. The conjecture on hyponormal operators following Corollary 3.7 is proved, and question 4 is answered negatively.

REFERENCES

1. H. O. Cordes, *On a class of C^* -algebras*, Math. Ann. **170** (1967), 283–313. MR 35 #749.
2. L. A. Fialkow, *A note on limits of unitarily equivalent partial isometries* (preprint).
3. R. Gellar and L. Page, *Limits of unitarily equivalent normal operators*, Duke Math. J. **41** (1974), 319–322. MR 49 #3581.
4. P. R. Halmos, *Quasitriangular operators*, Acta Sci. Math. (Szeged) **29** (1968), 283–293. MR 38 #2627.
5. ———, *A Hilbert space problem book*, Van Nostrand, Princeton, N.J., 1967. MR 34 #8178.
6. ———, *Limits of shifts*, Acta Sci. Math. (Szeged) **34** (1973), 131–139. MR 49 #3576.
7. H. Radjavi and P. Rosenthal, *Invariant subspaces*, Springer-Verlag, New York, 1973.
8. N. Salinas, *Reducing essential eigenvalues*, Duke Math. J. **40** (1973), 561–580.

DEPARTMENT OF MATHEMATICS, WESTERN MICHIGAN UNIVERSITY, KALAMAZOO, MICHIGAN 49008