

OPTIMAL STOCHASTIC SWITCHING AND THE DIRICHLET PROBLEM FOR THE BELLMAN EQUATION

BY

LAWRENCE C. EVANS AND AVNER FRIEDMAN¹

ABSTRACT. Let L^i be a sequence of second order elliptic operators in a bounded n -dimensional domain Ω , and let f^i be given functions. Consider the problem of finding a solution u to the Bellman equation $\sup_i (L^i u - f^i) = 0$ a.e. in Ω , subject to the Dirichlet boundary condition $u = 0$ on $\partial\Omega$. It is proved that, provided the leading coefficients of the L^i are constants, there exists a unique solution u of this problem, belonging to $W^{1,\infty}(\Omega) \cap W_{loc}^{2,\infty}(\Omega)$. The solution is obtained as a limit of solutions of certain weakly coupled systems of nonlinear elliptic equations; each component of the vector solution converges to u . Although the proof is entirely analytic, it is partially motivated by models of stochastic control. We solve also certain systems of variational inequalities corresponding to switching with cost.

1. Introduction. Consider a sequence of linear elliptic partial differential operators

$$L^k u \equiv - \sum_{i,j=1}^n a_{ij}^k(x) \frac{\partial^2 u}{\partial x_i \partial x_j} - \sum_{i=1}^n b_i^k(x) \frac{\partial u}{\partial x_i} + c^k(x) u \quad (k = 1, 2, \dots)$$

in a bounded domain $\Omega \subset R^n$. We assume:

$$\partial\Omega \in C^{2+\beta} \quad \text{for some } \beta > 0, \quad (1.1)$$

$$\sum_{i,j=1}^n a_{ij}^k(x) \xi_i \xi_j \geq \gamma |\xi|^2 \quad (x \in \Omega, \xi \in R^n, \gamma > 0), \quad (1.2)$$

$$|D^\alpha a_{ij}^k(x)|, |D^\alpha b_i^k(x)|, |D^\alpha c^k(x)| < C \quad (x \in \Omega, 1 \leq i, j \leq n; 0 \leq |\alpha| \leq 2, C > 0), \quad (1.3)$$

and

$$c^k(x) \geq c_0 \quad (x \in \Omega; c_0 > 0), \quad (1.4)$$

where γ, C, c_0 are constants independent of k . (The hypothesis $c_0 > 0$ is convenient, but $c_0 \geq 0$ is sufficient; see Remark 1, §6.)

Received by the editors April 16, 1978.

AMS (MOS) subject classifications (1970). Primary 35J60, 49A20; Secondary 35J25, 47H05, 47H15, 60H10, 93E20.

Key words and phrases. Bellman equation, elliptic operators, nonlinear elliptic equation, accretive operators, stochastic differential equations.

¹This work is partially supported by National Science Foundation Grants MCS 77-09152 and MC 575-21416 A01.

© 1979 American Mathematical Society
 0002-9947/79/0000-0417/\$07.25

Suppose further that the $f^k(x)$ are given functions for $x \in \Omega$, satisfying

$$|D^\alpha f^k(x)| \leq C \quad (x \in \Omega, 0 \leq |\alpha| \leq 2, C > 0), \tag{1.5}$$

where C is a constant independent of k .

In this paper we investigate the nonlinear partial differential equation

$$\sup_k (L^k u(x) - f^k(x)) = 0 \quad \text{a.e. for } x \in \Omega, \tag{1.6}$$

with the boundary condition

$$u = 0 \quad \text{on } \partial\Omega. \tag{1.7}$$

Equation (1.6) is called the *Bellman equation of dynamic programming*, arising in stochastic optimal control theory (see §7). It was studied by Krylov [13] and Nisio [14] in the case $\Omega = R^n$. Krylov proved (under assumptions similar to (1.2)–(1.5) and with the additional condition that c_0 is sufficiently large) that there exists a unique solution u in R^n such that, for some $\varphi(x) = (1 + \mu|x|^2)^{1/2}$, $\mu > 0$, $e^{-\varphi(x)}u(x)$ belongs to $W^{2,p}(R^n)$, $\forall p < \infty$. His proof was somewhat simplified by Nisio [14] (see also [3]). As is pointed out in [14], [3], the existence of a solution to the Bellman equation in a bounded domain is an open problem, except when $n = 2$.

In this paper we establish the existence of a unique solution u of (1.6), (1.7) satisfying $u \in W^{1,\infty}(\Omega) \cap W^{2,\infty}_{\text{loc}}(\Omega)$. We assume the conditions (1.1)–(1.5) and the additional restriction

$$a_{ij}^k \text{ are constants.} \tag{1.8}$$

Unlike Krylov’s approach, our proof does not use probabilistic methods; it is however partially motivated by probabilistic considerations. In fact, we approximate the solution of (1.6), (1.7) by the system of equations

$$\begin{aligned} L^k u^k + \beta_\varepsilon(u^k - u^{k+1}) &= f^k(x) \quad (1 \leq k \leq m, \quad x \in \Omega), \\ u^k &= 0 \quad \text{on } \partial\Omega, \end{aligned} \tag{1.9}$$

where $u^{m+1} \equiv u^1$. Here β_ε represents a “penalty” term: $\beta_\varepsilon(t) = 0$ if $t < 0$, $\beta_\varepsilon(t) \rightarrow \infty$ if $t > 0$, $\varepsilon \rightarrow 0$. Probabilistically, the solution component $u^k(x)$ represents the optimal cost starting at $x \in \Omega$ in state k of the same cost functional as in Krylov’s work plus an additional cost for every switching from one generator L^i to the next one L^{i+1} . We shall explain this model more precisely in §7.

In §2 we study the system (1.9) and prove that it has a unique classical solution. §3 comprises an a priori estimate on the $W^{1,\infty}(\Omega)$ -norm of the solution; the estimate does not depend on m and ε . In §4 we derive a priori estimate on the $W^{2,\infty}(\Omega_0)$ -norm of the solution of (1.9), $\bar{\Omega}_0 \subset \Omega$. It is only here that the condition (1.8) is used.

In §5 we take $\varepsilon \rightarrow 0$ in (1.9) and show that each $u^k = u^{k,\varepsilon}$ converges to the same function v^m , v^m solving the Bellman problem (1.6), (1.7) (where k

ranges over the indices $1, 2, \dots, m$). Finally, in §6 we prove that, as $m \rightarrow \infty$, v^m converges to the solution of (1.6), (1.7). These convergence arguments use nonlinear functional analytic methods and, in particular, certain accretive operator techniques.

In §7 we study the system

$$\begin{aligned} L^i u_i + \beta_\varepsilon(u_i - u_{i+1} - k_i) &= f_i & (x \in \Omega), \\ u_i &= 0 & \text{on } \partial\Omega \quad (1 < i < m) \end{aligned} \tag{1.10}$$

where the k_i are positive numbers, and also the limit case, as $\varepsilon \rightarrow 0$, a system of variational inequalities

$$\begin{aligned} L^i u_i + \beta(u_i - u_{i+1} - k_i) &\ni f^i & \text{a.e. in } \Omega, \\ u_i &= 0 & \text{on } \partial\Omega. \end{aligned} \tag{1.11}$$

(Here $\beta(t)$ is the maximal monotone graph: $\beta(t) = \{0\}$ if $t < 0$, $\beta(0) = [0, \infty]$.) We give probabilistic interpretation for these problems and prove that each component u_i of the solution of (1.11) converges to the solution v^m (of (1.6), (1.7) for L^1, \dots, L^m) as $(k_1, \dots, k_m) \rightarrow 0$.

We also show that for fixed and positive k_i , the solution of (1.10) converges to the solution of (1.11), as $\varepsilon \rightarrow 0$. This is proved under weaker assumptions than (1.3), (1.5) and without the restrictive condition (1.8).

In §8 we specialize to the case $m = 2$. It was proved by Brezis and Evans [5] that the corresponding Bellman equation

$$\begin{aligned} \max(L^1 u(x) - f^1(x), L^2 u(x) - f^2(x)) &= 0 & \text{a.e. in } \Omega, \\ u &= 0 & \text{on } \partial\Omega \end{aligned} \tag{1.12}$$

has a solution u in $C^{2+\alpha}(\Omega_0)$ for some $\alpha > 0$ and any domain $\Omega_0, \bar{\Omega}_0 \subset \Omega$. Here we show (without making the restriction (1.8)) that the solution u_1, u_2 of (1.11) with $m = 2$ satisfies: $u_i(x) \rightarrow u(x)$ as $(k_1, k_2) \rightarrow 0$.

NOTE ADDED IN PROOF. P. L. Lions (in work to appear) has recently removed the restriction hypothesis (1.8). His method gives $W^{2,\infty}(\Omega)$ estimates for the approximating system (2.3), (2.4) and is based on a nontrivial extension of our proof of Lemma 4.1.

2. Construction of approximate solutions. Let $\beta_\varepsilon(t)$ ($\varepsilon > 0, t$ real) be a C^∞ function in t such that

$$\begin{aligned} \beta_\varepsilon(t) &= 0 & \text{if } t < 0, \\ \beta_\varepsilon(t) &\rightarrow \infty & \text{if } t > 0, \quad \varepsilon \rightarrow 0, \\ \beta'_\varepsilon(t) &\geq 0, & \beta''_\varepsilon(t) \geq 0, \end{aligned} \tag{2.1}$$

and

$$|t\beta'_\varepsilon(t) - \beta_\varepsilon(t)| \leq C \quad (C \text{ constant}). \tag{2.2}$$

We can take, for example,

$$\beta_\varepsilon(t) = \frac{t - \varepsilon}{\varepsilon} \quad \text{if } 2\varepsilon < t < \infty,$$

$$= 0 \quad \text{if } t \leq 0,$$

and

$$0 < \beta_\varepsilon(t) < 1, \quad 0 \leq \beta'_\varepsilon(t) < \frac{1}{\varepsilon} \quad \text{if } 0 < t < 2\varepsilon.$$

Consider now the semilinear elliptic system

$$L^k u^k + \beta_\varepsilon(u^k - u^{k+1}) = f^k \quad \text{in } \Omega \quad (1 \leq k \leq m), \tag{2.3}$$

$$u^k = 0 \quad \text{on } \partial\Omega \tag{2.4}$$

where $u^{m+1} \equiv u^1$.

This system can be solved by several standard methods. It is most convenient for our purposes to invoke certain facts of nonlinear functional analysis, as these considerations are crucial for the convergence results of §5. We begin by recalling some definitions; for more details, see [1].

In any real Banach space X one defines the pairing

$$[x, y]_+ \equiv \inf_{\lambda > 0} \frac{\|x + \lambda y\| - \|x\|}{\lambda} \quad (x, y \in X).$$

It can be shown that

$$\lim_{\lambda \downarrow 0} \frac{\|x + \lambda y\| - \|x\|}{\lambda} \text{ exists and equals } [x, y]_+.$$

Note that

$$[\cdot, \cdot]_+ : X \times X \rightarrow R \text{ is upper semicontinuous.} \tag{2.5}$$

A nonlinear operator A with domain $D(A)$ in X and range in X is called *accretive* if

$$\|x - y\| \leq \|x - y + \lambda(Ax - Ay)\| \quad \forall x, y \text{ in } D(A), \quad \lambda > 0,$$

and *m-accretive* if, in addition, $R(I + \lambda A) = X$ for all $\lambda > 0$ (or, equivalently, for some $\lambda > 0$). A is accretive if and only if

$$[x - y, Ax - Ay]_+ \geq 0 \quad \forall x, y \text{ in } D(A). \tag{2.6}$$

If $X = C(\bar{\Omega})$, the brackets $[\cdot, \cdot]_+$ admit the representation

$$[f, g]_+ = \max_{\substack{y \in \bar{\Omega} \\ |f(y)| = \|f\|}} g(y) \cdot \text{sgn } f(y) \quad (f \not\equiv 0), \tag{2.7}$$

see, for example, Sato [16, p. 431] or Sinestrari [17, p. 22]. As we shall see, this characterization is useful for studying partial differential equations satisfying a maximum principle.

The following lemma is due to Frank Massey (unpublished).

LEMMA 2.1. *Let $X = X^1 \times \dots \times X^n$ be the product of real Banach spaces*

X^i with norm

$$\|x\| = \max_{1 \leq i \leq n} \|x^i\| \quad (x = (x^1, \dots, x^n)).$$

Denote by $[\cdot, \cdot]_+^i$ and $[\cdot, \cdot]_+$ the brackets in X^i and X , respectively. Then

$$[x, y]_+ = \max_{1 \leq i \leq n} \{ [x^i, y^i]_+^i; \|x^i\| = \|x\| \}.$$

PROOF. If $\|x^i\| = \|x\|$, then

$$\|x + \lambda y\| - \|x\| \geq \|x^i + \lambda y^i\| - \|x^i\|.$$

Dividing by λ and taking the infimum over λ , we get

$$[x, y]_+ \geq [x^i, y^i]_+^i.$$

This proves the inequality

$$[x, y]_+ \geq \max_i \{ [x^i, y^i]_+^i; \|x^i\| = \|x\| \}. \tag{2.8}$$

To prove the reverse, we take a sequence $\lambda_m > 0$ such that $\lambda_m \downarrow 0$ and

$$\|x + \lambda_m y\| = \|x^i + \lambda_m y^i\| \quad \text{for some fixed } i.$$

Then $\|x\| = \|x^i\|$ and

$$\|x + \lambda_m y\| - \|x\| = \|x^i + \lambda_m y^i\| - \|x^i\|.$$

Dividing by λ_m and taking $\lambda_m \downarrow 0$, we obtain the opposite inequality to (2.8).

□

We now take $X^i = C(\bar{\Omega})$, $X = X^1 \times \dots \times X^m$ and introduce the vector notation

$$u = \begin{bmatrix} u^1 \\ u^2 \\ \vdots \\ u^m \end{bmatrix}, \quad Lu = \begin{bmatrix} L^1 u^1 \\ L^2 u^2 \\ \vdots \\ L^m u^m \end{bmatrix}, \quad B_\epsilon u = \begin{bmatrix} \beta_\epsilon(u^1 - u^2) \\ \beta_\epsilon(u^2 - u^3) \\ \vdots \\ \beta_\epsilon(u^m - u^1) \end{bmatrix}.$$

Each operator L^i is defined on the set

$$D(L^i) = \{v \in W_0^{1,2}(\Omega) \cap W^{2,p}(\Omega); L^i v \in C(\bar{\Omega})\}$$

for any fixed $p > n$, and L is defined on $D(L) = D(L^1) \times \dots \times D(L^m)$. The operator B_ϵ is defined everywhere on X .

LEMMA 2.2. The operator $A_\epsilon = L - c_0 I + B_\epsilon$ with domain $D(L)$ is m -accretive in X .

PROOF. According to a standard perturbation theory [1] it suffices to prove that

$$B_\epsilon \text{ is Lipschitz and accretive} \tag{2.9}$$

and

$$L - c_0I \text{ is } m\text{-accretive.} \tag{2.10}$$

Since $0 \leq \beta'_\epsilon \leq 1/\epsilon$, B_ϵ is clearly Lipschitz. Furthermore, by Lemma 2.1,

$$\begin{aligned} & [u - \bar{u}, B_\epsilon u - B_\epsilon \bar{u}]_+ \\ &= \max\{ [u^i - \bar{u}^i, \beta_\epsilon(u^i - u^{i+1}) - \beta_\epsilon(\bar{u}^i - \bar{u}^{i+1})]_+, \\ & \qquad \qquad \qquad \|u^i - \bar{u}^i\| = \|u - \bar{u}\| \}. \end{aligned} \tag{2.11}$$

Take i such that $\|u^i - \bar{u}^i\| = \|u - \bar{u}\|$ and, without loss of generality, assume that

$$\|u^i - \bar{u}^i\| = u^i(x^0) - \bar{u}^i(x^0) \quad \text{for some } x^0 \in \bar{\Omega}.$$

Then

$$u^i(x^0) - \bar{u}^i(x^0) \geq u^{i+1}(x^0) - \bar{u}^{i+1}(x^0),$$

and so

$$u^i(x^0) - u^{i+1}(x^0) \geq \bar{u}^i(x^0) - \bar{u}^{i+1}(x^0).$$

Because of the monotonicity of β_ϵ , it follows that

$$\beta_\epsilon(u^i - u^{i+1}) \geq \beta_\epsilon(\bar{u}^i - \bar{u}^{i+1}) \quad \text{at } x^0.$$

Recalling (2.7), we conclude that the right-hand side of (2.11) is nonnegative. This proves (2.9).

The proof that $L^i - c_0I$ is accretive follows by the maximum principle, making use of (2.7). In fact, if $v \in C(\bar{\Omega}) \cap W_{loc}^{2,p}(\Omega)$ and v takes a positive maximum at a point $x^0 \in \Omega$, then by Bony [4] (see also [8])

$$\text{ess } \liminf_{x \rightarrow x^0} (L^i v(x) - c_0 v(x)) \geq 0. \tag{2.12}$$

Since, in particular, $L^i v - c_0 v$ is continuous, then $v(x^0)(L^i v(x^0) - c_0 v(x^0)) \geq 0$. Using this and (2.7), it is clear that $L^i - c_0I$ is accretive. Employing Lemma 2.1 we find that also $L - c_0I$ is accretive.

By the general theory of elliptic equations, each $L^i - c_0I$ is m -accretive; consequently $L - c_0I$ is also m -accretive. \square

THEOREM 2.3. *Let (1.1)–(1.5) hold. Then there exists a unique solution of (2.3), (2.4) with components u^k in $D(L^k)$; further, u^k belongs to $C^{2,\beta}(\bar{\Omega}) \cap C^{3,\theta}(\Omega_\theta)$ for any $0 < \theta < 1$, $\bar{\Omega}_\theta \subset \Omega$.*

The first part follows from the m -accretiveness of A_ϵ . The second part is a consequence of standard regularity results for elliptic equations. \square

3. $W^{1,\infty}$ estimates. In this section we derive uniform estimates on the solution $u = (u^1, \dots, u^m)$ of (2.3), (2.4) and on its first derivatives; these bounds will not depend on the parameters ϵ, m . The condition (1.8) will not be needed here.

LEMMA 3.1.

$$\max_k \|u^k\|_{L^\infty(\Omega)} \leq \frac{1}{c_0} \max_k \|f^k\|_{L^\infty(\Omega)}. \tag{3.1}$$

PROOF. We first prove that

$$u^k(x) \geq -\frac{1}{c_0} \max_k \|f^k\|_{L^\infty(\Omega)} \quad (x \in \Omega). \tag{3.2}$$

Denote by N_k the minimum of u^k in $\bar{\Omega}$ and choose j such that $N_j = \min N_k$. Without loss of generality we may suppose that $N_j < 0$. Let x^0 be a point in Ω such that $u^j(x^0) = N_j$. Then

$$L^j u^j(x^0) \leq c^j(x^0) u^j(x^0) = c^j(x^0) N_j \leq c_0 N_j,$$

$$\beta_\varepsilon(u^j(x^0) - u^{j+1}(x^0)) = 0$$

(since $u^j(x^0) < u^{j+1}(x^0)$). Consequently,

$$c_0 N_j \geq L^j u^j(x^0) + \beta_\varepsilon(u^j(x^0) - u^{j+1}(x^0)) = f^j(x^0),$$

so that

$$N_j \geq -\frac{1}{c_0} \|f_j\|_{L^\infty(\Omega)};$$

this proves (3.2). Similarly one estimates the u^k from above. \square

In what follows we shall denote by C a generic positive constant which is independent of m, ε .

LEMMA 3.2.

$$\|Du^k\|_{L^\infty(\partial\Omega)} \leq C \quad (1 \leq k \leq m). \tag{3.3}$$

PROOF. Let \bar{v}^k be the solution of

$$L^k \bar{v}^k = f^k \quad \text{in } \Omega,$$

$$\bar{v}^k = 0 \quad \text{on } \partial\Omega.$$

Since $L^k u^k < f^k$, the maximum principle implies that $u^k < \bar{v}^k$. It follows that

$$\frac{\partial u^k}{\partial \nu} < \frac{\partial \bar{v}^k}{\partial \nu} < C \quad \text{on } \partial\Omega, \tag{3.4}$$

where ν is the inner normal.

Next, let x^0 be a point on $\partial\Omega$ and let y^0 be the center of a ball B of radius R such that $\bar{B} \cap \bar{\Omega} = \{x^0\}$; R can be taken to be independent of x^0 . For sufficiently large positive constants p and D , the function

$$w(x) = D \left(\frac{1}{R^p} - \frac{1}{r^p} \right) \quad (r = |x - y^0|)$$

is a barrier at x^0 for each L^i ; that is, $w = 0$ at x^0 , $w > 0$ in $\bar{\Omega} \setminus \{x^0\}$, and $L^i w > 1$ (the constants p, D are independent of i).

Let $v^k \equiv -Cw$ for $1 < k < m$, for $C > \max_L \|f^k\|_{L^\infty(\Omega)}$. Then

$$\begin{aligned} L^k v^k + \beta_\varepsilon(v^k - v^{k+1}) &= g^k \quad \text{in } \Omega, \\ v^k &< 0 \quad \text{on } \partial\Omega, \end{aligned} \tag{3.5}$$

and $g^k = -CL^k w < f^k$. We claim that

$$v^k < u^k \quad \text{for all } k, x \in \Omega. \tag{3.6}$$

Indeed, otherwise choose j such that

$$\max_{\Omega} (v^k(x) - u^k(x)) \equiv M_k$$

is the largest when $k = j$, and let $x^* \in \Omega$ be such that

$$0 < M_j = v^j(x^*) - u^j(x^*).$$

At $x = x^*$, $L^j(v^j - u^j) > c^j(v^j - u^j) > 0$, $g^j - f^j < 0$, and consequently

$$\beta_\varepsilon(v^j - v^{j+1}) < \beta_\varepsilon(u^j - u^{j+1}).$$

Noting that $v^j \equiv v^{j+1}$, we deduce that $u^j(x^*) - u^{j+1}(x^*) > 0$. Hence

$$\begin{aligned} \max_{\Omega} (v^{j+1} - u^{j+1}) &> v^{j+1}(x^*) - u^{j+1}(x^*) > v^j(x^*) - u^j(x^*) \\ &= \max_{\Omega} (v^j - u^j), \end{aligned}$$

thus contradicting the definition of j .

Having proved (3.6), we now notice that $v^k = u^k = 0$ at x^0 . Hence

$$\frac{\partial u^k}{\partial \nu} > \frac{\partial v^k}{\partial \nu} = -C \frac{\partial w}{\partial \nu} \quad \text{at } x^0.$$

This, together with (3.4), complete the proof of (3.3). \square

LEMMA 3.3.

$$\|Du^k\|_{L^\infty(\Omega)} < C \quad (1 < k < m). \tag{3.7}$$

PROOF. Let λ be a positive number to be determined later on (independently of m, ε), and choose j such that

$$\max_{\Omega} [|\nabla u^j|^2 + \lambda(u^j)^2] > \max_{\Omega} [|\nabla u^k|^2 + \lambda(u^k)^2] \tag{3.8}$$

for all k . We take $j = 1$ for simplicity and write $u = u^1$, $L = L^1$, $a_{ij} = a_{ij}^1$, $b_i = c_i^1$, $c = c^1$, $f = f^1$ and $\beta = \beta_\varepsilon$. We also set $v = u^2$.

We shall denote partial derivatives by subscripts and use the summation convention.

We shall estimate the function

$$w = |\nabla u|^2 + \lambda u^2 = u_i u_i + \lambda u^2$$

by applying to it the maximum principle. First we compute

$$\begin{aligned} w_\mu &= 2u_i u_{i\mu} + 2\lambda u u_\mu, \\ w_{\mu\nu} &= 2u_{i\nu} u_{i\mu} + 2u_i u_{i\mu\nu} + 2\lambda u_\nu u_\mu + 2\lambda u u_{\mu\nu}. \end{aligned} \tag{3.9}$$

Differentiating the equation

$$Lu + \beta(u - v) = f \tag{3.10}$$

we obtain

$$Lu_i + \beta'(u - v)(u_i - v_i) = f_i + \tilde{D}^2u \quad (1 \leq i \leq n) \tag{3.11}$$

where

$$\tilde{D}^2u = \sum_{|\alpha| < 2} \sigma_\alpha D^\alpha u, \quad \sigma_\alpha \text{ bounded.}$$

We now compute Lw , by substituting from (3.9)

$$\begin{aligned} Lw &= -2a_{\mu\nu}u_{i\mu}u_{i\nu} - 2u_i a_{\mu\nu}u_{i\mu\nu} - 2\lambda a_{\mu\nu}u_\mu u_\nu \\ &\quad - 2\lambda u a_{\mu\nu}a_{\mu\nu} - 2u_i b_\mu u_{i\mu} - 2\lambda u b_\mu u_\mu + cu_i u_i + \lambda cu^2 \\ &\leq -2\gamma u_{i\mu}u_{i\mu} - 2\gamma \lambda u_\mu u_\mu + 2u_i Lu_i + 2\lambda u Lu. \end{aligned}$$

Recalling (3.10), (3.11), we get

$$\begin{aligned} Lw &\leq -2\gamma u_{ij}u_{ij} - 2\gamma \lambda u_i u_i + 2u_i f_i + 2u_i \tilde{D}^2u + 2\lambda u f \\ &\quad - 2u_i \beta'(u - v)(u_i - v_i) - 2\lambda u \beta(u - v). \end{aligned}$$

Since

$$|2u_i \tilde{D}^2u| \leq \gamma u_{ij}u_{ij} + Cu_i u_i,$$

and since $|u| \leq C$, we obtain, upon choosing λ to be sufficiently large (depending only on C)

$$Lw \leq C - J, \tag{3.12}$$

where

$$J = 2\lambda u \beta(u - v) + 2\beta'(u - v)u_i(u_i - v_i). \tag{3.13}$$

Suppose the maximum of w in $\bar{\Omega}$ is attained at a point $x^0 \in \Omega$. Then

$$u_i u_i + \lambda u^2 \geq v_i v_i + \lambda v^2 \quad \text{at } x = x^0 \tag{3.14}$$

(since (3.8) holds with $j = 1$). Next, because of (2.2),

$$J > 2\beta'(u - v)u_i(u_i - v_i) + 2\lambda u \beta'(u - v)(u - v) - C. \tag{3.15}$$

Noting that

$$\begin{aligned} 2u_i(u_i - v_i) &> u_i u_i - v_i v_i, \\ 2u(u - v) &> u^2 - v^2, \end{aligned}$$

we obtain

$$J > \beta'(u - v)(u_i u_i - v_i v_i + \lambda u^2 - \lambda v^2) - C.$$

Hence, by (3.14), $J > -C$ at $x = x^0$. Thus, by (3.12), $Lw(x^0) < C$. But since w attains its maximum in $\bar{\Omega}$ at x^0 , we must have

$$Lw(x^0) > c(x^0)w(x^0) > c_0 w(x^0).$$

Therefore $w(x^0) < C/c_0$.

If w attains its maximum only at boundary points, then the assertion $\max_{\bar{\Omega}} w \leq C$ follows from Lemma 3.2. This completes the proof of (3.7). \square

4. $W^{2,\infty}$ estimates. In this section we require also, in addition to (1.1)–(1.5), the condition (1.8). As before, C will denote a generic constant independent of ε, m .

LEMMA 4.1. *For any compact subdomain Ω_0 of Ω*

$$\|u^k\|_{W^{2,\infty}(\Omega_0)} \leq C \quad (1 \leq k \leq m) \tag{4.1}$$

where C depends on Ω_0 .

PROOF. Without loss of generality we may assume that the u^k belong to $C^4(\Omega)$. Indeed, otherwise we approximate L^i by $L^{i,q}$ with C^3 coefficients (say); the derivation of (4.1) for the corresponding solution $u^{k,q}$ (given below) shows that C depends only on the constants occurring in (1.2)–(1.5) and thus can be taken to be independent of q . Taking $q \rightarrow \infty$, the assertion (4.1) then follows.

Let ξ denote any direction and let ζ be a function in $C_0^\infty(\Omega)$ such that $\zeta = 1$ in Ω_0 , $\zeta > 0$ elsewhere. Consider the numbers

$$M_i = \max_{\bar{\Omega}} \left\{ \zeta^2 \left((u_{\xi\xi}^i)^+ \right)^2 + \lambda |\nabla u_i|^2 \right\} \quad (\lambda > 0)$$

and suppose for definiteness that

$$M_1 \geq M_i \quad \text{for all } 1 \leq i \leq k; \tag{4.2}$$

here λ is a positive number (independent of ε, m) to be determined later on. As in the proof of Lemma 3.3, we set $u = u^1$, $v = u^2$, $L = L^1$, $f = f^1$ and $\beta = \beta_1$.

Differentiating (3.10) twice with respect to ξ , we get, using (1.8),

$$Lu_{\xi\xi} + \beta'(u - v)(u_{\xi\xi} - v_{\xi\xi}) + \beta''(u - v)(u_\xi - v_\xi)^2 = f_{\xi\xi} + \tilde{D}^2u$$

where

$$\tilde{D}^2u = \sum_{|\alpha| < 2} \sigma_\alpha D^\alpha u, \quad \sigma_\alpha \text{ bounded.} \tag{4.3}$$

Since $\beta'' > 0$, we conclude that

$$Lu_{\xi\xi} + \beta'(u - v)(u_{\xi\xi} - v_{\xi\xi}) \leq f_{\xi\xi} + \tilde{D}^2u. \tag{4.4}$$

Consider now the function

$$w = \zeta^2 \left((u_{\xi\xi})^+ \right)^2 + \lambda |\nabla u|^2,$$

and suppose that it attains its maximum in $\bar{\Omega}$ at a point x^0 . If $x^0 \in \partial\Omega$ then

$$w(x^0) = \lambda |\nabla u(x^0)|^2 \leq C, \tag{4.5}$$

by Lemma 3.2. Suppose next that $x^0 \in \Omega$. If $u_{\xi\xi}(x^0) < 0$, then again (4.5) holds, by Lemma 3.3.

We shall now consider the case where $u_{\xi\xi}(x^0) > 0$. Then $u_{\xi\xi} > 0$ in a neighborhood G of x^0 , and thus

$$w = \zeta^2(u_{\xi\xi})^2 + \lambda|\nabla u|^2, \quad u_{\xi\xi} > 0 \text{ in } G.$$

We now compute

$$\begin{aligned} w_i &= 2u_{\xi\xi}u_{\xi\xi i}\zeta^2 + 2\zeta\zeta_i u_{\xi\xi}^2 + 2\lambda u_k u_{ki}, \\ w_{ij} &= 2u_{\xi\xi i}u_{\xi\xi j}\zeta^2 + 2u_{\xi\xi}u_{\xi\xi ij}\zeta^2 + 4(u_{\xi\xi}u_{\xi\xi i}\zeta\zeta_j + u_{\xi\xi}u_{\xi\xi j}\zeta\zeta_i) \\ &\quad + 2u_{\xi\xi}^2(\zeta\zeta_{ij} + \zeta_i\zeta_j) + 2\lambda u_{kj}u_{ki} + 2\lambda u_k u_{kij}. \end{aligned}$$

It follows easily that

$$\begin{aligned} -a_{ij}w_{ij} &\leq -2\zeta^2 a_{ij}u_{\xi\xi i}u_{\xi\xi j} - 2\zeta^2 u_{\xi\xi} a_{ij}u_{\xi\xi ij} \\ &\quad + C\zeta \sum_i |u_{\xi\xi i}u_{\xi\xi i}| + Cu_{\xi\xi}^2 - 2\lambda a_{ij}u_{ki}u_{kj} - 2\lambda u_k a_{ij}u_{kij}, \\ -b_i w_i &\leq 2\zeta^2 u_{\xi\xi} b_i u_{\xi\xi i} + C\zeta u_{\xi\xi}^2 + 2\lambda u_k b_i u_{ki}. \end{aligned}$$

Hence

$$\begin{aligned} Lw &\leq -2\gamma\zeta^2 u_{\xi\xi i}u_{\xi\xi i} + 2\zeta^2 u_{\xi\xi} Lu_{\xi\xi} \\ &\quad + \left(\gamma\zeta^2 u_{\xi\xi i}u_{\xi\xi i} + \frac{C}{\gamma} u_{\xi\xi}^2 \right) - 2\lambda\gamma u_{ki}u_{kj} + 2\lambda u_k Lu_k. \end{aligned}$$

We now specialize to $x = x^0$. Since $u_{\xi\xi}(x^0) > 0$, we can substitute $Lu_{\xi\xi}$ from (4.4) and still preserve the last inequality. Substituting also Lu_k from (3.11), we get

$$\begin{aligned} Lw &\leq -\gamma\zeta^2 u_{\xi\xi i}u_{\xi\xi i} + 2\zeta^2 u_{\xi\xi}(f_{\xi\xi} + \tilde{D}^2u - \beta'(u-v)(u_{\xi\xi} - v_{\xi\xi})) \\ &\quad + Cu_{\xi\xi}^2 - 2\lambda\gamma u_{ki}u_{ki} + 2\lambda u_k(f_k + \tilde{D}^2u - \beta'(u-v)(u_k - v_k)). \end{aligned}$$

Choosing λ sufficiently large and using also Lemma 3.3, we are led to the inequality

$$Lw \leq C - J \quad \text{at } x = x^0, \tag{4.6}$$

where

$$J = 2\beta'(u-v) [\zeta^2 u_{\xi\xi}(u_{\xi\xi} - v_{\xi\xi}) + \lambda u_k(u_k - v_k)].$$

Noting that $u_{\xi\xi}v_{\xi\xi} \leq u_{\xi\xi}^+v_{\xi\xi}^+$ at x^0 , we find that

$$J \geq \beta'(u-v) [\zeta^2 u_{\xi\xi}^2 - \zeta^2 (v_{\xi\xi}^+)^2 + \lambda u_k u_k - \lambda v_k v_k] \geq 0$$

where the last inequality is a consequence of (4.2).

Since w attains its maximum at x^0 ,

$$Lw(x^0) > c(x^0)w(x^0) \geq c_0w(x^0).$$

But since $Lw < C - J < C$ at x^0 , we deduce that $w(x^0) < C$. This completes

the proof that $\|w\|_{L^\infty(\Omega)} \leq C$. It follows that

$$\frac{\partial^2}{\partial \xi^2} u^k \leq C, \quad 1 \leq k \leq m, x \in \Omega_0, \quad (4.7)$$

for any direction ξ .

For a particular L^k , choose the coordinate system so that $a_{ij}^k = \delta_{ij}$. Then

$$\begin{aligned} -\Delta u^k &= L^k u_k + \sum b_i^k \frac{\partial u^k}{\partial x_i} - c^k u^k \\ &< f^k + \sum b_i^k \frac{\partial u^k}{\partial x_i} - c^k u^k \leq C. \end{aligned}$$

By (4.7) we also have

$$\frac{\partial^2 u^k}{\partial x_i^2} \leq C \quad \text{for each } i, x \in \Omega_0.$$

It follows that

$$\frac{\partial^2 u^k}{\partial x_i^2} \geq -(n-1)C, \quad x \in \Omega.$$

Since x_i can be any direction, the second mixed derivatives are also bounded. The proof of the lemma is complete. \square

5. Passage to the limit as $\varepsilon \rightarrow 0$. In this section we study the "truncated" Bellman equation

$$\max_{1 \leq k \leq m} (L^k u(x) - f^k(x)) = 0 \quad \text{a.e. in } \Omega, \quad (5.1)$$

with the boundary condition

$$u = 0 \quad \text{on } \partial\Omega. \quad (5.2)$$

THEOREM 5.1. *Let (1.1)–(1.5) and (1.8) hold. Then there exists a unique solution u of (5.1), (5.2) such that*

$$u \in W^{1,\infty}(\Omega) \cap W_{\text{loc}}^{2,\infty}(\Omega). \quad (5.3)$$

PROOF. Denote the solution of (2.3), (2.4) by $u_\varepsilon = (u_\varepsilon^1, u_\varepsilon^2, \dots, u_\varepsilon^m)$.

From Lemmas 3.3, 4.1 we have the uniform estimates

$$|Du_\varepsilon^k(x)| \leq C \quad (x \in \Omega), \quad (5.4)$$

$$|D^2 u_\varepsilon^k(x)| \leq C \quad (x \in \Omega_0) \quad (5.5)$$

for any compact domain Ω_0 in Ω . Take a sequence of ε 's (which we denote for simplicity again by ε) such that

$$u_\varepsilon^k \rightarrow v^k(x) \quad \text{uniformly in } \bar{\Omega}, \quad (5.6)$$

$$u_\varepsilon^k \rightarrow v^k(x) \quad \text{weakly in } W^{2,p}(\Omega_0) \quad (5.7)$$

for any $p < \infty$ and for any compact subdomain $\Omega_0 \subset \Omega$.

From the relation

$$\beta_\varepsilon(u_\varepsilon^k - u_\varepsilon^{k+1}) = f^k - L^k u_\varepsilon^k$$

and (5.4), (5.5) it follows that

$$\beta_\varepsilon(u_\varepsilon^k - u_\varepsilon^{k+1}) < C \quad \text{in } \Omega_0.$$

Noting that $\beta_\varepsilon(t) \rightarrow \infty$ if $t > 0, \varepsilon \rightarrow 0$, we deduce that

$$u_\varepsilon^k(x) - u_\varepsilon^{k+1}(x) \rightarrow 0 \quad \text{if } \varepsilon \rightarrow 0, x \in \Omega.$$

It follows that the v^k defined in (5.6) satisfy $v^1 = v^2 = \dots = v^m \equiv u$. We shall prove that

$$\max_{1 < k < m} (L^k u(x) - f^k(x)) = 0 \quad \text{a.e. in } \Omega. \tag{5.8}$$

We begin by noticing that $L^k u_\varepsilon^k < f^k$ and that the weak convergence (in (5.7)) preserves this inequality. Hence

$$L^k u(x) - f^k(x) < 0 \quad \text{a.e. in } \Omega,$$

so that also

$$\max_{1 < k < m} (L^k u(x) - f^k(x)) < 0 \quad \text{a.e. in } \Omega. \tag{5.9}$$

To prove the reverse inequality we employ an argument of Evans [9] which generalizes the Minty lemma to the space $C(\bar{\Omega})$. Using the notation of §2 we have, by the accretiveness of $L + B_\varepsilon$,

$$\begin{aligned} 0 < [u_\varepsilon - \tilde{\varphi}, (L + B_\varepsilon)u_\varepsilon - (L + B_\varepsilon)\tilde{\varphi}]_+ &= [u_\varepsilon - \tilde{\varphi}, f - (L + B_\varepsilon)\tilde{\varphi}]_+ \\ &= [u_\varepsilon - \tilde{\varphi}, f - L\tilde{\varphi}]_+, \end{aligned} \tag{5.10}$$

the last relation obtained by choosing $\tilde{\varphi}$ with components φ^i such that $\varphi^1 = \varphi^2 = \dots = \varphi^m \equiv \varphi$.

We take φ in $C_0^\infty(\Omega)$. Since $u_\varepsilon^k \rightarrow u$ uniformly in Ω , taking $\varepsilon \rightarrow 0$ in the last expression in (5.10) gives, according to (2.5) and Lemma 2.1,

$$0 < [\tilde{u} - \tilde{\varphi}, f - L\tilde{\varphi}]_+ = \max_{1 < k < m} [u - \varphi, f^k - L^k \varphi]_+ \tag{5.11}$$

where \tilde{u} is the vector with m equal components u .

By Lemma 2.2 of [9], for a.e. $x^0 \in \Omega$ there exists a sequence of functions $\varphi_j \in C_0^\infty(\Omega)$ such that

$$D^\alpha \varphi_j(x^0) \rightarrow D^\alpha u(x^0) \quad (0 < |\alpha| < 2), \tag{5.12}$$

$$\begin{aligned} \varphi_j(x^0) - u(x^0) = \|\varphi_j - u\|_{C(\bar{\Omega})} > \varphi_j(x) - u(x) \\ \forall x \in \Omega, \quad x \neq x^0. \end{aligned} \tag{5.13}$$

Taking $\varphi = \varphi_j$ in (5.11) and using (2.7), (5.13), we get

$$0 < \max_{1 < k < m} (L^k \varphi_j(x^0) - f^k(x^0)).$$

Taking $j \rightarrow \infty$ and using (5.12), we find that

$$\max_{1 < k < m} (L^k u(x^0) - f^k(x^0)) > 0 \quad \text{a.e.}$$

This completes the proof of (5.8).

The proof of uniqueness will be given in §7. \square

REMARK. The applicability of the accretive operator methods to the Bellman equation was first noted by Pliska [15].

6. Existence for the Bellman equation. We shall now consider the “full” Bellman equation

$$\sup_{1 < k < \infty} (L^k u(x) - f^k(x)) = 0 \quad \text{a.e. in } \Omega \tag{6.1}$$

with the boundary condition

$$u = 0 \quad \text{on } \partial\Omega. \tag{6.2}$$

THEOREM 6.1. *Let (1.1)–(1.5) and (1.8) hold. Then there exists a unique solution u of (6.1), (6.2) such that*

$$u \in W^{1,\infty}(\Omega) \cap W_{loc}^{2,\infty}(\Omega). \tag{6.3}$$

PROOF. Denote the solution of (5.1), (5.2) by u_m . Then

$$|Du_m(x)| < C \quad (x \in \Omega) \tag{6.4}$$

and, for any compact subdomain $\Omega_0 \subset \Omega$,

$$|D^2u_m(x)| < C \quad (x \in \Omega_0); \tag{6.5}$$

here C is a generic constant independent of m .

Let

$$A_m v(x) \equiv \max_{1 < k < m} (L^k v(x) - f^k(x)),$$

$$A v(x) \equiv \sup_{1 < k < \infty} (L^k v(x) - f^k(x)).$$

If $\varphi \in C_0^\infty(\Omega)$ then the $L^k \varphi$ are uniformly bounded and equicontinuous, so that as $m \rightarrow \infty$,

$$A_m \varphi(x) \rightarrow A \varphi(x) \quad \text{uniformly in } \bar{\Omega}. \tag{6.6}$$

Consider A_m as an operator in $X = L^\infty(\Omega)$ with the domain

$$D(A_m) = \{v \in W_0^{1,\infty}(\Omega) \cap W_{loc}^{2,\infty}(\Omega); A_m v(x) \in L^\infty(\Omega)\}.$$

LEMMA 6.2. A_m is accretive.

PROOF. We recall (see Sato [16, p. 433]) that the bracket $[,]_+$ in $X = L^\infty(\Omega)$ can be computed by

$$[f, g]_+ = \lim_{\epsilon \downarrow 0} \text{ess sup}_{\Omega(f, \epsilon)} g(x) \cdot \text{sgn } f(x) \quad (f \not\equiv 0) \tag{6.7}$$

where

$$\Omega(f, \varepsilon) \equiv \{x \in \Omega; |f(x)| > \|f\|_{L^\infty(\Omega)} - \varepsilon\}.$$

Now, if $v, \bar{v} \in D(A_m)$ then $v - \bar{v}$ is continuous in $\bar{\Omega}$; and without loss of generality we may assume that $v - \bar{v}$ takes a positive maximum $\|v - \bar{v}\|_{L^\infty(\Omega)}$ in $\bar{\Omega}$. We have proven that $[v - \bar{v}, A_m v - A_m \bar{v}]_+ > 0$, or, in view of (6.7), that

$$\lim_{\varepsilon \downarrow 0} \text{ess sup}_{G_\varepsilon} (A_m v(x) - A_m \bar{v}(x)) > 0 \tag{6.8}$$

where

$$G_\varepsilon = \left\{x; v(x) - \bar{v}(x) \geq \max_{\bar{\Omega}} |v - \bar{v}| - \varepsilon\right\}.$$

The maximum principle of Bony [4] (see (2.12)) implies that (6.8) holds if A_m is replaced by each L^k ; hence obviously (6.8) holds. \square

Using Lemma 6.2 we can now pass to the limit with $m \rightarrow \infty$ by an argument similar to that used in Theorem 5.1. Indeed, for any $\varphi \in C_0^\infty(\Omega)$,

$$0 < [u_m - \varphi, A_m u_m - A_m \varphi]_+ = [u_m - \varphi, -A_m \varphi]_+. \tag{6.9}$$

In view of (6.4), (6.5), there exists a subsequence of u_m , which we again denote by u_m , such that $u_m \rightarrow u$ uniformly in $\bar{\Omega}$, $Du_m \rightarrow Du$ in the weak star topology of $L^\infty(\Omega)$, $D^2 u_m \rightarrow D^2 u$ in the weak star topology of $L^\infty(\Omega_0)$ (for any $\Omega_0, \bar{\Omega}_0 \subset \Omega$).

Taking $m \rightarrow \infty$ in (6.9) we obtain, using (6.6) and (2.5), $[u - \varphi, -A\varphi]_+ > 0$. Choosing $\varphi = \varphi_j$ as in the proof of Theorem 5.1, we get $A\varphi_j(x^0) > 0$.

Now, since

$$D^\alpha \varphi_j(x^0) - D^\alpha u(x^0) \rightarrow 0 \quad (0 < |\alpha| < 2)$$

and the coefficients of L^k are uniformly bounded, we conclude that

$$A\varphi_j(x^0) - Au(x^0) \rightarrow 0.$$

It follows that $Au(x^0) > 0$.

Conversely, since $L^k u_m - f^k < 0$ a.e. in Ω , $k < m$, we obtain

$$L^k u - f^k \leq 0 \quad \text{a.e. in } \Omega.$$

Hence $Au \leq 0$ a.e. in Ω . We have thus proved that u is a solution of (6.1), (6.2) satisfying (6.3).

The uniqueness of the solution will be proved in §7. \square

REMARK 1. The assumption $c_0 > 0$ made in §§2-6 may be replaced by $c_0 \geq 0$. Indeed, if $c_0 = 0$, we observe that the maximum principle still applies to $L - \delta I$ provided δ is positive and sufficiently small. Hence Theorem 2.3 remains valid (with $L - \delta I + B_\varepsilon$ m -accretive). Next, (3.1) holds with $1/c_0$ replaced by some positive constant C . The remaining arguments now proceed as before. Thus, Theorem 6.1 remains true also when $c_0 = 0$.

REMARK 2. As a special case of Theorem 6.1 one can immediately establish the existence and uniqueness of a solution for the problem

$$\lambda u - F(u_{x_i x_j}) = f(x) \quad \text{in } \Omega, \quad (6.10)$$

$$u = 0 \quad \text{on } \partial\Omega \quad (6.11)$$

provided $\lambda > 0$ and $F(x_{ij})$ satisfies the following conditions:

- (a) $F: R^{n^2} \rightarrow R$ is convex and C^2 ,
- (b) $\sum F_{x_i x_j} \xi_i \xi_j \geq \gamma |\xi|^2$, $\gamma > 0$ (ellipticity),
- (c) $|F(x) - \sum F_{x_{ij}}(x) x_{ij}| \leq C \quad \forall x = (x_{ij})$.

Indeed, writing

$$F(x) = \max_{\xi} [F(\xi) + F_{x_{ij}}(\xi) \cdot (x_{ij} - \xi_{ij})], \quad \xi = (\xi_{ij}),$$

we may rewrite (6.10) as a Bellman equation with

$$L^{\xi} u \equiv \lambda u - \sum F_{x_{ij}}(\xi) \frac{\partial^2 u}{\partial x_i \partial x_j},$$

$$f^{\xi}(x) = f(x) + F(\xi) - \sum F_{x_{ij}}(\xi) \xi_{ij};$$

it suffices to let ξ range over a dense countable set in R^{n^2} .

7. Probabilistic interpretation and methods. Let k_1, \dots, k_m be positive numbers. Consider the system of m variational inequalities

$$L^i u_i < f^i \quad \text{a.e. in } \Omega,$$

$$u_i < k_i + u_{i+1} \quad \text{in } \Omega,$$

$$(L^i u_i - f^i)(u_i - k_i - u_{i+1}) = 0 \quad \text{a.e. in } \Omega, \quad 1 < i < m. \quad (7.1)$$

This system is the (formal) limit of the penalized problem

$$L^i u_i + \beta_{\epsilon}(u_i - k_i - u_{i+1}) = f^i \quad \text{in } \Omega,$$

$$u_i = 0 \quad \text{on } \partial\Omega, \quad 1 < i < m, \quad (7.2)$$

where $\beta_{\epsilon}(t)$ is the function constructed in §2.

We shall need the conditions

$$|a_{ij}^k|, |b_i^k|, |c^k|, |f^k| < C,$$

$$|Da_{ij}^k|, |Db_i^k|, |Dc^k| < C. \quad (7.3)$$

Under these assumptions and (1.1), (1.2), (1.4) and the additional condition that $|Df^k| < C$ one can establish the estimates of §3 for the solution $u_i = u_{i,\epsilon}$ of (7.2) (cf. the proof of Theorem 7.2 below). These estimates will provide a uniform modulus of continuity for solutions of (7.2) and these, in turn, imply as we shall see below uniform $W^{2,p}(\Omega)$ estimates (without using condition (1.8)).

Let us now interpret (7.1) probabilistically (cf. Bensoussan and Lions [2] for the case $m = 2$). This approach gives heuristic insight into the stochastic

control situation modeled by (7.1) (and (1.6), (1.7)). As a bonus we shall derive the $W^{2,p}(\Omega)$ estimate for solutions of (7.2) under somewhat weaker assumptions than those for the plan outlined above.

Consider the stochastic differential system

$$d\xi^i(t) = \sigma^i(\xi(t)) dw(t) + b^i(\xi(t)) dt$$

where $\sigma^i(x)$ is a symmetric matrix such that $\frac{1}{2}(\sigma^i(x))^2$ is the matrix $(a_{jk}(x))$ and $b^i(x)$ is the vector with components $b_j^i(x)$. Here $w(t)$ is an n -dimensional Brownian motion. Denote by \mathcal{F}_t the σ -field $\sigma(w(s), 0 \leq s \leq t)$.

For any sequence $\theta = (\theta_1, \theta_2, \dots)$ of \mathcal{F}_t stopping times $\theta_1 < \theta_2 < \dots < \theta_l < \dots$ ($\theta_l \uparrow \infty$ if $l \uparrow \infty$) we define the following process $\xi(t)$:

$$\xi(t) = \xi^j(t) \text{ if } \theta_{l_{m+j-1}} < t < \theta_{l_{m+j}} \quad (\theta_0 = 0)$$

where l is a nonnegative integer, and $\xi(t)$ is continuous at $t = \theta_i$ for all i . Thus, $\xi(t)$ switches cyclically from the stochastic system with generator L^i to the stochastic system with generator L^{i+1} ($L^{m+1} \equiv L^1$), and it starts with $\xi(t) = \xi^1(t)$ if $0 \leq t \leq \theta_1$, $\xi(0) = x$. We shall assume for simplicity that $c^i(x) \equiv \text{const} = \alpha > 0$.

Introduce the cost function

$$J_x^1(\theta) \equiv E_x \left[\int_0^T e^{-\alpha t} f(\xi(t)) dt + \sum_{i=0}^{\infty} \sum_{j=1}^m k_j e^{-\alpha \theta_{l_{m+j}}} I_{\theta_{l_{m+j}} < T} \right]$$

where T is the exit time from Ω and

$$f(\xi(t)) \equiv f^i(\xi^i(t)) \text{ if } \xi(t) = \xi^i(t). \tag{7.4}$$

This cost functional represents a running cost (per unit time) of $f^i(x)$ when the system is in state i , and a switching cost k_i for any transition from state i to state $i + 1$, the entire cost discounted with a factor α .

Set $u_1(x) \equiv \inf_{\theta} J_x^1(\theta)$. More generally we can define $\xi(t)$ which starts with $\xi(t) = \xi^i(t)$ for $0 \leq t \leq \theta_1$, and then proceed cyclically as before to change from any state j to state $j + 1$. We denote the corresponding cost function by $J_x^i(\theta)$, and set

$$u_i(x) = \inf_{\theta} J_x^i(\theta). \tag{7.5}$$

THEOREM 7.1. *Assume that (1.1), (1.2), (1.4), (1.5), and (7.3) hold. Then (u_1, \dots, u_m) forms a solution of (7.1), belonging to $C(\bar{\Omega}) \cap W^{2,p}(\Omega)$ for any $p < \infty$. Conversely, every solution of (7.1) which belongs to $C(\bar{\Omega}) \cap W_{loc}^{2,p}(\Omega)$ for some $p > n/2$ is given by (7.5) (and is therefore unique).*

PROOF. The second part of the theorem follows by a standard application of Ito's formula to functions in $C(\bar{\Omega}) \cap W_{loc}^{2,p}(\Omega)$ (cf. [3] [11]). Thus it remains to prove the existence of a solution of (7.1) which belongs to $W^{2,p}(\Omega)$ for all $p < \infty$.

We truncate the cost function $J_x^i(\theta)$ by restricting the number of θ_i 's to be at most N . Denote the corresponding cost function by $J_x^{i,N}(\theta)$ and set $u_i^N(x) = \inf_{\theta} J_x^{i,N}(\theta)$. Thus, if $N = 1$ then u_i^1 is simply a solution of the variational inequality

$$L^i u_i^1 \leq f^i, \quad u_i^1 \leq k_i, \quad (L^i u_i^1 - f^i)(u_i^1 - k_i) = 0,$$

that is, when $N = 1$ the obstacle for u_i^N is k_i .

Similarly, for any N , the obstacle for u_i^N is $k_i + u_{i+1}^{N-1}$. This in fact can be proved in a standard way by the principle of dynamic programming, using the strong Markov property.

It is well known (see, for instance, [3]) that if the obstacle is continuous then the solution of the variational inequality is also continuous. Hence by induction it follows that the u_i^N are all continuous functions in $\bar{\Omega}$.

We claim that

$$|u_i^N(x) - u_i(x)| \leq C/N \quad (x \in \bar{\Omega}) \tag{7.6}$$

where $u_i(x)$ is defined by (7.5).

Notice first that

$$u_i(x) \leq u_i^N(x) \tag{7.7}$$

since any cost $J_x^{i,N}(\theta)$ is also a cost $J_x^i(\theta)$ (with $\theta_j = \infty$ if $j > N$).

Next, in estimating $u_i(x)$ from below we may restrict ourselves only to "good" choices of stopping times θ_i . More specifically, since we may always choose not to switch, it is clear that we may restrict the θ_i to be such that

$$E_x \left[\sum_{lm+j>i} k_j e^{-\alpha\theta_{lm+j}} I_{\theta_{lm+j} < T} \right] \leq E_x \left[\int_{\theta_i}^T e^{-\alpha t} f(\xi(t)) dt \right].$$

Recalling that f is bounded, we conclude that in computing $u_i(x)$ it suffices to take the infimum on θ with θ_k satisfying:

$$\sum_{i>j} E_x [e^{-\alpha\theta_j} I_{\theta_j < T}] \leq C E_x [e^{-\alpha\theta} I_{\theta < T}] \tag{7.8}$$

for some C .

We may also restrict the θ_k to be such that $J_x^i(\theta) \leq C$ for some C sufficiently large. In view of (7.8) and the boundedness of f , this implies that

$$\sum_i E_x [e^{-\alpha\theta_i} I_{\theta_i < T}] \leq C.$$

Since the i th term is nonnegative and decreasing, it follows that

$$j E_x [e^{-\alpha\theta_j} I_{\theta_j < T}] \leq C. \tag{7.9}$$

Using (7.8), (7.9) it follows that $|J_x^i(\theta) - J_x^{i,N}(\tilde{\theta})| \leq C/N$ for any $\theta = (\theta_1, \theta_2, \dots)$, where $\tilde{\theta}$ is obtained from θ by dropping all the θ_j with $j > N$. It follows that $u_i^N(x) - u_i(x) \leq C/N$, and recalling (7.7), the assertion (7.6) follows.

Since $u_i^N(x)$ are continuous in $\bar{\Omega}$, (7.6) implies equicontinuity of the $u_i^N(x)$ and the continuity of $u_i(x)$ in $\bar{\Omega}$.

At each point $x^0 \in \Omega$ we cannot have $u_i(x^0) = k_i + u_{i+1}(x^0)$ for all $1 < i < m$. Hence there exist $\varepsilon_0 > 0, \delta > 0$ and some i such that

$$u_i(x) - k_i - u_{i+1}(x) < -\varepsilon_0 \quad \text{if } |x - x^0| < \delta.$$

It follows that $u_i^N(x) < k_i + u_{i+1}^{N-1} - \varepsilon_0/2$ if $|x - x^0| < \delta$ if N is sufficiently large, say $N \geq N_0$. Hence

$$L^i u_i^N = f^i \quad \text{if } |x - x^0| < \delta, \quad N \geq N_0.$$

Since u_i^N is uniformly bounded, the standard elliptic estimates give $\|u_i^N\|_{W^{2,p}(G_\delta)} \leq C$ for any $p < \infty, N \geq N_0$, where $G_\delta = \{x; x \in \Omega, |x - x^0| < \delta/2\}$ and C is a constant independent of N .

Considering $k_{i-1} + u_i^N$ as an obstacle in the variational inequality for u_{i-1}^{N+1} , we then deduce that $\|u_{i-1}^{N+1}\|_{W^{2,p}(G_\delta)} \leq C$ with a smaller δ . Proceeding in this manner step by step $m - 1$ times, we find that $\|u_{i-j}^{N+j}\|_{W^{2,p}(G_\delta)} \leq C$ ($1 < j < m - 1, N \geq N_0$). Recalling (7.6) we deduce that $\|u_j\|_{W^{2,p}(G_\delta)} \leq C$. Hence $u_j \in W^{2,p}(\Omega)$.

Finally, taking $N \rightarrow \infty$ in the variational inequalities for u_j^N ($1 < j < m$) we find that (u_1, \dots, u_m) is a solution of the system (7.1). \square

REMARK. Using the $C^{1,1}$ regularity of solutions of variational inequalities [6], the above proof establishes that $u_i \in C^{1,1}(\bar{\Omega})$ provided the f^i are Hölder continuous and a_{ij}^k, b_i^k are in $C^2(\bar{\Omega})$.

The probabilistic idea underlying the penalized problem (2.3), (2.4) is that $\beta_\varepsilon(u^k - u^{k+1})$ represents (in some heuristic sense) a penalty for switching from the stochastic system for ξ^k to the stochastic system for ξ^{k+1} . In the limit case of the Bellman equation, there is no cost for switching. In fact, Krylov [12] writes the solution $u(x)$ of the Bellman equation (in R^n) as the infimum of the cost functions

$$E_x \left[\int_0^\infty e^{-at} \sum_{i=1}^\infty (f^{v(t)}(\xi^{v(t)}(t)) I_{\sigma(t)=i}) dt \right] \tag{7.10}$$

where $v(t)$ is any nonanticipative function with range $1, 2, \dots$. If we restrict $v(t)$ to have the range $1, 2, \dots, m$ and to have only a countable number of discontinuities at times $\theta_1 < \theta_2 < \dots, \theta_i \uparrow \infty$, and if further $v(\theta_i + 0) = v(\theta_i - 0) + 1$ (if $v(\theta_i - 0) < m$; $v(\theta_i + 0) = 1$ if $v(\theta_i - 0) = m$), then this cost coincides with the cost $J_x^1(\theta)$ introduced above when $T = \infty$ and with no penalty for switching (i.e., $k_j = 0$ for all j).

It thus seems natural to try to obtain the solution u of the Bellman equation (5.1), (5.2) as the limit of the solution of (7.1) as $k = (k_1, \dots, k_m) \rightarrow$

0. In fact we have:

THEOREM 7.2. *Let (1.1)–(1.5) and (1.8) hold. Then there exists a unique solution $u^k = (u_1^k, \dots, u_m^k)$ of (7.1) with u_i^k in $W^{1,\infty}(\Omega) \cap W_{loc}^{2,\infty}(\Omega)$, and*

$$u_i^k(x) \rightarrow u(x) \text{ uniformly in } \bar{\Omega} \text{ as } k \rightarrow 0, \tag{7.11}$$

where u is the solution of (5.1), (5.2).

The proof is similar to the proof of Theorem 5.1. We first consider the system (7.2) and prove that it has a unique solution $(u_1^{k,\varepsilon}, \dots, u_m^{k,\varepsilon})$. Next we derive estimates on $u_i^{k,\varepsilon}$ analogous to those derived in §§3 and 4. In deriving the estimate on $Du_i^{k,\varepsilon}$ we get an inequality analogous to (3.15), with $\beta' = \beta'(u - v - k_1)$ and with an extra term $\hat{J} = 2\lambda u k_1 \beta'(u - v - k_1)$ on the right-hand side. Without loss of generality we may assume that $u > 0$ (otherwise we derive the gradient estimates for $u_i^{k,\varepsilon} + C$ instead of $u_i^{k,\varepsilon}$); consequently $\hat{J} > 0$, and we then continue as in §3. We can now take $\varepsilon \rightarrow 0$ to obtain the solution (u_1^k, \dots, u_m^k) of (7.1). Finally, passing to the limit with $k \rightarrow 0$ and arguing as in §5, we find that (7.11) is valid where u solves (5.1), (5.2). \square

We can actually prove in a similar way that $u_i^{k,\varepsilon} \rightarrow u$ if $k \rightarrow 0, \varepsilon \rightarrow 0$. The proof of Theorem 5.1 is in fact the case where $k = 0, \varepsilon \rightarrow 0$.

Uniqueness for the Bellman equation. The proof of the uniqueness asserted in Theorem 6.1 is similar to the proof of uniqueness for R^n given by Krylov [13] (see also [3, Chapter 4, §5]. One constructs a nearly optimal “feedback control” on a subset of Ω whose complement in Ω has a small measure. The crucial step is in being able to apply Ito’s formula for a function in $W_{loc}^{2,p}(\Omega) \cap C(\bar{\Omega})$ ($p > n$) for a process $d\xi(t) = \sigma(t) dw(t) + b(t) dt$ where $\sigma(t), b(t)$ are bounded and nonanticipative. This, in turn, is established by using Krylov’s inequality [12]

$$\left| E_x \left[\int_0^{T_0} \varphi(\xi(t)) dt \right] \right| \leq C \|\varphi\|_{L^p(\Omega_0)}$$

where Ω_0 is any domain with $\bar{\Omega}_0 \subset \Omega$ and T_0 is the exit time from Ω_0 ; C is a constant depending only on a bound on σ, b, σ^{-1} and the diameter of Ω .

One can represent the solution of (5.1), (5.2) in the form (we take for simplicity $c^i(x) \equiv \alpha$)

$$u(x) = \inf_{\theta} E_x \left[\int_0^T e^{-at} f(\xi(t)) dt \right] \tag{7.12}$$

where $f(\xi(t))$ is defined by (7.4), θ is any sequence of stopping times θ_i increasing to ∞ and $\xi(t)$ switches from each ξ^j to ξ^k according to the rule: $\xi^1, \xi^2, \xi^1, \xi^2, \xi^3, \xi^1, \xi^2, \xi^3, \xi^4, \xi^1, \dots$. This fact follows from the probabilistic interpretation of the solution of the “truncated” Bellman equation.

In the uniqueness proof for the Bellman equation one can choose a nearly optimal “feedback control” $v(t)$ on a subset of Ω whose complement has a

small measure as follows: $v(t)$ assigns the state $\xi^i(t)$ where $\xi(t)$ is in closed set $A_i \subset \Omega$, and the A_i are disjoint sets. Using such controls, we can easily give another proof of the assertion (7.12). \square

REMARK. Uniqueness for a solution of the truncated Bellman equation (5.1) is also immediate from the maximum principle of Bony [4].

We conclude this section with a generalization of Theorem 7.1 to the system of variational inequalities

$$L^i u_i + \sum_{j=1}^m \beta(u_i - u_j - k_{ij}) \ni f^i \quad \text{a.e. in } \Omega,$$

$$u_i = 0 \quad \text{on } \partial\Omega \quad (1 < i < m) \tag{7.13}$$

where $k_{ij} > 0$ if $i \neq j$, $k_{ii} = 0$.

We can again give a probabilistic interpretation of the solution (the switchings cost k_{ij} and need not be cyclic) and prove existence and uniqueness as for Theorem 7.1. Another way to prove existence is to derive estimates on the first derivatives of the solution of the penalized system

$$L^i u_i + \sum_{j=1}^m \beta_\epsilon(u_i - u_j - k_{ij}) = f^i \quad \text{in } \Omega,$$

$$u_i = 0 \quad \text{on } \partial\Omega \quad (1 < i < m) \tag{7.14}$$

by the method of §3 and then proceed as in the proof of Theorem 7.1 to derive $W^{2,p}$ estimates. [The method of §3 requires the additional assumption that $|Df^i| < C$.]

In establishing the $W^{2,p}$ estimates let us notice that for any $x^0 \in \Omega$ we may assume that $u_1 < u_2 < \dots < u_m$ at x^0 , and therefore

$$\beta_\epsilon(u_i - u_j - k_{ij}) = 0 \quad \text{if } j > i$$

in $G_\delta = \{|x - x^0| < \delta\} \cap \Omega$ for some $\delta > 0$ and all ϵ sufficiently small. Therefore

$$L^1 u_1 = f^1, \tag{7.15}$$

$$L^i u_i + \sum_{j=1}^{i-1} \beta_\epsilon(u_i - u_j - k_{ij}) = f^i \tag{7.16}$$

for $2 < i < m$.

Clearly $u_1 \in W^{2,p}_{loc}(G_\delta)$. We can now prove by induction that $u_i \in W^{2,p}_{loc}(G_\delta)$. Indeed, to pass from $i - 1$ to i , we set $\varphi_j = u_j + k_{ij}$ ($1 < j < i - 1$) and multiply (7.16) by $\zeta^p \beta_\epsilon^{p-1}(u_i - \varphi_k)$ ($1 < k < i$), where ζ is a cutoff function

with support in G_δ . By integration by parts,

$$\begin{aligned} & \int L^i u_i \zeta^p \beta_\epsilon^{p-1}(u_i - \varphi_k) \\ &= \int L^i(u_i - \varphi_k) \zeta^p \beta_\epsilon^{p-1}(u_i - \varphi_k) + \int L^i \varphi^k \cdot \zeta^p \beta_\epsilon^{p-1}(u_i - \varphi_k) \\ &> -C \int \zeta^{p-1} |\nabla \zeta| \beta_\epsilon^{p-1}(u_i - \varphi_k) \\ &\quad - \eta \int \zeta^p \beta_\epsilon^p(u_i - \varphi_k) - C(\eta) \int \zeta^p |L^i(\varphi_k)|^p \\ &> -C(\eta) - 2\eta \int \zeta^p \beta_\epsilon^p(u_i - \varphi_k) \quad (\forall \eta > 0). \end{aligned}$$

Since

$$\beta_\epsilon(u_i - \varphi_j) \beta_\epsilon^{p-1}(u_i - \varphi_k) > 0,$$

we easily get

$$\int \zeta^p \beta_\epsilon^p(u_i - \varphi_k) < C.$$

It follows that $L^i u_i \in L^p_{loc}(G_\delta)$, so that $u_i \in W^{2,p}_{loc}(G_\delta)$.

We have proved:

THEOREM 7.3. *Assume that (1.1), (1.2), (1.5) and (7.3) hold. Then there exists a unique solution of (7.13) whose components belong to $W^{2,p}(\Omega)$ for any $p < \infty$.*

8. The case of two operators. Consider now the system of two variational inequalities

$$\begin{aligned} L^1 u^1 + \beta(u^1 - u^2 - k_1) &\ni f^1 \quad \text{in } \Omega, \\ L^2 u^2 + \beta(u^2 - u^1 - k_2) &\ni f^2 \quad \text{in } \Omega, \\ u^1 = u^2 = 0 &\quad \text{on } \partial\Omega \end{aligned} \tag{8.1}$$

where $k_1 > 0, k_2 > 0, \beta(t) = \{0\}$ if $t < 0, \beta(0) = [0, \infty]$. Consider also the Bellman equation

$$\begin{aligned} \max\{L^1 u(x) - f^1(x), L^2 u(x) - f^2(x)\} &= 0 \quad \text{in } \Omega, \\ u &= 0 \quad \text{on } \partial\Omega. \end{aligned} \tag{8.2}$$

We assume, in addition to (1.1)–(1.3) that

$$f^i \in W^{1,p}(\Omega) \quad \text{for some } p > n, \tag{8.3}$$

$$c^1(x) > \alpha_0, \quad c^2(x) > \alpha_0, \tag{8.4}$$

where α_0 is sufficiently large depending on the coefficients of L^i and on Ω . Then [5] there exists a unique solution u of (8.2) in $H^3(\Omega) \cap H^1_0(\Omega)$, and it belongs to $C^{2,\alpha}(\Omega_0)$ for some $\alpha > 0$ and for any domain Ω_0 with $\bar{\Omega}_0 \subset \Omega$. Further, by [10], $u \in C(\bar{\Omega})$.

We denote the solution of (8.1) by (u_k^1, u_k^2) where $k = (k_1, k_2)$.

THEOREM 8.1. *If (1.1)–(1.3) and (8.3), (8.4) hold, then*

$$u_k^i(x) \rightarrow u(x) \text{ weakly in } H^2(\Omega) \text{ as } k \rightarrow 0. \tag{8.5}$$

This theorem was stated in §7 for a general system (see Theorem 7.2) in case α_0 is any positive number, but only where the coefficients $a_{ij}^k(x)$ are constants.

PROOF. We shall use below an inequality of Sobolevsky [18] (for proof see, for instance, [5]):

$$\int_{\Omega} L^1 v \cdot L^2 v \, dx \geq C \|v\|_{H^2(\Omega)}^2 \quad \forall v \in H^2(\Omega) \cap H_0^1(\Omega), \quad C > 0. \tag{8.6}$$

Clearly (8.1) implies

$$k_2 \leq u_k^1(x) - u_k^2(x) \leq k_1. \tag{8.7}$$

We shall now prove that

$$\|u_k^i\|_{H^2(\Omega)} \leq C \quad (i = 1, 2) \tag{8.8}$$

where C is a generic positive constant independent of k .

Set for simplicity $u_k^1 = u, u_k^2 = v, L^1 = L, L^2 = M, f^1 = f, f^2 = g$ and

$$\gamma_i = f^i - L^i u_k^i \in \beta(u^i - v^j - k_i) \quad (i, j = 1, 2; i \neq j).$$

Then (8.1) may be written in the form $Lu + \gamma_1 = f, Mv + \gamma_2 = g$ a.e. in Ω . It follows that

$$\int_{\Omega} Lu(Mu - Mv) + \int_{\Omega} \gamma_1(Mu - Mv) = \int_{\Omega} f(Mu - Mv), \tag{8.9}$$

$$\int_{\Omega} Mv(Lv - Lu) + \int_{\Omega} \gamma_2(Lv - Lu) = \int_{\Omega} g(Lv - Lu). \tag{8.10}$$

By the coercivity of M and L it follows that the second terms on the left-hand sides of (8.9), (8.10) are nonnegative (cf. Brezis-Strauss [7, Lemma 2]).

Adding (8.9), (8.10) and using the Sobolevsky inequality we get

$$C(\|u\|_{H^2}^2 + \|v\|_{H^2}^2) \leq \int_{\Omega} f(Mu - Mv) + \int_{\Omega} g(Lv - Lu) + 2 \int_{\Omega} MvLu. \tag{8.11}$$

Observe next that $\gamma_1 \gamma_2 = 0$ a.e., and so $(Lu - f)(Mv - g) = 0$ a.e. Hence

$$\int_{\Omega} MvLu \leq \varepsilon(\|u\|_{H^2}^2 + \|v\|_{H^2}^2) + \frac{C}{\varepsilon} \int_{\Omega} (f^2 + g^2).$$

Using this in (8.11), the assertion (8.8) follows.

We can now take a subsequence of (u_k, v_k) , which we again denote by (u_k, v_k) , such that (recall (8.7))

$$u_k, v_k \rightarrow w \text{ weakly in } H^2(\Omega). \tag{8.12}$$

Since $Lu_k < f, Mv_k < g$, we get $Lw < f, Mw < g$, so that

$$\Phi \equiv \max(Lw - f, Mw - g) < 0, \quad w \in H^2(\Omega) \cap H_0^1(\Omega). \quad (8.13)$$

We shall prove that $w = u$, where u is the solution of (8.2).

From the probabilistic interpretation of u_k, v_k we deduce that

$$u < u_k, \quad u < v_k; \quad (8.14)$$

therefore

$$u < w. \quad (8.15)$$

This we can also prove by the maximum principle. Indeed, if (8.14) is not true then without loss of generality we may assume that there exists a point $x^0 \in \Omega$ such that

$$(u - u_k)(x^0) = \max_{\bar{\Omega}}(u - u_k) > 0, \quad (8.16)$$

$$\max_{\bar{\Omega}}(u - u_k) > \max_{\bar{\Omega}}(u - v_k). \quad (8.17)$$

Then, by Bony [4],

$$f(x^0) \geq Lu(x^0) > \operatorname{ess\,lim\,sup}_{x \rightarrow x^0} Lu_k(x) = f(x^0) - \operatorname{ess\,lim\,inf}_{x \rightarrow x^0} \gamma_1,$$

so that $\operatorname{ess\,lim\,inf} \gamma_1 > 0$. Hence $u_k(x^0) > v_k(x^0)$. It follows that $(u - v_k)(x^0) > (u - u_k)(x^0)$, which contradicts (8.16), (8.17).

Having proved (8.15), it remains to prove that $w \geq u$. Notice, by (8.13), that

$$\max\{Lw - (f + \Phi), Mw - (g + \Phi)\} \equiv 0$$

and $f + \Phi < f, g + \Phi < g$. Hence it suffices to prove a comparison theorem for the Bellman equation, namely,

$$\text{if } f < \tilde{f}, g < \tilde{g} \text{ then } \tilde{u} < u \quad (8.18)$$

where \tilde{u} is the solution corresponding to \tilde{f}, \tilde{g} .

For smooth $f, \tilde{f}, g, \tilde{g}$ the proof follows by the probabilistic interpretation of u, \tilde{u} or by a simple maximum principle argument using the $C^{2+\alpha}(\Omega) \cap C(\bar{\Omega})$ regularity of u and \tilde{u} .

To prove (8.18) for general $f, \tilde{f}, g, \tilde{g}$, we approximate these functions by smooth functions and use the following lemma. \square

LEMMA 8.2. Let $\max(Lu_m - f_m, Mu_m - g_m) = 0$ a.e. in $\Omega, u_m \in H^2(\Omega) \cap H_0^1(\Omega)$. If $f_m \rightarrow f, g_m \rightarrow g$ in $L^2(\Omega)$, then $u_m \rightarrow u$ in $H^2(\Omega)$.

PROOF. We have $Lu_m + \beta(Mu_m - g_m) \ni f_m, Lu + \beta(Mu - g) \ni f$. Hence

$$L(u_m - u) + \gamma_m - \gamma = f_m - f \quad (8.19)$$

where $\gamma_m \in \beta(Mu_m - g_m), \gamma \in \beta(Mu - g)$. Multiplying (8.19) by $M(u_m - u) + g - g_m$ and integrating over Ω we obtain, after using the Sobolevsky

inequality,

$$C\|u_m - u\|_{H^2}^2 \leq \frac{C}{\varepsilon} (\|f_m - f\|_{L^2}^2 + \|g_m - g\|_{L^2}^2) + \varepsilon\|u_m - u\|_{H^2}^2 \\ - \int_{\Omega} (\gamma_m - \gamma)(M(u_m - u) + g - g_m).$$

Since the last integral is nonnegative (by the monotonicity of β), the assertion of the lemma follows. \square

REFERENCES

1. V. Barbu, *Nonlinear semigroups and differential equations in Banach spaces*, Noordhoff, Leyden, The Netherlands, 1976.
2. A. Bensoussan and J. L. Lions, *Contrôle impulsionnel et systèmes d'inéquations quasi variationnelles*, C. R. Acad. Sci. Paris **278** (1974), 747–751.
3. ———, *Applications des inéquations variationnelles en contrôle stochastique*, Dunod, Paris, 1978.
4. J.-M. Bony, *Principe du maximum dans les espaces de Sobolev*, C. R. Acad. Sci. Paris **265** (1967), 333–336.
5. H. Brézis and L. C. Evans, *A variational inequality approach to the Bellman-Dirichlet equation for two elliptic operators*, Arch. Rational Mech. Anal. (to appear).
6. H. Brézis and D. Kinderlehrer, *The smoothness of solutions to nonlinear variational inequalities*, Indiana Univ. Math. J. **9** (1974), 831–844.
7. H. Brézis and W. Strauss, *Semi-linear second order elliptic equations in L^1* , J. Math. Soc. Japan **25** (1973), 565–590.
8. M. Chicco, *Principio di massimo per soluzioni di equazioni ellittiche del secondo ordine di tipo Cordes*, Ann. Mat. Pura Appl. (4) **100** (1974), 239–258.
9. L. C. Evans, *A convergence theorem for solutions of nonlinear second order elliptic equations*, Indiana Univ. Math. J. **27** (1978), 875–887.
10. L. C. Evans and P. L. Lions, *Deux résultats de régularité pour le problème de Bellman-Dirichlet*, C. R. Acad. Sci. Paris Sér. A-B **286** (1978), 587–589.
11. A. Friedman, *Stochastic differential equations and applications*. vol. II, Academic Press, New York, 1976.
12. N. V. Krylov, *An inequality in the theory of stochastic integrals*, Theor. Probability Appl. **16** (1971), 438–448.
13. ———, *Control of a solution of a stochastic integral equation*, Theor. Probability Appl. **17** (1972), 114–130.
14. M. Nisio, *Remarks on stochastic optimal controls*, Japan J. Math. **1** (1975), 159–183.
15. S. Pliska, *A semigroup representation of the maximum expected reward vector in continuous parameter Markov decision theory*, SIAM J. Control **13** (1975), 1115–1129.
16. K. Sato, *On the generators of non-negative contraction semigroups in Banach spaces*, J. Math. Soc. Japan **20** (1968), 423–436.
17. E. Sinestrani, *Accretive differential operators*, Boll. Un. Mat. Ital. **13** (1976), 19–31.
18. P. Sobolevsky, *Equations with operators constituting an acute angle*, Dokl. Akad. Nauk SSSR **116** (1957), 754–757. (Russian)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KENTUCKY, LEXINGTON, KENTUCKY 40506

DEPARTMENT OF MATHEMATICS, NORTHWESTERN UNIVERSITY, EVANSTON, ILLINOIS 60201