

THE ATOMIC MODEL THEOREM AND TYPE OMITTING

DENIS R. HIRSCHFELDT, RICHARD A. SHORE, AND THEODORE A. SLAMAN

ABSTRACT. We investigate the complexity of several classical model theoretic theorems about prime and atomic models and omitting types. Some are provable in RCA_0 , and others are equivalent to ACA_0 . One, that every atomic theory has an atomic model, is not provable in RCA_0 but is incomparable with WKL_0 , more than Π_1^1 conservative over RCA_0 and strictly weaker than all the combinatorial principles of Hirschfeldt and Shore (2007) that are not Π_1^1 conservative over RCA_0 . A priority argument with Shore blocking shows that it is also Π_1^1 -conservative over $\text{B}\Sigma_2$. We also provide a theorem provable by a finite injury priority argument that is conservative over $\text{I}\Sigma_1$ but implies $\text{I}\Sigma_2$ over $\text{B}\Sigma_2$, and a type omitting theorem that is equivalent to the principle that for every X there is a set that is hyperimmune relative to X . Finally, we give a version of the atomic model theorem that is equivalent to the principle that for every X there is a set that is not recursive in X , and is thus in a sense the weakest possible natural principle not true in the ω -model consisting of the recursive sets.

1. INTRODUCTION

In this paper we investigate the complexity of a few classical model theoretic theorems about prime and atomic models and about omitting types in terms of both computational (recursion theoretic) and proof theoretic (reverse mathematical) calibrations and of the interplay between these approaches. The theorems of interest in such investigations are typically of the form $\forall A (\Theta(A) \rightarrow \exists B \Phi(A, B))$, where Θ and Φ are arithmetic and $A, B \in 2^{\mathbb{N}}$. Thus, from the recursion theoretic point of view, we want to bound or characterize the computational complexity of B given an A satisfying Θ (typically in terms of Turing degree or place in one of the standard arithmetic/analytic definability or jump hierarchies). From the reverse mathematics point of view we want to determine the axiom systems in which the theorem is provable (typically subsystems of second order arithmetic determined by the amount of comprehension assumed). Here, characterizations correspond to reversals in the sense that one proves (over some weak system) the axioms of one of the subsystems of second order arithmetic from the statements of the mathematical theorems being investigated.

Received by the editors July 25, 2007.

2000 *Mathematics Subject Classification*. Primary 03B30, 03C15, 03C50, 03C57, 03D45, 03F35.

The first author's research was partially supported by NSF Grants DMS-0200465 and DMS-0500590.

The second author's research was partially supported by NSF Grants DMS-0100035 and DMS-0554855.

The third author's research was partially supported by NSF Grants DMS-9988644 and DMS-0501167.

We briefly review the five standard systems of reverse mathematics. For completeness, we include systems stronger than arithmetical comprehension, but these will play no part in this paper. Details, general background, and results, as well as many examples of reversals, can be found in Simpson [32], the standard text on reverse mathematics. Each of the systems is given in the language of second order arithmetic, that is, the usual first order language of arithmetic augmented by set variables and the membership relation \in . Each contains the standard basic axioms for $+$, \cdot , and $<$ (which say that \mathbb{N} is an ordered semiring). In addition, they all include a form of induction that applies only to sets (that happen to exist):

$$(\mathbf{I}_0) \quad (0 \in X \wedge \forall n (n \in X \rightarrow n + 1 \in X)) \rightarrow \forall n (n \in X).$$

We call the system consisting of \mathbf{I}_0 and the basic axioms of ordered semirings \mathbf{P}_0 . All the five standard systems are defined by adding various types of set existence axioms to \mathbf{P}_0 . They also correspond to classical construction principles in recursion theory.

(RCA₀) Recursive Comprehension Axioms: This is a system just strong enough to prove the existence of the recursive sets but not of $0'$ nor indeed of any nonrecursive set. In addition to \mathbf{P}_0 its axioms include the schemes of Δ_1^0 comprehension and Σ_1^0 induction:

$$\begin{aligned} (\mathbf{\Delta}_1^0\text{-CA}_0) \quad & \forall n (\varphi(n) \leftrightarrow \psi(n)) \rightarrow \exists X \forall n (n \in X \leftrightarrow \varphi(n)) \text{ for all} \\ & \Sigma_1^0 \text{ formulas } \varphi \text{ and } \Pi_1^0 \text{ formulas } \psi \text{ in which } X \text{ is not free.} \\ (\mathbf{I\Sigma}_1) \quad & (\varphi(0) \wedge \forall n (\varphi(n) \rightarrow \varphi(n + 1))) \rightarrow \forall n \varphi(n) \text{ for all } \Sigma_1^0 \text{ for-} \\ & \text{mulas } \varphi. \end{aligned}$$

The next system says that every infinite binary tree has an infinite path. It is connected to the Low Basis Theorem (Jockusch and Soare [24]) of recursion theory, which says that every such tree has an infinite path whose jump is recursive in that of the tree itself.

(WKL₀) Weak König's Lemma: This system consists of \mathbf{RCA}_0 plus the statement that every infinite subtree of $2^{<\mathbb{N}}$ has an infinite path.

We next move up to arithmetic comprehension.

(ACA₀) Arithmetic Comprehension Axioms: This system consists of \mathbf{RCA}_0 plus the axioms $\exists X \forall n (n \in X \leftrightarrow \varphi(n))$ for every arithmetic formula φ in which X is not free.

In recursion theoretic terms, \mathbf{ACA}_0 proves the existence of $0'$ and, by relativization, it proves, and in fact is equivalent to, the existence of X' for every set X .

The next system corresponds to the existence of all (relativized) H -sets, i.e. the existence, for every set X , of the H_e^X (and so the hyperarithmetical hierarchy up to e) for each $e \in \mathcal{O}^X$, the hyperjump of X . It says that arithmetic comprehension can be iterated along any countable well order.

(ATR₀) Arithmetical Transfinite Recursion: This system consists of \mathbf{RCA}_0 plus the following axiom. If X is a set coding a well order $<_X$ with domain D and Y is a code for a set of arithmetic formulas $\varphi_x(z, Z)$ (indexed by $x \in D$) each with one free set variable and one free number variable, then there is a sequence $\langle K_x \mid x \in D \rangle$ of sets such that if y is the immediate successor of x in $<_X$, then $\forall n (n \in K_y \leftrightarrow \varphi_x(n, K_x))$, and if x is a limit point in $<_X$, then K_x is $\bigoplus \{K_y \mid y <_X x\}$.

The systems climbing up to full second order arithmetic (i.e. comprehension for all formulas) are classified by the syntactic level of the second order formulas for which we assume a comprehension axiom.

($\Pi_n^1\text{-CA}_0$) Π_n^1 Comprehension Axioms: $\exists X \forall k (k \in X \leftrightarrow \varphi(k))$ for every Π_n^1 formula φ in which X is not free.

The recursion theoretic equivalent of the simplest of these systems, $\Pi_1^1\text{-CA}_0$, is the existence of \mathcal{O}^X for every set X . Together with the four systems listed above, this system makes up the standard list of the axiomatic systems of reverse mathematics. Almost all theorems of classical mathematics whose proof theoretic complexities have been determined have turned out to be equivalent to one of them.

Below, we will also consider induction and bounding principles. In the definition of $\text{I}\Sigma_1$ above, if we replace Σ_1^0 formulas by Σ_k^0 or Π_k^0 formulas, we obtain the principles $\text{I}\Sigma_k$ and $\text{I}\Pi_k$, respectively. The principles $\text{I}\Sigma_k$ and $\text{I}\Pi_k$ are equivalent over RCA_0 (see Exercise II.3.12 in Simpson [32]). The Σ_k bounding principle is defined as follows.

($\mathbf{B}\Sigma_k$) $(\forall i < n)(\exists x)\varphi(x) \rightarrow (\exists u)(\forall i < n)(\exists x < u)\varphi(x)$ for all Σ_k^0 formulas φ .

Over RCA_0 , the principle $\mathbf{B}\Sigma_k$ is strictly between $\text{I}\Sigma_{k-1}$ and $\text{I}\Sigma_k$ (Paris and Kirby [30]; see also Hájek and Pudlák [16]).

The early connections between recursion theoretic ideas and methods on the one hand and reverse mathematics on the other typically involved recursive mathematics, diagonalization or finite injury arguments, and coding. Consider, for example, a theorem of the form $\forall A (\Theta(A) \rightarrow \exists B \Phi(A, B))$, where Θ and Φ are arithmetic. We call B a *solution* for the instance of the theorem specified by A if $\Theta(A) \rightarrow \Phi(A, B)$.

A construction of recursive mathematics that shows that there is a solution B recursive in any given A generally shows that the theorem is provable in RCA_0 . One that shows that B can be obtained arithmetically in A usually shows that the theorem is provable in ACA_0 . Standard forms of applications of recursion theoretic diagonalization or finite injury results to reverse mathematical calibrations are as follows:

- (1) If there is no solution B recursive in some given A , then the theorem is not provable in RCA_0 .
- (2) If there is no solution B that is low over A , i.e. $(B \oplus A)' \equiv_{\text{T}} A'$, then the theorem is not provable in WKL_0 .
- (3) If there is no solution B arithmetic in A , then the theorem is not provable in ACA_0 .

Coding methods tend to give reversals.

- (1) If, for any recursive tree T , there is a recursive instance of the theorem such that any solution codes a path through T , then the theorem usually implies WKL_0 .
- (2) If there is a recursive instance of the theorem such that any solution computes $0'$, then the theorem usually implies ACA_0 .
- (3) If there is a recursive instance of the theorem such that any solution computes \mathcal{O} , then the theorem usually implies $\Pi_1^1\text{-CA}_0$.

In this paper we will analyze several theorems that fall into this standard classification scheme and some that fall well outside. We are also particularly interested in reverse mathematical relationships derived from more complex recursion theoretic results and constructions.

In this paper all theories T in a language \mathcal{L} are deductively closed ($T \vdash \varphi \Rightarrow \varphi \in T$), complete ($T \vdash \varphi$ or $T \vdash \neg\varphi$ for every sentence φ of \mathcal{L}) and consistent (there is no formula φ such that $T \vdash \varphi$ and $T \vdash \neg\varphi$). All languages and models are countable. For the purposes of reverse mathematics, we include with a model \mathcal{A} of

T the function interpreting terms and the full satisfaction predicate for formulas with constants from the model. Formal definitions suitable for work in RCA_0 can be found in Simpson [32, Section II.8] along with, for example, a proof in RCA_0 that any complete consistent theory has a model in this sense.

Definition 1.1. A *partial type* of a theory T is a set of formulas in a fixed number of free variables that is consistent with T . We will reserve the unmodified term *type* for complete types, i.e., maximal partial types. A partial type Γ is *realized* in a model \mathcal{A} of T if there is an \vec{a} such that $\mathcal{A} \models \varphi(\vec{a})$ for every $\varphi \in \Gamma$. Otherwise, Γ is *omitted* in \mathcal{A} .

A formula $\varphi(x_1, \dots, x_n)$ of T is an *atom* of T if for each formula $\psi(x_1, \dots, x_n)$ we have $T \vdash \varphi \rightarrow \psi$ or $T \vdash \varphi \rightarrow \neg\psi$, but not both. A partial type Γ is *principal* if there is a formula φ consistent with T such that $T \vdash \varphi \rightarrow \psi$ for all $\psi \in \Gamma$. Thus a complete type is principal if and only if it contains an atom of T .

The theory T is *atomic* if, for every formula $\psi(x_1, \dots, x_n)$ consistent with T , there is an atom $\varphi(x_1, \dots, x_n)$ of T extending it, i.e. one such that $T \vdash \varphi \rightarrow \psi$. A model \mathcal{A} of T is *atomic* if every n -tuple from \mathcal{A} satisfies an atom of T , that is, every type realized in \mathcal{A} is principal. The model \mathcal{A} is *prime* if it can be elementarily embedded in every model of T .

The standard classical theorems about these notions (as in e.g. Chang and Keisler [1, 2.3]) are as follows:

- (1) *AMT* (Existence): T has an atomic model iff T is atomic.
- (2) *AUniqueness*: If \mathcal{A} and \mathcal{B} are atomic models of T , then $\mathcal{A} \cong \mathcal{B}$.
- (3) *PMT* (Existence): T has a prime model iff T is atomic.
- (4) *PUniqueness*: If \mathcal{A} and \mathcal{B} are prime models of T , then $\mathcal{A} \cong \mathcal{B}$.
- (5) *Equivalence*: A model of T is atomic iff it is prime.

Counting each biconditional as two implications, this list provides eight implications to analyze in terms of reverse mathematics. Three are easily seen in §2 to be provable in RCA_0 (the “only if” directions of AMT (1) and PMT (3) and the “if” direction of Equivalence (5)). We prove in §2 as well that three others are equivalent to ACA_0 (AUniqueness (2); the “only if” direction of PMT (3) and the “only if” direction of Equivalence (5)). Of the remaining two, the implication PUniqueness (4) is easily provable in ACA_0 , but its status in terms of reversals or connections to other systems remains completely open. Thus, when we refer below to principles considered in this paper, we always exclude PUniqueness. It is the “if” direction of AMT (1), which we call simply AMT, that turns out to be quite interesting.

(AMT) Atomic Model Theorem: Every complete atomic theory has an atomic model.

AMT is an exceptionally weak principle but nonetheless not provable in RCA_0 or even WKL_0 , as we will show below in Corollary 3.4. It turns out to be strictly weaker than most of the combinatorial principles studied in Hirschfeldt and Shore [21], which were themselves shown to be weaker than the classical Ramsey’s Theorem for pairs that has long been of interest both recursion theoretically and in reverse mathematics. We refer to Cholak, Jockusch and Slaman [2] for the best analysis of Ramsey’s Theorem itself.

Our general analysis and, in particular, our proof that AMT is implied by the weakest of these combinatorial principles (SADS, as defined below) are related to

various forcing constructions and priority arguments. Some of them supply more general conservation results; others touch on the issue of the role of $B\Sigma_2$ and $I\Sigma_2$ in (reverse) recursion theory. In particular, AMT is shown to be conservative over not only RCA_0 ($I\Sigma_1$) and $I\Sigma_2$ but over $B\Sigma_2$ as well. This last argument uses a finite injury argument with Shore blocking to carry out the relevant construction in $B\Sigma_2$. (We know of only two other examples of principles that are known to be conservative over $B\Sigma_2$: WKL_0 by Hájek [15] and COH by Chong, Slaman and Yang [3].) On the other hand, we are lead in §4 to a single statement about the existence of certain types of generics (a bit less than weakly 2-generics) that can be proven by a finite injury priority argument and that is conservative over both $I\Sigma_1$ and $I\Sigma_2$ but not over $B\Sigma_2$, as it implies $I\Sigma_2$ over $B\Sigma_2$. This seems to be the first example of such a theorem.

The existence of atomic models can be seen not only as a forcing/genericity construction but also (as is traditional in model theory) as a type omitting argument. In §5, we investigate the strength of another type omitting argument and show that it is equivalent (both recursion theoretically and in the sense of reverse mathematics) to the existence of hyperimmune sets. In §6, we give a weaker version of AMT that is equivalent to the existence of nonrecursive sets, and is thus in a sense the weakest possible natural principle not true in the ω -model consisting of the recursive sets. (An ω -model of a fragment of second order arithmetic is one whose first order part is standard, and hence is determined entirely by which subsets of ω it contains.)

To set the stage for our eventual analysis of AMT we define here many of the principles studied in [2] and [21] and summarize the relationships among them in Diagram 1 below. (Single arrows are implications; double arrows are strict implications and negated arrows represent known nonimplications. See [21] for attributions and references.) Each principle is to be understood as an addition to RCA_0 .

Definition 1.2. An n -coloring of $[\mathbb{N}]^k$, the unordered k -tuples (n_1, \dots, n_k) of natural numbers (listed by convention in increasing order), is a map $f : [\mathbb{N}]^k \rightarrow n$. A subset H of \mathbb{N} is *homogeneous* for the coloring f if H is infinite and $|f''[H]^k| = 1$ (i.e. all tuples from H have the same color). Unless otherwise stated all colorings will be 2-colorings of $[\mathbb{N}]^2$.

(RT₂²) Ramsey's Theorem for pairs: Every 2-coloring of $[\mathbb{N}]^2$ has a homogeneous set.

Definition 1.3. A coloring f of $[\mathbb{N}]^2$ is *stable* if $(\forall x)(\exists y)(\forall z > y)[f(x, y) = f(x, z)]$.

(SRT₂²) Stable Ramsey's Theorem for pairs: Every *stable* 2-coloring of $[\mathbb{N}]^2$ has a homogeneous set.

Definition 1.4. If $\vec{R} = \langle R_i \mid i \in \mathbb{N} \rangle$ is a sequence of sets, an infinite set S is \vec{R} -cohesive if $(\forall i)(\exists s)[(\forall j > s)(j \in S \rightarrow j \in R_i) \vee (\forall j > s)(j \in S \rightarrow j \notin R_i)]$.

(COH) Cohesive Principle: For every sequence $\vec{R} = \langle R_i \mid i \in \mathbb{N} \rangle$ there is an \vec{R} -cohesive set.

We will denote the n th partial recursive in A function (in some standard effective listing) by Φ_n^A .

(DNR) Diagonally Nonrecursive Principle: For every set A there is a function f that is diagonally nonrecursive relative to A , i.e. $\forall n \neg(f(n) = \Phi_n^A(n))$.

(CAC) Chain-AntiChain Principle: Every infinite partial order (P, \leq_P) has an infinite subset S that is either a *chain*, i.e. $(\forall x, y \in S)(x \leq_P y \vee y \leq_P x)$, or an *antichain*, i.e. $(\forall x, y \in S)(x \not\leq_P y \wedge y \not\leq_P x)$.

Definition 1.5 (RCA_0). An infinite linear order (L, \leq_L) in which all nonfirst elements have immediate predecessors and all nonlast ones have immediate successors has type

- ω if every element has finitely many predecessors;
- ω^* if every element has finitely many successors;
- $\omega + \omega^*$ if it does not have type ω or ω^* and every element has either finitely many predecessors or finitely many successors.

(ADS) Ascending or Descending Sequence: Every infinite linear order (L, \leq_L) has an infinite subset S that is either an ascending sequence, i.e. $(\forall s < t)(s, t \in S \rightarrow s <_L t)$, and so of order type ω , or a descending sequence, i.e. $(\forall s < t)(s, t \in S \rightarrow t <_L s)$, and so of order type ω^* .

(SADS) Stable ADS: Every linear order of type $\omega + \omega^*$ has a subset of order type ω or ω^* .

(CADS) Cohesive ADS: Every linear order has a subset S of order type ω , ω^* , or $\omega + \omega^*$.

Definition 1.6. A partial order \mathcal{P} is *stable* if either

$$(\forall i \in P)(\exists s)[(\forall j > s)(j \in P \rightarrow i <_P j) \vee (\forall j > s)(j \in P \rightarrow i \mid_P j)]$$

or

$$(\forall i \in P)(\exists s)[(\forall j > s)(j \in P \rightarrow i >_P j) \vee (\forall j > s)(j \in P \rightarrow i \mid_P j)].$$

(SCAC) Stable CAC: Every infinite stable partial order has an infinite chain or antichain.




Diagram 1

2. WITHIN RCA_0 , ACA_0 AND REVERSALS

We begin with the two facts provable in RCA_0 . The first is the “only if” direction of AMT (1) and requires simply the classical proof.

Proposition 2.1 (RCA_0). *If T has an atomic model, T is atomic.*

Proof. Let \mathcal{A} be an atomic model of T . Consider any formula $\psi(x_1, \dots, x_n)$ consistent with T . As T is complete, $T \vdash \exists x_1, \dots, x_n \psi(x_1, \dots, x_n)$, and so there are c_1, \dots, c_n in \mathcal{A} such that $\mathcal{A} \models \psi(c_1, \dots, c_n)$. As \mathcal{A} is atomic there is an atom $\varphi(x_1, \dots, x_n)$ of T such that $\mathcal{A} \models \varphi(c_1, \dots, c_n)$. Thus $\varphi(x_1, \dots, x_n)$ is the required atom of T extending ψ . \square

The second is the “if” direction of Equivalence (5) and uses the effective version of the omitting types theorem in the classical proof.

Proposition 2.2 (RCA_0). *Every prime model \mathcal{A} of T is atomic.*

Proof. Consider any n -tuple c_1, \dots, c_n of \mathcal{A} and the set S of formulas it satisfies, $S = \{\psi(x_1, \dots, x_n) \mid \mathcal{A} \models \psi(c_1, \dots, c_n)\}$. We wish to show that S contains an atom of T . If not, the classical omitting types theorem says that there is a model \mathcal{B} of T that fails to realize S , i.e. there are no elements d_1, \dots, d_n of \mathcal{B} such that $S = \{\psi(x_1, \dots, x_n) \mid \mathcal{B} \models \psi(d_1, \dots, d_n)\}$. In this situation there could not be an elementary embedding of \mathcal{A} into \mathcal{B} , for the desired contradiction. Thus we need only verify in RCA_0 that there is such a model \mathcal{B} . The usual effective version of the Henkin construction for the omitting types theorem, as for example in Harizanov [17, 6.1], provides in RCA_0 a complete consistent set of sentences in the language with Henkin constants in which every n -tuple d_1, \dots, d_n satisfies $\neg\psi(d_1, \dots, d_n)$ for some $\psi(d_1, \dots, d_n) \in S$. Simpson [32, II.8.6] then constructs the required model in RCA_0 . \square

These two propositions together obviously supply the proof in RCA_0 of the “only if” direction of PMT (3). We now prove that three other of the eight basic implications are equivalent to ACA_0 .

Theorem 2.3. *The following assertions are equivalent (over RCA_0):*

- (1) ACA_0 .
- (2) *Every atomic model of T is prime.*
- (3) *A Uniqueness.*
- (4) *Every atomic theory T has a prime model.*

Proof. The classical proofs of assertions (2)–(4) can be easily carried out in ACA_0 . We construct a theory T that will establish all the implications in the other direction. The language of T has unary predicates R_i and $R_{i,j}$ for $i, j \in \mathbb{N}$. The theory T is the deductive closure of the following set S of axioms.

Axioms for T :

1. Axioms asserting that the R_i define infinite, pairwise disjoint sets.
2. $R_{i,s}(x) \rightarrow R_i(x)$ for every $i, s \in \mathbb{N}$.
3. $R_{i,s}(x) \rightarrow R_{i,s+1}(x)$.
4. If $\Phi_i(i)$ does not converge in fewer than s many steps, then the axiom $\neg R_{i,s}(x)$ is included.

5. If $\Phi_i(i)$ converges in exactly s many steps, then there are axioms asserting that there are infinitely many x such that $R_{i,s}(x)$ and infinitely many x such that $\neg R_{i,s}(x)$.
6. If $\Phi(i)$ converges in exactly s many steps, then the axioms $\neg R_{i,s}(x) \rightarrow \neg R_{i,t}(x)$ are included for every $t > s$.

This set S of axioms is clearly recursive and so provably exists in RCA_0 . We claim that S admits quantifier elimination, i.e. for every formula $\psi(\bar{x})$ there is a quantifier free $\varphi(\bar{x})$ (possibly just the formal propositional symbols T, for true, or F, for false) with the same free variables such that $S \vdash \varphi(\bar{x}) \leftrightarrow \psi(\bar{x})$. This claim implies that S is complete, as any sentence is equivalent to T or F modulo S , and so its deductive closure is Δ_1 definable and provably exists in RCA_0 . This closure is our desired theory T .

To see that S admits quantifier elimination it suffices to present a recursive procedure that finds a quantifier free equivalent $\varphi(\bar{x})$ for a given existential formula $\exists \bar{y} \varphi(\bar{x}, \bar{y})$, as one can then carry out the usual induction (for the provable, in S , equivalence of the resulting formulas) in RCA_0 by IS_1 . By the usual applications of DeMorgan's laws and the distribution of \exists over \vee , we may assume φ is a conjunction of atomic formulas and negations of atomic formulas. As the language has only unary predicates the only atomic formulas are of the form $R_i(z)$ or $R_{i,s}(z)$ for $z \in \bar{x}\bar{y}$.

We begin with the following simple consistency checks. If any of them fail, the formula is inconsistent with S and its equivalent is F.

Consistency checks:

- (1) Is φ propositionally consistent, i.e. is there no occurrence of both $R(z)$ and $\neg R(z)$ for any relation R and variable z ?
- (2) Is φ consistent with axiom 1, i.e. does it not contain, for any z and $i \neq j$, both $R_i(z)$ and $R_j(z)$?
- (3) Is φ consistent with axiom 2, i.e. if it contains some $R_{i,s}(z)$, then does it not contain $\neg R_i(z)$?
- (4) Is φ consistent with axiom 3, i.e. if it contains some $R_{i,s}(z)$, then does it not contain $\neg R_{i,t}(z)$ for any $t > s$?
- (5) Is φ consistent with axiom 4, i.e. if $\Phi_i(i)$ does not converge in fewer than s many steps, then does $R_{i,s}(z)$ not occur in φ ?
- (6) Is φ consistent with axiom 6, i.e. if $\neg R_{i,s}(z)$ and $R_{i,t}(z)$ occur in φ with $s < t$, then does $\Phi_i(i)$ converge in exactly n steps with $s < n \leq t$?

If φ passes all these tests, then the required equivalent to $\exists \bar{y} \varphi(\bar{x}, \bar{y})$ is obtained by replacing all the atomic and negated atomic conjuncts about any $z \in \bar{y}$ with T. It is straightforward to see that the result is equivalent to $\exists \bar{y} \varphi(\bar{x}, \bar{y})$ modulo S . Thus S is complete, as any sentence is equivalent to T or F modulo S .

We now determine the atoms of T . As the language has only unary predicates, the formulas satisfied by an n -tuple are determined by the ones satisfied by the individual elements of the tuple. Thus it suffices to determine the quantifier free formulas with one free variable x that decide all atomic formulas about x . If $\Phi_i(i)$ is divergent, then $R_i(x)$ is such a formula since it implies $\neg R_{i,s}(x)$ for every s and also $\neg R_j(x)$ and $\neg R_{j,t}(x)$ for every $j \neq i$ and every t . If $\Phi_i(i)$ is convergent and converges in exactly s many steps, then $R_{i,s}(x)$ and $\neg R_{i,s}(x) \wedge R_i(x)$ are such

formulas. The first implies $R_i(x)$, implies $R_{i,t}(x)$ for every $t > s$ and $\neg R_{i,t}(x)$ for every $t < s$ and implies $\neg R_j(x)$ and $\neg R_{j,t}(x)$ for every $j \neq i$ and every t . The second one implies $\neg R_{i,t}(x)$ for every t and also $\neg R_j(x)$ and $\neg R_{j,t}(x)$ for every $j \neq i$ and every t . In any model \mathcal{A} of T any element that satisfies some R_i satisfies one of these atoms. Thus any model in which every element is in some R_i is atomic.

Finally, we claim that T is atomic. Again we need to consider only quantifier free formulas $\varphi(x)$ with one free variable, which we can take to be a conjunct of atomic formulas about x and negations of such formulas. (If we can extend all such consistent formulas to atoms, we can extend any consistent disjunction of them to an atom and so any formula to an atom, as required.) If φ fails one of the above consistency checks, it is inconsistent with T and so of no concern. Otherwise, if it contains some $R_i(x)$ or $R_{i,s}(x)$ as a conjunct, then we can obviously extend it to one of the atoms described above. Otherwise, it contains only negated atomic formulas. In this case the atom $R_i(x)$ extends φ for any i larger than all j such that R_j or any $R_{j,t}$ appears in φ and $\Phi_j(j)$ is divergent.

To deduce ACA_0 from each of assertions (2)–(4) of the theorem, we build two atomic models \mathcal{A}, \mathcal{B} of T such that if there is a model \mathcal{C} with embeddings into both \mathcal{A} and \mathcal{B} , then $0'$ is recursive in \mathcal{C} and the embeddings.

The domains of both \mathcal{A} and \mathcal{B} are \mathbb{N} , and in each the interpretation of R_i is $\{\langle i, n \rangle \mid n \in \mathbb{N}\}$. To define $R_{i,s}$ first see whether $\Phi_i(i)$ is convergent in at most s steps. If not, then $R_{i,s}$ is empty. If so, say it converges at step $t \leq s$. In \mathcal{A} we put $\{i\} \times [0, t] \subseteq R_{i,t}$, split the rest of $i \times \mathbb{N}$ into two recursive infinite pieces, and put one into $R_{i,t}$ and the other into $\neg R_{i,t}$. In \mathcal{B} we put $\{i\} \times [0, t] \subseteq \neg R_{i,t}$ and then also split the rest of $i \times \mathbb{N}$ into two recursive infinite pieces, putting one into $R_{i,t}$ and the other into $\neg R_{i,t}$. Clearly \mathcal{A} and \mathcal{B} are recursive models of T and so provably exist in RCA_0 .

Suppose now that \mathcal{C} is also a model of T and there are embeddings $f : \mathcal{C} \rightarrow \mathcal{A}$ and $g : \mathcal{C} \rightarrow \mathcal{B}$. We claim that we can compute $0'$ from these models and embeddings, i.e. that it is Δ_1 in them and so provably exists in RCA_0 . Given i we find a $c \in \mathcal{C}$ such that $\mathcal{C} \models R_i(c)$. (One exists by our first axioms for T .) We then get $f(c)$. If $\Phi_i(i)$ converges in at most $f(c)$ many steps, then, of course, $i \in 0'$. If not, we get $g(c)$. Again if $\Phi_i(i)$ converges in at most $g(c)$ many steps, then, of course, $i \in 0'$. If not, then we claim that $\Phi_i(i)$ diverges and so $i \notin 0'$. Otherwise, by our assumptions, $\Phi_i(i)$ converges in exactly t many steps for some $t > f(c)$. Thus by our construction of \mathcal{A} , we have $\{i\} \times [0, t] \subseteq R_{i,t}^{\mathcal{A}}$ and so, in particular, $\mathcal{A} \models R_{i,t}(\langle i, f(c) \rangle)$. As f and g are embeddings, $\mathcal{B} \models R_{i,t}(\langle i, g(c) \rangle)$, and so by the construction of \mathcal{B} we have $g(c) > t$ for the desired contradiction.

Finally, we argue that each assertion (2)–(4) of the theorem provides a model \mathcal{C} of T and embeddings into both \mathcal{A} and \mathcal{B} . For (2), note that \mathcal{A} is atomic and so prime by (2). Thus there is an embedding of \mathcal{A} into \mathcal{B} , and \mathcal{A} itself fulfills the requirements on \mathcal{C} . Similarly, assuming (3) there is an isomorphism between \mathcal{A} and \mathcal{B} , and so either will fulfill the requirements on \mathcal{C} . As for (4), we have shown that T is atomic, and so (4) implies that it has a prime model \mathcal{C} which by definition is embeddable into both \mathcal{A} and \mathcal{B} , as required. \square

Of course, by Theorems 2.2 and 2.3(3), $\text{ACA}_0 \vdash \text{PUniqueness}$, but the reversal is open. We now turn to the remaining implication AMT and its relation to other principles.

3. AMT: NONIMPLICATIONS

The reverse mathematically most interesting theorem of our group on prime and atomic models is (the existence part of) AMT, that every atomic theory has an atomic model. Along with those studied in [21], it is another example of a mathematical existence theorem weaker than ACA_0 and incomparable with WKL_0 . We begin our analysis by using some of the recursion theoretic results of Csima, Hirschfeldt, Knight and Soare [9], Csima [8], Hirschfeldt and Shore [21] and others to draw reverse mathematical conclusions of the form that AMT does not imply various other principles.

Theorem 3.1 (Csima [8]). *Every atomic decidable T (i.e. the set of sentences in T is recursive) has an atomic prime model \mathcal{A} with a low complete diagram, i.e. if D is the set of sentences in the language of T with constants added for each element of \mathcal{A} true in \mathcal{A} , then $D' \equiv_T \mathbf{0}'$.*

Definition 3.2 (Csima, Hirschfeldt, Knight and Soare [9]). A degree \mathbf{a} is *prime bounding* if every decidable atomic T has an atomic model recursive in \mathbf{a} , i.e. its atomic diagram is recursive in \mathbf{a} .

Theorem 3.3 (Csima, Hirschfeldt, Knight and Soare [9]). *A degree $\mathbf{a} \leq \mathbf{0}'$ is prime bounding iff $\mathbf{a} \in \bar{\mathbf{L}}_2$, i.e. $\mathbf{a}'' > \mathbf{0}''$.*

Corollary 3.4. $\text{WKL}_0 \not\leq \text{AMT}$.

Proof. Essentially by the low basis theorem (Jockusch and Soare [24]) as shown in Simpson [32, proof of Theorem VIII.2.17], there is a low degree \mathbf{d} , i.e. $\mathbf{d}' = \mathbf{0}'$, bounding a standard model \mathcal{M} of WKL_0 , i.e. every set in \mathcal{M} is recursive in \mathbf{d} . On the other hand, Theorem 3.3 says that no low degree can bound a standard model of AMT. \square

We now want to combine and slightly improve the two previous theorems to get the reverse nonimplication. As we will need the notion of the tree of Henkin constructions of models of a theory for this and later proofs, we first make it precise with the following definition.

Definition 3.5 (RCA_0). Let T be a (complete, consistent, deductively closed) theory in a language \mathcal{L} . Let \mathcal{L}' be the extension of \mathcal{L} obtained by adding on countably many new (Henkin) constants c_i , and let φ_i be an enumeration of all the sentences of \mathcal{L}' . The *tree \mathcal{F} of all possible standard Henkin constructions of models of T* is defined by recursion. Each node $\sigma \in 2^{\mathbb{N}}$ in \mathcal{F} will also be labeled by a set S_σ of sentences of \mathcal{L}' consistent with T associated with it. We begin by putting \emptyset into \mathcal{F} and setting $S_\emptyset = \emptyset$. If $\sigma \in \mathcal{F}$ and $|\sigma| = n$, we ask whether $S_\sigma \cup \{\varphi_n\}$ is consistent with T . If so, we put $\sigma \hat{\ } 1$ into \mathcal{F} . If φ_n is of the form $\exists x\psi(x)$ we choose the least i such that c_i does not appear in S_σ and let $S_{\sigma \hat{\ } 1} = S_\sigma \cup \{\varphi_n, \psi(c_i)\}$; otherwise $S_{\sigma \hat{\ } 1} = S_\sigma \cup \{\varphi_n\}$. If $\neg\varphi_n$ is consistent with T , we put $\sigma \hat{\ } 0$ into \mathcal{F} and let $S_{\sigma \hat{\ } 0} = S_\sigma \cup \{\neg\varphi_n\}$.

Remark 3.6. Even in RCA_0 , the usual arguments show that at least one of φ and $\neg\varphi$ must be consistent with $T \cup S_\sigma$, and if $\sigma \hat{\ } i \in \mathcal{F}$, then $S_{\sigma \hat{\ } i}$ is consistent with T . If P is any (infinite) path through \mathcal{F} , then $S_P = T \cup \{S_\sigma \mid \sigma \in P\}$ is consistent and recursively defines a model \mathcal{M}_P of T (in RCA_0) by the usual Henkin argument. (The elements are the c_i for i least such that there is some c_j with $c_i = c_j \in S_P$.)

The functions and relations are defined in the standard fashion.) The recursion theoretic version of this fact is that the tree \mathcal{F} is recursive in T and each \mathcal{M}_P is recursive in $T \oplus P$. This consideration gives rise to a tree \mathcal{S} (with the same nodes as \mathcal{F}) labeled with D_σ , subsets of the complete diagrams of Henkin models of T . Of course, \mathcal{F} and \mathcal{S} trivially have infinitely many splits along every path (determined by various constants being equal or not if by nothing else). Thus if T is decidable (recursive), \mathcal{F} and \mathcal{S} are recursively isomorphic to the full binary tree $2^{<\mathbb{N}}$.

Proposition 3.7. *If T is atomic and decidable and $\mathbf{a} \notin \mathbf{GL}_2$, i.e. $(\mathbf{a} \vee \mathbf{0}')' < \mathbf{a}''$, then T has an atomic model \mathcal{B} with complete diagram of degree $\mathbf{b} \leq \mathbf{a}$ with $\mathbf{b} \in \mathbf{GL}_1$, i.e. $\mathbf{b} \vee \mathbf{0}' = \mathbf{b}'$.*

Proof. As T is decidable, the tree \mathcal{F} of all possible standard Henkin constructions of models of T and the associated tree \mathcal{S} of Henkin models of T are recursive and recursively isomorphic to $2^{<\mathbb{N}}$. As always, the sets of nodes that decide each membership question of the jump of the model (given that it lies on \mathcal{S}) are dense. Any path through \mathcal{S} that meets each of these dense sets is in \mathbf{GL}_1 . (To decide whether e is in the jump just find a node σ such that either $\Phi_e^{D_\sigma}(e)$ is convergent or σ has no extension τ (on \mathcal{F}) with $\Phi_e^{D_\tau}(e)$ convergent.) As T is atomic, the set of nodes that make a given n -tuple c_1, \dots, c_n of Henkin constants satisfy an atom is also dense. (At any node there is some formula $\psi(x_1, \dots, x_n)$ that contains all the facts determined so far about c_1, \dots, c_n . By atomicity of T , that formula $\psi(x_1, \dots, x_n)$ can be extended to an atom $\varphi(x_1, \dots, x_n)$. One then simply finds an extension of the given node on \mathcal{S} whose label includes the formula $\varphi(c_1, \dots, c_n)$.) Thus all we need to do is to produce a path through \mathcal{S} that is recursive in \mathbf{a} and meets all these dense sets of conditions. The corresponding Henkin model of T is then atomic, and its complete theory is recursive in \mathbf{a} and in \mathbf{GL}_1 as required.

The crucial point is that the function going from one of these dense sets and a node σ to an extension τ in the dense set is uniformly recursive in $0'$. (For deciding the jump this is standard. For atomicity, just note that being an atom is a Π_1 property of a formula.) We combine these operations into a single function f recursive in $0'$ for which we can use the hypothesis that $\mathbf{a} \notin \mathbf{GL}_2$. We define a function f recursive in $0'$ by recursion. Given $f(n)$, we let $f(n+1)$ be the least $m > f(n)$ such that

- (1) every σ of length at most $f(n)$ that has any extension τ on \mathcal{S} such that $\Phi_e^{D_\tau}(e) \downarrow$ for any $e \leq n$ has one at level m and
- (2) for every tuple \bar{a} of Henkin constants with code below $f(n)$ and every σ on \mathcal{S} of length at most $f(n)$,
 - (a) σ has an extension τ on \mathcal{F} of length less than m whose label has a formula isolating the type of \bar{a} and
 - (b) letting τ be such an extension of shortest length, for every $\rho \supseteq \sigma$ such that $|\rho| \leq |\tau|$, if ρ does not have a formula isolating the type of \bar{a} , then ρ has a split in the type of \bar{a} (as given by D_ρ) appearing on (the labels of) \mathcal{S} below level m .

Now as $\mathbf{a} \notin \mathbf{GL}_2$, there is a g recursive in \mathbf{a} that is strictly increasing and not dominated by f (see Lerman, [26, Corollary IV.3.4]).

We now construct a path through \mathcal{S} recursive in g that meets all the specified dense sets. Inductively assume that at stage n we have σ on \mathcal{S} of length at most

$\min\{f(n), g(n)\}$. We find the least e or \bar{a} below n (in a priority ordering where we order e before the tuple \bar{a} with code e) such that

- (1) $\Phi_e^{D\sigma}(e)$ diverges, but $\Phi_e^{D\tau}(e)$ converges for some $\tau \supseteq \sigma$ of length at most $g(n+1)$ or
- (2) there is a split in the type of \bar{a} above σ below level $g(n+1)$ in \mathcal{S} , but for some $\tau \supseteq \sigma$ of length less than $g(n+1)$ there is no split of the type of \bar{a} in \mathcal{S} above τ below level $g(n+1)$.

Choose a τ of minimal length satisfying the required condition for this least requirement. Let the $(n+1)^{\text{st}}$ step in constructing our desired path be τ . Note that we maintain our inductive hypothesis in that $|\tau| \leq f(n+1), g(n+1)$. The point here is that τ is by construction of length less than $g(n+1)$, and if $f(n+1) \leq g(n+1)$, then it is of length less than $f(n+1)$ by the definition of f and our inductive hypothesis that $|\sigma| < f(n)$.

Now we argue that every requirement is satisfied. Say deciding $\Phi_e^M(e)$ is the least one not satisfied and all higher priority ones have been satisfied by node ρ constructed at stage k . At some $n > k$ we are at a node σ of length at most $\min\{f(n), g(n)\}$ and $g(n+1) > f(n+1)$. If there is any extension τ of σ such that $\Phi_e^{D\tau}(e) \downarrow$, then there is one of length at most $f(n+1) < g(n+1)$, and so we find it at stage n and take the first one as the next element of our path and satisfy this requirement. Otherwise, there is no such extension and we also decide the jump at e , as required.

If \bar{a} is the least tuple with a nonatomic type in the model we built, again choose an $n > \bar{a}$ such that $g(n+1) > f(n+1)$ and all higher priority requirements have been satisfied by the node σ of length at most $\min\{f(n), g(n)\}$ produced at stage n (i.e. all $\Phi_e^{D\sigma}(e)$ have been decided and all $\bar{b} < \bar{a}$ have had their types isolated by σ). At stage $n+1$ our computation sees that σ does not isolate the type of \bar{a} (by our assumption that it is not isolated in our model and the definition of f). If we search for a $\tau \supseteq \sigma$ that has no extensions splitting the type of \bar{a} in the labels below level $g(n+1)$, the one of shortest length that we find actually isolates the type of \bar{a} by the definition of f . We choose such one as the next node on our tree. \square

Corollary 3.8. *Every degree $\mathbf{a} \notin \mathbf{GL}_2$ is prime bounding and, in fact, bounds a standard model of AMT all of whose sets are recursive in sets below \mathbf{a} which are also in \mathbf{GL}_1 .*

Proof. Given any $\mathbf{a} \notin \mathbf{GL}_2$ we can build a standard model of AMT bounded by \mathbf{a} by iterating Proposition 3.7 to dovetail and produce a class of sets closed under AMT (i.e. for any atomic theory in the class there is an atomic model in the class), with each member of the class recursive in \mathbf{a} and indeed recursive in some \mathbf{GL}_1 degree below \mathbf{a} . To be more precise, we construct three sequences T_i, \mathcal{A}_i and \mathbf{b}_i such that for each i , we have $T_i \leq_T \mathbf{b}_i \leq \mathbf{b}_{i+1} \leq \mathbf{a}$ and $\mathbf{b}_i \in \mathbf{GL}_1$, so that every complete atomic theory recursive in any \mathbf{b}_j occurs as some T_k and \mathcal{A}_k is an atomic model of T_k whose complete diagram is recursive in \mathbf{b}_{k+1} . Say we have $T_i \leq_T \mathbf{b}_i \in \mathbf{GL}_1$ and $\mathbf{b}_i \leq \mathbf{a}$. Since $\mathbf{b}_i \in \mathbf{GL}_1$, we have $(\mathbf{a} \vee \mathbf{b}'_i)' = (\mathbf{a} \vee (\mathbf{b}_i \vee \mathbf{0}'))'$. As $\mathbf{b}_i \leq \mathbf{a}$, we have $(\mathbf{a} \vee (\mathbf{b}_i \vee \mathbf{0}'))' = (\mathbf{a} \vee \mathbf{0}')'$. As $\mathbf{a} \notin \mathbf{GL}_2$, we have $(\mathbf{a} \vee \mathbf{0}')' < \mathbf{a}''$. Thus $(\mathbf{a} \vee \mathbf{b}'_i)' < \mathbf{a}''$, so $\mathbf{a} \notin \mathbf{GL}_2^{\mathbf{b}_i}$. By Proposition 3.7 (relativized to \mathbf{b}_i) we can get an atomic model \mathcal{A}_i of T_i with its complete diagram recursive in \mathbf{a} and $\mathbf{GL}_1^{\mathbf{b}_i}$. We let \mathbf{b}_{i+1} be the degree of \mathcal{A}_i and note that it is in \mathbf{GL}_1 as $\mathbf{b}'_{i+1} = (\mathbf{b}_{i+1} \vee \mathbf{b}'_i) = (\mathbf{b}_{i+1} \vee \mathbf{b}_i \vee \mathbf{0}') = \mathbf{b}_{i+1} \vee \mathbf{0}'$, as required. \square

Corollary 3.9. *AMT does not imply WKL_0 .*

Proof. No incomplete r.e. degree can bound a model of WKL_0 by Jockusch and Soare [24]. On the other hand, by Corollary 3.8 every r.e. degree not in \mathbf{L}_2 bounds one. \square

We now turn to the relations between AMT and the combinatorial principles ranging in strength from RT_2^2 to ADS and SADS studied in [2] and [21] and defined in §1. We begin with the recursion theoretic arguments for nonimplications.

Corollary 3.10. *AMT does not imply RT_2^2 , SRT_2^2 , CAC, COH or CADS.*

Proof. RT_2^2 and CAC have recursive instances with no Δ_2 solution by Jockusch [23] and Herrmann [19]. COH has recursive instances whose only Δ_2 solutions are high (i.e. with jump $0''$) by Cooper [7] and Jockusch and Stephan [25], and so does CADS, since it has the same ω -models as COH (Hirschfeldt and Shore [21]). SRT_2^2 has recursive instances with no low (i.e. with jump $0'$) solution (Downey, Hirschfeldt, Lempp and Solomon [10]). As above, however, every r.e. degree not in \mathbf{L}_2 bounds a model of AMT in which every degree is low. \square

A more subtle degree theoretic argument shows that AMT does not imply SADS.

Theorem 3.11 (Hirschfeldt and Shore [21]). *For any r.e. A, B there is a recursive order of type $\omega + \omega^*$ such that, if C is a suborder of type ω or ω^* , then either $B \leq_T A \oplus C$ or $A \leq_T B \oplus C$.*

Corollary 3.12. *AMT $\not\leq$ SADS.*

Proof. By standard results there are high r.e. \mathbf{h} , \mathbf{a} and \mathbf{b} such that $\mathbf{a} \not\leq \mathbf{b} \vee \mathbf{h}$ and $\mathbf{b} \not\leq \mathbf{a} \vee \mathbf{h}$. (For example, by Robinson [31] as in Soare [34, VIII.4.8], every finite usl can be embedded in every interval of r.e. degrees, and we can take such an embedding with the bottom of the interval high.) By Corollary 3.8, there is a model of AMT with all sets recursive in \mathbf{h} . \square

AMT also does not imply any of the other principles studied in [21] that are below ACA_0 and incomparable with WKL_0 . In fact, we have a very strong conservation result similar to that for COH in [21] that implies almost all of these independence results over AMT.

The basic situation here is that we start with a countable model \mathcal{M} of RCA_0 and a notion of forcing \mathcal{P} defined over \mathcal{M} . Moreover, if G is sufficiently generic over \mathcal{P} (usually 1-generic plus satisfying a specific list of requirements that guarantee that if we force an instance of a Σ_1^0 (or Π_1^0) formula, then there is a least instance that can be forced, and we do so), we can show that $\mathbf{I}\Sigma_1$ holds for formulas over \mathcal{M} with G as an added set parameter. (This typically relies on a sufficiently simple definition of forcing for one quantifier sentences.) In this situation, \mathcal{M} is an ω -submodel of the model $\mathcal{M}[G]$ obtained by adding onto the sets of \mathcal{M} all sets definable over \mathcal{M} by Δ_1^0 formulas with G as an added set parameter, and $\mathcal{M}[G]$ is itself a model of RCA_0 (Friedman [12], or see Simpson [32] or [2, §6]), where a submodel \mathcal{A} of \mathcal{B} is an ω -submodel if \mathcal{A} has the same first-order part as \mathcal{B} . (Warning: in Cholak, Jockusch and Slaman [2], $\mathcal{M}[G]$ is first officially defined as the model of $\mathbf{I}\Sigma_1$ obtained by adding on just G to the sets of \mathcal{M} . What we generally want, however, is its extension to a model of RCA_0 .)

Here we are interested in AMT, and the relevant notion of forcing is standard Cohen forcing on the appropriate tree. Note that if G is Cohen 1-generic (i.e. every Π_1^0 formula or its negation is forced and the conditions specific to forcing $\mathbb{I}\Sigma_1$ mentioned above are met), then adding G to \mathcal{M} preserves $\mathbb{I}\Sigma_1$, and so $\mathcal{M}[G] \models \text{RCA}_0$.

Theorem 3.13. *Let \mathcal{M} be a countable model of RCA_0 and let $\Phi(A, B)$ be a Σ_3^0 predicate of two set variables such that for some fixed $A \in \mathcal{M}$ there is no $B \in \mathcal{M}$ with $\mathcal{M} \models \Phi(A, B)$. If G is Cohen 2-generic over \mathcal{M} , then there is no $B \in \mathcal{M}[G]$ with $\mathcal{M}[G] \models \Phi(A, B)$.*

Proof. Suppose that $\Phi(A, B) = \exists x \forall y \exists z \varphi(A, B, x, y, z)$ where φ is Δ_0^0 and, for the sake of a contradiction, that there is a $B \in \mathcal{M}[G]$ such that $\mathcal{M}[G] \models \Phi(A, B)$. Now \mathcal{M} is an ω -submodel of $\mathcal{M}[G]$, and B is of the form $\Phi_e^{G \oplus C}$ for some Turing functional Φ_e and set C in \mathcal{M} , so for some $x \in \mathcal{M}$,

$$\mathcal{M}[G] \models \Phi_e^{G \oplus C} \text{ is a total characteristic function and } \forall y \exists z \varphi(A, \Phi_e^{G \oplus C}, x, y, z).$$

As this whole formula is equivalent (even in RCA_0) to a $\Pi_2^0(G)$ (over \mathcal{M}) formula $\forall y \exists z \Psi(G, C, A, x, y, z)$ and G is 2-generic, there is a condition $p \in \mathcal{M}$ such that $p \Vdash \forall y \exists z \Psi(G, C, A, x, y, z)$.

By the general definition of forcing this means that for every y in \mathcal{M} and every condition $p' \leq p$, there is a $z \in \mathcal{M}$ and a condition $p'' \leq p'$ such that $p'' \Vdash \Psi(G, C, A, x, y, z)$. As forcing a Δ_0^0 formula is itself a Δ_0 relation, we can now define by recursion the function f such that $f(0) = p$ and $f(n+1) \Vdash \exists z \Psi(G, C, A, x, n, z)$. As \mathcal{M} is a model of RCA_0 , this function is an element of \mathcal{M} and so gives the characteristic function of a set $H \in \mathcal{M}$, which by construction satisfies $\forall y \exists z \Psi(H, C, A, x, y, z)$. Finally, if $D = \Phi_e^{H \oplus C}$, then $D \in \mathcal{M}$ and $\mathcal{M} \models \forall y \exists z \varphi(A, D, x, y, z)$ for the desired contradiction. \square

We now want to observe that adding a Cohen 2-generic adds an atomic model of each atomic theory $T \in \mathcal{M}$. (In fact, even less than meeting all uniformly Π_1^0 sequences of dense sets suffices for AMT, as we will see in §4.) Iterating the forcing will then provide our conservation result.

Proposition 3.14. *Let \mathcal{M} be a countable model of RCA_0 and let T be an atomic theory in \mathcal{M} . If G is Cohen 2-generic over \mathcal{M} , then there is an $A \in \mathcal{M}[G]$ that is an atomic model of T in $\mathcal{M}[G]$.*

Proof. Consider again the tree \mathcal{S} in \mathcal{M} of all standard Henkin constructions of models of T . For each initial segment of Henkin constants \bar{c} the set of nodes σ of \mathcal{S} that include an atom φ about \bar{c} is dense by the assumption that T is atomic. Moreover, this set is Π_1 , as that is the complexity of the statement that φ is an atom of T . Thus the path through \mathcal{S} determined by G meets every such set and so produces an atomic model of T , as required. \square

We now have our conservation and nonimplication results. In the terminology of Simpson [32, Definition VII.2.28], a model \mathcal{M} is a *restricted β -submodel* of a model \mathcal{M}' if \mathcal{M} is an ω -submodel of \mathcal{M}' and for every sentence of the form $\exists X \psi$ where ψ is Π_2^0 with parameters in \mathcal{M} , we have $\mathcal{M} \models \exists X \psi$ if and only if $\mathcal{M}' \models \exists X \psi$. (This condition says that a subtree of $\mathbb{N}^{<\mathbb{N}}$ in \mathcal{M} has a path in \mathcal{M} if and only if it has a path in \mathcal{M}' , whence the terminology.)

Corollary 3.15. *AMT is restricted $(r)\Pi_2^1$ conservative over RCA_0 , i.e. conservative over RCA_0 for sentences of the form $\forall A(\Theta(A) \rightarrow \exists B\Phi(A, B))$, where Θ is arithmetic and Φ is Σ_3^0 . Furthermore, every model of RCA_0 is a restricted β -submodel of a model of AMT.*

Proof. Consider any sentence of the specified form and any model \mathcal{M} of RCA_0 not satisfying the sentence. Then there is a set A of \mathcal{M} such that $\mathcal{M} \models \Theta(A)$ for which there is no B in \mathcal{M} such that $\mathcal{M} \models \Phi(A, B)$. Construct a sequence $\langle G_i \mid i \in \omega \rangle$ of subsets of $\mathbb{N}^{\mathcal{M}}$ such that G_{i+1} is Cohen 2-generic over $\mathcal{M}[G_1] \dots [G_i]$. Let $\mathcal{M}' = \bigcup \{ \mathcal{M}[G_0] \dots [G_i] \mid i \in \omega \}$. By the results mentioned above, $\mathcal{M}[G_0] \dots [G_i] \models RCA_0$ for each i , and, by Proposition 3.14, every atomic theory T in $\mathcal{M}[G_0] \dots [G_i]$ has an atomic model in $\mathcal{M}[G_0] \dots [G_i][G_{i+1}]$. Thus $\mathcal{M}' \models RCA_0 + AMT$ as the first order part of every element of this ascending sequence of models is the same. By induction (on true ω) our theorem shows that there is no $B \in \mathcal{M}[G_0] \dots [G_i]$ such that $\mathcal{M}[G_0] \dots [G_i] \models \Phi(A, B)$. Again, as each successive model in this list is an ω -extension of the preceding one and \mathcal{M}' is just their union, we see that while $\mathcal{M}' \models \Theta(A)$, there is no $B \in \mathcal{M}'$ such that $\mathcal{M}' \models \Phi(A, B)$. Thus our original sentence is not a theorem of $RCA_0 + AMT$, as required.

The version of the corollary in terms of restricted β -extensions follows by taking Θ to be empty and Φ to be Π_2^0 . Our argument shows that if $\mathcal{M}' \models \exists B\Phi$, then $\mathcal{M} \models \exists B\Phi$. The other direction follows from the fact that \mathcal{M} is an ω -submodel of \mathcal{M}' . □

We note that this result also provides the usual Π_1^1 conservativity result, since any Π_1^1 sentence $\forall A\Psi(A)$ is equivalent to $\forall A(\neg\Psi(A) \rightarrow \exists B(0 \neq 0))$. Moreover, the result is the best possible one of this form, as AMT itself is a sentence of the form $\forall A(\Theta(A) \rightarrow \exists B\Phi(A, B))$ with Θ arithmetic and Φ a Π_3^0 formula. It is, however, strong enough to show that any principle that asserts, for example, the existence of an infinite set satisfying some recursive condition such as being homogeneous, a path through a tree, or a chain or antichain in a linear or partial order cannot be implied by AMT, as all of these statements are ones in which Φ is Π_2^0 .

Corollary 3.16. *None of the following principles are implied by AMT: RT_2^2 , SRT_2^2 , WKL_0 , DNR , CAC , ADS , $SADS$.*

By the argument of Hirschfeldt and Shore [21] using Mathias forcing we can also add COH so that $AMT + COH$ has the same properties specified in Corollaries 3.15 and 3.16 for AMT alone.

Note that by [21] CAC , ADS and $SADS$ are also incomparable with WKL_0 .

4. AMT, SADS AND Σ_2

We now place AMT with respect to the combinatorial principles of [21] by showing that it is implied by the weakest of them that is not conservative over RCA_0 : SADS. (Note that by Corollary 3.16 this implication is strict.) We also place it more precisely in the purely proof theoretic hierarchy by showing that it is Π_1^1 -conservative over $B\Sigma_2$. The proof of the implication uses a finite injury priority argument that must be carried out in RCA_0 (and so only with $I\Sigma_1$) by using Shore blocking. The situation is more subtle than might first appear, as similar constructions provide priority arguments for slightly stronger conclusions that, provably, cannot be carried out in RCA_0 . The conservation proof also uses a priority argument. We will discuss these issues after proving our implication from SADS.

Theorem 4.1. $RCA_0 \vdash SADS \rightarrow AMT$.

Proof. We work in RCA_0 and begin with an atomic theory T . We again have the tree \mathcal{S} of standard Henkin constructions of models of T (with Henkin constants c_i). We want to recursively define an ordering $<_L$ on \mathbb{N} of type $\omega + \omega^*$ such that recursively in any subsequence R of type ω or ω^* we can construct an infinite path P through \mathcal{S} that corresponds to an atomic model \mathcal{A} of T . Looking ahead to the construction of the path and model, we define a monotonic recursive operation Φ taking finite sequences of numbers to finite paths in \mathcal{S} .

We define $\Phi(x_1, \dots, x_n) = \langle \sigma_1, \dots, \sigma_n \rangle$ to be an increasing sequence of nodes in \mathcal{S} by recursion on n . We let σ_1 be the least node σ on \mathcal{S} (in some standard ordering of nodes of type ω) that mentions c_0 and such that in a (standardized) search of x_1 many steps we have not found a witness to the assertion that σ is not an atom about c_0 . (Remember that the nodes on \mathcal{S} correspond to the conjunction of sets of formulas about the Henkin constants. This conjunction, viewed as a sentence about any particular set of constants (by existentially quantifying out the rest), may or may not be an atom of T . To see that it is not an atom, it suffices to find two incompatible extensions both of which are consistent with T . As T is complete, this is a recursive test of the candidate extensions.) We now proceed by induction. Given σ_i we let σ_{i+1} be the least σ (in the standard ordering) extending σ_i on \mathcal{S} that mentions all of c_0, \dots, c_{x_i} such that in x_{i+1} many steps we have not found a witness to the assertion that σ is not an atom about c_0, \dots, c_{x_i} .

Our goal in defining our order L is to guarantee that any ω or ω^* subsequence will infinitely often have successive elements far enough apart so that applying Φ will, for each i , produce at some point a correct witness that some σ_n is an atom for c_0, \dots, c_i . If so, Φ applied to any such sequence will produce our desired path through \mathcal{S} generating an atomic model of T .

The construction of L will involve a movable marker procedure with two sets of markers $d_{i,s}$ and $u_{i,s}$ that remain in two disjoint parts D_s and U_s of L with every member of D_s being L -less than every member of U_s . The intuition is that the limit D of D_s is the ω part of L (the down side) and the limit U of U_s is the ω^* part (the up side). The $d_{i,s}$ and $u_{i,s}$ will also stabilize to numbers d_i and u_i that will be in D and U and ordered by $<_L$ in the end in ω and ω^* orderings, respectively. We have requirements \mathcal{D}_i and \mathcal{U}_i with a standard priority ordering giving \mathcal{D}_i priority over \mathcal{D}_j for $j > i$ and over \mathcal{U}_j for $j \geq i$, and \mathcal{U}_i priority over \mathcal{U}_j for $j > i$.

We say that the requirement \mathcal{D}_i needs attention at stage s if there exists a sequence $x_1, \dots, x_n, d_{i,s}$ of elements of D_s increasing in both L and natural order such that $\Phi(x_1, \dots, x_n, d_{i,s}) = \langle \sigma_1, \dots, \sigma_{n+1} \rangle$, and by stage s we have seen a witness that σ_{n+1} does not supply an atom about c_0, \dots, c_{x_n} . Similarly, we say that \mathcal{U}_i needs attention at stage s if there is a sequence $x_1, \dots, x_n, u_{i,s}$ of elements of U_s decreasing in L and increasing in the natural order such that $\Phi(x_1, \dots, x_n, u_{i,s}) = \langle \sigma_1, \dots, \sigma_{n+1} \rangle$, and by stage s we have seen a witness that σ_{n+1} does not supply an atom about c_0, \dots, c_{x_n} .

At stage s let r be the smallest number not yet appearing in $L_s = D_s \cup U_s$. Suppose the highest priority requirement needing attention at s is \mathcal{D}_i . We move all numbers in D_s that are L greater than or equal to $d_{i,s}$ to be the L smallest members of U_{s+1} (preserving the L order already defined on them). We let $d_{i,s+1} = r$ and insert it as the L largest element of D_{s+1} . All $d_{j,s+1}$ become undefined for $j > i$, as do all $u_{j,s+1}$ for $j \geq i$. Similarly, if the highest priority requirement needing

attention at s is \mathcal{U}_i , we move all numbers in U_s that are L smaller than or equal to $u_{i,s}$ to be the L largest members of D_{s+1} (preserving the L order already defined on them). We let $u_{i,s+1} = r$ and insert it as the L smallest element of U_{s+1} . All $u_{j,s+1}$ become undefined for $j > i$, as do all $d_{j,s+1}$ for $j > i$. If no requirement needs attention, we add r as the L largest member of D_{s+1} , let it be $d_{j,s+1}$ for the least j such that $d_{j,s}$ is undefined, and add $r + 1$ as the L smallest member of U_{s+1} and let it be $u_{j,s+1}$ for the least j such that $u_{j,s}$ is undefined. We then go on to the next stage.

We first claim that every n is eventually in L_s and from some stage onward is always in D_s or always in U_s . As for entering L_s , as we always put in the least number not yet in L_s at every stage, $n \in L_{n+1}$. Once n is in D_s or U_s , it can change to the other at $t > s$ only when we act for some requirement. By $\text{I}\Sigma_1$ there is a requirement of highest priority for which we ever act that moves n . Say it is \mathcal{D}_i and it acts to move n at stage t . This means that $d_{i,t} \leq_L n$, we move n to U_{t+1} and all markers of lower priority become undefined. No action for U_j of higher priority ever moves n back to D_s by our choice of \mathcal{D}_i . Any marker of lower priority is appointed after t and assigned to a number that, by construction, is L -less than n . Moving such a lower priority marker for its requirement, by construction, does not move any number L -above it. Thus n never leaves U_s after t . The argument if the highest priority requirement moving n is \mathcal{U}_i is symmetric.

Once n has stopped moving at t , all numbers entering L after t are L -above n (if it is in D_t) or all are L -below n (if it is in U_t). Thus n has either finitely many L predecessors or finitely many L successors and is therefore of type $\omega + \omega^*$, as desired. We now apply SADS to get an infinite ascending or descending sequence $R = \langle x_i \rangle$ in L . Without loss of generality, assume it is ascending in L and in natural order. Let $P = \Phi(\langle x_i \rangle)$ be the path in \mathcal{S} determined by R . We claim that it includes an atom for each initial segment c_0, \dots, c_n of Henkin constants and so provides the required atomic model of T . We need an important fact about the construction.

Lemma 4.2. *If we act for a requirement \mathcal{D}_i at s and never act for any requirement of higher priority again, then we eventually stop acting for \mathcal{D}_i and so $d_{i,t}$ eventually remains fixed. Moreover, if $d_{i,s}$ never changes after t , then, for any sequence $x_1, \dots, x_n, d_{i,t}$ that is increasing in both L and natural order, σ_{n+1} includes an atom about c_0, \dots, c_{x_n} , where $\Phi(x_1, \dots, x_n, d_{i,t}) = \langle \sigma_1, \dots, \sigma_{n+1} \rangle$.*

Similarly, if we act for a requirement \mathcal{U}_i at s and never act for any requirement of higher priority again, then we eventually stop acting for \mathcal{U}_i and so $u_{i,t}$ eventually remains fixed. Moreover, if $u_{i,s}$ never changes after t , then, for any sequence $x_1, \dots, x_n, u_{i,t}$ that is decreasing in L and increasing in natural order, σ_{n+1} includes an atom about c_0, \dots, c_{x_n} , where $\Phi(x_1, \dots, x_n, u_{i,t}) = \langle \sigma_1, \dots, \sigma_{n+1} \rangle$.

Proof. We prove the first part of the lemma. The proof of the second part is symmetric.

Suppose we act for \mathcal{D}_i at s . We then appoint some r as $d_{i,s+1}$. We now act for \mathcal{D}_i again only if we discover a new witness as described in the definition of needing attention. Since we never act for any requirement of higher priority the numbers L -below $d_{i,s+1}$ are fixed in D_t forever. We then act again for \mathcal{D}_i at $t > s$ only when we discover some new witness that some σ is not an atom, where σ is the final value of $\Phi(x_1, \dots, x_n, d_{i,t})$ for some increasing (in both L and natural order) sequence x_1, \dots, x_n of numbers L -below $d_{i,t}$. Each such sequence x_1, \dots, x_n produces a fixed $\sigma(x_1, \dots, x_n)$ as the next-to-last value under Φ . The only variations arise from

trying to find the first node $\tau(x_1, \dots, x_n)$ extending $\sigma(x_1, \dots, x_n)$ that includes an atom about c_1, \dots, c_{x_n} . The set of such $\sigma(x_1, \dots, x_n)$ is finite and each has such an atom extending it by the assumption that T is atomic. We can avoid the use of $B\Sigma_2$ that might naturally be invoked in order to find a common bound on all such extensions by a trick peculiar to our setting. We simply relabel each $\sigma(x_1, \dots, x_n)$ so as to replace the constants c_1, \dots, c_{x_n} by new free variables that are distinct from the ones used for all the other sequences. The other constants appearing are existentially quantified out as usual. This gives a formula with x_n many free variables for each sequence x_1, \dots, x_n , with all the sets of variables distinct for different sequences. We can now consider the conjunction of all these formulas as a single formula of many variables. (Note that this conjunction is consistent with T . Each conjunct is consistent with T and so the existential closure of each conjunct is a theorem of T . As all the variables are distinct, the conjunction of these existential closures is the existential closure of the conjunct which is therefore itself consistent with T .) As T is atomic this formula can be extended to an atom Ψ . If in Ψ we existentially quantify out the variables not free in the relabeled $\sigma(x_1, \dots, x_n)$, we have an atom extending $\sigma(x_1, \dots, x_n)$. Thus there is a single bound on atoms extending all the relevant $\sigma(x_1, \dots, x_n)$, and so by $I\Sigma_1$ there is a first one for each and a bound on the witnesses needed to show that all smaller candidates are not such atoms. Once we are above such a stage, no change occurs in $d_{i,t}$ and its value must also be above the stage at which the correct extensions are found for each $\sigma(x_1, \dots, x_n)$. After such a stage \mathcal{D}_i will never again need attention. In particular, this stage is large enough so that the search incorporated in Φ produces a σ_{n+1} including an atom about c_1, \dots, c_{x_n} . \square

Returning to our verification that P determines an atomic model of T , we choose any $m = x_{j+1} \in R$ so that x_j is larger than n and let t be the last stage at which m is put into D_t . If it is moved into D_t (say by the action of some \mathcal{U}_i), then it is part of a block of numbers moved, all of which are L -below $u_{i,t}$. Otherwise, it is just put into D_t on its own as some $d_{i,t}$ at t and never moves. In the second case, by the lemma applied to \mathcal{D}_i , if $\Phi(x_1, \dots, x_j, m) = \langle \sigma_1, \dots, \sigma_{j+1} \rangle$, then σ_{j+1} includes an atom about c_0, \dots, c_{x_j} and so one about c_0, \dots, c_n , as required. In the first case, no \mathcal{D}_k of higher priority than \mathcal{U}_i can act after t , as that would move m . The highest priority requirement that acts at a $t' \geq t$ must be a \mathcal{U}_j for $j \leq i$. Let \mathcal{U}_j be that requirement and (by the lemma) let v be the last stage at which it acts. At v , the marker $u_{j,v}$ and a block of numbers are moved from U_v to U_{v+1} . They include m (if $v = t$) and otherwise are all L -above m . At $v + 1$ no requirement can need attention, as all of lower priority than \mathcal{U}_j have had their markers undefined at v and no higher priority ones can act by our assumptions. Thus we insert some number r as the L -greatest element of D_{v+1} and let it be $d_{j+1,v+1}$. By the lemma there is a stage $w \geq v + 1$ at which we last act for \mathcal{D}_{j+1} . At w we appoint some r' as $d_{j+1,w}$, and it too is L -above m . Let m' be the last element of R in $[m, r')$ and m'' be the first element of R greater than or equal to r' . By the lemma, if $\Phi(x_1, \dots, x_j, m', r') = \langle \sigma_1, \dots, \sigma_{j+2} \rangle$, then σ_{j+2} includes an atom about $c_1, \dots, c_{m'}$. As $m'' \geq r'$, the search to produce the last element of $\Phi(x_1, \dots, x_j, m', m'')$ begins with the same σ_{j+1} and runs for at least as many steps as the one that produced the correct answer in r' many steps. Thus it too produces the desired atoms, and as x_1, \dots, x_j, m', m'' is an initial segment of R , the sequence $\Phi(R)$ contains an atom about $c_1, \dots, c_{m'}$ and so, of course, about c_1, \dots, c_n , as required. \square

The above argument is a finite injury argument that we carried out in RCA_0 and so only using $\text{I}\Sigma_1$. The amount of induction needed for various types of priority arguments has been extensively studied as reverse recursion theory but primarily in the setting of first order arithmetic (where one talks about r.e. or Δ_2 sets in terms of their indices). Good basic references here include Chong and Yang [4] and [5]. A number of finite injury arguments have been carried out in $\text{I}\Sigma_1$, such as the Friedberg-Muchnik theorem and even the Sacks splitting theorem. The general situation described by Groszek and Slaman [14] is that ones with recursive bounds on the number of injuries (as in the Friedberg-Muchnik type arguments) can be carried out in $\text{I}\Sigma_1$. Ones with unbounded (but still finite) injuries (as in the Sacks splitting theorem) can at times be carried out in $\text{I}\Sigma_1$ or $\text{B}\Sigma_2$. However, the ability to carry out all such arguments implies $\text{I}\Sigma_2$ over $\text{B}\Sigma_2$. On the other hand, until now there did not seem to be a known example of a single theorem of this type that cannot be carried out in $\text{B}\Sigma_2$.

Our theorem about SADS and atomic models involves a finite injury priority argument as does the one in [21] building low solutions to SADS. One could also frame the basic construction of an atomic model of an atomic theory as an argument of this sort. Viewed as a forcing argument, the general construction of an atomic model involves meeting a uniformly Π_1^0 set of dense sets (the atoms along the tree of the Henkin construction). Our construction in Theorem 4.1 gets by with $\text{I}\Sigma_1$ because of two special properties of the argument. The first corresponds to the technique developed in α -recursion theory [33] and latter called Shore blocking in applications to reverse recursion theory in models of first order arithmetic (as in Mytilinaios [29]). It relies on the fact that producing an atom for the single n -tuple c_1, \dots, c_n automatically produces one for all j -tuples from the set $\{c_1, \dots, c_n\}$ for all $j \leq n$. Thus at any stage s we can take care of all requirements of this form up to s by a single action. The second use of such a special property allowed us to avoid $\text{B}\Sigma_2$ in the argument that action for each single requirement was bounded by combining finitely many formulas into one before finding an atomic extension.

If we move on to the general case of meeting arbitrary uniformly Π_1^0 collections of dense sets we actually find a theorem proven by a priority argument of the unbounded but finite type that requires $\text{I}\Sigma_2$ in the sense of reverse mathematics. The theorem is that for any uniformly Π_1^0 collection of sets D_i , each of which is dense in $2^{<\mathbb{N}}$, there is a set G that meets each of them, i.e. $\forall i \exists s (G \upharpoonright s \in D_i)$, and is low, i.e. G' is Δ_2 . This theorem is easily proved assuming $\text{I}\Sigma_2$ by a standard finite injury argument. The number of times we act to get into a single D_i is finite but, in general, unbounded. (Each lowness requirement acts at most once after it has highest priority.) We denote the formal version of this theorem by $\Pi_1^0\text{G}$ (for Π_1^0 Generic) and claim that it implies $\text{I}\Sigma_2$ over the base theory $\text{B}\Sigma_2$. To remain in keeping with the rest of this paper we formulate our result in second order arithmetic over RCA_0 , but it is not difficult to translate it to the more usual setting of first order arithmetic.

($\Pi_1^0\text{G}$) For any uniformly Π_1^0 collection of sets D_i , each of which is dense in $2^{<\mathbb{N}}$, there is a G such that $\forall i \exists s (G \upharpoonright s \in D_i)$.

Theorem 4.3. $\text{RCA}_0 + \text{B}\Sigma_2 \vdash \Pi_1^0\text{G} \rightarrow \text{I}\Sigma_2$.

Proof. Let (M, \mathcal{S}) be a model of $\text{RCA}_0 + \text{B}\Sigma_2$ in which each uniformly Π_1^0 collection of dense subsets of $2^{<\mathbb{N}}$ has a G meeting all its members. Suppose for the sake of contradiction that $\text{I}\Sigma_2$ fails in M . Then there is a Σ_2 cut \mathcal{I} in M , i.e. $x \in \mathcal{I} \Leftrightarrow$

$\exists y \forall z \varphi(x, y, z)$, and \mathcal{I} is an initial segment of M closed under successor. The dense sets D_i we consider correspond to the actions of getting a gap large enough to correctly recognize one more witness for membership in \mathcal{I} (and also in a D_i). To be precise we let

$$D_i = \{ \sigma \in 2^{<\mathbb{N}} \mid \text{the last two elements of } \sigma \text{ are } w \text{ and } \langle i, x, y \rangle \\ \wedge \exists y' < y \forall z \varphi(x, y', z) \wedge \forall y' \leq w \exists z' < y (\neg \varphi(x, y', z')) \}.$$

Clearly the D_i are uniformly Π_1^0 . As \mathcal{I} is a cut and so not a member of \mathcal{S} , there is no bound on the least witnesses y such that $\forall z \varphi(x, y, z)$ for $x \in \mathcal{I}$, while by $\text{I}\Sigma_1$ every $x \in \mathcal{I}$ has such a least witness y , and by $\text{B}\Sigma_2$ there is a bound on the z' needed to show that y is in fact the least witness for x . Thus each D_i is dense, for if we consider any τ with last element w , there is an $x \in \mathcal{I}$ whose least witness y is larger than w and a $z > y$ such that counterexamples for all $y' \leq w$ can be found below z . So if we let σ extend τ by adding on as its first new element $\langle i, x, z \rangle$, we have our extension of τ in D_i , as required.

Let G be as in $\Pi_1^0\text{G}$ and let a be any number above all those in \mathcal{I} . By our assumption $\forall i < a \exists s (G \upharpoonright s \in D_i)$. As being in D_i is a Π_1 property uniformly, by $\text{B}\Sigma_2$ there is a t such that $\forall i < a \exists s < t (G \upharpoonright s \in D_i)$. Moreover, the set $R = \{s < t \mid \exists i < a (G \upharpoonright s \in D_i)\}$ is M -finite (i.e. coded in M by a number), as is the relation

$$S = \{ \langle i, x \rangle \mid i, x < a \wedge \exists s < t (G \upharpoonright s \in D_i) \\ \wedge \exists y (\text{the last element of } G \upharpoonright s \text{ is } \langle i, x, y \rangle) \}$$

by $\text{B}\Sigma_2$. Finally, we claim S is, in fact, a one-one function from a into \mathcal{I} and so into a proper subset of a for the desired contradiction. We have already seen that for every $i < a$ there is an $x \in \mathcal{I}$ such that $\langle i, x \rangle \in S$. Suppose then that we have $i \neq j$ both below a and $r, s < t$ such that $G \upharpoonright r \in D_i$ and $G \upharpoonright s \in D_j$, the next-to-last and last elements of $G \upharpoonright r$ are v and $\langle i, x_i, y_i \rangle$, and the next-to-last and last elements of $G \upharpoonright s$ are w and $\langle j, x_j, y_j \rangle$. Without loss of generality, we assume that $r < s$, and so $v < \langle i, x_i, y_i \rangle \leq w < \langle j, x_j, y_j \rangle$. (As $i \neq j$, we know that $G \upharpoonright r$ and $G \upharpoonright s$ have different last elements, and so $r \neq s$.) If $x_i = x_j = x$, we would have an immediate contradiction, as the definitions of D_i and D_j would say that $\exists y' < y_i \forall z \varphi(x, y', z)$ and $\forall y' \leq w \exists z' < y (\neg \varphi(x, y', z'))$, respectively, but $y_i \leq w$. \square

Note that $\Pi_1^0\text{G}$ is obviously implied by the assumption of the existence of sets Cohen 2-generic with respect to any given set. The addition of such a set by standard Cohen forcing preserves $\text{I}\Sigma_1$ and $\text{I}\Sigma_2$ (if true in the ground model). Thus the arguments of Theorem 3.13 and Corollary 3.15 show that $\Pi_1^0\text{G}$, while not conservative over $\text{B}\Sigma_2$, is $\text{r-}\Pi_2^1$ conservative over both $\text{I}\Sigma_1$ and $\text{I}\Sigma_2$ as is AMT.

To conclude our analysis, we show that retreating to AMT gives us Π_1^1 -conservativity over $\text{B}\Sigma_2$ as well as $\text{I}\Sigma_1$ and $\text{I}\Sigma_2$. The only other known examples of mathematical principles conservative over all three theories are WKL_0 (Hájek [15]) and COH (Chong, Slaman and Yang [3]). Here too we use a priority argument with Shore blocking.

Theorem 4.4. *Let \mathcal{M} be a model of $\text{B}\Sigma_2$ and T be an atomic theory in \mathcal{M} . Let \mathcal{F} be the tree of Henkin constructions of models of T in \mathcal{M} (as in Definition 3.5). Then there is a path P through \mathcal{F} such that $\mathcal{M}[P] \models \text{B}\Sigma_2$ and the model of T given recursively in P as in Remark 3.6 is atomic.*

Proof. Our goals for P are that it preserve $B\Sigma_2$ and that the associated formulas S_σ for $\sigma \in P$ (when extraneous new constants are replaced by new free variables and existentially quantified) include atoms for each initial segment c_1, \dots, c_n of Henkin constants. As auxiliary requirements we will make P be 1-generic (for every e there is a $\sigma \subseteq P$ such that $\Phi_e^\sigma(e) \downarrow$ or $\forall \tau \supseteq \sigma (\Phi_e^\tau(e) \uparrow)$) and preserve IS_1 . Our construction will be a movable marker one that produces P as a limit of approximations σ_s so that P will be Δ_2 with convergence at all initial segments σ : $(\forall \sigma \in 2^{<\mathbb{N}})[\sigma \subseteq P \Leftrightarrow \exists s \forall t > s (\sigma \subseteq \sigma_t) \Leftrightarrow \forall s \exists t > s (\sigma \subseteq \sigma_t)]$. Thus we will have $\Pi_1^P \subseteq \Sigma_2$: $\Phi_e^P(e) \uparrow \Leftrightarrow (\exists \sigma)(\sigma \subseteq P \wedge \forall \tau \supseteq \sigma (\Phi_e^\tau(e) \uparrow))$. In particular, any Σ_2^P formula in $\mathcal{M}[P]$ will then be equivalent to one that is Σ_2 in \mathcal{M} , and so $B\Sigma_2$ will hold in $\mathcal{M}[P]$ since it holds in \mathcal{M} .

We have the following requirements:

$R_{\langle e, n \rangle}$: Decide the e th Σ_1^0 sentence with parameter n , i.e. get a $\sigma \subseteq P$ such that $\Phi_e^\sigma(n) \downarrow$ or $\forall \tau \supseteq \sigma (\Phi_e^\tau(n) \uparrow)$. If we satisfy the $R_{\langle e, n \rangle}$ corresponding to a Σ_1^0 formula $\exists x \psi(x, n)$ by forcing convergence, we act to preserve IS_1 by guaranteeing that there is a least m such that $\exists x \psi(x, m)$ by getting a $\sigma \subseteq P$ such that $\Phi_e^\sigma(n) \downarrow$ and, for every $k < m$ and $\tau \supseteq \sigma$, we have $\Phi_e^\tau(k) \uparrow$.

A_e : There is a $\sigma \subseteq P$ such that S_σ implies an atom for c_1, \dots, c_e .

At each stage s of the construction we have an approximation σ_s to P and movable markers $q_{i,s}$ (which may be of type R or A) for an initial segment of i 's indicating the place $(\sigma_s \upharpoonright q_{i,s})$ at which we think we have satisfied Q_j ($Q = R$ or A) for $j < q_{i-1,s}$. At stage s we define whether a marker $q_{i,s}$ needs attention and the associated action for $q_{i,s}$ by cases.

$Q = R$: First, q_i needs attention if there exists an $\langle e, n \rangle < q_{i-1,s}$ such that $\Phi_e^{\sigma_s \upharpoonright q_{i,s}}(n) \uparrow$ and a $\tau \supseteq \sigma_s \upharpoonright q_{i,s}$ with $|\tau| < s$ such that $\Phi_e^\tau(n) \downarrow$. If so, we find the least such $\langle e, n \rangle$ and the least τ for it and set $\sigma_{s+1} = \tau \hat{\ } 0^k$ where k is chosen so that $|\sigma_{s+1}| = s$; then $q_{i,s+1} = s$. If not, but if there is an $\langle e, n \rangle < q_{i-1,s}$ such that $\Phi_e^{\sigma_s \upharpoonright q_{i,s}}(n) \downarrow$, a $k < n$ and $\tau \supseteq \sigma \upharpoonright q_{i,s}$ such that $\Phi_e^\tau(k) \downarrow$, but no $l \leq k$ such that $\Phi_e^{\sigma_s \upharpoonright q_{i,s}}(l)[s] \downarrow$, then q_i also needs attention. In this case, we find the least such $\langle e, n \rangle$ and the least k and τ for it and set $\sigma_{s+1} = \tau \hat{\ } 0^k$ where k is chosen so that $|\sigma_{s+1}| = s$; then $q_{i,s+1} = s$.

$Q = A$: If we see in s many steps that $S_{\sigma_s \upharpoonright q_{i,s}}$ does not imply an atom about $c_1, \dots, c_{q_{i-1,s}}$, then q_i needs attention. If so, we let $\tau \supseteq \sigma_s \upharpoonright q_{i-1,s}$ be least such that in s many steps we do not find a witness that S_τ does not include an atom about $c_1, \dots, c_{q_{i-1,s}}$ and set $\sigma_{s+1} = \tau \hat{\ } 0^k$ where k is chosen so that $|\sigma_{s+1}| = s$; then $q_{i,s+1} = s$.

In each case, all markers q_j for $j > i$ become undefined and all with $j < i$ remain as they were at s .

Of course, at stage s we act for the least q_i that needs attention. If no marker now defined requires attention we let $\sigma_{s+1} = \sigma_s \hat{\ } 0$ and set $q_{i,s+1} = |\sigma_{s+1}|$ for the least i for which $q_{i,s}$ is undefined.

As in the proof of Theorem 4.1 and as usual in blocking arguments, it suffices to show that if, for all $j < i$, the $q_{j,s}$ have reached their final positions q_j by stage t , then $q_{i,s}$ eventually stops moving (with final value q_i) and so P is well defined as the union of the σ_{q_i} . (The q_i are cofinal in \mathcal{M} by the construction, as they are the stage at which $q_{i,s}$ is last changed, and if none are changed at s , then a new one larger than all the others is added to the list of defined markers.) Moreover, we argue that the associated requirements are all satisfied. We have two cases.

$Q = A$: By our assumption that T is atomic there is a $\tau \supseteq \sigma \upharpoonright q_{i-1}$ such that S_τ includes an atom for $c_1, \dots, c_{q_{i-1}}$. There is a least one by $\text{I}\Sigma_1$ (in the form of the equivalent principle III_1) and by $\text{B}\Sigma_1$ a stage by which all smaller $\rho \supseteq \sigma_{q_{i-1}}$ have been seen to not do so. At the first such stage s after q_{i-1} has settled down, we set $\sigma_s = \tau \hat{\ } 0^k$ for some k and $q_{i,s} = s$ and never change $q_{i,s}$ or $\sigma_s \upharpoonright q_{i,s}$ again.

$Q = R$: Once $q_{i-1,s}$ has settled down, $q_{i,s}$ moves monotonically upward. It moves only for the sake of some $\langle e, n \rangle < q_{i-1}$ as in the first case of the description of the action of R , or some $\langle e, n \rangle < q_{i-1}$ and $k < n$ as in the second case of that description. In any event, it moves at most once for each such $\langle e, n \rangle$ or pair of $\langle e, n \rangle$ and k . The set of $\langle e, n \rangle < q_{i-1}$ and pairs of $\langle e, n \rangle < q_i$ and $k < n$ for which $q_{i,s}$ moves is Σ_1 and so by $\text{I}\Sigma_1$ finite in \mathcal{M} , and the set of stages at which it moves is then bounded as well. Once $q_{i,s}$ has settled down to its final value q_i , the value of $\sigma_s \upharpoonright q_i$ never changes either, by construction. Finally, it is clear from the definition of markers of type R requiring attention and the associated action that we satisfy the requirements: for every $\langle e, n \rangle < q_{i-1}$ either $\Phi_e^{P \upharpoonright q_i}(n) \downarrow$ or $(\forall \tau \supseteq P \upharpoonright q_i)(\Phi_e^\tau(n) \uparrow)$, and if $\Phi_e^{P \upharpoonright q_i}(n) \downarrow$, then there is a $k \leq n$ such that $\Phi_e^{P \upharpoonright q_i}(k) \downarrow$, while for every $l < k$ and $\tau \supseteq P \upharpoonright q_i$, we have $\Phi_e^\tau(l) \uparrow$.

Thus for each i in an initial segment of \mathcal{M} the $q_{i,s}$ settle down to a sequence q_i cofinal in \mathcal{M} and we have satisfied all the R requirements. Once q_i has stopped moving $\sigma \upharpoonright q_i$ never changes either, and so P settles down in the way required by the argument given at the beginning of this proof to show that $\mathcal{M}[P] \models \text{B}\Sigma_2$. Moreover, the model defined in $\mathcal{M}[P]$ by S_P as in Remark 3.6 is an atomic model of T , as we have satisfied the A requirements. \square

The usual iteration of this construction gives an ω -extension of \mathcal{M} that satisfies both $\text{B}\Sigma_2$ and AMT , and so we have our final result here.

Corollary 4.5. *AMT is Π_1^1 -conservative over $\text{B}\Sigma_2$.*

For any complete decidable atomic theory T , let \mathcal{S} be the tree of standard Henkin constructions of models of T . For any tuple of constants \vec{c} , the set of nodes of T that are atoms about \vec{c} is dense, and these sets are uniformly Π_1^0 when taken over all tuples of constants. Thus $\Pi_1^0\text{G}$ implies AMT . On the other hand, Theorem 4.3 and Corollary 4.5 show that the converse does not hold. However, Conidis [6] has shown that every ω -model of AMT is also a model of $\Pi_1^0\text{G}$, so the use of non- ω -models is essential in establishing this nonimplication. As, like AMT , $\Pi_1^0\text{G}$ can be forced by iterating the addition of Cohen 2-generics, it is also $\text{r-}\Pi_2^1$ conservative over RCA_0 by Theorem 3.13. Thus it (even with the addition of COH as in Corollary 3.16) does not imply any of WKL_0 , RT_2^2 , SRT_2^2 , CAC , ADS , SADS or DNR (even for ω -models). As we remarked above, it is easy in the standard model to meet any uniformly Π_1^0 collection of dense sets with a low set, so the degree theoretic properties mentioned in Corollary 3.10 show that it does not imply COH or CADS either, even for ω -models.

We do not know whether RT_2^2 implies $\Pi_1^0\text{G}$. It is straightforward to check that the construction used in Conidis' proof mentioned in the previous paragraph can be carried out and verified in $\text{RCA}_0 + \text{I}\Sigma_2$, and hence $\text{RCA}_0 + \text{I}\Sigma_2 \vdash \text{AMT} \rightarrow \Pi_1^0\text{G}$. Since RT_2^2 implies AMT , we have that $\text{RT}_2^2 + \text{I}\Sigma_2$ implies $\Pi_1^0\text{G}$. On the other hand, by Hirst [22], RT_2^2 implies $\text{B}\Sigma_2$, so by Theorem 4.3, $\text{RT}_2^2 + \Pi_1^0\text{G}$ implies $\text{I}\Sigma_2$. Thus the question of whether RT_2^2 implies $\Pi_1^0\text{G}$ is equivalent to the question of whether RT_2^2 implies $\text{I}\Sigma_2$, which is the first part of Question 13.7 in Cholak, Jockusch and

Slaman [2]. Of course, this equivalence remains true if we replace RT_2^2 by any of the principles considered in [21] such as SRT_2^2 , CAC and ADS that lie between RT_2^2 and AMT and imply $B\Sigma_2$.

5. TYPE OMITTING AND HYPERIMMUNITY

In this section we consider a version of the Omitting Types Theorem that has close connections with the Atomic Model Theorem. We begin with a recursion theoretic analysis, and later derive some reverse mathematical consequences. The classical Omitting Types Theorem states that if T is a complete theory and S is a countable set of nonprincipal types of T , then there is a model of T omitting all the types in S . Millar [28] proved the following effective version of this fact.

Theorem 5.1 (Millar [28]). *Let T be a complete decidable theory. Let S_0 be a recursive set of complete types of T . Let S_1 be a recursive set of nonprincipal partial types of T . There is a decidable model of T omitting all nonprincipal types in S_0 and all partial types in S_1 .*

Millar's proof involves an effective Henkin construction with additional requirements of two kinds: For each partial n -type Γ in S_1 and each n -tuple of constants \vec{c} , there is a requirement that \vec{c} should not realize Γ . Also, for each n -type Γ in S_0 and each n -tuple of constants \vec{c} , there is a requirement that if Γ is nonprincipal, then \vec{c} should not have type Γ . At a given stage of the construction, we can satisfy a requirement of the first kind simply by extending what we have currently said about \vec{c} to disagree with Γ , which we know we can do because we are guaranteed that Γ is nonprincipal.

To satisfy requirements of the second kind, we need to be a little more careful. At a given stage, when deciding whether to add a sentence φ or its negation to the elementary diagram of the model we are building, we find the strongest priority requirement (if any) that would be satisfied by adding φ or its negation, and proceed accordingly. We can then easily argue that, for any given requirement R , once all previous requirements have been satisfied, R will also eventually be satisfied.

This argument is by induction, of course, but the construction involves no injuries. Once a requirement acts, it is permanently satisfied, even if stronger requirements act later on. So we can cast our argument as a Σ_1^0 induction as follows. Given a requirement R , let F be the (finite) set of all stronger requirements that ever act. By the inductive hypothesis, there is a stage s such that every requirement in F has acted by stage s . Now we know that no requirement stronger than R will ever act after stage s , so if the type mentioned in R is nonprincipal, then R will eventually act and be satisfied.

Thus Millar's proof carries through in RCA_0 , and hence RCA_0 proves the following statement: Let T be a complete theory. Let S_0 be a set of complete types of T . Let S_1 be a set of nonprincipal partial types of T . There is a model of T omitting all nonprincipal types in S_0 and all partial types in S_1 .

On the other hand, Millar [27] also showed that there is a sense in which the Omitting Types Theorem cannot be made fully effective, as a corollary to the following result on atomic models.

Theorem 5.2 (Goncharov and Nurtazin [13]; Millar [27]). *There is a complete decidable atomic theory T such that all types of T are recursive but T has no recursive atomic model.*

Since the set of all partial types of such a theory T has a recursive listing, we have the following corollary.

Corollary 5.3 (Millar [27]). *There is a complete decidable theory T and a recursive set S of partial types of T such that no decidable model of T omits all nonprincipal partial types in S .*

Improving a result of Csima [8], Hirschfeldt [20] showed that Theorem 5.2 cannot be improved, in the following sense.

Theorem 5.4 (Hirschfeldt [20]). *Let T be a complete decidable atomic theory such that all types of T are recursive, and let $\mathbf{d} > \mathbf{0}$. Then T has a \mathbf{d} -decidable atomic model.*

We will consider a version of this result in the context of reverse mathematics in the next section. It is natural to ask whether a similar result holds for the kind of type omitting in Corollary 5.3. Csima [8] obtained the following partial result.

Theorem 5.5 (Csima [8]). *Let T be a complete decidable theory, let S be a recursive set of partial types of T , and let \mathbf{d} be such that $\mathbf{0} < \mathbf{d} \leq \mathbf{0}'$. Then there is a \mathbf{d} -decidable model of T that omits all nonprincipal partial types in S .*

However, outside the Δ_2 degrees, the situation for type omitting is different from that for atomic models. As we now show, the degrees that can always carry out the kind of type omitting in Corollary 5.3 are exactly the hyperimmune degrees. Recall that a degree is *hyperimmune* if it contains a total function f that is not dominated by any recursive function; that is, for each total recursive function g , there are infinitely many n such that $f(n) > g(n)$. If a degree is not hyperimmune, then it is *hyperimmune-free*.

Theorem 5.6.

1. *Let T be a complete decidable theory, let L be a recursive set of partial types of T , and let \mathbf{d} be a hyperimmune degree. Then there is a \mathbf{d} -decidable model of T that omits all nonprincipal partial types in L .*
2. *There is a complete decidable theory T and a recursive set L of partial types of T such that any countable model of T that omits all nonprincipal partial types in L has hyperimmune degree.*

Proof. 1. Since \mathbf{d} is hyperimmune, we can fix a \mathbf{d} -recursive function f that is not dominated by any recursive function. We use the tree \mathcal{S} of Henkin constructions of models of T as in Definition 3.5 and Remark 3.6 so that we think of our \mathbf{d} -decidable model as a path in \mathcal{S} .

Thus it is enough to show that there is an f -recursive path P of \mathcal{S} such that for every nonprincipal $\Gamma \in L$ and every tuple \vec{c} of new constants of the same arity as Γ , the following requirement is met:

$$R_{\Gamma, \vec{c}} : \text{if } \Gamma \text{ is nonprincipal, then there is a } \gamma \in \Gamma \text{ with } \neg\gamma(\vec{c}) \in \theta^P.$$

We give these requirements a priority ordering in the usual way.

We build $P = \bigcup \sigma_i$, where we will have $\sigma_0 \subset \sigma_1 \subset \dots$ and $|\sigma_i| = i$. At each stage in the construction, each requirement may have a target string. Begin by letting σ_0 be the empty sequence, and declaring that no requirement has a target.

At stage i , given $\sigma_i \in \mathcal{S}$, define σ_{i+1} as follows. Say that $R_{\Gamma, \vec{c}}$ is satisfied if there is a $\gamma \in \Gamma$ with $\neg\gamma(\vec{c}) \in S_{\sigma_i}$. Once a requirement is satisfied, cancel its target. Suppose that $R_{\Gamma, \vec{c}}$ is not satisfied and does not have a target, and there is a $\tau \in \mathcal{S}$

with S_τ propositionally extending S_{σ_i} such that $|\tau| < f(i)$ and $\neg\gamma(\vec{c}) \in S_\tau$ for some $\gamma \in \Gamma$. Then declare the least such τ (in some fixed ordering of $2^{<\mathbb{N}}$) to be the target of $R_{\Gamma, \vec{c}}$.

Let $R_{\Gamma, \vec{c}}$ be the strongest among the first i requirements that is not satisfied and has a target, and let τ be its target. If there is no such requirement, let τ be the least immediate successor of σ_i on \mathcal{S} . Let σ_{i+1} be the immediate successor of σ_i extended by τ . Cancel the targets of all requirements weaker than $R_{\Gamma, \vec{c}}$.

Clearly P is an f -recursive path of \mathcal{P} . We show by induction that each requirement is satisfied. Assume by induction that there is a stage s after which no requirement stronger than $R_{\Gamma, \vec{c}}$ has a target. If there is a stage $t \geq s$ at which $R_{\Gamma, \vec{c}}$ has a target τ , then P extends τ , which, by the way targets are chosen, implies that $R_{\Gamma, \vec{c}}$ is eventually satisfied. Thus it is enough to assume for a contradiction that Γ is nonprincipal and never satisfied, and show that there is a stage $t \geq s$ at which $R_{\Gamma, \vec{c}}$ has a target.

Define the recursive function g as follows. Given n , let M_n be the set of all $\sigma \in \mathcal{S}$ such that $|\sigma| = n$. For each $\sigma \in M_n$, let τ_σ be the least extension of σ on \mathcal{S} such that $\neg\gamma(\vec{c}) \in S_{\tau_\sigma}$ for some $\gamma \in \Gamma$. Such a τ_σ must exist, since otherwise we would have $T \cup S_\sigma \vdash \gamma(\vec{c})$ for all $\gamma \in \Gamma$, contradicting the assumption that Γ is nonprincipal. Let $g(n) = \max\{|\tau_\sigma| : \sigma \in M_n\}$. Now g is a total recursive function, so there is a $t > s$ such that $f(t) > g(t)$. Since $\sigma_t \in M_t$, there is a τ extending σ_t such that $|\tau| < f(t)$ and $\neg\gamma(\vec{c}) \in S_\tau$ for some $\gamma \in \Gamma$. Since we are assuming $R_{\Gamma, \vec{c}}$ is not satisfied at stage t , this means that $R_{\Gamma, \vec{c}}$ acquires a target at stage t , as desired.

2. Let \mathcal{T} be a subtree of $2^{\mathbb{N}}$ with no dead ends. A *partial path* of \mathcal{T} is a collection G of pairs $\langle n, i \rangle$ with $n \in \mathbb{N}$ and $i < 2$, such that there is a path P of \mathcal{T} with $P(n) = i$ for all $\langle n, i \rangle \in G$. We call any such path P a *completion* of G . We can think of a path P of \mathcal{T} as a partial path by identifying P with $\{\langle n, P(n) \rangle : n \in \mathbb{N}\}$. A partial path G is *isolated* if there is a $\sigma \in \mathcal{T}$ such that every path of \mathcal{T} extending σ is a completion of G .

We translate \mathcal{T} into a theory T in the language with infinitely many unary relation symbols U_0, U_1, \dots as follows. Let Δ be the set consisting of the following axioms:

- (1) For each $\sigma \in \mathcal{T}$ and each n ,

$$\exists^{\geq n} x \left(\bigwedge_{\sigma(i)=1} U_i(x) \wedge \bigwedge_{\sigma(i)=0} \neg U_i(x) \right).$$

- (2) For each $\sigma \notin \mathcal{T}$,

$$\neg \exists x \left(\bigwedge_{\sigma(i)=1} U_i(x) \wedge \bigwedge_{\sigma(i)=0} \neg U_i(x) \right).$$

Let T be the theory axiomatized by Δ . It is straightforward to show that T is a complete theory, and is decidable if \mathcal{T} is recursive. Furthermore, the partial paths of \mathcal{T} correspond to partial 1-types of T in a natural way, and a partial path of \mathcal{T} is isolated if and only if the corresponding partial type of T is principal. Finally, a model of T corresponds to a choice of a dense sequence of paths of \mathcal{T} (possibly with repetitions).

Thus it is enough to build a recursive subtree \mathcal{T} of $2^{\mathbb{N}}$ with no dead ends, and a uniformly recursive collection C of partial paths of \mathcal{T} , such that any countable dense sequence of paths of \mathcal{T} of hyperimmune-free degree must contain an extension of some nonisolated element of C . Since there are continuum many hyperimmune-free degrees, we cannot play directly against such sequences of paths. Instead, we will build \mathcal{T} and define a functional Ψ to satisfy the requirements

R_e : if M is a countable dense sequence of paths of \mathcal{T} and
 either Ψ^M is not total or Φ_e dominates Ψ^M , then
 M contains an extension of some nonisolated element of C .

We reserve the portion of \mathcal{T} above $0^e 1$ for satisfying R_e . The functional Ψ is defined by letting $\Psi^M(n)$ be the least k such that for each $e \leq n$, there is a $P \in M$ extending $0^e 1$, and for the first such P (in the order of M), there are $m_0 < m_1 < \dots < m_{n+1} < k$ with $P(m_i) = 1$ for each $i \leq n$. We will ensure that every path of \mathcal{T} of the form $0^e 1 \sigma 0^\omega$ is nonisolated and is in C , which will imply that if Ψ^M is not total, then either M is not dense or M contains an extension of some nonisolated element of C .

Above $0^e 1$, we act as follows, defining \mathcal{T} above this node and defining an element D_e of C . Whenever we declare a string to be in \mathcal{T} , we of course also put all of its substrings in \mathcal{T} . Except for this requirement, any string we do not declare to be in \mathcal{T} is not in \mathcal{T} . To explain the reasoning behind our construction, consider a countable dense sequence M of paths of \mathcal{T} such that Ψ^M is total and Φ_e dominates Ψ^M , and let P be the first element of M extending $0^e 1$. Our task will be to define \mathcal{T} and C so that P extends D_e , while making D_e nonisolated.

We begin by putting $\langle i, 0 \rangle$ for $i < e$ and $\langle e, 1 \rangle$ into D_e . This action ensures that any path extending D_e must extend $0^e 1$. We then declare that $0^e 1 0^n$ is in \mathcal{T} for all n , and put $0^e 1 0^\omega$ into C . For each s such that $\Phi_e(e)[s] \uparrow$ and each $n < s$, we declare that $0^e 1 0^n 1^{s-n}$ is in \mathcal{T} . Suppose there is a least stage t_0 such that $\Phi_e(e)[t_0] \downarrow$. By the usual conventions, $\Phi_e(e) < t_0$. Thus, by the definition of Ψ , the path P must extend $0^e 1 0^n 1$ for some $n < t_0$. So for each such n we declare that the string $0^e 1 0^n 1^{t_0-n}$ is in \mathcal{T} , and we call these strings the *level 1 active nodes above $0^e 1$* . For each $n \geq t_0$, we declare $0^e 1 0^n 1^k$ to be in \mathcal{T} for all k . Notice that for every path Q extending a level 1 active node above $0^e 1$ (including the path P), we have $Q(e + t_0) = 1$, while for every other path Q extending $0^e 1$, we have $Q(e + t_0) = 0$. We now put $\langle e + t_0, 1 \rangle$ into D_e , thus ensuring that any path extending D_e must extend some level 1 active node above $0^e 1$.

We now work only above the level 1 active nodes above $0^e 1$. Above these nodes, we repeat a version of the above construction. More precisely, given a level m active node σ above $0^e 1$, we proceed as follows. (We will have already defined t_0, \dots, t_{m-1} so that $|\sigma| = e + t_0 + \dots + t_{m-1}$.) We declare that $\sigma 0^n$ is in \mathcal{T} for all n and put $\sigma 0^\omega$ into C . For each s such that $\Phi_e(e + t_0 + \dots + t_{m-1})[s] \uparrow$ and each $n < s$, we declare that $\sigma 0^n 1^{s-n}$ is in \mathcal{T} . If there is a least stage t_m such that $\Phi_e(e + t_0 + \dots + t_{m-1})[t_m] \downarrow$, then we proceed as follows. For each $n < t_1$, we declare that the string $\sigma 0^n 1^{t_m-n}$ is in \mathcal{T} , and we call these strings the *level $m + 1$ active nodes above $0^e 1$* . For each $n \geq t_m$, we declare $\sigma 0^n 1^k$ to be in \mathcal{T} for all k . Finally, we put $\langle e + t_0 + \dots + t_m, 1 \rangle$ into D_e . As argued above for the $m = 1$ case, for every path Q extending a level m active node above $0^e 1$ (including the path P)

we have $Q(e + t_0 + \dots + t_m) = 1$, while for every other path Q extending $0^e 1$ we have $Q(e + t_0 + \dots + t_m) = 0$.

We have completed the definition of \mathcal{T} and C . Now suppose that M is a countable dense sequence of paths of \mathcal{T} and has hyperimmune-free degree. If Ψ^M is not total, then there is a path P in M of the form $0^e 1 \sigma 0^\omega$ for some e and σ . But every such path is put into C during the construction and is made to be nonisolated. Thus in this case M contains a path extending a nonisolated element of C . On the other hand, if Ψ^M is total, then there is some e such that Φ_e dominates Ψ^M . Let P be the first element of M extending $0^e 1$. As argued during the construction, for every m , we have $P(e + t_0 + \dots + t_m) = 1$. Since $D_e = \{\langle i, 0 \rangle : i < e\} \cup \{\langle e, 1 \rangle\} \cup \{\langle e + t_0 + \dots + t_m, 1 \rangle : m \in \mathbb{N}\}$, we see that P extends D_e . Furthermore, D_e is nonisolated, since for every path Q extending D_e and every $\sigma \subset Q$, there is a $\tau \supset \sigma$ that is not extended by any node active at any level, which implies that no path extending τ can also extend D_e . Thus in this case also, M contains a path extending a nonisolated element of C . \square

The previous results lead us to consider the following principle.

(OPT) Omitting Partial Types: Let T be a complete theory and let S be a set of partial types of T . Then there is a model of T that omits all nonprincipal partial types in S .

Since there is an ω -model of WKL_0 consisting entirely of sets of hyperimmune-free degree (see Simpson [32]), part 2 of Theorem 5.6 implies that $\text{WKL}_0 \not\vdash \text{OPT}$.

Iterating Theorem 5.5 and dovetailing in the usual way (also using the density of the r.e. degrees), we see that every r.e. degree bounds an ω -model of OPT. This fact means that OPT does not imply any of the principles mentioned above, except those that follow from RCA_0 and possibly $\text{I}\Sigma_2$ and $\text{B}\Sigma_2$. As we will see below, AMT implies OPT, and hence OPT also does not imply $\text{I}\Sigma_2$ or $\text{B}\Sigma_2$.

The reverse mathematical analogue of Theorem 5.6 also holds. Let HYP be the statement that for every set X there is a degree that is hyperimmune relative to X ; in other words, there is a function not dominated by any X -recursive function. (See Definition VII.1.4 of Simpson [32] for a definition of relative recursiveness in RCA_0 .)

Theorem 5.7. $\text{RCA}_0 \vdash \text{OPT} \leftrightarrow \text{HYP}$.

Proof. It is straightforward to check that the proof of part 2 of Theorem 5.6 relativizes to any set X and can be carried out in RCA_0 .

The proof of part 1 of Theorem 5.6 can also be carried out in RCA_0 . As usual, the only worry is the induction required to prove that every requirement is satisfied. Notice, however, that while a requirement's target may be canceled by a stronger priority requirement, once a requirement is satisfied, it can no longer be injured. Therefore we can argue by Σ_1^0 induction as follows.

Fix a requirement $R_{\Gamma, \vec{c}}$ and let F be the (finite) set of all stronger requirements that are ever satisfied. By induction, there is a stage s by which all requirements in F have been satisfied. If Γ is not principal, then, by the same argument as before, there is a stage $i > s$ at which there is a $\tau \in \mathcal{S}$ with S_τ propositionally extending S_{σ_i} such that $|\tau| < f(i)$ and $\neg\gamma(\vec{c}) \in S_\tau$ for some $\gamma \in \Gamma$. We claim that no requirement stronger than $R_{\Gamma, \vec{c}}$ acquires a target at a stage $\geq i$. Indeed, suppose otherwise and let $R_{\Gamma', \vec{c}'}$ be the strongest requirement that acquires a target at a stage $\geq i$. Then this target is never canceled, and hence $R_{\Gamma', \vec{c}'}$ is eventually satisfied. But no

requirement stronger than $R_{\Gamma, \bar{c}}$ is satisfied after stage s , so we have a contradiction. Thus, if $R_{\Gamma, \bar{c}}$ does not have a target at stage i , then it acquires one at that stage. This target is never canceled, so $R_{\Gamma, \bar{c}}$ is eventually satisfied. \square

Corollary 5.8. $RCA_0 \vdash AMT \rightarrow OPT$.

Proof. The theory built in part 2 of Theorem 5.6 is atomic, and hence AMT also implies HYP. \square

Corollary 5.9. $RCA_0 \vdash COH \rightarrow OPT$.

Proof. Jockusch and Stephan [25] proved that every p-cohesive set (i.e., one that is cohesive for the collection of primitive recursive sets) has hyperimmune degree. Their proof goes through a degree theoretic characterization of the p-cohesive sets, so we give a more direct proof in RCA_0 .

Let s_n^e be the least s such that $\Phi_e \upharpoonright n + 1[s] \downarrow$, if such an s exists. So $s_n^e \leq s_{n+1}^e$. We assume the convention that $\Phi_e(n) \leq s_n^e$. Let $r_0^e = s_0^e$ and $r_{n+1}^e = s_{\sum_{i=0}^n r_i^e}^e$.

For each e , let S_e be the set whose characteristic function is given by the sequence $0^{r_0^e} 1^{r_1^e} 0^{r_2^e} 1^{r_3^e} 0^{r_4^e} 1^{r_5^e} \dots$ if Φ_e is total, and $0^{r_0^e} 1^{r_1^e} 0^{r_2^e} 1^{r_3^e} \dots (1-i)^{r_{n-1}^e} i^\omega$ if n is the least number such that r_n^e is undefined, and $i = n \bmod 2$. It is easy to see that the sequence S_0, S_1, \dots can be defined in RCA_0 . Let C be cohesive for this sequence.

Let f be the C -recursive function defined by letting $f(n)$ be the least m such that the sequence giving the characteristic function of C has n zeroes and n ones among its first m bits. We claim that f is not dominated by any recursive function. To show this, let Φ_e be total. Suppose C is almost contained in S_e (the case where C is almost contained in the complement of S_e being symmetric). It follows from the definition of S_e that for any m we can find $y > x > m$ such that $x, y \in C$, there is no number strictly between x and y in C , and for some n we have $x < \sum_{i=0}^n r_i^e$ and $y \geq \sum_{i=0}^{n+1} r_i^e$. Then $f(x) > y > r_{n+1}^e = s_{\sum_{i=0}^n r_i^e}^e \geq s_x^e \geq \Phi_e(x)$. Since we can take m as large as we want, Φ_e does not dominate f .

Thus C has hyperimmune degree. This proof is obviously relativizable, so COH implies HYP, and hence OPT. \square

6. A WEAK FORM OF AMT

In this section we consider the effect of adding effectiveness conditions on the types of a theory to AMT.

It follows from Theorem 5.1 that a complete decidable atomic theory whose types are uniformly recursive has a decidable atomic model. Theorem 5.2 shows that this result no longer holds if we weaken the hypothesis by assuming only that the types of T are recursive, but not necessarily uniformly so. On the other hand, Theorem 5.4 shows that it does come very close to holding under this weaker hypothesis. Recall that Theorem 5.4 says that if T is a complete decidable atomic theory such that all types of T are recursive, and $\mathbf{d} > \mathbf{0}$, then T has a \mathbf{d} -decidable atomic model. This fact leads us to consider the following weak version of AMT, which captures the hypothesis of Theorem 5.4 without explicit reference to recursiveness.

We say that partial types Γ and Δ of a theory T are *equivalent* if they imply the same formulas over T . Note that if Γ is a complete type equivalent to a partial type Δ , then Γ is recursive in Δ . We say that a set S is a *subenumeration* of the types of a theory T if for every type Γ of T there is an i such that $\{\varphi \mid \langle i, \varphi \rangle\} \in S$

is equivalent to Γ . If the types of T have a subenumeration, then we say that they are *subenumerable*.

(AST) Atomic model theorem with Subenumerable Types: Let T be a complete atomic theory whose types are subenumerable. Then T has an atomic model.

Suppose the types of T are recursive. Let ψ_0, ψ_1, \dots enumerate the formulas in the language of T . Let $S = \{ \langle e, \bigwedge_{i \leq s} \psi_n \rangle \mid \Phi_e(n)[s] \downarrow = 1 \}$. Then S is recursive and is a subenumeration of the types of T . Thus, by Theorem 5.2, AST does not hold in the ω -model consisting of the recursive sets, so $\text{RCA}_0 \not\vdash \text{AST}$. Theorem 5.4 leads us to expect that AST may indeed be equivalent to the existence of relatively nonrecursive sets. We now show that this is the case, which is remarkable given the fact that the statement of AST does not use any recursion theoretic notions. We begin with the following recursion theoretic result.

Theorem 6.1 (Goncharov and Nurtazin [13]; Harrington [18]). *Let T be a complete decidable atomic theory whose principal types are uniformly recursive. Then T has a decidable atomic model.*

This result is proven by a priority argument. It has the following analogue in reverse mathematics which we prove by a similar argument using Shore blocking.

Theorem 6.2 (RCA_0). *Let T be a complete atomic theory such that there is a listing of the principal types of T . Then T has an atomic model.*

Proof. Let $\Gamma_0, \Gamma_1, \dots$ be the principal types of T . Let \mathcal{S} be the tree of standard Henkin constructions of models of T , with Henkin constants c_i . For $\sigma \in \mathcal{S}$, let S_σ be the set of sentences associated with σ as in Definition 3.5. For $\sigma \in \mathcal{S}$ and $n \in \mathbb{N}$, let $t(\sigma, n)$ be the least t such that $\exists x_{n+1}, \dots, x_m \bigwedge_{\varphi \in S_\sigma} \varphi[c_0/x_0, \dots, c_m/x_m] \in \Gamma_t$, where c_m is the largest constant mentioned in S_σ . For a path P of \mathcal{S} , let $t(P, n) = \lim_k t(P \upharpoonright k, n)$, which of course can be infinite. Note that if $t(P, n) < \infty$, then c_0, \dots, c_n has type $\Gamma_{t(P, n)}$ in the model \mathcal{M} of T determined by P . Thus, if $t(P, n) < \infty$ for all n , then \mathcal{M} is atomic.

We define a path P of \mathcal{S} as follows. Begin at the root node of \mathcal{S} , and define $n_{i,0} = i$ for all $i \in \mathbb{N}$. Suppose we have reached a node σ of length k . If σ has a single successor $\sigma \hat{\ } j$, then proceed to that successor and let $n_{i,k+1} = n_{i,k}$ for all i . Note that in this case $t(\sigma \hat{\ } j, n) = t(\sigma, n)$ for all n .

If σ has two successors, then for $j \in \{0, 1\}$, let $d_{\sigma \hat{\ } j}$ be the least $i \leq k$ such that $t(\sigma \hat{\ } j, n_{i,k}) > t(\sigma, n_{i,k})$, or $d_{\sigma \hat{\ } j} = \infty$ if there is no such i . If $d_{\sigma \hat{\ } 0} = d_{\sigma \hat{\ } 1}$, then let $j = 0$. (Note that this case can happen only if $d_{\sigma \hat{\ } 0} = d_{\sigma \hat{\ } 1} = \infty$.) Otherwise, let j be such that $d_{\sigma \hat{\ } j} > d_{\sigma \hat{\ } (1-j)}$. Proceed to $\sigma \hat{\ } j$. For $i < d_{\sigma \hat{\ } j}$, let $n_{i,k+1} = n_{i,k}$. For $i \geq d_{\sigma \hat{\ } j}$, let $n_{i,k+1} = k + i$.

Having thus defined P , let $n_i = \lim_k n_{i,k}$, which can be infinite. We claim that the finite values of n_i are unbounded. In other words, for each n there is an i such that $n < n_i < \infty$.

Suppose otherwise, and let n be such that every n_i is either less than n or infinite. Let e be least such that $n_{e,k} > n$ for some k . Such an e exists by Σ_1^0 -induction (and the fact that there exist i, k such that $n_{i,k} > n$). Let $i < e$. Then $i < t(P \upharpoonright k+1, n_i)$ for all $k \geq n$, so $t(P \upharpoonright k+1, n_i) = t(P \upharpoonright k, n_i)$ for all $k \geq n$, and hence $t(P, n_i) < \infty$. Let α_{n_e-1} be an atom in $\Gamma_{t(P, n_e)}$. From α_{n_e-1} we can obtain atoms α_i in $\Gamma_{t(P, n_i)}$ for each $i < e$. Let m be such that $S_{P \upharpoonright m}$ contains $\alpha_i[x_0/c_0, \dots, x_{n_i}/c_{n_i}]$ for all $i < e$. Note that for all $\tau \succ S_{P \upharpoonright m}$ on \mathcal{S} and all $i < e$, we must have $t(\tau, n_i) = t(P, n_i)$.

Since $n_e > n$, we have $n_e = \infty$. So there is an $l \geq n$ such that $d_{P \upharpoonright l+1} \leq e$. Let $\tau = P \upharpoonright l \wedge (1 - P(l))$. By the construction of P we have $d_\tau < d_{P \upharpoonright l+1} \leq e$, so for some $i < e$ we have $t(\tau, n_i) > t(P \upharpoonright l, n_i) = t(P, n_i)$, contradicting the last statement of the previous paragraph.

Thus the finite values of n_i are unbounded. So given n there is an i such $n < n_i < \infty$. For all sufficiently large k we must have $d_{P \upharpoonright k} > i$, and hence $t(P \upharpoonright k+1, n_i) = t(P \upharpoonright k, n_i)$, which implies that $t(P \upharpoonright k+1, n) = t(P \upharpoonright k, n)$. So $t(P, n) < \infty$. Thus the model of T determined by P is atomic. \square

We are now ready to show that AST is equivalent to the existence of relatively nonrecursive sets.

Theorem 6.3. $RCA_0 \vdash AST \leftrightarrow \forall X \exists Y (Y \not\leq_T X)$.

Proof. First assume AST and fix a set X . The relativized form of the component of the proof of Theorem 5.2 provided as the proof of Theorem 5.2 of Csima, Hirschfeldt, Knight, and Soare [9] builds an X -recursive subtree \mathcal{T} of $2^{<\mathbb{N}}$ with no dead ends, all of whose paths are X -recursive, but such that there is no X -recursive listing of the isolated paths of \mathcal{T} . It is straightforward to check that this construction can be carried out in RCA_0 . Translate \mathcal{T} into a tree T as in the proof of Theorem 5.6. The 1-types of T correspond to the paths of \mathcal{T} . But because the language of T has only unary relations, any n -type is simply a union of 1-types. It follows that T is atomic and its types are X -recursive. By the same argument as in the paragraph preceding this theorem, we can X -recursively define a subenumeration of the types of T . By AST, T has an atomic model \mathcal{M} . We can form the set of all 1-types realized in \mathcal{M} , that is, all principal 1-types of T . From this set we get a set consisting exactly of the isolated paths of \mathcal{T} . This set is not X -recursive, so $\exists Y (Y \not\leq_T X)$.

Now assume $\forall X \exists Y (Y \not\leq_T X)$ and let T be a complete atomic theory whose types have a subenumeration S . Let $D \not\leq_T S$. For each n , let \mathcal{T}_n be the tree of n -types of T , which we can think of as a subtree of $2^{<\omega}$. Let \mathcal{T} be the result of gluing together the \mathcal{T}_n into a single tree by placing a copy of \mathcal{T}_n with root $0^n 1$ for each n . Then \mathcal{T} has no dead ends and the isolated paths of \mathcal{T} are dense. Furthermore, each path of \mathcal{T} (other than 0^ω) is a type of T equivalent to one of the rows of S , and hence is recursive in S . So for each path P of \mathcal{T} we have $D \not\leq_T P$. It is easy to check that the proof of Theorem 2.1 of Hirschfeldt [20] can now be carried out in RCA_0 to show that there is a D -recursive listing of the isolated paths of \mathcal{T} , and hence a D -recursive listing of the principal types of T . It follows from Theorem 6.2 that T has an atomic model. \square

Thus AST is implied by all principles mentioned above other than the ones that follow from RCA_0 , $I\Sigma_2$ and $B\Sigma_2$, and implies none of the principles mentioned above other than the ones that follow from RCA_0 . Indeed, AST is the weakest reasonable principle that does not hold in the ω -model consisting of the recursive sets.

7. SUMMARY AND OPEN QUESTIONS

We summarize most of our main results in Diagram 2. As before, double arrows are strict implications and negated arrows are nonimplications. We conclude with three open questions.

Question 7.1. Does COH (or CADS) imply AMT over RCA_0 ?

Question 7.2. Does CADS imply OPT over RCA_0 ?

Question 7.3. Is AMT $\text{r-}\Pi_2^1$ conservative over $\text{B}\Sigma_2$?




Diagram 2

ACKNOWLEDGEMENTS

The authors thank the Institute for Mathematical Sciences of the National University of Singapore for its generous support during the period July 18 – August 15, 2005, when part of this research was done.

REFERENCES

- [1] Chang, C. C.; Keisler, H. J. [1973], *Model Theory*, North-Holland, Amsterdam. MR0409165 (53:12927)
- [2] Cholak, P. A.; Jockusch, Jr., C. G.; Slaman, T. A. [2001], On the strength of Ramsey's Theorem for pairs. *J. Symbolic Logic* **66**, 1–55. MR1825173 (2002c:03094)
- [3] Chong, C. T.; Slaman, T. A.; Yang, Y. [ta], On the strength of Ramsey's theorem for pairs. In preparation.
- [4] Chong, C. T.; Yang, Y. [1998], Recursion theory in weak fragments of Peano arithmetic: A study of cuts. *Proc. Sixth Asian Logic Conference, Beijing 1996*, World Scientific, Singapore, 47–65. MR1789730 (2001g:03111)

- [5] Chong, C. T.; Yang, Y. [2000], Computability theory in arithmetic: Provability, structure and techniques. *Computability Theory and Its Applications: Current Trends and Open Problems*, P. Cholak, S. Lempp, M. Lerman and R. A. Shore eds., *Contemporary Mathematics* **257**, AMS, Providence, RI, 73–82. MR1770735 (2001d:03098)
- [6] Conidis, C. [2008], Classifying model-theoretic properties, *J. Symbolic Logic*, **73** 885–905. MR2444274
- [7] Cooper, S. B. [1972], Jump equivalence of the Δ_2^0 hyperhyperimmune sets. *J. Symbolic Logic* **37**, 598–600. MR0360240 (50:12690)
- [8] Csimá, B. F. [2004], Degree spectra of prime models. *J. Symbolic Logic* **69**, 430–442. MR2058182 (2005d:03067)
- [9] Csimá, B. F.; Hirschfeldt, D. R.; Knight, J.; Soare, R. I. [2004], Bounding prime models. *J. Symbolic Logic* **69**, 1117–1142. MR2135658 (2005m:03065)
- [10] Downey, R.; Hirschfeldt, D. R.; Lempp, S.; Solomon, R. [2001], A Δ_2^0 set with no infinite low subset in either it or its complement. *J. Symbolic Logic* **66**, 1371–1381. MR1856748 (2002i:03046)
- [11] Ershov, Yu. L.; Goncharov, S. S.; Nerode, A.; Remmel, J. B. (eds.); Marek, V. W. (assoc. ed.) [1998], *Handbook of Recursive Mathematics*, 2 vols., Elsevier, New York.
- [12] Friedman, H. [1976], Systems of second order arithmetic with restricted induction I (abstract). *J. Symbolic Logic* **41**, 557–558.
- [13] Goncharov, S. S.; Nurtazin, A. T. [1973], Constructive models of complete decidable theories. *Algebra Logic* **12**, 67–77. MR0398816 (53:2667)
- [14] Groszek, M.; Slaman, T. A. [nd], Foundations of the priority method I: finite and infinite injury. Preprint.
- [15] Hájek, P. [1993], Interpretability and fragments of arithmetic. *Arithmetic, Proof Theory, and Computational Complexity (Prague, 1991)*, *Oxford Logic Guides* **23**, Oxford Univ. Press, New York, 185–196. MR1236462 (94f:03066)
- [16] Hájek, P.; Pudlák, P. [1998], *Metamathematics of First-Order Arithmetic*, second printing. *Perspect. Math. Logic*, Springer-Verlag, Berlin. MR1748522 (2000m:03003)
- [17] Harizanov, V. S. [1998], Pure computable model theory, in Ershov et al. [1998], vol. 1, 3–114. MR1673621 (2000f:03108)
- [18] Harrington, L. [1974], Recursively presentable prime models. *J. Symbolic Logic* **39**, 305–309. MR0351804 (50:4292)
- [19] Herrmann, E. [2001], Infinite chains and antichains in computable partial orderings. *J. Symbolic Logic* **66**, 923–934. MR1833487 (2003e:03082)
- [20] Hirschfeldt, D. R. [2006], Computable trees, prime models, and relative decidability. *Proc. Amer. Math. Soc.* **134**, 1495–1498. MR2199197 (2007b:03055)
- [21] Hirschfeldt, D. R.; Shore, R. A. [2007], Combinatorial principles weaker than Ramsey’s Theorem for pairs. *J. Symbolic Logic* **72**, 171–206. MR2298478 (2007m:03115)
- [22] Hirst, J. L. [1987], *Combinatorics in Subsystems of Second Order Arithmetic*, Ph.D. Dissertation, The Pennsylvania State University.
- [23] Jockusch, Jr., C. G. [1972], Ramsey’s Theorem and recursion theory. *J. Symbolic Logic* **37**, 268–280. MR0376319 (51:12495)
- [24] Jockusch, Jr., C. G.; Soare, R. I. [1972], Π_1^0 classes and degrees of theories. *Trans. Amer. Math. Soc.* **173**, 33–56. MR0316227 (47:4775)
- [25] Jockusch, Jr., C. G.; Stephan, F. [1993], A cohesive set which is not high. *Math. Log. Quart.* **39**, 515–530 (correction in *Math. Log. Quart.* **43**, 569). MR1270396 (95d:03078)
- [26] Lerman, M. [1983], *Degrees of Unsolvability*, Springer-Verlag, Berlin. MR708718 (85h:03044)
- [27] Millar, T. S. [1978], Foundations of recursive model theory. *Ann. Math. Logic* **13**, 45–72. MR482430 (80a:03051)
- [28] Millar, T. S. [1983], Omitting types, type spectrums, and decidability. *J. Symbolic Logic* **48**, 171–181. MR693259 (85a:03049)
- [29] Mytilinaios, M. E. [1989], Finite injury and Σ_1 -induction. *J. Symbolic Logic* **54**, 212–221. MR987320 (90i:03067a)
- [30] Paris, J. B.; Kirby, L. A. S. [1978], Σ_n -collection schemas in arithmetic. *Logic Colloquium ’77*, *Stud. Logic Foundations Math.* **96**, North-Holland, Amsterdam-New York, 199–209. MR519815 (81e:03056)
- [31] Robinson, R. W. [1971], Interpolation and embedding in the recursively enumerable degrees. *Ann. Math. (2)* **93**, 285–314. MR0274286 (43:51)

- [32] Simpson, S. G. [1999], *Subsystems of Second Order Arithmetic, Perspect. Math. Logic*, Springer-Verlag, Berlin. MR1723993 (2001i:03126)
- [33] Shore, Richard A., *Splitting an α -recursively enumerable set*. Trans. Amer. Math. Soc. **204** (1975), 65-77. MR0379154 (52:60)
- [34] Soare, R. I. [1987], *Recursively Enumerable Sets and Degrees*, Springer-Verlag, Berlin. MR882921 (88m:03003)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, ILLINOIS 60637
E-mail address: `drh@math.uchicago.edu`

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY, ITHACA, NEW YORK 14853
E-mail address: `shore@math.cornell.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, BERKELEY, CALIFORNIA 94720
E-mail address: `slaman@math.berkeley.edu`