

ORBIT FULL GROUPS FOR LOCALLY COMPACT GROUPS

A. CARDERI AND F. LE MAÎTRE

ABSTRACT. We show that the topological rank of an orbit full group generated by an ergodic, probability measure-preserving free action of a non-discrete unimodular locally compact Polish group is two. For this, we use the existence of a cross section and show that for a locally compact Polish group, the full group generated by any dense subgroup is dense in the orbit full group of the action of the group.

We prove that the orbit full group of a free action of a locally compact Polish group is extremely amenable if and only if the acting group is amenable, using the fact that the full group generates the von Neumann algebra of the action.

CONTENTS

Introduction	2321
1. Orbit equivalence in the locally compact case	2323
2. Dense subgroups in orbit full groups	2330
3. Topological rank of orbit full groups	2337
4. The orbit full group as a unitary group	2337
5. Extreme amenability of orbit full groups	2340
Appendix A. Haar measures for equivalence relations	2342
Acknowledgments	2347
References	2347

INTRODUCTION

The full group of a measure-preserving action of a countable group on a standard probability space (X, μ) is the group of measure-preserving transformations which preserve every orbit. It is a (complete) invariant of orbit equivalence for the action and has a natural Polish topology induced by the uniform metric $d_u(S, T) := \mu(\{x \in X : S(x) \neq T(x)\})$. This group topology encodes many interesting properties of the action. For example Giordano and Pestov proved in [GP07] that if the group acts freely, then the full group is extremely amenable if and only if the acting group is amenable. Another example was provided by the second-named author in [LM14]: the cost of the action is very closely related to the topological rank of the full group, that is, the minimum number of generators needed to generate a dense subgroup.

Received by the editors January 31, 2016, and, in revised form, May 24, 2016.

2010 *Mathematics Subject Classification.* Primary 37A15, 37A20; Secondary 22D10, 46L10.

The authors were partially supported by Projet ANR-14-CE25-0004 GAMME.

The first author was partially supported by ERC Consolidator Grant No. 681207.

Now let G be a Polish group and consider a measure-preserving G -action on a standard probability space (X, μ) . In [CM16], we initiated the study of a Polish group topology on the associated *orbit full group*

$$[\mathcal{R}_G] = \{T \in \text{Aut}(X, \mu) : \forall x \in X, T(x) \in G \cdot x\}$$

which coincides with the uniform topology when G is a countable discrete group. In this work, we want to analyze orbit full groups associated to free actions of second countable locally compact groups, which most of the time we will suppose to be unimodular.

This study is motivated by the fact that for actions of locally compact groups, orbit full groups are still complete invariants of orbit equivalence [CM16, Thm. 3.26], so their topological properties should reflect properties of the associated equivalence relation. Moreover, these orbit full groups are better behaved since they arise naturally as unitary groups (see Section 4) and “preserve density” as follows.

Theorem A (see Theorem 2.1). *For every measure-preserving action of a locally compact Polish group G on a probability space (X, μ) and for every dense subgroup $H \subset G$, the orbit full group $[\mathcal{R}_H]$ is dense in $[\mathcal{R}_G]$.*

The above theorem is false for general actions of Polish groups: an example of Kolmogorov gives a measure-preserving action of the bijection group of the integers \mathfrak{S}_∞ such that whenever $H \leq \mathfrak{S}_\infty$ is a dense countable group, the full group $[\mathcal{R}_H]$ is *not* dense in $[\mathcal{R}_{\mathfrak{S}_\infty}]$ (see [CM16, Ex. 3.14]).

We will actually prove that Theorem A holds for every *suitable* action of a Polish group (see Definition 2.2 and Theorem 2.3)). This notion was introduced by Becker in [Bec13], where he proved that any measure-preserving action of a locally compact Polish group is suitable. On the other hand, general Polish groups can have actions which are suitable and actions which are not. As an example, the standard Bernoulli shift of \mathfrak{S}_∞ on $[0, 1]^\mathbb{N}$ is suitable (see Example 2.4).

Using Theorem A, we can then show that orbit full groups associated to actions of locally compact non-compact and non-discrete Polish groups contain a dense 2-generated subgroup. This is in sharp contrast with the discrete case where the topological rank reflects the cost of the equivalence relation and thus can be equal to any integer $n \geq 2$. Our result also shows that cost cannot provide a rich invariant of orbit equivalence for actions of non-discrete locally compact groups (see Remark 1.21).

Theorem B (see Theorem 3.1). *Let G be a locally compact unimodular non-discrete non-compact Polish group. For every measure-preserving, essentially free and ergodic action of G , there is a dense G_δ of couples (T, U) in $[\mathcal{R}_G]^2$ which generate a dense free subgroup of $[\mathcal{R}_G]$ acting freely. In particular, the topological rank of $[\mathcal{R}_G]$ is 2.*

A key tool in the proof of Theorem B is a well-known result of Forrest, namely the existence of a *cross section* for actions of locally compact groups [For74]. This will roughly provide a countable group Γ such that the cost 1 group $\mathbb{Z} \times \Gamma$ is a “measurable dense subgroup” of G . We can then use the results of [LM14] along with Theorem A to find a dense 2-generated subgroup.

Remark. Theorem B is also true in the case G is compact acting ergodically on (X, μ) . Indeed in this case the action is essentially transitive and $[\mathcal{R}_G] = \text{Aut}(X, \mu)$ which has a dense G_δ of couples of topological generators inducing a free action of

the free group on two generators by results of Prasad [Pra81, Pra83]. We do not know whether Theorem B holds for non-discrete Polish groups in general, even in the case of suitable actions.

In this work, we also extend a result of Giordano and Pestov, Theorem 5.7 of [GP07], that says that a full group of a free, ergodic action of a countable group is *extremely amenable* if and only if the acting countable group is amenable.

Before stating the theorem, let us recall that a group is *extremely amenable* if every action of the group on a compact space admits a fixed point. The first example of an extremely amenable group was given by Christensen and Herer [CH75]. Since then several examples of extremely amenable groups have been found such as the unitary group of a separable Hilbert space [GM83] or the group of measure-preserving bijections of a standard probability space [GP02].

Theorem C. *Let G be a locally compact second countable unimodular group acting freely and ergodically on (X, μ) . Then the full group of G is extremely amenable if and only if G is amenable.*

The proof of the direct implication in the theorem is an easy adaptation of Giordano and Pestov's arguments to the locally compact case, using cross sections.

For the other direction, we follow a different path and use von Neumann algebras. We first prove that the von Neumann algebra of the action $G \curvearrowright (X, \mu)$ is generated by the full group $[\mathcal{R}_G]$; see Proposition 4.3. We use this to show that if $[\mathcal{R}_G]$ is extremely amenable, then the von Neumann algebra of the action is amenable and therefore the acting group is amenable.

1. ORBIT EQUIVALENCE IN THE LOCALLY COMPACT CASE

1.1. Measure-preserving actions. Whenever a group G acts on a set X and $x \in X$, we denote by $G_x \leq G$ the stabilizer of x . The **free part** of an action $G \curvearrowright X$ is the G -invariant set of all $x \in X$ such that $G_x = \{e\}$.

A standard probability space is a probability space (X, \mathcal{B}, μ) such that (X, \mathcal{B}) is a standard Borel space and μ is a Borel non-atomic probability measure. All such probability spaces are isomorphic; see [Kec95, Thm. 17.41]. A subset A of X is a **Borel set** if it belongs to the σ -algebra \mathcal{B} . It is called a (Lebesgue-) **measurable set** if it belongs to the μ -completion of \mathcal{B} . From now on, we will drop the \mathcal{B} and fix a standard probability space (X, μ) .

Whenever G is a Polish group, a **Borel G -action** is a Borel action map $\alpha : G \times X \rightarrow X$. As usual, we will often drop the letter α and let $g \cdot x := \alpha(g, x)$ for every $g \in G$ and $x \in X$. The following lemma is well known; for a proof see [MRV13, Lem. 10].

Lemma 1.1. *Let G be a locally compact Polish¹ group, and consider a Borel G -action on a standard Borel space X . Then the free part of the G -action is a Borel subset of X .*

We denote by $\text{Aut}(X, \mu)$ the group of all measure-preserving Borel bijections of (X, μ) , where we identify two such bijections if they coincide on a full measure subset of X . It is equipped with the **weak topology**, defined to be the coarsest group

¹Recall that a locally compact group is Polish if and only if it is second countable (see [Kec95, Theorem 5.3]).

topology which makes the maps $T \in \text{Aut}(X, \mu) \mapsto \mu(T(A) \triangle A)$ continuous for every Borel set A . This turns $\text{Aut}(X, \mu)$ into a Polish group (see e.g. [Kec10, I.1.(B)]).

A **measure-preserving** G -action on (X, μ) is a Borel G -action on X such that for every $g \in G$ and every Borel $A \subseteq X$, one has $\mu(gA) = \mu(A)$. If G is a group, a near- G -action on (X, μ) is a homomorphism $G \rightarrow \text{Aut}(X, \mu)$. Every measure-preserving action induces a near-action, and a near-action is the same as an action by μ -preserving automorphisms on the measure algebra of (X, μ) .

The following lemma is well known when G is locally compact (see for instance [OW87, Lem. II.1.1]). We include a simple proof which works for all Polish groups.

Lemma 1.2. *Every measure-preserving action α of a Polish group G on (X, μ) induces a continuous near-action $\rho_\alpha : G \rightarrow \text{Aut}(X, \mu)$.*

Proof. By Pettis' Lemma (see [BK96, Thm. 1.2.6]), we only need to check that ρ_α is a Borel map. By definition of the weak topology on $\text{Aut}(X, \mu)$ it is enough to show that for every Borel subset A of X and every $\epsilon > 0$, the set

$$B := \{g \in G : \mu(g(A) \triangle A) < \epsilon\}$$

is Borel. For this, observe that since the action is Borel, the subset $\Gamma := \{(g, x) \in X \times G : x \in g(A)\}$ is Borel and hence $\Gamma_A := \Gamma \triangle (A \times G)$ is also Borel. This implies that the map

$$M : g \mapsto \mu(\{x \in X : x \in A \triangle g(A)\})$$

is also Borel. So we can conclude observing that $B = M^{-1}([0, \epsilon])$. \square

For locally compact Polish groups, measure-preserving actions and continuous near-actions are in one-to-one correspondence.

Theorem 1.3 (Mackey, [Mac62]). *Let G be a locally compact Polish group and let (X, μ) be a standard probability space. Then for every continuous homomorphism $\rho : G \rightarrow \text{Aut}(X, \mu)$ there exists a measure-preserving action α of G on (X, μ) such that the induced homomorphism $\rho_\alpha : G \rightarrow \text{Aut}(X, \mu)$ is equal to ρ .*

Moreover if α and β are two measure-preserving actions of G such that the induced homomorphisms ρ_α and ρ_β are equal, then there is a Borel G -invariant subset $A \subset X$ of full measure such that $\alpha|_A = \beta|_A$.

Remark 1.4. The above result is in sharp contrast with the following situation which was uncovered by Glasner, Tsirelson and Weiss: if G is a Levy Polish group, every measure-preserving G -action is trivial but G can still have interesting continuous near-actions. Examples include $\text{Aut}(X, \mu)$ itself or the orthogonal group of an infinite-dimensional Hilbert space; see [GTW05].

If G is locally compact and Polish, then we will call the measure-preserving action associated to a near-action $\rho : G \rightarrow \text{Aut}(X, \mu)$ a realization of the near-action. Let us recall two important definitions.

Definition 1.5. A measure-preserving action of a Polish group G on the probability measure space (X, μ) is

- **essentially free** if the free part of the action has full measure, that is, if there is a measurable subset of full measure $A \subset X$ such that for every $x \in A$ and every $g \in G$, we have that $gx \neq x$;
- **ergodic** if every Borel subset $A \subset X$ which is **almost G -invariant** (i.e. for all $g \in G$ we have $\mu(A \triangle g(A)) = 0$) has measure 0 or 1.

Remark 1.6.

- In the definition of essential freeness, one can actually assume that A is G -invariant and Borel by Lemma 1.1.
- There are actions of compact groups such that for every $g \in G$, the set $\{x \in X : gx = x\}$ has measure 0 but which are not essentially free.
- Mackey's Theorem implies that if G is locally compact and Polish, then if a realization of an action is essentially free, then all Borel realizations are essentially free.
- Another easy consequence of Mackey's Theorem 1.3 is that a measurable action of a locally compact Polish group is ergodic if and only if every Borel subset $A \subset X$ which is G -invariant (i.e. for all $g \in G$, we have $g(A) = A$) has measure 0 or 1 (see [Mac62, Theorem 3]). This is not true for Polish groups in general, as witnessed by Kolmogorov's example [Dan00, Example 9].

Every locally compact Polish group admits an essentially free measure-preserving action (see e.g. Proposition 1.2 in [AEG94]). We will now give two concrete examples of measure-preserving actions.

Example 1.7. Suppose that G is totally disconnected and non-compact. By van Dantzig's Theorem, there exists a chain $(K_n)_{n \in \mathbb{N}}$ of compact open subgroups of G such that $\bigcap_n K_n = \{1_G\}$. We now let G act by permutations on the countable set $\bigsqcup_n G/K_n$. The associated Bernoulli shift on $[0, 1]^{\bigsqcup_n G/K_n}$ is essentially free since the G -action on $\bigsqcup_n G/K_n$ is faithful, and ergodic because every G -orbit on $\bigsqcup_n G/K_n$ is infinite.

Example 1.8. Suppose that G has a lattice $\Gamma < G$, let λ be a Haar measure on G and let D be a fundamental domain of the right Γ -action on G . Then any probability measure-preserving action of Γ on (X, μ) induces a measure-preserving action of G on $(X \times D, \mu \times \lambda|_D)$; see Definition 4.2.21 in [Zim84].

We will see in Section 1.3 that all actions of any locally compact Polish group can be decomposed as a product equivalence relation analogous to the previous example.

1.2. Orbit full groups. Let us start by recalling Dye's definition of full groups [Dye59]. A subgroup $\mathbb{G} \leq \text{Aut}(X, \mu)$ is **full** if whenever (A_n) is a partition of a full measure subset of (X, μ) and $(T_n)_{n \in \mathbb{N}}$ is a sequence of elements of \mathbb{G} such that $(T_n(A_n))_{n \in \mathbb{N}}$ is a partition of a full measure subset of X , the new element $T \in \text{Aut}(X, \mu)$ defined by

$$T(x) = T_n(x) \quad \text{for all } x \in A_n$$

actually belongs to \mathbb{G} . A full group is **ergodic** if every almost \mathbb{G} -invariant Borel subset of X has measure 0 or 1. Given a group $G \leq \text{Aut}(X, \mu)$, there is a smallest full group containing G , denoted by $[G]$. If the corresponding G -almost action is ergodic, then $[G]$ is ergodic. The following proposition is well known in the case of full groups of ergodic measure-preserving equivalence relations. Its proof in the general case can be found in [Dye59, Lem. 3.2].

Proposition 1.9. *Let $\mathbb{G} \leq \text{Aut}(X, \mu)$ be an ergodic full group. Then for any $A, B \subseteq X$ of same measure, there is $T \in \mathbb{G}$ such that $T(A) = B$ up to measure zero.*

For a measure-preserving action of a Polish group G on X , we will denote by \mathcal{R}_G the *orbit equivalence relation*

$$\mathcal{R}_G := \{(x, y) \in X \times X : \text{there exists } g \in G \text{ such that } g \cdot x = y\}.$$

We recall now the definition of orbit full groups and their Polish topology (see [CM16] for more details and proofs).

Definition 1.10. Let G be a Polish group. The **orbit full group** of a probability measure-preserving action of G on (X, μ) is the group

$$[\mathcal{R}_G] := \{T \in \text{Aut}(X, \mu) : T(x) \in G \cdot x \text{ for almost every } x \in X\}.$$

Let d_G be a compatible, right-invariant and bounded metric on G . We denote by $L^0(X, \mu, G)$ the space of measurable functions from X to G which we equip with the metric

$$\tilde{d}_G(f, g) := \int_X d_G(f(x), g(x)) d\mu(x).$$

The topology induced by this metric only depends on the topology of G . It is a Polish topology called the topology of *convergence in measure*.

For a probability measure-preserving action of G on (X, μ) , for every measurable subset $A \subset X$ and measurable function $f : A \rightarrow G$, we define

$$\Phi(f) : A \rightarrow X, \text{ by } \Phi(f)(x) = f(x)x,$$

and we put $\widetilde{[\mathcal{R}_G]} := \Phi^{-1}([\mathcal{R}_G])$.

The Polish space $\widetilde{[\mathcal{R}_G]}$ equipped with the product $f \cdot g(x) = f(g(x)x)g(x)$ becomes a Polish group for the topology of convergence in measure. Moreover the map $\Phi : \widetilde{[\mathcal{R}_G]} \rightarrow [\mathcal{R}_G]$ is a group homomorphism with respect to this product.

The topology of **convergence in measure** on $[\mathcal{R}_G]$ is the quotient topology induced by Φ and we proved in Theorem 1 of [CM16] that it is a Polish group topology.

Remark 1.11. If the action of G is essentially free, then the map Φ is a bijection, so the convergence in measure on $[\mathcal{R}_G]$ is given by the metric \tilde{d}_G .

We also recall that full groups of locally compact Polish groups are complete invariants of orbit equivalence.

Definition 1.12. A probability measure-preserving action of the group G on (X, μ) is **orbit equivalent** to a probability measure-preserving action of the group H on (Y, ν) if there exists a subset of full measure $A \subset X$ and a measure-preserving Borel injection $\varphi : A \rightarrow Y$ such that

$$\varphi \times \varphi(\mathcal{R}_G \cap (A \times A)) = \mathcal{R}_H \cap (\varphi(A) \times \varphi(A)).$$

We will also say that the equivalence relations \mathcal{R}_G and \mathcal{R}_H are *isomorphic up to measure zero*. We recall the following theorem.

Theorem 1.13 ([CM16, Thm. 3.26]). *Let G and H be locally compact Polish groups acting on the probability space (X, μ) preserving the measure. Then the actions are orbit equivalent if and only if the associated orbit full groups are isomorphic.*

1.3. Cross sections and product decomposition. We now present the most important property of measure-preserving actions of locally compact Polish groups: the existence of a *cross section*.

Definition 1.14. Consider an essentially free, measure-preserving action of a locally compact Polish group G on a standard probability space (X, μ) . A Borel subset $Y \subset X$ is a **cross section** of the action if there exists a neighborhood of the identity $U \subset G$ such that the map $\theta : U \times Y \rightarrow X$ defined by $\theta(u, y) := uy$ is injective and such that $\mu(X \setminus GY) = 0$.

The existence of cross sections was proved by Forrest in [For74, Proposition 2.10] in the more general context of *non-singular* actions. For a more recent proof, we invite the reader to read Theorem 4.2 of [KPV15]. The following theorem is essentially a version of [For74, Proposition 2.13] in the context of a measure-preserving action of a unimodular locally compact group.

Theorem 1.15. *Let G be a unimodular locally compact non-compact and non-discrete Polish group, denote by λ a Haar measure on G . Consider a measure-preserving, essentially free and ergodic action of G on the standard probability space (X, μ) .*

- (i) *There exists a standard probability space (Y, ν) and a countable group Γ acting on (Y, ν) by measure-preserving transformations such that the action of G is orbit equivalent to the product action of $\mathbb{S}^1 \times \Gamma$ on $(\mathbb{S}^1 \times Y, L \times \mu)$, where \mathbb{S}^1 is the circle group² acting on itself by translation and L is its normalized Lebesgue measure.*
- (ii) *Identifying $G \times X$ with \mathcal{R}_G via the map $(g, x) \mapsto (g, g \cdot x)$, one can choose an orbit equivalence map $\Theta : (\mathbb{S}^1 \times Y, L \times \nu) \rightarrow (X, \mu)$ such that the induced map between equivalence relations*

$$\Theta \times \Theta : (\mathcal{R}_{\mathbb{S}^1} \times \mathcal{R}_\Gamma, L \times L \times \tilde{\nu}) \rightarrow (\mathcal{R}_G, \Lambda \times \mu)$$

is measure-preserving, where $\tilde{\nu}$ is the σ -finite measure induced by ν on \mathcal{R}_Γ via integration of the counting measure of the fibers.

- (iii) *The group G is amenable if and only if the orbit equivalence relation induced by Γ on (Y, ν) is amenable.*

Proof. In this proof we will use the notation and conventions of Proposition 4.3 of [KPV15]. Let $Y \subset X$ be a cross section and let $U \subset G$ be a neighborhood of the identity as in Definition 1.14. We consider the restriction of \mathcal{R}_G to Y ,

$$\mathcal{R} := \{(y, y') \in Y \times Y : \exists g \in G, y' = gy\}.$$

By [KPV15, Proposition 4.3.1], \mathcal{R} is a Borel, countable equivalence relation on Y . Define $\Psi : G \times X \rightarrow \mathcal{R}_G$ by $\Psi(g, x) = (gx, x)$ and observe that since the action is free, Ψ is a bijection. Denote by Ψ_Y the restriction of Ψ to $G \times Y$ and put $Z = \Psi_Y(G \times Y)$. Observe that the projection $\pi : Z \rightarrow X$ on the first coordinate is countable-to-one hence we can define a measure η on Z by integrating with respect to μ the counting measure over the projection π . By definition we have $\mu(UY) = \eta(\Psi_Y(U \times Y))$. Put $\text{covol}(Y) := \lambda(U)/\mu(UY)$. As explained in the proof of [KPV15, Proposition 4.3.2], the unicity of the Haar measure on G implies the existence of a probability measure ν on Y such that $\Psi_*(\lambda \times (\nu/\text{covol}(Y))) = \eta$.

²Actually, any infinite compact metrizable group will do; the point is that the orbit equivalence relation associated to $\mathbb{S}^1 \curvearrowright \mathbb{S}^1$ is transitive.

Moreover by [KPV15, Proposition 4.3], we know that

- (1) the probability measure ν is \mathcal{R} -invariant,
- (2) (\mathcal{R}, ν) is ergodic if and only if the action of G is ergodic,
- (3) (\mathcal{R}, ν) has infinite orbits almost everywhere if and only if G is non-compact,
- (4) (\mathcal{R}, ν) is amenable if and only if G is.

By property (3) above, we deduce that (Y, ν) is diffuse. Moreover since \mathcal{R} is countable and measure-preserving, Feldman and Moore's result ([FM77, Theorem 1]) gives us a measure-preserving action of a countable group Γ on (Y, ν) which induces the equivalence relation \mathcal{R} .

Up to taking an open subset of U , we may assume that $\mu(U \cdot Y) = \frac{1}{K}$ for some integer $K \in \mathbb{N}$. Set $A = U \cdot Y$. By ergodicity of G , we can find $T \in [\mathcal{R}_G]$ of order K such that $\{A, T(A), \dots, T^{K-1}(A)\}$ is a partition of a full measure subset of X . Let us denote by c the counting measure on $\mathbb{Z}/K\mathbb{Z}$ and consider the equivalence relation \mathcal{S}' on $(\mathbb{Z}/K\mathbb{Z} \times U \times Y, c \times \lambda_U \times \nu)$ defined by

$$(k, u, y) \mathcal{S}'(k', u', y') \text{ if } y \mathcal{R} y'.$$

The measure-preserving map

$$\begin{aligned} \Theta' : (\mathbb{Z}/K\mathbb{Z} \times U \times Y, c \times \lambda_U \times \nu) &\rightarrow (X, \mu) \\ \Theta'(k, u, y) &:= T^k(u \cdot y), \end{aligned}$$

defines an orbit equivalence between \mathcal{S}' and \mathcal{R}_G . Denote by L the normalized Lebesgue measure on the circle \mathbb{S}^1 and fix a measure-preserving isomorphism

$$\alpha : (\mathbb{S}^1, L) \rightarrow (\mathbb{Z}/K\mathbb{Z} \times U, c \times \lambda_U).$$

Let \mathcal{S} be the equivalence relation induced by the action of $\Gamma \times \mathbb{S}^1$ on $Y \times \mathbb{S}^1$ where Γ acts on Γ and \mathbb{S}^1 acts on itself by translation. Observe that α induces an orbit equivalence between \mathcal{S}' and \mathcal{S} , which combined with Θ' gives an orbit equivalence Θ between \mathcal{S} and \mathcal{R}_G and hence (i) is proved.

Now (ii) can be deduced by an easy computation; it is also a direct application of the uniqueness of the Haar measure on \mathcal{R}_G (see Theorem A.13). Condition (iii) follows from property (4). \square

1.4. Weak orbit equivalence versus orbit equivalence. Let \mathcal{R} be a Borel equivalent relation on X and $A \subseteq X$ be a Borel subset with positive measure. The **restriction** of \mathcal{R} to A is the equivalence relation $\mathcal{R} \cap (A \times A)$ on the standard probability space (A, μ_A) where the measure μ_A is defined by: for all Borel $B \subseteq A$, $\mu_A(B) = \frac{\mu(B)}{\mu(A)}$. Let us recall two important definitions.

Definition 1.16. Let Γ and Λ be two countable groups.

- The groups Γ and Λ are **measure equivalent** if there exists a standard σ -finite measured space (Ω, m) and commuting measure-preserving actions of Γ and Λ on (Ω, m) which are essentially free and admit a fundamental domain with finite measure.
- The groups Γ and Λ are **weakly orbit equivalent** if Γ and Λ admit measure-preserving essentially free ergodic actions on probability spaces (X, μ) and (Y, ν) such that there exist measurable subsets $A \subset X$ and $B \subset Y$ such that \mathcal{R}_Γ restricted to A is orbit equivalent to \mathcal{R}_Λ restricted to B . The quantity $\mu(A)/\nu(B)$ is called the *coupling constant* of the weak orbit equivalence.

Remark 1.17. An application of the ergodic decomposition theorem yields that one can drop the ergodicity assumption in the definition of weak orbit equivalence.

Furman proved in [Fur99b] that two countable groups are measure equivalent if and only if they are weakly orbit equivalent. Let us now study these notions for non-discrete locally compact groups.

Definition 1.18. Let G and H be two Polish locally compact groups.

- The groups G and H are **orbit equivalent** if they admit ergodic measure-preserving essentially free actions which are orbit equivalent.
- The groups G and H are **weakly orbit equivalent** if they admit ergodic measure-preserving essentially free actions on (X, μ) such that there exist positive measurable subsets $A \subset X$ and $B \subset X$ such that \mathcal{R}_G restricted to A is orbit equivalent to \mathcal{R}_H restricted to B .

Note that a non-discrete locally compact group is never weakly orbit equivalent to a discrete one.

Lemma 1.19. *Let G be a non-discrete locally compact Polish group acting essentially freely ergodically in a measure-preserving manner on (X, μ) . Then for every Borel subset $A \subseteq X$ of positive measure, \mathcal{R}_G is orbit equivalent to its restriction to A .*

Proof. By Theorem 1.15, we can find a standard probability space (Y, ν) and a countable group Γ acting on (Y, ν) by measure-preserving transformations such that the action $\mathbb{S}^1 \times \Gamma$ on $(\mathbb{S}^1 \times Y, L \times \mu)$ is orbit equivalent to \mathcal{R}_G . It thus suffices to show that for every measurable $A \subseteq \mathbb{S}^1 \times Y$ of positive measure, $\mathcal{R}_{\mathbb{S}^1 \times \Gamma} = \mathcal{R}_{\mathbb{S}^1} \times \mathcal{R}_\Gamma$ is orbit equivalent to its restriction to (A, μ_A) .

So let A be a subset of $\mathbb{S}^1 \times Y$ of positive measure. Let \tilde{A} be a Borel subset of \mathbb{S}^1 with measure $\mu(A)$. Then since $\mathcal{R}_{\mathbb{S}^1}$ is transitive it is orbit equivalent to its restriction to \tilde{A} , which in turn yields that $\mathcal{R}_{\mathbb{S}^1} \times \mathcal{R}_\Gamma$ is orbit equivalent to its restriction to $\tilde{A} \times Y$.

Since the G -action is ergodic, there exists $\varphi \in [\mathcal{R}_{\mathbb{S}^1 \times \Gamma}]$ which maps a full measure subset of $\tilde{A} \times Y$ to a full measure subset of A (see Proposition 1.9), so that the restrictions of $\mathcal{R}_{\mathbb{S}^1 \times \Gamma}$ to A and $\tilde{A} \times Y$ are orbit equivalent. We conclude that $\mathcal{R}_{\mathbb{S}^1 \times \Gamma}$ is orbit equivalent to its restriction to A , hence the same conclusion holds for \mathcal{R}_G . \square

Theorem 1.20. *Let G and H be two non-discrete locally compact Polish groups. Then G and H are orbit equivalent if and only if they are weakly orbit equivalent.*

Proof. The direct implication is by definition and the converse is a straightforward application of the previous lemma. \square

Remark 1.21. Any reasonable definition of cost for equivalence relations coming from measure-preserving actions of non-discrete locally compact groups would have to satisfy an induction formula for restrictions to subsets. So by the previous theorem it could not be invariant under orbit equivalence, but only under *Haar-measure-preserving* orbit equivalence (see Appendix A for a precise definition of the Haar measure in our setup).

Cost for locally compact groups would nevertheless provide three distinct orbit equivalence classes as it does for countable groups up to weak orbit equivalence: cost 1, finite cost greater than 1, and infinite cost.

This is in complete analogy with the recent theory of L^2 Betti numbers for locally compact groups; see [Pet13] and [KPV15]. In order to define the L^2 Betti numbers, one has to fix a Haar measure and then depend on it linearly. However when there are at least two L^2 Betti numbers finite and non-zero, the ratios do not depend on the choice of the Haar measure.

The following proposition is surely well known to experts, but we were not able to find it in the literature. It guarantees that orbit equivalence for locally compact non-discrete groups is at least as complicated as measure equivalence for countable groups.

Proposition 1.22. *Two countable groups Γ and Λ are measure equivalent if and only if the locally compact groups $\Gamma \times \mathbb{S}^1$ and $\Lambda \times \mathbb{S}^1$ are orbit equivalent.*

Proof. Suppose that $\Gamma \times \mathbb{S}^1$ admits an action on the probability space (X, μ) which is orbit equivalent to an action of $\Lambda \times \mathbb{S}^1$. Then we can let Γ act on $\mathcal{R}_{\Gamma \times \mathbb{S}^1}$ on the left and we can let Λ act on the right via the orbit equivalence. These two actions commute and in both cases a fundamental domain is given by $X \times \mathbb{S}^1$ which has finite measure.

Suppose now that the groups Γ and Λ are measure equivalent; then by [Fur99a, Lem. 2.2.2] and [Fur99b, Lem. 3.2] they are weakly orbit equivalent. Then $\Gamma \times \mathbb{S}^1$ and $\Lambda \times \mathbb{S}^1$ are weakly orbit equivalent, hence orbit equivalent by Theorem 1.20. \square

Note that the groups $\Gamma \times \mathbb{S}^1$ and $\Lambda \times \mathbb{S}^1$ are unimodular, so Theorem A.13 applies. Therefore any orbit equivalence between them sends the Lebesgue measure of the circle group to a multiple of the Lebesgue measure of the other circle. This constant is the coupling constant of the induced measure equivalence between Γ and Λ .

Corollary 1.23. *There are uncountably many non-discrete locally compact groups up to orbit equivalence.*

Proof. This follows from the fact that there are uncountably many countable groups up to measure equivalence (see the paragraph preceding **P_{ME}15*** in [Gab05]). \square

2. DENSE SUBGROUPS IN ORBIT FULL GROUPS

The aim of this section is to prove the following theorem.

Theorem 2.1. *Let G be a locally compact Polish group. For every measure-preserving action of G on the probability space (X, μ) and for every dense subgroup $H \subset G$, we have that $[\mathcal{R}_H]$ is dense in $[\mathcal{R}_G]$.*

Theorem 2.1 will be a crucial tool to compute the topological rank³ of an orbit full group (see the next section).

2.1. Suitable actions. We will prove Theorem 2.1 under a weaker hypothesis in the context of Polish group actions. Recall however that Kolmogorov's example ([Dan00, Example 9]) shows that Theorem 2.1 does not generalize to arbitrary measure-preserving actions of Polish groups. Indeed there is a Borel probability measure on $\{0, 1\}^{\mathbb{N}}$ such that the full group generated by the finitely supported permutations is not dense in the orbit full group of the Polish group of all permutations of \mathbb{N} acting by shift on $\{0, 1\}^{\mathbb{N}}$. The fact is that this action is not *suitable*.

³Recall that the topological rank of a topological group is the minimum of the rank of a countable dense subgroup.

Definition 2.2 (Becker, [Bec13, Definition 1.2.7]). Let G be a Polish group. A Borel measure-preserving action of G on the probability space (X, μ) is **suitable** if for all Borel subsets $A, B \subset X$ of positive measure, one of the following two conditions holds:

- (1) for any open neighborhood of the identity $O \subset G$, there is $g \in O$ such that $\mu(A \cap gB) > 0$;
- (2) there are Borel subsets $A' \subset A$ and $B' \subset B$ of full measure in A and B and an open neighborhood O of the identity in G such that $(OA') \cap B' = \emptyset$.

We will prove the following.

Theorem 2.3. *Let G be a Polish group. For every Borel, measure-preserving, suitable action of G on the probability space (X, μ) and for every dense subgroup $H \subset G$, the orbit full group $[\mathcal{R}_H]$ is dense in $[\mathcal{R}_G]$.*

Becker proved in Theorem 1.2.9 of [Bec13], that all measure-preserving actions of locally compact Polish groups are suitable, so Theorem 2.3 implies Theorem 2.1. We end this section by giving a different example to which our results apply.

Example 2.4. The standard Bernoulli shift $\mathfrak{S}_\infty \curvearrowright ([0, 1]^\mathbb{N}, \lambda^{\otimes \mathbb{N}})$ is a suitable action, where λ is the Lebesgue measure.

Proof. Set $X := [0, 1]^\mathbb{N}$ and $\mu := \lambda^{\otimes \mathbb{N}}$ and let $A, B \subseteq X$ be Borel subsets of positive measure. Suppose that (1) does not hold, and let O be a neighborhood of the identity such that

$$\text{for all } g \in O, \mu(A \cap gB) = 0.$$

By shrinking O if necessary, we may assume that there exists $N \in \mathbb{N}$ such that O is the subgroup of \mathfrak{S}_∞ consisting of all the permutations which fix pointwise the set $\{0, \dots, N-1\}$. We endow $Y := [0, 1]^{\{0, \dots, N-1\}}$ with the measure $\eta := \lambda^{\otimes N}$ and $Z := [0, 1]^{\mathbb{N} \setminus \{0, \dots, N-1\}}$ with the measure $\nu := \lambda^{\mathbb{N} \setminus \{0, \dots, N-1\}}$. Observe that $(X, \mu) = (Y, \eta) \times (Z, \nu)$.

Let $\tau \in O$ be a bijection whose set of fixed points is $\{0, \dots, N-1\}$, and which has only one non-trivial orbit. Then its Bernoulli shift on (Z, ν) is ergodic since it is conjugate to the \mathbb{Z} -shift on $([0, 1]^\mathbb{Z}, \lambda^{\otimes \mathbb{Z}})$. Moreover the ergodic decomposition of its Bernoulli shift on (X, μ) is given by $(\mu_y)_{y \in Y}$ where μ_y is the probability measure on X defined by $\mu_y := \delta_y \otimes \nu$.

Since $\mu(A \cap \tau^k B) = 0$ for all $k \in \mathbb{Z}$ and every μ_y is τ -ergodic, there is a full measure Borel subset Y' of Y such that for all $y \in Y'$ we have that $\mu_y(A) = 0$ whenever $\mu_y(B) > 0$. Let Y_0 be the Borel set of $y \in Y'$ such that $\mu_y(B) > 0$, and put $Y_1 := Y' \setminus Y_0$. The sets $A' := (Y_1 \times Z) \cap A$ and $B' := (Y_0 \times Z) \cap B$ witness that (2) holds. \square

Remark 2.5. In the above example, since the countable group $\mathfrak{S}_{(\infty)}$ of finitely supported permutations is amenable, the full group it generates is extremely amenable for the uniform topology by [GP07, Thm. 5.7]. Since the uniform topology refines the topology of convergence in measure and $\mathfrak{S}_{(\infty)}$ is dense in \mathfrak{S}_∞ , Theorem 2.3 yields that $[\mathcal{R}_{\mathfrak{S}_\infty}]$ is extremely amenable. It would be interesting to understand for

which non-locally compact closed subgroups $G \leq \mathfrak{S}_\infty$ the orbit full group associated to the standard Bernoulli shift is extremely amenable.

2.2. An equivalent statement. From now on, we will use the notation of Section 1.2. For every Borel, measure-preserving action of G on the probability space (X, μ) , we denote by $[\widetilde{G}]_D \subset [\widetilde{\mathcal{R}_G}]$ the subset of functions with countable (essential) image and we put $[G]_D := \Phi([\widetilde{G}]_D)$. Note that $[G]_D$ is the smallest full group containing the image of G inside $\text{Aut}(X, \mu)$.

Theorem 2.3 follows from the following weaker theorem, which is also important in its own right.

Theorem 2.6. *Let G be a Polish group. For every Borel, measure-preserving, suitable action of G on the probability space (X, μ) , we have that $[\widetilde{G}]_D \subset [\widetilde{\mathcal{R}_G}]$ is a dense subgroup.*

Before deducing Theorem 2.3 from the above result, we need the following lemma which will be used several times.

Lemma 2.7. *Let $A \subseteq X$ and let $f : A \rightarrow G$ be a function with countable image. If the map $\Phi(f) : A \rightarrow X$ defined by $\Phi(f)(x) = f(x)x$ is injective, then there exists $f' \in [\widetilde{G}]_D$ which extends f .*

Proof. Let Γ be the group generated by the range of the function f . Since $\Phi(f)$ is injective, the map $\Phi(f)$ is an element of the pseudo-full group⁴ of Γ . The elements of the pseudo-full group of Γ preserve the \mathcal{R}_Γ -conditional measure and any two sets having the same \mathcal{R}_Γ -conditional measure can be sent to one another by an element of the pseudo-full group of Γ (see [LM15, Sec. 2.1] for details). Therefore there is an element $T \in [G]_D$ which extends $\Phi(f)$. By lifting such a T to $[\widetilde{G}]_D$ where f was not defined, we obtain $f' \in [\widetilde{G}]_D$ which extends f . \square

Proof of Theorem 2.3. Let G be a Polish group and let H be a dense subgroup. Consider a Borel, measure-preserving suitable action of G on the probability space (X, μ) . By Theorem 2.6, we only need to prove that $[\widetilde{H}]_D \subset [\widetilde{G}]_D$ is dense.

Fix a compatible, right-invariant metric d_G on G bounded by 1, fix $\varepsilon > 0$ and take $f \in [\widetilde{\mathcal{R}_G}]_D$. There are $k \in \mathbb{N}$, a finite subset $\{g_1, \dots, g_k\} \subset G$ and a finite partition $\{A_0, \dots, A_k\}$ of X such that $\mu(A_0) \leq \varepsilon/2$ and for every $i \geq 1$, we have $f(A_i) = \{g_i\}$. By density and weak-continuity of the action, there exists $\{h_1, \dots, h_k\} \subset H$ such that for every $i \in \{1, \dots, k\}$, we have that $d_G(g_i, h_i) \leq \varepsilon$ and $\mu(g_i(A_i)\Delta h_i(A_i)) \leq \varepsilon/2k$. Put

$$B_i := h_i^{-1}(g_i(A_i) \cap h_i(A_i)) \subset A_i, \quad B := \bigcup_{i=1}^k B_i,$$

and observe that $\mu(B) \geq 1 - \varepsilon$. Consider the map $f' : B \rightarrow H$ defined by $f'(x) = h_i$ whenever $x \in B_i$. Then f has finite range, and since the subsets $h_i B_i$ are disjoint, $\Phi(f)$ is injective. Lemma 2.7 allows us to extend f' to $f'' \in [\widetilde{G}]_D$. We clearly have $\tilde{d}_G(f, f'') \leq 2\varepsilon$, which ends the proof. \square

⁴The pseudo-full group of the countable group Γ acting on (X, μ) is defined to be the set of Borel partial maps whose graph is a subset of \mathcal{R}_Γ .

2.3. Proof of Theorem 2.6.

Definition 2.8. Fix $f \in \widetilde{[\mathcal{R}_G]}$ and a neighborhood of the identity $N \subset G$. We say that a couple (A, g) is (N) -good if

- (1) $A \subset X$ is a measurable subset of positive measure and $g : A \rightarrow G$ is a measurable function with countable image,
- (2) for every $x \in A$, we have $f(x)g(x)^{-1} \in N$,
- (3) the map $\Phi(g) : A \rightarrow X$ defined by $\Phi(g)(x) = g(x)x$ is injective.

We note that for a fixed $f \in \widetilde{[\mathcal{R}_G]}$ the existence of a good couple is not a trivial fact. Indeed, we will use the hypothesis that the action is suitable only to show the existence of such couples.

The proof of the theorem will be a measurable version of the Hall's Marriage Theorem and it will follow the same strategy as Hudson's in [Hud93]. For a fixed f as in Definition 2.8, using Zorn's Lemma, we will construct for every $\varepsilon > 0$ and neighborhood of the identity $N \subset G$ a good couple such that $\mu(A) > 1 - \varepsilon$ in three steps.

Step 1. In the first step (and only in this one), we will use the hypothesis that the action is suitable.

Proposition 2.9. Let $f \in \widetilde{[\mathcal{R}_G]}$ and let $N \subset G$ be a neighborhood of the identity. For every $B \subset X$ of positive measure, there is a good couple (A, g) such that $A \subset B$ has positive measure and $\Phi(g)(A) \subset \Phi(f)(B)$.

Proof. Consider a neighborhood of the identity $O \subset G$ such that $O = O^{-1}$ and $O^2 \subset N$. Let $f(x_0)$ be an element of the support of the pushforward measure $f_*\mu|_B$ and put $A_O := B \cap f^{-1}(Of(x_0))$. For every neighborhood of the identity O' in G , set $C_{O'} := B \cap f^{-1}(O'f(x_0))$. Note that $C_{O'} \subset A_O$, whenever $O' \subset O$. By definition of the support of $f_*\mu|_B$ the Borel set $C_{O'}$ has positive measure.

Let us show that condition (2) of Definition 2.2 is not satisfied for the two Borel sets $\Phi(f)(A_O)$ and $f(x_0)A_O$. Indeed, $\Phi(f)^{-1}$ and $f(x_0)^{-1}$ are measure-preserving so if condition (2) holds, then there is a full measure subset $A' \subseteq A_O$ such that $\Phi(f)(A')$ and $O'f(x_0)A'$ are disjoint. This is a contradiction because $\Phi(f)(A')$ and $O'f(x_0)A'$ contain $\Phi(f)(A' \cap C_{O'})$ which has positive measure.

Since the action is suitable, (1) of Definition 2.2 has to hold. So there is $h \in O$ such that

$$\mu(\Phi(f)(A_O) \cap hf(x_0)A_O) > 0.$$

Set $A := A_O \cap f(x_0)^{-1}h^{-1}\Phi(f)(A_O)$ and for every $x \in A$ put $g(x) := hf(x_0)$. The couple (A, g) is good, because for every $x \in A$ we have that $f(x)f(x_0)^{-1} \in O$ and

$$f(x)g(x)^{-1} = f(x)f(x_0)^{-1}h \in O^2 \subset N. \quad \square$$

Step 2. For a neighborhood N of the identity in G and $\varepsilon > 0$, we now define the order on the family of N -good couples associated to a function $f \in \widetilde{[\mathcal{R}_G]}$.

Definition 2.10. Let (A_1, g_1) and (A_2, g_2) be two good couples. We say that $(A_1, g_1) \prec (A_2, g_2)$ if $A_2 \supseteq A_1$ almost everywhere and if

$$\mu(\{x \in A_1 : g_1(x) \neq g_2(x)\}) \leq \frac{1}{\varepsilon}(\mu(A_2) - \mu(A_1)).$$

The factor ε^{-1} is the scaling constant which appears in the end of the proof of Proposition 2.12. We identify two good couples (A_1, g_1) and (A_2, g_2) if $\mu(A_1 \triangle A_2) = 0$ and for almost all $x \in A_1 \cap A_2$, $g_1(x) = g_2(x)$. This makes the relation \prec antisymmetric. It is moreover clearly reflexive.

Lemma 2.11. *The relation \prec is an order relation on the set of good couples.*

Proof. The only fact left to prove is that \prec is transitive. For this suppose that

$$(A_1, g_1) \prec (A_2, g_2) \prec (A_3, g_3);$$

then

$$\{x \in A_1 : g_1(x) \neq g_3(x)\} \subset \{x \in A_1 : g_1(x) \neq g_2(x)\} \cup \{x \in A_2 : g_2(x) \neq g_3(x)\},$$

so we get

$$\begin{aligned} \mu(\{x \in A_1 : g_1(x) \neq g_3(x)\}) &\leq \mu(\{x \in A_1 : g_1(x) \neq g_2(x)\}) + \mu(\{x \in A_2 : g_2(x) \neq g_3(x)\}) \\ &\leq \frac{1}{\varepsilon}(\mu(A_2) - \mu(A_1)) + \frac{1}{\varepsilon}(\mu(A_3) - \mu(A_2)) \\ &= \frac{1}{\varepsilon}(\mu(A_3) - \mu(A_1)). \end{aligned} \quad \square$$

The following proposition is the core of the proof of Theorem 2.6.

Proposition 2.12. *For every good couple (A, g) with $\mu(A) < 1 - \varepsilon$, there exists a good couple (A', g') such that $(A, g) \prec (A', g')$ and $\mu(A' \setminus A) > 0$.*

We would like to say that for every good couple (A, g) there is $B \subset X \setminus A$ such that $\Phi(f)(B) \cap \Phi(g)(A) = \emptyset$. When this is the case, we can conclude using Proposition 2.9. The problem is that this is not always possible, but it is possible in a finite number of steps.




FIGURE 1. In the figure $\Phi(f)$ acts vertically and $\Phi(g)$ acts diagonally. Since $\Phi(f)(X \setminus A) \subset \Phi(g)(A)$, we cannot use Proposition 2.9 directly.

Proof. We first show that there exists $k \in \mathbb{N}$ with $k \leq 1/\varepsilon$ and there are two sequences $\{D_i\}_{i \leq k}$ and $\{E_i\}_{i \leq k}$ of measurable subsets of X of positive measure such that (see Figure 1)

- (1) the $\{D_i\}_{i \leq k}$ are pairwise disjoint as are the $\{E_i\}_{i \leq k}$,
- (2) $D_1 \subset X \setminus A$ and $D_i \subset A$ for $i > 1$,
- (3) $E_k \subset X \setminus \Phi(g)(A)$ and $E_i \subset \Phi(g)(A)$ for $i < k$,
- (4) $\Phi(f)(D_i) = E_i$ and $E_{i-1} = \Phi(g)(D_i)$ for all $i \in \{2, \dots, k\}$.

Set $B_1 := X \setminus A$ and $C_1 := \Phi(f)(B_1)$. For $i \geq 2$ define recursively

$$B_i := \Phi(g)^{-1}(C_{i-1} \cap \Phi(g)(A)) \quad \text{and} \quad C_i := \Phi(f)(B_i).$$

Observe that $\{B_i\}_i$ are pairwise disjoint as are the $\{C_i\}_i$. Suppose now that for $l \geq 1$, we have that $C_i \subset \Phi(g)(A)$ for all $i \leq l$. Since $\Phi(g)$ and $\Phi(f)$ preserve the measure, we have that $\mu(C_i) = \mu(B_1)$ for all $i \leq l$ and hence we have that $l\mu(B_1) \leq 1 - \mu(B_1)$. By hypothesis $\mu(B_1) \geq \varepsilon$, so $l \leq 1/\varepsilon - 1$. Therefore there exists $k \leq 1/\varepsilon$, such that C_k is not contained in $\Phi(g)(A)$ and $C_i \subset \Phi(g)(A)$ for every $i < k$. Put $E_k := C_k \setminus \Phi(g)(A)$ and set $D_k := \Phi(f)^{-1}(E_k)$. Observe that $D_k \subset B_k$ and define recursively $E_i := \Phi(g)(D_{i+1})$ and $D_i := \Phi(f)^{-1}(E_i)$.

Now we will use Proposition 2.9 to extend g to a subset B_1 of D_1 . If $k > 1$, then $E_1 = \Phi(f)(D_1) \subset \Phi(g)(A)$ and so we cannot directly apply Proposition 2.9. Instead, we will modify g in a subset B_i of each D_i (with values in E_i) for $i > 1$ to obtain a larger good couple (A', g') with $A' = A \cup B_1$ and $\Phi(g)(A' \setminus B_k) = \Phi(g)(A)$ and $\Phi(g')(B_k) \subset E_k \subset X \setminus \Phi(g)(A)$.

By Proposition 2.9, there exists a good couple (A_1, g_1) such that $A_1 \subset D_1$ and $\Phi(g_1)(A_1) \subset \Phi(f)(A_1) \subset E_1$. For $i \in \{2, \dots, k\}$, whenever A_{i-1} is defined, we set

$$A'_i := \Phi(g)^{-1}(\Phi(g_{i-1})(A_{i-1})) \subset \Phi(g)^{-1}(E_{i-1}) = D_i.$$

For every i such that A'_i is defined, Proposition 2.9 implies that there is a good couple (A_i, g_i) such that $A_i \subset A'_i$ is non-negligible and $\Phi(g_i)(A_i) \subset \Phi(f)(A'_i) \subset E_i$. Put $B_k := A_k$. For $i \in \{1, \dots, k-1\}$, we define recursively $B_i := \Phi(g_i)^{-1}(\Phi(g)(B_{i+1}))$.

Set $A' := A \cup B_1$ and define

$$g'(x) := \begin{cases} g(x) & \text{if } x \in A \setminus \bigcup_{i \geq 2} B_i, \\ g_i(x) & \text{if } x \in B_i. \end{cases}$$

By construction, $\Phi(g') : A' \rightarrow X$ is injective and preserves the measure. Moreover (A', g') is obtained by *cutting and pasting* N -good couples, so it is an N -good couple. Let us finally check that $(A, g) \prec (A', g')$. Clearly we have $A' \supset A$ and $\mu(A' \setminus A) = \mu(B_1) > 0$. Moreover

$$\mu(\{x \in A : g(x) \neq g'(x)\}) \leq \mu\left(\bigcup_{i \geq 2} B_i\right) \leq k\mu(B_1) \leq \frac{1}{\varepsilon}(\mu(A') - \mu(A)). \quad \square$$

Step 3. We verify now that we can apply Zorn's Lemma to the set of good couples.

Proposition 2.13. *Every chain for \prec has an upper bound.*

Proof. Let us assume for the moment that $\{(A_n, g_n)\}_n$ is a countable chain of good couples. For every $n \in \mathbb{N}$ set

$$B_n := \{x \in A_n : g_n(x) \neq g_{n+1}(x)\}, \quad C_n := \bigcap_{k \geq n} B_k, \quad \text{and} \quad A := \bigcup_n C_n.$$

Clearly $A \subset \bigcup_n A_n$ and we now check that the two measurable subsets have the same measure. In fact, since $\{A_n\}_n$ and $\{C_n\}_n$ are increasing sequences, for every $\eta > 0$, there is $K \in \mathbb{N}$ such that

$$\mu\left(\bigcup_n A_n\right) - \mu(A_K) < \eta \quad \text{and} \quad \mu\left(\bigcup_n C_n\right) - \mu(C_K) < \eta,$$

hence we have

$$\begin{aligned} \mu\left(\bigcup_n A_n\right) - \mu(A) &\leq 2\eta + \mu(A_K) - \mu(C_K) = 2\eta + \mu(A_K \setminus C_K) \\ &= 2\eta + \mu(A_K \cap (\bigcup_{k \geq K} X \setminus B_k)) = 2\eta + \mu\left(\bigcup_{k \geq K} A_K \setminus B_k\right) \\ &\leq 2\eta + \sum_{k \geq K} \mu(A_k \setminus B_k) \leq 2\eta + \frac{1}{\varepsilon} \sum_{k \geq K} \mu(A_{k+1} \setminus A_k) \\ &\leq 2\eta + \frac{1}{\varepsilon} \mu\left(\bigcup_{k \geq K+1} A_k \setminus A_K\right) \leq 2\eta + \frac{\eta}{\varepsilon}. \end{aligned}$$

As η is arbitrarily small, we get that $A = \bigcup_n A_n$ almost everywhere. For $x \in C_n$, observe that $g_n(x) = g_{n+j}(x)$ for every $j \geq 0$. We define

$$g(x) := g_n(x) \quad \text{if } x \in C_n.$$

The couple (A, g) is obtained by cutting and pasting N -good couples so the couple is N -good. Moreover $A \supseteq \bigcup_n A_n$ almost everywhere and for every $n \in \mathbb{N}$, we have

$$\mu(x \in A_n : g_n(x) \neq g(x)) \leq \mu(A_n \setminus C_n) \leq \frac{1}{\varepsilon} \sum_{k \geq n} \mu(A_{k+1} - A_k) = \frac{1}{\varepsilon} (\mu(A) - \mu(A_n)).$$

Therefore the couple (A, g) is an upper bound for the countable chain. Consider now an arbitrary chain $\{(A_c, g_c)\}_{c \in C}$ and set $\lambda = \sup_{c \in C} \mu(A_c)$. If there is a good couple (A_c, g_c) such that $\mu(A_c) = \lambda$, then this couple is an upper bound of the chain and there is nothing to prove. Suppose that this is not the case and consider a subsequence $\{(A_n, g_n)\}_{n \in \mathbb{N}}$ of the chain such that $\lim_n \mu(A_n) = \lambda$. Let (A, g) be an upper bound for this sequence. Given any element of the chain (A_c, g_c) there exists n such that $\mu(A_c) \leq \mu(A_n)$ and hence $(A_c, g_c) \prec (A_n, g_n) \prec (A, \varphi)$. \square

End of the proof of Theorem 2.6. Let $f \in \widetilde{[\mathcal{R}_G]}$. By definition of the topology of convergence in measure, a base of neighborhoods of f is given by the open sets

$$\mathcal{U}_{\varepsilon, N} := \left\{ g \in \widetilde{[\mathcal{R}_G]} : \mu(\{x \in X : g(x) \in Nf(x)\}) > 1 - \varepsilon \right\},$$

where $\varepsilon > 0$ and $N \subset G$ is a neighborhood of the identity. For every neighborhood of the identity $N \subset G$, Proposition 2.9 implies that the set of good couples for f is not empty. For $\varepsilon > 0$, Proposition 2.13 tells us that there is a maximal good couple (A, g) . The maximality of the couple and Proposition 2.12 imply that $\mu(A) \geq 1 - \varepsilon$. So by Lemma 2.7 there is $g' \in \widetilde{[\mathcal{R}_G]}_D$ such that $g' \in \mathcal{U}_{\varepsilon, N}$.

3. TOPOLOGICAL RANK OF ORBIT FULL GROUPS

We now use Theorem 2.1 to show that the topological rank of orbit full groups associated to free measure-preserving actions of unimodular locally compact groups is equal to two.

Theorem 3.1. *Let G be a locally compact unimodular non-discrete and non-compact Polish group. For every measure-preserving, essentially free and ergodic action of G , there is a dense G_δ of couples (T, U) in $[\mathcal{R}_G]^2$ which generate a dense free subgroup of $[\mathcal{R}_G]$ acting freely. In particular, the topological rank of $[\mathcal{R}_G]$ is 2.*

Proof. Let G be a locally compact unimodular non-discrete and non-compact Polish group. Suppose that G acts on the probability space (X, μ) preserving the measure, essentially freely and ergodically. Let us denote by F_2 the free group on two generators. Since $[\mathcal{R}_G]$ is a Polish group, a standard argument shows that the set

$$\left\{ (T, U) \in [\mathcal{R}_G]^2 : \overline{\langle T, U \rangle} = [\mathcal{R}_G] \text{ and } \langle T, U \rangle \cong F_2 \right\}$$

is G_δ (see e.g. [Pra81, Pra83]) so we have only to prove that it is dense.

By Theorem 1.15, there exists a (not necessarily free) action of a countable group Γ on a measure space (Y, ν) such that \mathcal{R}_G is orbit equivalent to the product action of $\mathbb{S}^1 \times \Gamma$ on $\mathbb{S}^1 \times Y$. Fix a copy of \mathbb{Z} in \mathbb{S}^1 generated by an irrational rotation; then $\mathbb{Z} \times \Gamma$ is dense in $\mathbb{S}^1 \times \Gamma$. By Theorem 2.1, we have that $[\mathcal{R}_{\mathbb{Z} \times \Gamma}]$ is dense in $[\mathcal{R}_G]$.

The equivalence relation $\mathcal{R}_{\mathbb{Z} \times \Gamma}$ has cost 1, by Proposition VI.23 of [Gab00] (note that the proof only uses that Γ_1 acts freely). So we can apply Theorem 1.7 in [LM15] to get the existence of an aperiodic $T \in [\mathcal{R}_{\mathbb{Z} \times \Gamma}]$ such that

$$\left\{ U \in [\mathcal{R}_{\mathbb{Z} \times \Gamma}] : \overline{\langle T, U \rangle}^{d_u} = [\mathcal{R}_{\mathbb{Z} \times \Gamma}] \text{ and } \langle T, U \rangle \cong F_2 \right\} \subset [\mathcal{R}_{\mathbb{Z} \times \Gamma}]$$

is a dense subset of $[\mathcal{R}_{\mathbb{Z} \times \Gamma}]$ with respect to the uniform topology. This concludes the proof since by Theorem 4.4 of [CM16], the conjugacy class of T is dense in $[\mathcal{R}_G]$ for the topology of convergence in measure. \square

4. THE ORBIT FULL GROUP AS A UNITARY GROUP

In this section, we study the relationship between orbit full groups arising from measure-preserving free actions of locally compact groups and the associated von Neumann algebra. Throughout this section, G will be a locally compact second countable unimodular group which we equip with a left and right invariant Haar measure m .

Let us recall the crossed product construction. See the first chapter of [vD78] for more about this. Note however that our left von Neumann algebra is the right von Neumann algebra in Van Daele's book.

Definition 4.1. Let G be a locally compact Polish group. For a measure-preserving free action of G on the probability space (X, μ) , the **crossed product** $L^\infty(X, \mu) \rtimes G$ is the von Neumann algebra on $L^2(G \times X, m \times \mu)$ generated by

- the set of unitary operators $\{\lambda_h \times \kappa_h\}_{h \in G}$ where $h \mapsto \kappa_h$ is the Koopman representation of G on $L^2(X, \mu)$ and $h \mapsto \lambda_h$ is the left regular representation,
- the abelian algebra $L^\infty(X, \mu)$ which acts on functions $\xi \in L^2(G \times X, \mu)$ by multiplication: for all $f \in L^\infty(X, \mu)$, we let $f\xi(g, x) = f(x)\xi(g, x)$.

We will show that this von Neumann algebra is generated by the orbit full group $[\mathcal{R}_G]$, which can be seen as a unitary group as follows. Recall that since we assume that the action of G is essentially free, the full group $[\mathcal{R}_G]$ is isomorphic as a Polish group to $\widetilde{[\mathcal{R}_G]}$, as explained in Section 1.2.

Definition 4.2. Consider the near-action of the full group $\widetilde{[\mathcal{R}_G]}$ on $(G \times X, m \times \mu)$ given by

$$g \cdot (h, x) = (g(x)h, g(x)x) \quad \text{for all } g \in \widetilde{[\mathcal{R}_G]}, h \in G, x \in X$$

and denote by π the associated Koopman representation on $L^2(G \times X)$. That is, for every $f \in L^2(G \times X)$ and $g \in \widetilde{[\mathcal{R}_G]}$, we have

$$\pi(g) \cdot f(h, x) = f(g(x)^{-1}h, g(x)^{-1}x).$$

Proposition 4.3. Let G be a unimodular non-compact locally compact Polish group acting freely on (X, μ) .

- (1) The map π is a continuous embedding of $[\mathcal{R}_G]$ into $\mathcal{U}(M)$.
- (2) The full group of \mathcal{R}_G consists of the intersection of $\text{Aut}(G \times X, m \times \mu)$ with $\mathcal{U}(G \ltimes L^\infty(X))$, seeing both as subgroups of $\mathcal{U}(L^2(G \times X, m \times \mu))$. In particular, it is a closed subgroup of $\mathcal{U}(L^2(G \times X, m \times \mu))$.
- (3) The full group generates the von Neumann algebra, that is, $\pi([\mathcal{R}_G])'' = L^\infty(X, \mu) \rtimes G$.

Proof. (1) First, observe that the action of $\widetilde{[\mathcal{R}_G]}$ on $G \times X$ is measure-preserving, so that π is a unitary representation.

Let us now see why $\pi(G) \subseteq \mathcal{U}(M)$. For this, note that the commutant of $M = L^\infty(X) \rtimes G$ is generated by the operators \tilde{f} for $f \in L^\infty(X)' = L^\infty(X)$ (acting by $(\tilde{f}\xi)(h, x) = f(h^{-1}x)\xi(h, x)$) and the operators $1 \times \rho_g$ where ρ_g is the right regular representation; see [vD78, Thm. 3.12]. For $g \in [\mathcal{R}_G]$, we have

$$\begin{aligned} \pi(g)\tilde{f}\xi(h, x) &= (\tilde{f}\xi)(g(x)^{-1}h, g(x)^{-1}x) \\ &= f(h^{-1}x)\xi(g(x)^{-1}h, g(x)^{-1}x) \\ &= \tilde{f}\pi(g)\xi(h, x) \end{aligned}$$

and we also have for $h' \in G$

$$\begin{aligned} \pi(g)\rho_{h'}\xi(h, x) &= (\rho_{h'}\xi)(g(x)^{-1}h, g(x)^{-1}x) \\ &= \xi(g(x)^{-1}hh', g(x)^{-1}x) \\ &= \rho_{h'}\pi(g)\xi(h, x), \end{aligned}$$

which concludes the proof.

(2) We will prove that every $T \in \text{Aut}(G \times X, m \times \mu)$ which commutes with the operators \tilde{f} and ρ_h is in the image of the full group $[\mathcal{R}_G]$.

Fix such a T and define maps t_1, t_2 by the equation $T(g, x) = (t_1(g, x)g, t_2(g, x))$. Since T commutes with ρ_h , we obtain that for almost all $(g, x) \in G \times X$,

$$(t_1(gh, x)gh, t_2(gh, x)) = (t_1(g, x)gh, t_2(g, x)).$$

In particular by Fubini's Theorem for almost all $x \in X$ and almost all $(g, h) \in G^2$ we have $t_1(gh, x) = t_1(g, x)$ so that $t_1(\cdot, x)$ is equal to a constant $g(x)$ up to a set of measure zero.

The above equation also yields $t_2(gh, x) = t_2(g, x)$ for almost all $(g, x) \in G \times X$ so again by Fubini's Theorem $t_2(\cdot, x)$ is equal to a constant $t(x)$ up to a set of measure zero.

So for almost all $g \in G$ and $x \in X$, one has $T(g, x) = (g(x)g, t(x))$. Now we observe that since T preserves the measure, $t : X \rightarrow X$ also does: if A is a Borel subset of G such that $m(A) = 1$ and $B \subseteq X$ is any Borel set, then

$$\begin{aligned} \mu(B) &= m \times \mu(A \times B) \\ &= m \times \mu(T^{-1}(A \times B)) \\ &= m \times \mu(\{(h, x) : g(x)h \in A \text{ and } t(x) \in B\}) \\ &= m \times \mu(\{(h, x) : h \in g(x)^{-1}A \text{ and } x \in t^{-1}(B)\}) \\ &= \int_X m(g(x)^{-1}A) \chi_{t^{-1}(B)}(x) d\mu(x) \\ &= m(A) \int_X \chi_{t^{-1}(B)}(x) d\mu(x) \\ &= \mu(t^{-1}(B)). \end{aligned}$$

Finally, we exploit the hypothesis that T commutes with the operators \tilde{f} for all $f \in L^\infty(X)$ which means

$$f(h^{-1}g(x)^{-1}t(x))\xi(g(x)h, t(x)) = f(h^{-1}x)\xi(g(x)h, t(x))$$

for almost all $(h, x) \in G \times X$. Since this is true for every f and ξ , we must have that $g(x)x = t(x)$ almost surely and hence T is in the image of the full group $[\mathcal{R}_G]$.

(3) Since $\lambda \times \kappa(G)$ is already a subgroup of $\pi([\mathcal{R}_G])$, by definition of the crossed product it suffices to show that $\pi([\mathcal{R}_G])''$ contains $L^\infty(X)$. For this, it is enough to show that for every $A \subset X$ the multiplication operator $\tilde{\chi}_A$ belongs to $\pi([\mathcal{R}_G])''$.

By Theorem 1.15, we may assume that $X = \mathbb{S}_1 \times Y$, and that $\mathcal{R}_G = \mathcal{R}_{\mathbb{S}_1 \times \Gamma}$, where $\mathbb{S}^1 \times \Gamma$ acts via a product action. Since G is non-compact, Γ has infinite orbits, but recall that the action is not necessarily free.

Let \mathcal{R}_Γ be the equivalence relation of the action of Γ on Y and $\tilde{\mathcal{R}}_\Gamma$ the equivalence relation of the action of Γ on $\mathbb{S}^1 \times Y$ obtained by making Γ act trivially on \mathbb{S}^1 . Observe that $\tilde{\mathcal{R}}_\Gamma = \mathcal{R}_\Gamma \times \mathbb{S}^1$ as measure spaces and by Theorem 1.15(ii), we have a measure-preserving isomorphism between $(G \times X, m \times \mu)$ and $(\mathcal{R}_\Gamma \times \mathbb{S}^1 \times \mathbb{S}^1, \tilde{\nu} \times L \times L)$.

By a well-known result of Dye (see e.g. [Kec10, Thm. 3.5]), we can choose an aperiodic element $T \in [\tilde{\mathcal{R}}_\Gamma] \subset [\mathcal{R}_{\Gamma \times \mathbb{S}^1}]$. Now let $T_{X \setminus A}$ be the first return map induced by T on $X \setminus A$, which belongs to $[\tilde{\mathcal{R}}_\Gamma]$.

Claim. *The sequence $(T_{X \setminus A}^n)_{n \in \mathbb{N}}$ tends to $\tilde{\chi}_A$ weakly as operators on the Hilbert space $L^2(\tilde{\mathcal{R}}_\Gamma)$.*

Proof of Claim. We denote by $[[\tilde{\mathcal{R}}_\Gamma]]$ the pseudo-full group of the action of Γ , that is, the pseudo-group of partial measure-preserving isomorphisms of X whose graph is contained in $\tilde{\mathcal{R}}_\Gamma$. For $\varphi \in [[\tilde{\mathcal{R}}_\Gamma]]$, we denote by $\text{dom}(\varphi) \subset X$ the domain of φ , that is, the Borel subset of X on which φ is defined. Given $\varphi \in [[\tilde{\mathcal{R}}_\Gamma]]$, we let $\chi_\varphi \in L^2(\tilde{\mathcal{R}}_\Gamma)$ be the characteristic function of the graph of φ . Note that $\tilde{\chi}_A \chi_\varphi = \chi_{\varphi|_A}$ where $\varphi|_A$ is the restriction of φ to $A \cap \text{dom}(\varphi)$.

Since the Hilbert space $L^2(\tilde{\mathcal{R}}_\Gamma)$ is spanned by the functions χ_φ where $\varphi \in [[\tilde{\mathcal{R}}_\Gamma]]$, we need to show that if $\varphi, \psi \in [[\tilde{\mathcal{R}}_\Gamma]]$, then

$$\left\langle T_{X \setminus A}^n \chi_\varphi, \chi_\psi \right\rangle \rightarrow \langle \tilde{\chi}_A \chi_\varphi, \chi_\psi \rangle.$$

By definition, this is equivalent to

$$\mu(\{x \in D : T_{X \setminus A}^n \varphi(x) = \psi(x)\}) \rightarrow \mu(\{x \in D \cap A : \varphi(x) = \psi(x)\}),$$

where $D := \text{dom}(\varphi) \cap \text{dom}(\psi)$.

So take φ, ψ in $[[\tilde{\mathcal{R}}_\Gamma]]$. Since the restriction of $T_{X \setminus A}$ to $X \setminus A$ is aperiodic, for all $x \in D \cap X \setminus A$ there is at most one $n \in \mathbb{N}$ such that $T_{X \setminus A}^n \varphi(x) = \psi(x)$. So the sets

$$B_n := \{x \in D \cap X \setminus A : T_{X \setminus A}^n \varphi(x) = \psi(x)\}$$

are all disjoint, hence their measure tends to 0. Moreover since $T_{X \setminus A}(x) = x$ for all $x \in A$, we have

$$\{x \in D \cap A : T_{X \setminus A}^n \varphi(x) = \psi(x)\} = \{x \in D \cap A : \varphi(x) = \psi(x)\}.$$

We conclude that

$$\mu(\{x \in D : T_{X \setminus A}^n \varphi(x) = \psi(x)\}) \rightarrow \mu(\{x \in D \cap A : \varphi(x) = \psi(x)\})$$

as desired. \square

Since $\mathcal{R}_{\Gamma \times \mathbb{S}^1} = \mathcal{R}_\Gamma \times \mathbb{S}^1 \times \mathbb{S}^1$ and $\tilde{\mathcal{R}}_\Gamma = \mathcal{R}_\Gamma \times \mathbb{S}^1$, we deduce that $(T_{X \setminus A}^n)_{n \in \mathbb{N}}$ tends to $\tilde{\chi}_A$ weakly as operators on the Hilbert space $L^2(\mathcal{R}_{\Gamma \times \mathbb{S}^1}) \cong L^2(\mathcal{R}_G)$. Therefore the sequence $(\pi(T_{X \setminus A}^n))_{n \in \mathbb{N}}$ tends to $\tilde{\chi}_A$ weakly in $\pi([\mathcal{R}_G])''$. \square

5. EXTREME AMENABILITY OF ORBIT FULL GROUPS

Let us recall that a Polish group is **extremely amenable** if whenever it acts continuously on a compact space, the action has a fixed point. It is **amenable** if whenever it acts continuously by affine transformations on a compact subset of a locally convex topological vector space, then the action has a fixed point.

The aim of this section is to extend Theorem 5.7 of Giordano and Pestov [GP07] to the locally compact setting.

Theorem 5.1. *Let G be a locally compact non-compact unimodular Polish group. Suppose that G acts freely on the probability space (X, μ) preserving the probability measure. Then the following are equivalent:*

- (i) G is amenable.
- (ii) $[\mathcal{R}_G]$ is amenable.
- (iii) $[\mathcal{R}_G]$ is extremely amenable.

Before we prove the theorem, let us recall the following useful well-known result which follows from Remark 5.3.29(2) and Corollary 6.2.12 of [ADR00].

Theorem 5.2. *Let G be a locally compact non-compact unimodular Polish group. Suppose that G acts freely on the probability space (X, μ) preserving the probability measure. Then G is amenable if and only if the crossed product $L^\infty(X, \mu) \rtimes G$ is injective.*

We will also need the following lemma, which provides basic extremely amenable orbit full groups.

Lemma 5.3. *Let G be a compact metrizable group acting freely on a standard probability space (X, μ) . Then the associated orbit full group is extremely amenable.*

Proof. As a consequence of [Var63, Thm. 3.2], we may view X as a Borel G -invariant subspace of a compact continuous G -space K . By [Gao09, Prop. 3.4.6], there is a Borel transversal⁵ for the G -action on K , in particular there is a Borel transversal Y for the G -action on X .

Let π be the Borel map which takes every $x \in X$ to the only $y \in Y$ such that $y \in G \cdot x$, and equip Y with the pushforward measure $\nu := \pi_*\mu$. Let λ be the Haar probability measure on G . By uniqueness of the Haar measure, the Borel G -equivariant bijection

$$\begin{aligned} \Phi : (G \times Y, \lambda \times \nu) &\rightarrow (X, \mu) \\ (g, y) &\mapsto g \cdot y \end{aligned}$$

is measure-preserving. Moreover, under the identification of X with $G \times Y$, the orbit full group becomes the group $L^0(Y, \nu, \text{Aut}(G, \lambda))$ equipped with the topology of convergence in measure. We now have two cases to consider:

- G is discrete hence finite, in which case (Y, ν) has to be non-atomic and $\text{Aut}(G, \lambda)$ is a finite permutation group, in particular it is a compact group. Then by a result of Glasner (see [Pes06, Thm. 4.2.2]), the group $L^0(Y, \nu, \text{Aut}(G, \lambda))$ is extremely amenable.
- G is non-discrete, in which case $\text{Aut}(G, \lambda)$ is extremely amenable by a result of Giordano and Pestov (see [Pes06, Thm. 4.5.15]), which implies that $L^0(Y, \nu, \text{Aut}(G, \lambda))$ also is.

In either case, we see that the orbit full group $L^0(Y, \nu, \text{Aut}(G, \lambda))$ is extremely amenable as desired. \square

Proof of Theorem 5.1. Clearly (iii) \Rightarrow (ii), so we will only have to show that (i) \Rightarrow (iii) and that (ii) \Rightarrow (i).

(i) \Rightarrow (iii): Suppose the group G is amenable. By Theorem 1.15, we can assume that X decomposes as a product $(Y \times \mathbb{S}^1, \nu \times \lambda)$, and that $\mathcal{R}_G = \mathcal{R} \times (\mathbb{S}^1 \times \mathbb{S}^1)$ where \mathcal{R} is a measure-preserving countable aperiodic amenable equivalence relation. By Connes-Feldman-Weiss' Theorem [CFW81], we can actually assume that $\mathcal{R} = \mathcal{R}_\Gamma$ where $\Gamma := \bigoplus_{n \in \mathbb{N}} \mathbb{Z}/2\mathbb{Z}$ is acting freely on (Y, ν) .

Let then $H := \Gamma \times \mathbb{S}^1$; then we have a natural H -action on $Y \times \mathbb{S}^1$ which induces the same equivalence relation as G . We thus only have to show that $[\mathcal{R}_H]$ is extremely amenable. The group H is naturally written as an increasing union of compact groups $K_n := (\bigoplus_{k \leq n} \mathbb{Z}/2\mathbb{Z}) \times \mathbb{S}^1$.

Note that the reunion $\bigcup_n \widetilde{[K_n]_D}$ is dense in $\widetilde{[H]_D}$. In fact given for every $f \in \widetilde{[H]_D}$ and every ε , there exists $N > 0$ such that

$$A := \{x \in X : f(x) \in K_N\}$$

has measure larger than $1 - \varepsilon$. Therefore by Lemma 2.7 we can extend f_A , the restriction of f to A , to an element of $\widetilde{[K_N]_D}$ which is close to f .

By Theorem 2.6 the group $[H]_D$ is dense in $[\mathcal{R}_H]$, so the union $\bigcup_n [K_n]$ is dense in $[\mathcal{R}_H]$. Observe now that $\bigcup_n [K_n]_D \subseteq \bigcup_n [\mathcal{R}_{K_n}]$ and hence $\bigcup_n [\mathcal{R}_{K_n}]$ is dense in $[\mathcal{R}_H]$. Finally observe that the full groups $[\mathcal{R}_{K_n}]$ are extremely amenable by Lemma 5.3.

⁵A Borel transversal is a Borel subset which intersects every G -orbit at exactly one point.

So $[\mathcal{R}_H]$ contains an increasing sequence of extremely amenable subgroups, whose union is dense, therefore $[\mathcal{R}_H] = [\mathcal{R}_G]$ is extremely amenable.

(ii) \Rightarrow (i): Suppose the full group $[\mathcal{R}_G]$ is amenable. By Theorem 5.2, the amenability of G is equivalent to the injectivity of the crossed product $L^\infty(X, \mu) \rtimes G$. Moreover by the celebrated result of Connes [Con76, Thm. 6], the injectivity of a von Neumann algebra $M \subseteq \mathcal{B}(\mathcal{H})$ is equivalent to Schwartz's property (P), which means that whenever $x \in \mathcal{B}(\mathcal{H})$, the closed convex hull of $(uxu^*)_{u \in \mathcal{U}(M)}$ intersects the commutant of M . It thus suffices to prove that $L^\infty(X) \rtimes G$ has Schwartz's property (P).

To this end, let $x \in \mathcal{B}(\mathcal{H})$. Then the convex closed hull K of $(uxu^*)_{u \in \mathcal{U}(L^\infty(X, \mu) \rtimes G)}$ is a weakly compact convex set on which $[\mathcal{R}_G]$ acts continuously by conjugation. Since $[\mathcal{R}_G]$ is amenable and this action is the restriction of a linear hence affine action, there exists $x_0 \in K$ which is fixed by the conjugation action. This means that $x_0 \in \pi([\mathcal{R}_G])' = (\pi([\mathcal{R}_G])'')'$ so x_0 belongs to the commutant of $L^\infty(X) \rtimes G$ by item (3) of Proposition 4.3, which concludes the proof. \square

The proof of (ii) \Rightarrow (i) actually shows that the von Neumann algebra generated by an amenable unitary group is injective. Note that de la Harpe proved that a von Neumann algebra is injective if and only if its unitary group is amenable [dlH79]. Our proof of (ii) \Rightarrow (i) is essentially a reformulation of his.

APPENDIX A. HAAR MEASURES FOR EQUIVALENCE RELATIONS

The content of this appendix is standard and can be carried out in a much more general setting (see [ADR00]). However, extracting the statements we need can be difficult, so we give complete proofs for which we claim no originality.

A.1. Invariant Haar systems. When $G = \Gamma$ is a countable discrete group, the first-coordinate projection $\pi : \mathcal{R}_\Gamma \rightarrow X$ has countable fibers, which allows us to define a **Haar measure** M on \mathcal{R}_Γ by integrating the counting measure over the fibers: for all Borel $A \subseteq \mathcal{R}_\Gamma$,

$$M(A) = \int_X |\pi^{-1}(\{x\}) \cap A| d\mu(x).$$

The definition of a Haar measure in a more general context of locally compact groups is however more complicated.

Definition A.1. Let \mathcal{R} be a Borel equivalence relation on (X, μ) . An **invariant Haar system** on \mathcal{R} is a family $(m_x)_{x \in X}$ of Borel measures on X which satisfy the following properties:

- (1) (*invariance*) There is a full measure subset X' of X such that for all $(x, y) \in \mathcal{R} \cap (X' \times X')$, $m_x = m_y$.
- (2) For all $x \in X$, m_x is non-trivial and supported on $[x]_\mathcal{R}$ (i.e. $m_x(X \setminus [x]_\mathcal{R}) = 0$ and $m_x([x]_\mathcal{R}) > 0$).
- (3) (*measurability*) For all Borel $A \subseteq \mathcal{R}$, the map $x \mapsto m_x(A_x)$ is Borel, where $A_x := \{y \in X : (x, y) \in A\}$.
- (4) (*σ -finiteness*) There exists an exhausting increasing sequence of Borel subsets (A_n) of \mathcal{R} such that for all $n \in \mathbb{N}$, one has $\int_X m_x((A_n)_x) d\mu(x) < +\infty$.
- (5) For all full measure subsets X' of X , one has $m_x(X \setminus X') = 0$ for μ -almost all $x \in X$.

The couple $(\mathcal{R}, (m_x)_{x \in X})$ is then called a **measured equivalence relation**.

Remark A.2. Note that condition (5) allows one to transport a Haar system on \mathcal{R} to a Haar system on \mathcal{R}' whenever \mathcal{R} and \mathcal{R}' are orbit equivalent.

Example A.3. Suppose \mathcal{R} is a Borel countable non-singular equivalence relation on (X, μ) . Then an invariant Haar system on \mathcal{R} is given by letting m_x be the counting measure on $[x]_{\mathcal{R}}$.

Example A.4. Suppose G is a locally compact Polish group with right Haar measure λ . Then given an essentially free measure-preserving G -action on (X, μ) , one can endow \mathcal{R}_G with an invariant Haar system $(\lambda_x)_{x \in X}$ given by the natural identification $g \mapsto g \cdot x$ between (G, λ) and $[x]_{\mathcal{R}}$. In other words, given a Borel subset A of G and $x \in X$, we set $\lambda_x(A \cdot x) := \lambda(A)$. Note that such an identification only makes sense when x belongs to the free part of the action, so when x does not belong to it we define λ_x to be the Dirac measure on x .

Let us check that the field of measure $(\lambda_x)_{x \in X}$ is an invariant Haar system. For a Borel subset A of G , we have

$$\lambda_{gx}(A \cdot x) = \lambda_{gx}(Ag^{-1}g \cdot x) = \lambda(Ag^{-1}) = \lambda(A) = \lambda_x(A \cdot x),$$

so condition (1) is satisfied. One can easily check that conditions (2), (3) and (4) are satisfied, while (5) is a consequence of the Fubini Theorem and the fact that the G -action preserves the measure: if X' has full measure in X , then for almost all $x \in X$, for λ -almost $g \in G$ one has $g \cdot x \in X'$.

Note that when G is discrete, this definition of the Haar measure coincides with the previous one.

Remark A.5. Actually, as the expert reader knows, one can define a Haar measure on \mathcal{R}_G regardless of the freeness of the G -action, whenever G is locally compact Polish. But since the construction of the measure is significantly more complicated, and since we will only deal with non-free actions when G is discrete, we chose not to present this more general setting.

Example A.6. If $(\mathcal{R}_1, (m_x^1)_{x \in X})$ and $(\mathcal{R}_2, (m_y^2)_{y \in Y})$ are measured equivalence relations on (X, μ) and (Y, ν) respectively, then $(\mathcal{R}_1 \times \mathcal{R}_2, (m_x^1 \times m_y^2)_{(x,y) \in X \times Y})$ is a measured equivalence relation on $(X \times Y, \mu \times \nu)$. A particular case of interest to us is when \mathcal{R}_1 is the transitive equivalence relation and \mathcal{R}_2 is a countable measure-preserving equivalence relation. Indeed by Theorem 1.15 every measured equivalence relation arising from a free action of a non-discrete unimodular locally compact group is of this form.

Remark A.7. There can be a lot of different invariant Haar systems on an equivalence relation \mathcal{R} , even in the ergodic case. For instance, if \mathcal{R} is the transitive equivalence relation on (X, μ) , then any choice of Borel σ -finite measure ν on X which is absolutely continuous with respect to μ yields an invariant Haar system $(m_x)_{x \in X}$ given by $m_x = \nu$. In the next section, we will add a condition which yields uniqueness: unimodularity.

By Weil's Theorem, a Polish group which admits a right-invariant measure is locally compact. Similarly, the existence of an invariant Haar system on an equivalence relation forces the acting group to be locally compact. Let a Polish group G act freely on (X, μ) , and suppose that there exists an invariant Haar system $(m_x)_{x \in X}$ on \mathcal{R}_G . Then we can define a natural right-invariant measure on G as

$$\lambda(A) := \int_X m_x(A \cdot x) d\mu(x) \text{ for } A \subset G.$$

This measure is not always σ -finite,⁶ but we now show how this can be circumvented.

Theorem A.8. *Let G be a Polish group acting freely on (X, μ) in a measure-preserving manner. If \mathcal{R}_G has an invariant Haar system, then G is locally compact.*

Proof. We will show that there exists a non-trivial right-*quasi*-invariant Borel probability measure on G . This implies that G is locally compact by Mackey's Theorem [Mac57, Thm. 7.1].

Let (m_x) be an invariant Haar system on \mathcal{R}_G , let (A_n) be a partition of \mathcal{R}_G into Borel sets of finite measure. We define a new Haar system (η_x) of *probability measures* on \mathcal{R}_G by putting, for every $x \in X$ and Borel $A \subseteq X$,

$$\eta_x(A) := \sum_{n=0}^{\infty} \frac{1}{2^{n+1}} \frac{m_x((A \cap A_n)_x)}{m_x((A_n)_x)}.$$

Then our Haar system satisfies all the axioms of invariant Haar systems except of course invariance (condition (1)) , which can be replaced by

(1') (*quasi-invariance*) There is a full measure subset X' of X such that for all $(x, y) \in \mathcal{R} \cap (X' \times X')$, $[\eta_x] = [\eta_y]$.

As before, we can integrate the Haar system to obtain a probability measure on G :

$$\lambda(A) := \int_X \eta_x(A \cdot x) d\mu(x) \text{ for } A \subset G.$$

To complete the proof, we will show that λ is quasi-invariant with respect to the right multiplication. For this, suppose that $\lambda(A) = 0$; then by definition for almost all $x \in X$ one has $\eta_x(A \cdot x) = 0$ which implies by (1') that for every $g \in G$ and almost all $x \in X$, $\eta_{gx}(A \cdot x) = 0$. Since moreover we have that $g_*\mu = \mu$, we can conclude the proof:

$$\lambda(Ag) = \int_X \eta_x(Agx) d\mu(x) = \int_X \eta_{g^{-1}x}(Ax) d\mu(x) = 0. \quad \square$$

A.2. Unimodularity. For a measured equivalence relation \mathcal{R} on (X, μ) , the **pre-orbit full group** $[\mathcal{R}]_B$ is the group of all Borel bijections $T : X \rightarrow X$ which preserve μ , and such that for all $x \in X$, one has $(x, T(x)) \in \mathcal{R}$. The pre-orbit full group has two natural actions on \mathcal{R} :

- the **left action** defined by $l_T(x, y) = (T(x), y)$ for all $(x, y) \in \mathcal{R}$ and
- the **right action** defined by $r_T(x, y) = (x, T(y))$ for all $(x, y) \in \mathcal{R}$.

These two actions are conjugated by the **flip** σ defined by $\sigma(x, y) := (y, x)$.

For $A \subset \mathcal{R}$, as in the last section, we put $A_x = \{y \in X : (x, y) \in A\}$. Every invariant Haar system $(m_x)_{x \in X}$ allows us to equip \mathcal{R} with a natural measure M defined as follows:

$$M(A) := \int_X m_x(A_x) d\mu(x) \quad \text{for every } A \subseteq \mathcal{R} \text{ Borel.}$$

Note that condition (4) on (m_x) corresponds to the σ -finiteness of (\mathcal{R}, M) .

Lemma A.9. *The left action of the pre-full group on \mathcal{R} preserves M .*

⁶Let $\Gamma = \mathbb{Z}/2\mathbb{Z}$ act on $[0, 1]$ via $T : x \mapsto (1 - x)$ and take a T -invariant function $f : [0, 1] \rightarrow [0, +\infty[$ which is not integrable; then $m_x = f(x)(\delta_x + \delta_{T(x)})$ is an invariant Haar system but the associated measure on $\mathbb{Z}/2\mathbb{Z}$ is infinite.

Proof. For all Borel $A \subseteq X$ and all $T \in [\mathcal{R}]_B$, one has

$$\begin{aligned} M(l_TA) &= \int_X m_x((l_TA)_x) d\mu(x) \\ &= \int_X m_{T^{-1}(x)}((l_TA)_x) d\mu(x) \\ &= \int_X m_{T^{-1}(x)}((A)_{T^{-1}(x)}) d\mu(x) \\ &= \int_X m_x(A_x) d\mu(x) = M(A), \end{aligned}$$

so the measure M is preserved by the left action of the pre-orbit full group. \square

Denote by $\text{Aut}(\mathcal{R}, M)$ the group of measure-preserving Borel bijections of \mathcal{R} , two such bijections being identified up to measure zero. Then the left action defines a morphism $[\mathcal{R}]_B \rightarrow \text{Aut}(\mathcal{R}, M)$ which factors through the orbit full group $[\mathcal{R}]$. So we have obtained a measure-preserving near-action of the orbit full group on (\mathcal{R}, M) .

Definition A.10. An invariant Haar system (m_x) on a Borel equivalence relation \mathcal{R} is called **unimodular** if the flip preserves M .

As the name suggests, free actions of unimodular locally compact groups give rise to unimodular Haar systems.

Proposition A.11. *Let G be a unimodular locally compact group acting essentially freely on (X, μ) and let λ be a Haar measure on G . Then the associated invariant Haar system $(\lambda_x)_{x \in X}$ on \mathcal{R}_G given by Example A.4 is unimodular.*

Proof. By Lemma 1.1 we may assume that G acts freely. Let $\Phi : X \times G \rightarrow \mathcal{R}_G$ be the Borel identification given by $\Phi(x, g) := (x, g \cdot x)$. By definition, the measure M on \mathcal{R}_G obtained by $M(A) = \int_X \lambda_x(A_x) d\mu(x)$ is just the product measure, $\Phi_*(\mu \otimes \lambda) = M$. Therefore in order to show that $(\lambda_x)_x$ is unimodular, we need to show that the map

$$\Psi := \Phi \circ \sigma \circ \Phi^{-1} : (X \times G, \mu \otimes \lambda) \rightarrow (X \times G, \mu \otimes \lambda)$$

is measure-preserving. Observe that $\Psi(x, g) = (gx, g^{-1})$. For a set C , let χ_C denote its characteristic function. Let $A \subseteq X$ and $B \subseteq G$ be Borel sets; then we have

$$\begin{aligned} \Psi_*(\mu \otimes \lambda)(A \times B) &= \int_{X \times G} \chi_{A \times B}(g \cdot x, g^{-1}) d\mu \otimes \lambda(g, x) \\ &= \int_{X \times G} \chi_A(g \cdot x) \chi_B(g^{-1}) d\mu \otimes \lambda(g, x) \\ &= \int_G \chi_B(g^{-1}) \left(\int_X \chi_A(g \cdot x) d\mu(x) \right) d\lambda(g) \\ &= \int_G \chi_B(g^{-1}) \mu(A) d\lambda(g) \\ &= \lambda(B) \mu(A), \end{aligned}$$

where the last three equalities are respectively consequences of Fubini's Theorem, the fact that G preserves the measure and the unimodularity of G . By uniqueness of the product measure, we conclude that $\Psi_*(\mu \times \lambda) = \mu \times \lambda$ as desired. \square

Remark A.12. Let G be a unimodular locally compact Polish group acting essentially freely on (X, μ) , let λ be a Haar measure on G and let (λ_x) be the associated unimodular invariant Haar system on \mathcal{R}_G . Then by the above proposition the right $[\mathcal{R}_G]$ -action on (\mathcal{R}_G, M) gives an embedding $[\mathcal{R}_G] \hookrightarrow L^0(X, \mu, \text{Aut}(G, \lambda))$ (in other words, the full group acts on every \mathcal{R}_G -class in a measure-preserving manner).

Since embeddings of Polish groups have closed range (see [Gao09, Prop. 2.2.1]), we can deduce that if (Y, ν) is a standard σ -finite space, then the group $L^0(X, \mu, \text{Aut}(Y, \nu))$ contains as a closed subgroup every orbit full group arising from a measure-preserving free action of a non-discrete unimodular Polish locally compact group. For a similar statement in the discrete case, see [KLM15, Prop. 13].

Theorem A.13. *Let G be a Polish group acting freely on (X, μ) . If there is a unimodular invariant Haar system $(m_x)_{x \in X}$ on \mathcal{R}_G , then G is locally compact unimodular.*

If the action is moreover ergodic, then there exists a constant $c > 0$ such that for almost all $x \in X$, one has $m_x = c\lambda_x$, where λ_x is the invariant Haar system associated to a fixed Haar measure λ on G .

Proof. First note that by Theorem A.8, G has to be locally compact. We fix a left Haar measure λ on G . For every x in the free part of the action, consider the G -equivariant bijective Borel map $\phi_x : [x]_{\mathcal{R}} \rightarrow G$ defined by $\phi_x(y) \cdot x = y$; then the pushforward measure $\eta_x := (\phi_x)_* m_x$ is a σ -finite measure on G .

Since the right action of the orbit full group $[\mathcal{R}]$ on (\mathcal{R}, M) is conjugate to the left action by the flip, unimodularity yields that the right action of $[\mathcal{R}]$ on \mathcal{R} preserves M . In particular, the right action of G on \mathcal{R} preserves M , so for a fixed $g \in G$, and any $A \subseteq \mathcal{R}$ we have

$$\int_X m_x(A_x) d\mu(x) = \int_X m_x((r(g)A)_x) d\mu(x) = \int_X m_x(gA_x) d\mu(x).$$

By the uniqueness of disintegration, this implies that for almost all $x \in X$, $g_* m_x = m_x$. Then by Fubini's Theorem, for almost all $x \in X$ and λ -almost all $g \in G$, $g_* m_x = m_x$. Since $\phi_x : [x]_{\mathcal{R}} \rightarrow G$ is left G -equivariant, this implies that for almost all $x \in X$, there is a full measure subgroup of G which preserves η_x when acting on the left. But every full measure subgroup of G equates G (see e.g. [Zim84, Prop. B.1]), so for almost all $x \in X$, one has that η_x is a Borel σ -finite left-invariant measure on G . By uniqueness of the Haar measure, we conclude that for almost all $x \in X$, the measure η_x is a multiple of λ . Let X_0 be the full measure G -invariant Borel set of $x \in X$ such that η_x is a multiple of λ .

Fix a Borel subset K of G such that $\lambda(K) = 1$. For all $x \in X_0$, we let $c_x = \eta_x(K)$; then $x \mapsto c_x$ is Borel and we have $\eta_x = c_x \lambda$. Moreover for all $(g, x) \in G \times X_0$,

$$c_{gx} = \eta_{gx}(K) = m_{gx}(Kg x) = m_x(Kgx) = \eta_x(Kg),$$

so that $c_{gx} = \Delta(g)c_x$, where Δ is the modular function on G .

Let $g \in G$, let $a > 0$ be an essential value of the function $x \mapsto c_x$ and consider the set of positive measure $A := \{x \in X : a/2 < c_x < 3a/2\}$. By Poincaré's Recurrence Theorem for almost all $x \in A$ there is an infinite subset $S_x \subset \mathbb{N}$ such that $g^k x \in A$ for every $k \in S_x$. So for $x \in A$ we have that

$$a/2 < \Delta(g^k)c_x = \Delta(g)^k c_x < 3a/2 \quad \text{for all } k \in S_x,$$

which implies that $\Delta(g) = 1$, and we conclude that G is unimodular.

Therefore $c_{gx} = c_x$ for all $x \in X_0$: the function $x \mapsto c_x$ is G -invariant. So whenever the G -action is ergodic, $x \mapsto c_x$ is a.s. constant, which yields the second part of the theorem. \square

Let us point out that when the acting group G is already known to be locally compact, the freeness hypothesis above can be replaced by almost freeness, since we know by Lemma 1.1 that the free part of the action is a Borel set and invariant Haar systems restrict well to full measure Borel subsets. So unimodular locally compact groups form a closed class under orbit equivalence among locally compact groups.

ACKNOWLEDGMENTS

Both authors would like to thank Damien Gaboriau, Henrik Petersen, Sven Raum and Todor Tsankov for many useful discussions around this topic. We are especially grateful to Sven Raum for explaining to us the proof of item (3) of Proposition 4.3 in the discrete case. We thank the anonymous referee for the detailed report which significantly improved the exposition.

REFERENCES

- [ADR00] C. Anantharaman-Delaroche and J. Renault, *Amenable groupoids*, Monographies de L'Enseignement Mathématique [Monographs of L'Enseignement Mathématique], vol. 36, L'Enseignement Mathématique, Geneva, 2000. With a foreword by Georges Skandalis and Appendix B by E. Germain. MR1799683
- [AEG94] Scot Adams, George A. Elliott, and Thierry Giordano, *Amenable actions of groups*, Trans. Amer. Math. Soc. **344** (1994), no. 2, 803–822, DOI 10.2307/2154508. MR1250814
- [Bec13] Howard Becker, *Cocycles and continuity*, Trans. Amer. Math. Soc. **365** (2013), no. 2, 671–719, DOI 10.1090/S0002-9947-2012-05570-X. MR2995370
- [BK96] Howard Becker and Alexander S. Kechris, *The descriptive set theory of Polish group actions*, London Mathematical Society Lecture Note Series, vol. 232, Cambridge University Press, Cambridge, 1996. MR1425877
- [CFW81] A. Connes, J. Feldman, and B. Weiss, *An amenable equivalence relation is generated by a single transformation*, Ergodic Theory Dynamical Systems **1** (1981), no. 4, 431–450 (1982). MR662736
- [CH75] Wojciech Herer and Jens Peter Reus Christensen, *On the existence of pathological submeasures and the construction of exotic topological groups*, Math. Ann. **213** (1975), 203–210, DOI 10.1007/BF01350870. MR0412369
- [CM16] Alessandro Carderi and François Le Maître, *More Polish full groups*, Topology Appl. **202** (2016), 80–105, DOI 10.1016/j.topol.2015.12.065. MR3464151
- [Con76] A. Connes, *Classification of injective factors. Cases II_1 , II_∞ , III_λ , $\lambda \neq 1$* , Ann. of Math. (2) **104** (1976), no. 1, 73–115, DOI 10.2307/1971057. MR0454659
- [Dan00] Alexandre I. Danilenko, *Point realization of Boolean actions of countable inductive limits of locally compact groups* (English, with English, Russian and Ukrainian summaries), Mat. Fiz. Anal. Geom. **7** (2000), no. 1, 35–48. MR1760945
- [dlH79] P. de la Harpe, *Moyennabilité du groupe unitaire et propriété P de Schwartz des algèbres de von Neumann* (French), Algèbres d'opérateurs (Sém., Les Plans-sur-Bex, 1978), Lecture Notes in Math., vol. 725, Springer, Berlin, 1979, pp. 220–227. MR548116
- [Dye59] H. A. Dye, *On groups of measure preserving transformation. I*, Amer. J. Math. **81** (1959), 119–159, DOI 10.2307/2372852. MR0131516
- [FM77] Jacob Feldman and Calvin C. Moore, *Ergodic equivalence relations, cohomology, and von Neumann algebras. I*, Trans. Amer. Math. Soc. **234** (1977), no. 2, 289–324, DOI 10.2307/1997924. MR0578656
- [For74] Peter Forrest, *On the virtual groups defined by ergodic actions of R^n and \mathbf{Z}^n* , Advances in Math. **14** (1974), 271–308, DOI 10.1016/0001-8708(74)90033-4. MR0417388

- [Fur99a] Alex Furman, *Gromov's measure equivalence and rigidity of higher rank lattices*, Ann. of Math. (2) **150** (1999), no. 3, 1059–1081, DOI 10.2307/121062. MR1740986
- [Fur99b] Alex Furman, *Orbit equivalence rigidity*, Ann. of Math. (2) **150** (1999), no. 3, 1083–1108, DOI 10.2307/121063. MR1740985
- [Gab00] Damien Gaboriau, *Coût des relations d'équivalence et des groupes* (French, with English summary), Invent. Math. **139** (2000), no. 1, 41–98, DOI 10.1007/s002229900019. MR1728876
- [Gab05] D. Gaboriau, *Examples of groups that are measure equivalent to the free group*, Ergodic Theory Dynam. Systems **25** (2005), no. 6, 1809–1827, DOI 10.1017/S0143385705000258. MR2183295
- [Gao09] Su Gao, *Invariant descriptive set theory*, Pure and Applied Mathematics (Boca Raton), vol. 293, CRC Press, Boca Raton, FL, 2009. MR2455198
- [GM83] M. Gromov and V. D. Milman, *A topological application of the isoperimetric inequality*, Amer. J. Math. **105** (1983), no. 4, 843–854, DOI 10.2307/2374298. MR708367
- [GP02] Thierry Giordano and Vladimir Pestov, *Some extremely amenable groups* (English, with English and French summaries), C. R. Math. Acad. Sci. Paris **334** (2002), no. 4, 273–278, DOI 10.1016/S1631-073X(02)02218-5. MR1891002
- [GP07] Thierry Giordano and Vladimir Pestov, *Some extremely amenable groups related to operator algebras and ergodic theory*, J. Inst. Math. Jussieu **6** (2007), no. 2, 279–315, DOI 10.1017/S1474748006000090. MR2311665
- [GTW05] E. Glasner, B. Tsirelson, and B. Weiss, *The automorphism group of the Gaussian measure cannot act pointwise*, Israel J. Math. **148** (2005), 305–329, DOI 10.1007/BF02775441. Probability in mathematics. MR2191233
- [Hud93] Steven M. Hudson, *A marriage theorem with Lebesgue measure*, J. Combin. Theory Ser. A **62** (1993), no. 2, 234–251, DOI 10.1016/0097-3165(93)90046-B. MR1207735
- [Kec95] Alexander S. Kechris, *Classical descriptive set theory*, Graduate Texts in Mathematics, vol. 156, Springer-Verlag, New York, 1995. MR1321597
- [Kec10] Alexander S. Kechris, *Global aspects of ergodic group actions*, Mathematical Surveys and Monographs, vol. 160, American Mathematical Society, Providence, RI, 2010. MR2583950
- [KLM15] Adriane Kaïchouh and François Le Maître, *Connected Polish groups with ample generics*, Bull. Lond. Math. Soc. **47** (2015), no. 6, 996–1009, DOI 10.1112/blms/bdv078. MR3431579
- [KPV15] David Kyed, Henrik Densing Petersen, and Stefaan Vaes, *L^2 -Betti numbers of locally compact groups and their cross section equivalence relations*, Trans. Amer. Math. Soc. **367** (2015), no. 7, 4917–4956, DOI 10.1090/S0002-9947-2015-06449-6. MR3335405
- [LM14] François Le Maître, *The number of topological generators for full groups of ergodic equivalence relations*, Invent. Math. **198** (2014), no. 2, 261–268, DOI 10.1007/s00222-014-0503-6. MR3274561
- [LM15] François Le Maître, *On full groups of non-ergodic probability-measure-preserving equivalence relations*, Ergodic Theory Dynam. Systems **36** (2016), no. 7, 2218–2245, DOI 10.1017/etds.2015.20. MR3568978
- [Mac57] George W. Mackey, *Borel structure in groups and their duals*, Trans. Amer. Math. Soc. **85** (1957), 134–165, DOI 10.2307/1992966. MR0089999
- [Mac62] George W. Mackey, *Point realizations of transformation groups*, Illinois J. Math. **6** (1962), 327–335. MR0143874
- [MRV13] Niels Meesschaert, Sven Raum, and Stefaan Vaes, *Stable orbit equivalence of Bernoulli actions of free groups and isomorphism of some of their factor actions*, Expo. Math. **31** (2013), no. 3, 274–294, DOI 10.1016/j.exmath.2012.08.012. MR3108102
- [OW87] Donald S. Ornstein and Benjamin Weiss, *Entropy and isomorphism theorems for actions of amenable groups*, J. Analyse Math. **48** (1987), 1–141, DOI 10.1007/BF02790325. MR910005
- [Pes06] Vladimir Pestov, *Dynamics of infinite-dimensional groups*, University Lecture Series, vol. 40, American Mathematical Society, Providence, RI, 2006. The Ramsey-Dvoretzky-Milman phenomenon; Revised edition of *Dynamics of infinite-dimensional groups and Ramsey-type phenomena* [Inst. Mat. Pura. Apl. (IMPA), Rio de Janeiro, 2005; MR2164572]. MR2277969

- [Pet13] Henrik Densing Petersen, *L^2 -Betti numbers of locally compact groups* (English, with English and French summaries), C. R. Math. Acad. Sci. Paris **351** (2013), no. 9–10, 339–342, DOI 10.1016/j.crma.2013.05.013. MR3072156
- [Pra81] V. S. Prasad, *Generating dense subgroups of measure preserving transformations*, Proc. Amer. Math. Soc. **83** (1981), no. 2, 286–288, DOI 10.2307/2043512. MR624915
- [Pra83] V. S. Prasad, *Sous-groupes libres et sous-ensembles indépendants de transformations préservant la mesure* (French), Measure theory and its applications (Sherbrooke, Que., 1982), Lecture Notes in Math., vol. 1033, Springer, Berlin, 1983, pp. 276–282, DOI 10.1007/BFb0099865. MR729543
- [Var63] V. S. Varadarajan, *Groups of automorphisms of Borel spaces*, Trans. Amer. Math. Soc. **109** (1963), 191–220, DOI 10.2307/1993903. MR0159923
- [vD78] A. van Daele, *Continuous crossed products and type III von Neumann algebras*, London Mathematical Society Lecture Note Series, vol. 31, Cambridge University Press, Cambridge-New York, 1978. MR500089
- [Zim84] Robert J. Zimmer, *Ergodic theory and semisimple groups*, Monographs in Mathematics, vol. 81, Birkhäuser Verlag, Basel, 1984. MR776417

INSTITUT FÜR GEOMETRIE, TECHNISCHE UNIVERSITÄT DRESDEN, 01062 DRESDEN, GERMANY
E-mail address: `alessandro.carderi@gmail.com`

INSTITUT DE MATHÉMATIQUES DE JUSSIEU-PRG, UNIVERSITÉ PARIS DIDEROT, SORBONNE PARIS
CITÉ, 75205 PARIS CEDEX 13, FRANCE
E-mail address: `francois.le-maitre@imj-prg.fr`