

American Mathematical Society

Colloquium Publications

Volume 36

Topological Dynamics

Walter Helbig Gottschalk
Gustav Arnold Hedlund



TOPOLOGICAL DYNAMICS

2000 *Mathematics Subject Classification*. Primary 54–XX.

Library of Congress Card Number 55-12710
ISBN 0-8218-1036-7
ISSN 0065-9258

Copying and reprinting. Individual readers of this publication, and nonprofit libraries acting for them, are permitted to make fair use of the material, such as to copy a chapter for use in teaching or research. Permission is granted to quote brief passages from this publication in reviews, provided the customary acknowledgment of the source is given.

Republication, systematic copying, or multiple reproduction of any material in this publication is permitted only under license from the American Mathematical Society. Requests for such permission should be addressed to the Acquisitions Department, American Mathematical Society, 201 Charles Street, Providence, Rhode Island 02904-2294, USA. Requests can also be made by e-mail to reprint-permission@ams.org.

Copyright © 1955 by the American Mathematical Society
Reprinted with corrections and an appendix, 1968
Printed in the United States of America.

The American Mathematical Society retains all rights
except those granted to the United States Government.

This book may not be reproduced in any form without the permission of the publisher


⊗ The paper used in this book is acid-free and falls within the guidelines
established to ensure permanence and durability.

Visit the AMS home page at <http://www.ams.org/>

10 09 08 07 06 10 09 08 07 06 05

TOPOLOGICAL DYNAMICS

WALTER HELBIG GOTTSCHALK
AND GUSTAV ARNOLD HEDLUND



This page intentionally left blank

PREFACE

By *topological dynamics* we mean the study of transformation groups with respect to those topological properties whose prototype occurred in classical dynamics. Thus the word "topological" in the phrase "topological dynamics" has reference to mathematical content and the word "dynamics" in the phrase has primary reference to historical origin.

Topological dynamics owes its origin to the classic work of Henri Poincaré and G. D. Birkhoff. It was Poincaré who first formulated and solved problems of dynamics as problems in topology. Birkhoff contributed fundamental concepts to topological dynamics and was the first to undertake its systematic development.

In the classic sense, a dynamical system is a system of ordinary differential equations with at least sufficient conditions imposed to insure continuity and uniqueness of the solutions. As such, a dynamical system defines a (one-parameter or continuous) flow in a space. A large body of results for flows which are of interest for classical dynamics has been developed, since the time of Poincaré, without reference to the fact that the flows arise from differential equations. The extension of these results from flows to transformation groups has been the work of recent years. These extensions and the concomitant developments are set forth in this book.

Part One contains the general theory. Part Two contains notable examples of flows which have contributed to the general theory of topological dynamics and which in turn have been illuminated by the general theory of topological dynamics.

In addition to the present Colloquium volume, the only books which contain extensive related developments are G. D. Birkhoff [2, Chapter 7], Niemytzki and Stepanoff [1, Chapter 4 of the 1st edition, Chapter 5 of the 2nd edition] and G. T. Whyburn [1, Chapter 12]. The contents of this volume meet but do not significantly overlap a forthcoming book by Montgomery and Zippin.

The authors wish to express their appreciation to the American Mathematical Society for the opportunity to publish this work. They also extend thanks to Yale University and the Institute for Advanced Study for financial aid in the preparation of the manuscript. The second named author extends to the American Mathematical Society his thanks for the invitation to give the Colloquium Lectures in which some aspects of the subject were discussed. Some of his work has been supported by the United States Air Force through the Office of Scientific Research of the Air Research and Development Command.

PHILADELPHIA, PENNSYLVANIA
NEW HAVEN, CONNECTICUT
July, 1954

This page intentionally left blank

CONVENTIONS AND NOTATIONS

Each of the two parts of the book is divided into sections and each section into paragraphs. Cross references are to paragraphs. 4.6 is the sixth paragraph of section 4. In general, a paragraph is either a definition, lemma, theorem or remark. A "remark" is a statement, the proof of which is left to the reader. These proofs are not always trivial, however.

References to the literature are, in general, given in the last paragraph of each section. Numbers in brackets following an author's name refer to the bibliography at the end of the book. Where there is joint authorship, the number given refers to the article or book as listed under the first named author.

An elementary knowledge of set theory, topology, uniform spaces and topological groups is assumed. Such can be gained by reading the appropriate sections of Bourbaki [1, 2, 3]. With a few exceptions to be noted, the notations used are standard and a separate listing seemed unnecessary.

Unless the contrary is specifically indicated, groups are taken to be multiplicative. Topological groups are not assumed to be necessarily separated (Hausdorff). The additive group of integers will be denoted by \mathcal{I} and the additive group of reals by \mathcal{R} .

Contrary to customary usage, the function or transformation sign is usually placed on the right. That is, if X and Y are sets, f denotes a transformation of X into Y and $x \in X$, then xf denotes the unique element of Y determined by x and f .

In connection with uniform spaces, the term *index* is used to denote an element of the filter defining the uniform structure, thus replacing the term *entourage* as used by Bourbaki [2]. In keeping with the notation for the value of a function, if X is a uniform space, α is an index of X and $x \in X$, then $x\alpha$ denotes the set of all $y \in X$ such that $(x, y) \in \alpha$. Unless the contrary is stated, a uniform space is not necessarily separated.

This page intentionally left blank

TABLE OF CONTENTS

SECTION	PAGE
1. TRANSFORMATION GROUPS	1
2. ORBIT-CLOSURE PARTITIONS	12
3. RECURSION	20
4. ALMOST PERIODICITY	31
5. REGULAR ALMOST PERIODICITY	49
6. REPLETE SEMIGROUPS	57
7. RECURRENCE	64
8. INCOMPRESSIBILITY	69
9. TRANSITIVITY	73
10. ASYMPTOTICITY	81
11. FUNCTION SPACES	90
12. SYMBOLIC DYNAMICS	102
13. GEODESIC FLOWS OF MANIFOLDS OF CONSTANT NEGATIVE CURVATURE	114
14. CYLINDER FLOWS AND A PLANAR FLOW	133
APPENDIX	143
BIBLIOGRAPHY	159
INDEX	165

This page intentionally left blank

APPENDIX²

MINIMAL SETS: AN INTRODUCTION TO TOPOLOGICAL DYNAMICS¹

W. H. GOTTSCHALK

The notion of minimal set is centrally located in topological dynamics. Topological dynamics may be defined as the study of transformation groups with respect to those properties, wholly or largely topological in nature, whose prototype occurred in classical dynamics.

Henri Poincaré was the first to introduce topological notions and methods in dynamics, that is, the study of ordinary differential equations. G. D. Birkhoff was the first to undertake the systematic development of topological dynamics, indicating its essentially abstract character and making fundamental contributions. Birkhoff's first paper on the subject appeared in 1912 [6, pp. 654–672]; Birkhoff's paper contains the first definition of minimal sets, some theorems about them, and some examples of them. Most of Birkhoff's work in topological dynamics from the point of view of general theory is to be found in Chapter 7 of his Colloquium volume *Dynamical systems* published in 1927 [5]. The Colloquium volume *Analytical topology* by G. T. Whyburn [27], published in 1942, contains related developments in its Chapter 12. The Russian book, *Qualitative theory of differential equations* by Nemyckii and Stepanov [22], first edition in 1947 and second edition in 1949, contains a chapter devoted to topological dynamics; an English translation of this book has recently been announced by Princeton Press. The treatments of topological dynamics in the above books are all from the points of view of a single transformation or a one-parameter group of transformations. The Colloquium volume *Topological dynamics* by Hedlund and myself [14], published in 1955, is concerned mainly with general transformation groups.

A *topological transformation group*, or *transformation group* for short, is defined to be an ordered triple (X, T, π) such that the following axioms are satisfied:

(A0) (STIPULATIVE AXIOM). X is a topological space, called the *phase space*; T is a topological group, called the *phase group*; and π is a map of the cartesian product $X \times T$ into X , called the *phase map*.

An address delivered before the Washington meeting of the Society on October 26, 1957, by invitation of the Committee to Select Hour Speakers for Eastern Sectional Meetings; received by the editors April 10, 1958.

¹ This address was prepared while the author was under contract No. AF 18(600)-1116 of the Air Force Office of Scientific Research. Reproduction in whole or in part is permitted for any purpose of the United States Government.

² Reprinted from the Bulletin of the American Mathematical Society, Vol. 64, No. 6, November, 1958, pp. 336-351.

[Let us for the moment use the multiplicative notation for the phase group T , let e denote the identity element of T , and let the value of π at the point (x, t) of $X \times T$ be denoted by xt .]

(A1) (IDENTITY AXIOM). $xe = x$ for all $x \in X$.

(A2) (HOMOMORPHISM AXIOM). $(xt)s = x(ts)$ for all $x \in X$ and all $t, s \in T$.

(A3) (CONTINUITY AXIOM). π is continuous.

Consider a given transformation group (X, T, π) . The phase map $(x, t) \rightarrow xt$ determines two kinds of maps when one of the variables x, t is replaced by a constant. Thus, for fixed $t \in T$, the map $x \rightarrow xt$ is a homeomorphism π^t of the phase space X onto itself which is called a *transition*. Again, for fixed $x \in X$, the map $t \rightarrow xt$ is a continuous map π_x of the phase group T into the phase space X which is called a *motion*. The set of all transitions is a group of homeomorphisms of the phase space X onto itself such that the transition π^e induced by the identity element e of T is the identity homeomorphism of the phase space X and such that the map $t \rightarrow \pi^t$ is a group homomorphism. Conversely, a group of homeomorphisms of X onto itself, suitably topologized, gives rise to a transformation group.

An *intrinsic* property of the transformation group (X, T, π) is defined to be a property which is describable in terms of the topology of the phase space X , the topology and group structure of the phase group T , and the phase map π . Topological dynamics is concerned with intrinsic properties of transformation groups with particular reference to those properties which first arose in classical dynamics.

Among the examples which have been studied in the past, the most frequently occurring phase groups are the additive group \mathcal{I} of integers with the discrete topology and the additive group \mathcal{R} of real numbers with its usual topology. In either case, the transformation group is called a *flow*. If the phase group T is \mathcal{I} , then the transformation group is called a *discrete* flow. If the phase group T is \mathcal{R} , then the transformation group is called a *continuous* flow. Here the word "continuous" apparently refers to the "real number continuum".

To see how transformation groups in the role of continuous flows appeared in classical dynamics, consider a system of n first order ordinary differential equations of the form

$$\frac{dx_i}{dt} = f_i(x_1, \dots, x_n) \quad (i = 1, \dots, n)$$

where the functions f_i may be defined on a region X of euclidean n -

space or they may be defined locally throughout an n -dimensional differentiable manifold X . Impose enough conditions to insure the existence, uniqueness, and continuity of solutions for all real values of the "time" t and through all points of X . Then there exists exactly one continuous flow in X whose motions are the solutions of the system of differential equations. To illustrate with a very simple example, the continuous flow in the plane determined by the system

$$\frac{dx_1}{dt} = x_1, \quad \frac{dx_2}{dt} = -x_2$$

has as phase map $\pi((x_1, x_2), t) = (x_1 e^t, x_2 e^{-t})$.

A *dynamical system* may be defined as a continuous flow in a differentiable manifold induced by an autonomous system of first order ordinary differential equations, as above. Of course, every dynamical system is a continuous flow. To what extent and under what conditions a continuous flow is a dynamical system appears to be a largely unsolved problem. Let me say that this usage of the term "dynamical system" is not at all universal. More likely than not it is often used as synonymous with "continuous flow" or even "discrete flow."

The two kinds of flows, discrete and continuous, are closely related. For example, a continuous flow determines many discrete flows by taking cyclic subgroups of \mathcal{G} . Conversely, a discrete flow determines a continuous flow when the phase space X is extended to the cartesian product of X and the closed unit interval, and the bases of the cylinder thus obtained are identified according to the transition induced by the integer 1, that is, $(x, 1)$ and $(x1, 0)$ are identified. The problem of imbedding a discrete flow in a continuous flow without alteration of the phase space appears to be largely unsolved.

A discrete flow determines and is determined by the transition induced by the integer 1. Thus, we may redefine a discrete flow more economically as an ordered pair (X, ϕ) where X is a topological space and ϕ is a homeomorphism of X onto X . For the sake of simplicity, I wish to confine most of my remaining remarks to flows, and in particular to discrete flows. Moreover, the phase space will be assumed to be compact metric for the same reason. It is to be pointed out that the present theory of topological dynamics contains results on transformation groups acting upon topological spaces and uniform spaces which specialize to give most of the known facts about flows on metric spaces.

Let X be a nonvacuous compact metric space with metric ρ and let ϕ be a homeomorphism of X onto X .

The phase space X is nonvacuous closed and invariant, that is, $X\phi = X$. In case X has these properties minimally it is said that X is a *minimal set*. The definition may be phrased for a subset of X . A subset M of X is said to be *minimal under ϕ* provided that M is nonvacuous closed and invariant, that is $M\phi = M$, and no proper subset of M has all these properties. Minimal sets may be characterized in various ways. If $x \in X$, then the *orbit of x* , denoted $O(x)$, is defined to be $\{x\phi^n \mid n \in \mathcal{I}\}$ where \mathcal{I} is the set of all integers; and the *orbit-closure of x* , denoted $\bar{O}(x)$, is defined to be the closure of the orbit $O(x)$ of x . A subset M of X is minimal if and only if M is nonvacuous and M is the orbit-closure of each of its points.

By the axiom of choice, there always exists at least one minimal subset of X [14, p. 15].

Now recursive properties enter into consideration. A point x of the phase space is said to be *almost periodic under ϕ* and ϕ is said to be *almost periodic at x* provided that if U is a neighborhood of x , then there exists a relatively dense subset A of \mathcal{I} such that $x\phi^n \in U$ for all $n \in A$. A subset A of \mathcal{I} is called *relatively dense* in case the gaps of A are bounded. Another way of saying the same thing is that $\mathcal{I} = A + K = \{a + k \mid a \in A \& k \in K\}$ for some finite subset K of \mathcal{I} . We may call the set K a *bond* of the set A . A periodic point is almost periodic, but not conversely. It has been remarked that a periodic point returns to itself every hour on the hour; but an almost periodic point returns to a neighborhood every hour within the hour.

The basic connection between minimal sets and almost periodic points is this: The orbit-closure of a point x is minimal if and only if the point x is almost periodic [14, p. 31]. Consequently, every point of a minimal set is almost periodic, and there always exists at least one almost periodic point. This, of course, is in strong contrast to the situation with respect to periodic points.

Since different minimal subsets of X are necessarily disjoint, we may conclude that the class of all orbit-closures is a partition of the phase space if and only if ϕ is pointwise almost periodic, that is to say, ϕ is almost periodic at each point of X . The partition of orbit-closures, when it exists, is necessarily star-open in the sense that the saturation or star of every open set is again open. In a compact metric space, which we are here considering, this is equivalent to lower semi-continuity in another terminology. It is however not always the case that the partition of orbit-closures is star-closed in the sense that the star of every closed set is itself closed, or equivalently, upper semi-continuous. This condition may be characterized by the following recursive property. The map ϕ is said to be *weakly almost periodic on*

X and the phase space X is said to be *weakly almost periodic under ϕ* provided that if ϵ is a positive real number, then there exists a finite subset K_ϵ of \mathcal{S} such that $x \in X$ implies the existence of a (necessarily relatively dense) subset A_x of \mathcal{S} such that $\mathcal{S} = A_x + K_\epsilon$ and $x\phi^n \in N_\epsilon(x)$ for all $n \in A_x$. Here the bond K_ϵ depends only upon ϵ , but the individual sets of return A_x depend also upon x . To summarize, the class of all orbit-closures is a star-closed partition of the phase space if and only if the map ϕ is weakly almost periodic [14, p. 34]. In particular, ϕ is weakly almost periodic on each minimal set.

We are now in the middle of the spectrum of recursive properties. A weaker recursive property than almost periodic may be mentioned first of all. A point x of the phase space is said to be *recurrent* provided that if U is a neighborhood of x , then there exists an extensive subset A of \mathcal{S} such that $x\phi^n \in U$ for all $n \in A$. Here “extensive” means containing a sequence diverging to $-\infty$ and a sequence diverging to $+\infty$. In other words a point is recurrent provided that it returns (or recurs) to a neighborhood infinitely often in the past and infinitely often in the future. The term recurrent as it is here used coincides with its meaning in the Poincaré Recurrence Theorem (1899), which seems to be one of the earliest appearances in the literature of a theorem on a recursive property different from periodic. In general, pointwise recurrent does not imply pointwise almost periodic [12]. However, if the phase space is also zero-dimensional, then pointwise recurrent implies weakly almost periodic [14, p. 65].

Corresponding to any recursive property there is also the property of regionally recursive. For example, ϕ is *regionally recurrent* provided that if $x \in X$ and if U is a neighborhood of x , then there exists an extensive subset A of \mathcal{S} such that $U \cap U\phi^n \neq \emptyset$ for every $n \in A$. The *center* of a discrete flow (X, ϕ) is defined to the greatest invariant subset on which ϕ is regionally recurrent. The center is also characterized as the closure of the set of recurrent points. Since every almost periodic point is also recurrent, the center is necessarily nonvacuous. The center has the property that the relative sojourn of every point of the phase space in a neighborhood of the center is equal to unity. If the flow has an invariant measure which is positive and finite for nonvacuous open sets, then the flow is regionally recurrent and the center is the entire phase space. This is the case for conservative dynamical systems [14; 22].

Let us examine a few stronger recursive properties. The map ϕ is said to be *almost periodic on X* and the phase space X is said to be *almost periodic under ϕ* provided that if $\epsilon > 0$, then there exists a relatively dense subset A of \mathcal{S} such that $x\phi^n \in N_\epsilon(x)$ for all $x \in X$ and

all $n \in A$. Almost periodic implies weakly almost periodic which in turn implies pointwise almost periodic; no converse holds. The almost periodicity of the map ϕ may be characterized as follows. These four statements are pairwise equivalent: (1) ϕ is almost periodic; (2) the set of powers of ϕ , namely $\{\phi^n \mid n \in \mathcal{J}\}$, is equicontinuous; (3) the set of powers of ϕ has compact closure where the ambient space is the group of all homeomorphisms of X onto X with the usual topology; (4) there exists a compatible metric of X which makes ϕ an isometry [14, p. 37].

A still stronger recursive property is regularly almost periodic (G. T. Whyburn 1942; P. A. Smith 1941 related) [27; 24]. This is like almost periodic except that the set of return A is taken to be a nontrivial subgroup of \mathcal{J} . We have then the properties: regularly almost periodic point, pointwise regularly almost periodic map, and regularly almost periodic map. We mention a few results in this context. The map ϕ is regularly almost periodic if and only if ϕ is both pointwise regularly almost periodic and weakly almost periodic. If ϕ is pointwise regularly almost periodic, then every orbit-closure is a zero-dimensional regularly almost periodic minimal set [14, p. 49f.]. That a regularly almost periodic homeomorphism on a manifold is necessarily periodic has been conjectured by P. A. Smith (1941) [24], but so far has been proved only for the two-dimensional case [for example, 14, p. 56].

By definition, an *inheritance* theorem is a theorem of the form " ϕ has property P if and only if ϕ^n has the property P " where n is a preassigned nonzero integer. The inheritance theorem holds for recurrent and almost periodic points as well as for certain other recursive properties of a point. Actually, a very general inheritance theorem can be proved which yields most known results for recursive points [14, p. 26]. No general inheritance theorem is known for such properties as regionally recurrent and weakly almost periodic although they too inherit, at least under certain conditions [14, pp. 67 and 35].

In this connection we may mention the fact that a connected minimal set under ϕ is also minimal under ϕ^n ($n \neq 0$) [14, p. 16].

Two discrete flows (X, ϕ) and (Y, ψ) are said to be *isomorphic* provided there exists a homeomorphism h of X onto Y such that $\phi h = h\psi$.

Let us consider now the problems: (1) (Construction problem.) To construct all minimal sets systematically; (2) (Classification problem.) To classify all minimal sets according to isomorphism type. At the present, only partial answers can be given to these questions, even for discrete flows on compact metric spaces.

In the spectrum of isomorphism types of minimal sets, those which are almost periodic seem to constitute an extreme case, for the following reason. A *monothetic group* is by definition a compact group which contains a dense cyclic subgroup, a generator of a dense cyclic subgroup being called a generator of the monothetic group. Now the almost periodic minimal sets under discrete flows are coextensive with the monothetic groups in the following sense. If X is a monothetic group with generator a , then X is a minimal set under the homeomorphism ϕ of X onto X defined by $x \rightarrow xa$. Conversely, if (X, ϕ) is an almost periodic minimal set, then there exists a group structure in X which makes X a monothetic group and such that the map ϕ is translation by a generator a , that is $x\phi = xa$ for all $x \in X$ [14, p. 39]. Since the monothetic groups can be constructed in their entirety as the character groups of the subgroups of the discrete circle group (Anzai and Kakutani 1943) [2], the construction problem for almost periodic minimal sets under discrete flows has a reasonably definite answer. It is not clear to me whether the corresponding classification problem can also be answered by the present theory of topological groups. It is known (Halmos and Samelson 1942; Anzai and Kakutani 1943) [15; 2] that every compact connected separable abelian group is monothetic. For example, the n -toral groups K^n where n is a positive integer and the infinite toral group $K^{\mathcal{J}}$ are monothetic.

The n -adic groups are monothetic and are topologically the Cantor discontinuum. Let us describe geometrically a minimal set given by the dyadic group and a generator. A similar construction is available for the general case.

If E is the disjoint union of segments (meaning closed line segments) E_1, \dots, E_r , let E^* denote the union of segments obtained by deleting the open middle third of each of the segments E_1, \dots, E_r . Let S be a segment of length 1. Define X_0, X_1, \dots inductively as follows: $X_0 = S, X_{n+1} = X_n^*$ ($n = 0, 1, 2, \dots$). Define $X = \bigcap_{n=0}^{\infty} X_n$. The space X is the Cantor discontinuum. Denote the 2^n disjoint segments which make up X_n ($n = 0, 1, 2, \dots$) by $S(n, m)$ ($0 \leq m \leq 2^n - 1$) where $S(0, 0) = S, S(n+1, m)$ is the initial third of $S(n, m)$, and $S(n+1, m+2^{n-1})$ is the terminal third of $S(n, m)$. For each nonnegative integer n permute the segments $S(n, m)$ ($0 \leq m \leq 2^{n-1}$) cyclicly according to the second coordinate m . These permutations induce a homeomorphism ϕ of X onto itself. More exactly, if $x \in X$, then for each nonnegative integer n we have that $x \in S(n, m_n)$ for exactly one integer m_n with $0 \leq m_n \leq 2^n - 1$ and $x\phi = \bigcap_{n=0}^{\infty} S(n, m_n + 1 \pmod{2^n})$. It follows that (X, ϕ) is a regularly almost periodic mini-

mal set which we may call the dyadic minimal set. To see the group structure of X , draw the horizontal segments $S(1, 0)$ and $S(1, 1)$ under the horizontal segment $S(0, 0)$, draw downward arrows to $S(1, 0)$ and $S(1, 1)$ from $S(0, 0)$, and label the arrows 0 and 1 from left to right. Continue this process. A point of X is uniquely represented by a downward chain of arrows and therefore by a sequence of 0's and 1's. The sequences are added coordinatewise with (possibly infinite) carry-over. The generator of the dyadic group X which corresponds to ϕ is given by the sequence $(1, 0, 0, 0, \dots)$. The n -adic minimal sets are all regularly almost periodic.

In the case of the circle (1-sphere) as a minimal set under a discrete flow, the isomorphism type is characterized by an irrational number r between 0 and 1, the Poincaré rotation number, and the minimal set is isomorphic to the spin of a circle through an angle which is $r\pi$ [for example, 17]. Actually, the discrete flows on a circle appear to be presently unclassified. Let us look at one. Let $\dots, E_{-1}, E_0, E_1, \dots$ be a disjoint bisequence of closed arcs on a circle K such that $\bigcup_{n \in \mathcal{G}} E_n$ is dense in K and such that the cyclic order of $\dots, E_{-1}, E_0, E_1, \dots$ on K agrees with the cyclic order of $\dots, xr^{-1}, x, xr, \dots$ on a circle when the circle is rotated through one radian, x being a point of the circle and r denoting the rotation. Map E_n ($n \in \mathcal{G}$) homeomorphically onto E_{n+1} in the evident manner. This defines a homeomorphism of $\bigcup_{n \in \mathcal{G}} E_n$ onto itself. Since both this homomorphism and its inverse are uniformly continuous, it can be extended uniquely to a homeomorphism ϕ of K onto K . The complement of $\bigcup_{n \in \mathcal{G}} \text{int } E_n$ is a minimal set under ϕ which is the Cantor discontinuum and which is not almost periodic. The endpoints of the E_n ($n \in \mathcal{G}$) constitute a pair of doubly asymptotic orbits, that is $\rho(x\phi^n, y\phi^n)$ goes to zero with $1/n$ where x and y are the endpoints of the arc E_0 .

The almost periodic minimal sets are homogeneous since their phase spaces are groups. Is every minimal set homogeneous? The answer is no (Floyd 1949) [11]. Floyd's example may be described in geometric language (University of Pennsylvania dissertation of Joseph Auslander 1957) [3].

If E is a box $A \times B$ where A and B are segments, define E^* to be $(A_1 \times B_1) \cup (A_3 \times B) \cup (A_5 \times B_2)$ where A_1, \dots, A_5 are the consecutive equal fifths of A and B_1, B_2 are the consecutive equal halves of B . If D is a disjoint union of boxes, define D^* to be the union of the E^* where E ranges over the boxes in D . Start with the unit box X_0 and define

$$X_{n+1} = X_n^* \quad (n = 0, 1, 2, \dots), \quad X = \bigcap_{n=0}^{+\infty} X_n.$$

The boxes in X_n ($n=0, 1, 2, \dots$) are permuted just like the segments are permuted in the construction of the triadic minimal set. For example, the first three stages have boxes numbered as follows from left to right: 0; 0, 1, 2; 0, 3, 6, 1, 4, 7, 3, 5, 8. The process gives rise to a homeomorphism ϕ of X onto X . It is seen that X consists of point-components and segments, X is zero-dimensional at point-components, X is one-dimensional on segments, X is therefore not homogeneous, X is minimal under ϕ , ϕ is not almost periodic, ϕ is regularly almost periodic at some points of X but not at all points, ϕ is not recurrent and no orbits are unilaterally asymptotic.

There is also an example (F. B. Jones 1949) [14, p. 139 f.] of a minimal set under a discrete flow whose phase space is a plane one-dimensional continuum which is locally connected at some points and not locally connected at other points.

We turn now to another class of examples which we shall call *shifting flows*. Consider a nonvacuous compact metric space E and form its cartesian power $X = E^{\mathcal{g}}$ with exponent \mathcal{g} where \mathcal{g} is the set of all integers. An element of X is simply a function on \mathcal{g} to E . Provide X with its product topology, or equivalently, its point-open topology. Define a homeomorphism σ of X onto X as follows: $(x_n | n \in \mathcal{g})\sigma = (x_{n+1} | n \in \mathcal{g})$ where $x = (x_n | n \in \mathcal{g}) \in X$. The map σ is customarily called the *shift* transformation. A point x of X may be denoted by a so-called *symbolic trajectory*.

$$\begin{array}{c} \downarrow \\ \dots x_{-1} \ x_0 x_1 \dots \end{array}$$

with *index* (the arrow) which denotes the value of the point x at 0. The shift transformation changes the indexed symbolic trajectory simply by shifting the index to the next symbol on the right. Thus

$$\begin{array}{c} x \ x\sigma \\ \downarrow \ \downarrow \\ \dots x_{-2} \ x_{-1} \ x_0 \ x_1 \ x_2 \dots \end{array}$$

The notational device of a symbolic trajectory helps very considerably in the study of shifting flows.

A *symbolic flow* is defined to be a shifting flow $(E^{\mathcal{g}}, \sigma)$ where E is a finite set with more than one element. The phase space $E^{\mathcal{g}}$ of a symbolic flow $(E^{\mathcal{g}}, \sigma)$ is the Cantor discontinuum. *Symbolic dynamics* is defined to be the study of symbolic flows; it is therefore largely combinatorial in nature. A large number of papers have been concerned with symbolic dynamics [14, p. 113]. The symbolic flows are descrip-

tive of geodesic flows over certain surfaces of constant negative curvature. Marston Morse (1921) [20] defined a symbolic trajectory as follows, where $E = \{a, b\}$ with $a \neq b$. The elements a and b are called *duals* of each other. Consider the 1-block a ; dualize to form the 1-block b and suffix it; we now have a 2-block $a b$. Consider the 2-block $a b$; dualize to form the 2-block $b a$ and suffix it; we now have a 4-block $a b b a$. The next step gives the 8-block $a b b a b a a b$. The process is continued to form a ray. This ray is then reflected to the left to form a trajectory which may be indexed at any symbol. This indexed trajectory is an example of an almost periodic point under σ which is not periodic. Consequently, Morse showed for the first time that there exist everywhere nonlocally connected compact minimal sets under certain geodesic flows.

A characteristic property of symbolic flows is *expansive*. A discrete flow (X, ϕ) is said to be *expansive* provided there exists a positive real number ϵ such that if x and y are different points of the phase space X , then $\rho(x\phi^n, y\phi^n) > \epsilon$ for some integer n . It may be seen that every symbolic flow is expansive and, conversely, if a discrete flow on a zero-dimensional compact metric space is expansive, then it may be imbedded in a symbolic flow. Observe that expansive is a strong negation of isometric, or equivalently, almost periodic. Consequently, the expansive minimal sets appear to be at the other end of the spectrum from almost periodic minimal sets. It has been proved (Schwartzmann 1952; Utz 1950 related) [23; 26; 14, p. 87] that every expansive discrete flow possesses a pair of points which is positively asymptotic and a pair of points which is negatively asymptotic. This means that $x \neq y$ and $\rho(x\phi^n, y\phi^n) \rightarrow 0$ as $n \rightarrow +\infty$ or $n \rightarrow -\infty$, respectively. The asymptotic points of a minimal set are at least partially characteristic of the minimal set.

In the shifting flow $(E^{\mathcal{G}}, \sigma)$ we may take E to be the closed unit interval J and obtain the Hilbert cube $J^{\mathcal{G}}$ as phase space. We may also take E to be the circle group K and obtain the infinite toral group $K^{\mathcal{G}}$ as phase space; in this case the shift transformation is a group automorphism. An example (R. F. Williams 1955) [28] of an expansive discrete flow on a continuum may be imbedded in the flow $(K^{\mathcal{G}}, \sigma)$. It is described as the set of all bisequences $(z_n/n \in \mathcal{G})$ such that $z_n = z_{n+1}^2$ for all $n \in \mathcal{G}$; it is therefore the 2-solenoidal group and the map is a group automorphism. The shifting flows in the Hilbert cube, the infinite torus, and other spaces, appear to be worthy of further study. So far as I am aware, shifting flows have been systematically investigated only in the case of symbolic flows.

I wish now to describe briefly some recent unpublished work [3,

related] tending toward the classification of locally almost periodic minimal sets. Given a discrete flow (X, ϕ) . The map ϕ is said to be *locally almost periodic* provided that if $x \in X$ and if U is a neighborhood of x , then there exists a neighborhood V of x and a relatively dense subset A of \mathcal{I} such that $V\phi^n \subset U$ for all $n \in A$. Two points x and y of X are said to be *proximal* provided that if $\epsilon > 0$, then $\rho(x\phi^n, y\phi^n) < \epsilon$ for some $n \in \mathcal{I}$. Assume now that X is compact metric and ϕ is locally almost periodic. The relation proximal is a closed equivalence relation in X . Consider now the star-closed partition space X^* determined by the relation proximal. The discrete flow induced on X^* is almost periodic. Suppose also that X is minimal. Then X^* is minimal. Hence X^* is an almost periodic minimal set and thus a monothetic group, called the *structure group* of the original locally almost periodic minimal set. To illustrate, the structure group of Floyd's example is the triadic group, and the structure group of Jones' example in the circle group.

Just a few words now about continuous flows. In general, the definitions and theorems mentioned before apply to continuous flows as well as discrete flows. There are some exceptions, however. A minimal set under a continuous flow is necessarily connected; not so for a discrete flow. Another exception appears in the theorem (A. A. Markov 1931) [19; 14, p. 14] that a finite-dimensional minimal set under a continuous flow is a Cantor-manifold and hence has the same dimension at each point. This is not true for discrete flows: witness the example of Floyd.

The examples of continuous flows which have been most extensively studied are the geodesic flows over surfaces of constant negative curvature. Geodesic flows are defined for Riemannian manifolds of arbitrary dimension but let us think of the 2-dimensional case. Let S be a surface of class C^2 provided with a Riemannian metric of class C^2 and which is complete in the sense that each geodesic arc in S can be extended to infinite length in both directions. The phase space X consists of the unit contravariant vectors on S which we may call the unitangents (unitangent = unit tangent). Let $x \in X$ and let $t \in \mathbb{R}$. Consider the geodesic g in S which has x as a tangent. If t is positive, measure t units of arc length along g in the same direction as x from the base point of x and take the unitangent to g at the new point to be the image of x under the transition π^t . Likewise for negative t . There results a continuous flow which is called the *geodesic flow over S* . Now X is a 3-dimensional manifold which is the bundle space of a fibre bundle with the circle as fiber and the surface S as base space. The geodesic flows are dynamical systems which are determined by the Euler equations.

We state one theorem on geodesic flows which is now classical. If S is a complete 2-dimensional analytic Riemannian manifold of constant negative curvature and of finite area, then the geodesic flow over S is regionally mixing and its periodic points are dense in the space of unitangents on S . This theorem is proved again in the Colloquium volume *Topological dynamics* [14, p. 131]. A continuous flow (X, \mathcal{R}, π) is said to be *regionally mixing* provided that if U and V are nonvacuous open subsets of X , then there exists a positive real number s such that $Ut \cap V \neq \emptyset$ for $t \in \mathcal{R}$ with $|t| \geq s$.

Closely related to the geodesic flows are the horocycle flows over surfaces of constant negative curvature. They have the same phase spaces as geodesic flows but different transitions and motions. It has been shown (Hedlund 1936) [16] that for each integer $p > 1$ there exists a closed orientable surface S of constant negative curvature and of genus p such that the space X of unitangents on S is a regionally mixing minimal set under the horocycle flow over S . Here the phase spaces X are compact 3-dimensional manifolds, are not almost periodic under the horocycle flows, and, even more, are not recurrent (since regionally mixing). A continuous flow (X, \mathcal{R}, π) is said to be *recurrent* provided that if $\epsilon > 0$, then there exists an extensive subset A of \mathcal{R} such that $xt \in N_\epsilon(x)$ for all $x \in X$ and all $t \in A$.

As remarked previously, every discrete flow (X, ϕ) gives rise to a continuous flow as follows: take the cartesian product of X and the closed unit interval, and identify bases according to the map ϕ . The capped n -adic minimal sets produce the n -solenoidal minimal sets. All of these minimal sets are regularly almost periodic, and their structure groups are the n -adic groups and the n -solenoidal groups.

Consider the space X of all continuous functions f on the line group \mathcal{R} to \mathcal{R} and provide X with its compact-open topology. Define a continuous flow in X by translation of the function f : $(f(r) | r \in \mathcal{R}) \rightarrow (f(r+t) | r \in \mathcal{R})$ defines π^t for each $t \in \mathcal{R}$. It is known (Bebutov 1940) [4] that every continuous flow in a compact metric space with at most one fixed point is imbeddable in the above flow. The minimal sets under continuous flows therefore all appear in the Bebutov example. A perspicuous proof of his theorem is much to be desired. From a topological point of view, the Bebutov example has a favorable property, namely, the orbit-closure of a point f is compact if and only if f is bounded and uniformly continuous.

The consideration of function spaces leads us to the almost periodic functions of Harold Bohr. Consider the space X of all bounded uniformly continuous functions on \mathcal{R} to \mathcal{R} and provide X with its uniform topology. Define a continuous flow in X by translation of the functions. An element of X is an almost periodic function in the sense

of Bohr if and only if it is an almost periodic point under the flow. This statement shows the consistency of the terminology almost periodic. A certain fraction of the theory of almost periodic functions can be presented under the theory of continuous flows or, more generally, transformation groups, [14, pp. 40–48]. It seems that future developments will encompass more. Recent work of Tornehave (1954) [25] is suggestive in this connection.

In the foregoing discussion of flows there are several features which suggest a broader and more general development in the context of transformation groups. Some of these features are: (1) analogous theories for discrete flows and continuous flows (the famous dictum of E. H. Moore is brought to mind); (2) the study of flows themselves leads to topological groups; (3) the study of flows leads to almost periodic functions and the theory of almost periodic functions includes functions on groups.

The Colloquium volume *Topological dynamics* [14] is written from this more general point of view. Let us mention briefly two representative notions of recursion for transformation groups. Let (X, T) be a transformation group. A subset A of the phase group T is said to be *syndetic* provided that $T = AK$ for some compact subset K of T . The notion syndetic replaces the notion relatively dense for sets in \mathcal{S} and \mathcal{R} . A point x of the phase space X is said to be *almost periodic* provided that if U is a neighborhood of x , then $xA = \{xt \mid t \in A\} \subset U$ for some syndetic subset A of T . A semigroup P in T is said to be *replete* provided that P contains some bilateral translate of each compact subset of T . A subset A of T is said to be *extensive* provided that A intersects every replete semigroup in T . A point x of X is said to be *recurrent* provided that if U is a neighborhood of x , then $xA \subset U$ for some extensive subset A of T . When T is \mathcal{S} or \mathcal{R} , then these notions of recursion reduce to the customary ones for flows [14].

For the remaining moments permit me to pose a few more questions which are presently unanswered so far as I know. Questions stated for flows on metric spaces are usually meaningful also for transformation groups on more general spaces.

(1) What compact metric spaces can be minimal sets under a discrete flow? Under a continuous flow? The universal curve of Sierpinski? The universal curve of Menger? A lens space? What polyhedra? Can they be nonorientable? About all that is known for polyhedra is that, in the continuous flow case, the Euler characteristic has to vanish. This follows from the fact that a minimal set with more than one point cannot have the fixed point property [1, p. 532]. The only closed surfaces with vanishing Euler characteristic are the torus

and the Klein bottle. The torus is a minimal set under both discrete flows and continuous flows. Can the Klein bottle be a minimal set? What can be said about the homology and homotopy groups of a minimal set?

(2) More generally, what subsets of a given phase space can be minimal subsets? Does there exist a discrete flow in euclidean 3-space such that some orbit-closure is the necklace of Antoine? If so, can it be minimal? In an n -dimensional manifold it is known that a minimal subset is either the whole manifold or of dimension less than n [14, p. 14]. If the dimension is equal to $n-1$, then is it necessarily an $(n-1)$ -torus? (Question of R. W. Bass.) Bass has pointed out that a theorem of Kodaira and Abe (1940) [18] shows the answer to be affirmative for an almost periodic minimal set under a continuous flow even when merely the phase space is imbeddable in euclidean n -space.

(3) Let X be a compact uniform space which is minimal under a homeomorphism ϕ . In general, X need not be metrizable since every compact connected separable abelian group is monothetic (for example, K^J where K is the circle group and J is the unit interval). What other conditions will guarantee that X be metrizable? One instance is that ϕ be expansive (Bryant 1955) [7; 9]. Another is that ϕ be regularly almost periodic.

(4) Distal means without distinct proximal points. Is every distal minimal set necessarily almost periodic? It is known that distal and locally almost periodic are equivalent to almost periodic, even without minimality [13].

(5) The study of geodesic flows and horocycle flows now requires rather much geometry and analysis. Is it possible, in some sense, to axiomatize these flows so that they are more accessible to immediate study?

(6) In many examples of geodesic flows, symbolic flows, and horocycle flows the following properties occur together: regionally mixing and dense periodic or almost periodic points. Does regionally mixing plus some auxiliary hypothesis imply that the almost periodic points are dense?

(7) What spaces can carry an expansive homeomorphism (Question of Utz and others)? The n -cells? May such spaces be locally connected?

(8) The roles of connected, locally connected, locally euclidean, and other kinds of connectivity, in question of existence and implication?

(9) When von Neumann defined his almost periodic functions on a

group, he generalized Bochner's characterization of the Bohr almost periodic functions. This led to the notion of left and right almost periodic functions which Maak subsequently proved equivalent. (Incidentally, the theorems of topological dynamics can be quoted to establish this.) When the original Bohr definition is generalized directly to functions on topological groups, a similar dichotomy occurs but the answer as to their equivalence in this case is not clear. If the two kinds are actually equivalent, then this new definition leads again to von Neumann's functions. (See [14, p. 40 f.] for references and full statements.)

(10) There are a number of theorems in topological dynamics whose hypothesis and conclusion are meaningful for nonmetric spaces, say compact or locally compact uniform spaces, and yet the only known proof makes use of a category argument valid for metric spaces or at least first-countable spaces [14]. The question is to find a replacement for the category arguments, either in particular cases or in general, which permit the removal of countability assumptions such as metric or first-countable. Recent work of Robert Ellis (1957) [10] is very interesting and suggestive in this connection.

(11) From an abstract point of view, there is an equivalence between transformation groups and spaces of functions on groups to spaces. Briefly, the space X of functions on a group T to a space Y determines a transformation group (X, T) by translation on the functions; and a transformation group (X, T, π) determines the space $\{\pi_x | x \in X\}$ of motions in the space Z of all functions on T to X . These two constructions are not symmetric because the first produces a relatively complicated space X from a relatively simple space Y , and the second produces a more complicated space Z from an already complicated space X . To avoid the difficulty, a theorem is needed for transformation groups which extends the Bebutov theorem on a universal flow for continuous flows.

More questions are stated in [21].

REFERENCES

1. Paul Alexandroff and Heinz Hopf, *Topologie I*, Die Grundlehren der mathematischen Wissenschaften, vol. 45, Berlin, 1935; Ann Arbor, 1945.
2. Hirotada Anzai and S. Kakutani, *Bohr compactifications of a locally compact abelian group II*, Proceedings of the Imperial Academy, Tokyo, vol. 19 (1943) pp. 533-539.
3. Joseph Auslander, *Mean-L-stable systems*, Dissertation, University of Pennsylvania, 1957.
4. M. Bebutov, *Sur les systèmes dynamiques dans l'espace des fonctions continues*, Comptes Rendus, Doklady Akad. Nauk SSSR, vol. 27 (1940) pp. 904-906.
5. G. D. Birkhoff, *Dynamical systems*, Amer. Math. Soc. Colloquium Publications, vol. 9, New York, 1927.

6. ———, *Collected mathematical papers*, 3 vols., American Mathematical Society, New York, 1950.
7. B. F. Bryant, *Unstable self-homeomorphisms of a compact space*, Dissertation, Vanderbilt University, 1954.
8. ———, *A note on unstable homeomorphisms*, Amer. Math. Monthly vol. 61 (1954) p. 509 (abstract).
9. ———, *A note on uniform spaces*, Amer. Math. Monthly, vol. 62 (1955) p. 529 (abstract).
10. Robert Ellis, *Locally compact transformation groups*, Duke Math. J. vol. 24 (1957) pp. 119–126.
11. E. E. Floyd, *A nonhomogeneous minimal set*, Bull. Amer. Math. Soc. vol. 55 (1949) pp. 957–960.
12. W. H. Gottschalk, *Almost periodic points with respect to transformation semi-groups*, Ann. of Math. vol. 47 (1946) pp. 762–766.
13. ———, *Characterizations of almost periodic transformation groups*, Proc. Amer. Math. Soc. vol. 7 (1956) pp. 709–712.
14. W. H. Gottschalk and G. A. Hedlund, *Topological dynamics*, Amer. Math. Soc. Colloquium Publications, vol. 36, Providence, 1955.
15. P. R. Halmos and H. Samelson, *On monothetic groups*, Proc. Nat. Acad. Sci. U.S.A., vol. 28 (1942) pp. 254–258.
16. G. A. Hedlund, *Fuchsian groups and transitive horocycles*, Duke Math. J. vol. 2 (1936) pp. 530–542.
17. E. R. van Kampen, *The topological transformations of a simple closed curve into itself*, Amer. J. Math. vol. 57 (1935) pp. 142–152.
18. Kunihiko Kodaira and Makoto Abe, *Über zusammenhängende kompakte abelsche Gruppen*, Proceedings of the Imperial Academy, Tokyo, vol. 16 (1940) pp. 167–172.
19. A. A. Markoff, *Sur une propriété générale des ensembles minimaux de M. Birkhoff*, C. R. Acad. Sci. Paris vol. 193 (1931) pp. 823–825.
20. Marston Morse, *Recurrent geodesics on a surface of negative curvature*, Trans. Amer. Math. Soc. vol. 22 (1921) pp. 84–100.
21. V. V. Nemyckiĭ, *Topological problems of the theory of dynamical systems*, Uspehi Mat. Nauk (N.S.) vol. 4 (1949) pp. 91–152 (Russian); Amer. Math. Soc. Translation, number 103.
22. V. V. Nemyckiĭ and V. V. Stepanov, *Qualitative theory of differential equations*, Moscow-Leningrad, 1947; 2d ed., 1949, Russian; English translation of 2d ed., Princeton University Press, to appear in 1959.
23. Sol Schwartzman, *On transformation groups*, Dissertation, Yale University, 1952.
24. P. A. Smith, *Periodic and nearly periodic transformations*, Lectures in Topology, Ann Arbor, 1941, pp. 159–190.
25. Hans Tornehave, *On almost periodic movements*, Det Kongelige Danske Videnskabernes Selskab, Matematisk-fysiske Meddelelser, vol. 28, no. 13, 1954.
26. W. R. Utz, *Unstable homeomorphisms*, Proc. Amer. Math. Soc. vol. 1 (1950) pp. 769–774.
27. G. T. Whyburn, *Analytic topology*, Amer. Math. Soc. Colloquium Publications, vol. 28, New York, 1942.
28. R. F. Williams, *A note on unstable homeomorphisms*, Proc. Amer. Math. Soc. vol. 6 (1955) pp. 308–309.

UNIVERSITY OF PENNSYLVANIA

BIBLIOGRAPHY

ANZAI, HIROTADA

1. (With S. Kakutani) *Bohr compactifications of a locally compact abelian group I*, Proceedings of the Imperial Academy, Tokyo, vol. 19 (1943), pp. 476–480.
2. (With S. Kakutani) *Bohr compactifications of a locally compact abelian group II*, Proceedings of the Imperial Academy, Tokyo, vol. 19 (1943), pp. 533–539.

ARENS, RICHARD

1. *A topology for spaces of transformations*, Annals of Mathematics, vol. 47 (1946), pp. 480–495.
2. *Topologies for homeomorphism groups*, American Journal of Mathematics, vol. 68 (1946), pp. 593–610.

AYRES, W. L.

1. *On transformations having periodic properties*, Fundamenta Mathematicae, vol. 33 (1939), pp. 95–103.

BAUM, J. D.

1. *An equicontinuity condition for transformation groups*, Proceedings of the American Mathematical Society, vol. 4 (1953), pp. 656–662.

BEBUTOFF, M.

1. *Sur les systèmes dynamiques dans l'espace des fonctions continues*, Comptes Rendus (Doklady) de l'Académie des Sciences de l'URSS, vol. 27 (1940), pp. 904–906.

BELTRAMI, E.

1. *Teoria fondamentale degli spazii di curvatura costante*, Annali di Matematica, ser. 2, vol. 2, pp. 232–255.

BESICOVITCH, A. S.

1. *A problem on topological transformations of the plane*, Fundamenta Mathematicae, vol. 28 (1937), pp. 61–65.
2. *A problem on topological transformations of the plane. II*, Proceedings of the Cambridge Philosophical Society, vol. 47 (1951), pp. 38–45.

BIANCHI, L.

1. *Vorlesungen über Differentialgeometrie*, Leipzig, 1899.

BIRKHOFF, G. D.

1. *Collected Mathematical Papers*, vols. 1, 2, 3, New York, 1950.
2. *Dynamical Systems*, American Mathematical Society Colloquium Publications, vol. 9, 1927.

BOCHNER, S.

1. *Beiträge zur Theorie der fastperiodischen Funktionen I*, Mathematische Annalen, vol. 96 (1926), pp. 119–147.

BOHR, HARALD

1. *Collected Mathematical Works*, vols. 1, 2, 3, Copenhagen, 1952.

BOURBAKI, N.

Éléments de mathématique, Première Partie, *Les structures fondamentales de l'analyse*.

1. Livre I, *Théorie des ensembles (Fascicule de résultats)*, Actualités scientifiques et industrielle 846, Paris, 1939.
2. Livre III, *Topologie générale*, Chapitre I, *Structures topologiques*, & Chapitre II, *Structures uniformes*, Actualités scientifiques et industrielles 858, Paris, 1940.
3. Livre III, *Topologie générale*, Chapitre III, *Groupes topologiques (Théorie élémentaire)*, Actualités scientifiques et industrielles 916, Paris, 1942.
4. Livre III, *Topologie générale*, Chapitre X, *Espaces fonctionnels*, Actualités scientifiques et industrielles 1084, Paris, 1949.

CAMERON, R. H.

1. *Almost periodic transformations*, Transactions of the American Mathematical Society, vol. 36 (1934), pp. 276–291.

CARATHÉODORY, C.

1. *Über den Wiederkehrsatz von Poincaré*, Sitzungsberichte der Preussischen Akademie der Wissenschaften, 1919, pp. 580–584.

ELLIS, ROBERT

1. *Continuity and homeomorphism groups*, Proceedings of the American Mathematical Society, vol. 4 (1953), pp. 969–973.

ENGEL, F.

1. See Lie [1].

ERDÖS, P.

1. (With A. H. Stone) *Some remarks on almost periodic transformations*, Bulletin of the American Mathematical Society, vol. 51 (1945), pp. 126–130.

FLOYD, E. E.

1. *A nonhomogeneous minimal set*, Bulletin of the American Mathematical Society, vol. 55 (1949), pp. 957–960.

FORD, L. R.

1. *Automorphic Functions*, New York, 1929.

FORT, M. K., Jr.

1. *A note on equicontinuity*, Bulletin of the American Mathematical Society, vol. 55 (1949), pp. 1098–1100.

FRANKLIN, PHILIP

1. *Almost periodic recurrent motions*, Mathematische Zeitschrift, vol. 30 (1929), pp. 325–331.

FRICKE, R.

1. (With F. Klein) *Automorphe Functionen*, vol. I, Leipzig, 1897.

GARCIA, MARIANO

1. (With G. A. Hedlund) *The structure of minimal sets*, Bulletin of the American Mathematical Society, vol. 54 (1948), pp. 954–964.

GOTTSCHALK, W. H.

1. *An investigation of continuous mappings with almost periodic properties*, Dissertation, University of Virginia, 1944.
2. *Powers of homeomorphisms with almost periodic properties*, Bulletin of the American Mathematical Society, vol. 50 (1944), pp. 222–227.
3. *Orbit-closure decompositions and almost periodic properties*, Bulletin of the American Mathematical Society, vol. 50 (1944), pp. 915–919.
4. *A note on pointwise nonwandering transformations*, Bulletin of the American Mathematical Society, vol. 52 (1946), pp. 488–489.
5. (With G. A. Hedlund) *Recursive properties of transformation groups*, Bulletin of the American Mathematical Society, vol. 52 (1946), pp. 637–641.
6. *Almost periodic points with respect to transformation semi-groups*, Annals of Mathematics, vol. 47 (1946), pp. 762–766.
7. *Almost periodicity, equi-continuity and total boundedness*, Bulletin of the American Mathematical Society, vol. 52 (1946), pp. 633–636.
8. *Recursive properties of transformation groups II*, Bulletin of the American Mathematical Society, vol. 54 (1948), pp. 381–383.
9. *Transitivity and equicontinuity*, Bulletin of the American Mathematical Society, vol. 54 (1948), pp. 982–984.
10. (With G. A. Hedlund) *The dynamics of transformation groups*, Transactions of the American Mathematical Society, vol. 65 (1949), pp. 348–359.
11. *The extremum law*, Proceedings of the American Mathematical Society, vol. 3 (1952), p. 631.

HALL, D. W.

1. (With J. L. Kelley) *Periodic types of transformations*, Duke Mathematical Journal, vol. 8 (1941), pp. 625-630.

HALMOS, P. R.

1. (With H. Samelson) *On monothetic groups*, Proceedings of the National Academy of Sciences of the United States, vol. 28 (1942), pp. 254-258.
2. *Measure Theory*, New York, 1950.
3. (With H. E. Vaughan) *The marriage problem*, American Journal of Mathematics, vol. 72 (1950), pp. 214-215.

HARTMAN, PHILIP

1. (With Aurel Wintner) *Integrability in the large and dynamical stability*, American Journal of Mathematics, vol. 65 (1943), pp. 273-278.

HAUSDORFF, FELIX

1. *Grundzüge der Mengenlehre*, 1st ed., reprinted, Chelsea, New York, 1949.

HEDLUND, G. A.

1. See Morse [3].
2. *The dynamics of geodesic flows*, Bulletin of the American Mathematical Society, vol. 45 (1939), pp. 241-260.
3. See Morse [4].
4. *Sturmian minimal sets*, American Journal of Mathematics, vol. 66 (1944), pp. 605-620.
5. See Gottschalk [5].
6. See Garcia [1].
7. See Gottschalk [10].

HILMY, HEINRICH

1. *Sur une propriété des ensembles minima*, Comptes Rendus (Doklady) de l'Académie des Sciences de l'URSS, vol. 14 (1937), pp. 261-262.
2. *Sur la théorie des ensembles quasi-minimaux*, Comptes Rendus (Doklady) de l'Académie des Sciences de l'URSS, vol. 15 (1937), pp. 113-116.
3. *Sur les mouvements des systèmes dynamiques qui admettent "l'incompressibilité" des domaines*, American Journal of Mathematics, vol. 59 (1937), pp. 803-808.
4. *Sur les théorèmes de récurrence dans la dynamique générale*, American Journal of Mathematics, vol. 61 (1939), pp. 149-160.

HOPF, E.

1. *Zwei Sätze über den wahrscheinlichen Verlauf der Bewegungen dynamischer Systeme*, Mathematische Annalen, vol. 103 (1930), pp. 710-719.
2. *Ergodentheorie*, Berlin, 1937.
3. *Statistik der geodätischen Linien in Mannigfaltigkeiten negativer Krümmung*, Berichte über die Verhandlungen der sächsischen Akademie der Wissenschaften zu Leipzig, Mathematisch-Physischen Klasse, vol. 91 (1939), pp. 261-304.

HOPF, H.

1. *Zum Clifford-Kleinschen Raumproblem*, Mathematische Annalen, vol. 95 (1926), pp. 313-339.
2. (With W. Rinow) *Ueber den Begriff der vollständigen differentialgeometrische Fläche*, Commentarii Mathematici Helvetici, vol. 3 (1931), pp. 209-225.

HUREWICZ, WITOLD

1. (With Henry Wallman) *Dimension Theory*, Princeton Mathematical Series No. 4, 1941.

KAKUTANI, S.

1. See Anzai [1].
2. See Anzai [2].

KELLEY, J. L.

1. See Hall [1].

VON KERÉKJÁRTÓ, B.

1. *Sur les similitudes de l'espace*, Comptes Rendus de l'Académie des Sciences Paris, vol. 198 (1934), p. 1345.

KLEIN, F.

1. See Fricke [1].

KOEBE, P.

1. *Riemannsche Mannigfaltigkeiten und nicht euklidische Raumformen*, Sitzungsberichte der Preussischen Akademie der Wissenschaften, (1927), pp. 164–196; (1928), pp. 345–442; (1929), pp. 414–457; (1930), pp. 304–364, 504–541; (1931), pp. 506–534.

KURATOWSKI, CASIMIR

1. *Topologie I*, Warsaw, 1933.

KUROSCH, A.

1. *Theory of Groups*, Moscow, 1944, (Russian).

LIE, SOPHUS

1. (With F. Engel) *Theorie der Transformationgruppen*, Leipzig, 1888.

LÖBEL, F.

1. *Die überall regulären unbegrenzten Flächen fester Krümmung*, Dissertation, Tübingen, 1927.
2. *Beispiele geschlossener dreidimensionaler Clifford-Kleinscher Räume negativer Krümmung*, Berichte über die Verhandlungen der sächsischen Akademie der Wissenschaften zu Leipzig, Mathematisch-Physischen Klasse, vol. 83 (1931), pp. 167–174.

MAAK, WILHELM

1. *Eine neue Definition der fastperiodischen Funktionen*, Abhandlungen aus dem Mathematischen Seminar der Hansischen Universität, vol. 11 (1936), pp. 240–244.
2. *Abstrakte fastperiodische Funktionen*, Abhandlungen aus dem Mathematischen Seminar der Hansischen Universität, vol. 11 (1936), pp. 367–380.

MAIER, A. G.

1. *On central trajectories and a problem of Birkhoff*, Matematičeskii Sbornik (N. S.), vol. 26 (68), (1950), pp. 266–290.

MARKOFF, A. A.

1. *Sur une propriété générale des ensembles minimaux de M. Birkhoff*, Comptes Rendus de l'Académie des Sciences, Paris, vol. 193 (1931), pp. 823–825.
2. *Stabilität im Liapounoffschen Sinne und Fastperiodizität*, Mathematische Zeitschrift, vol. 36 (1933), pp. 708–738.

MONTGOMERY, DEANE

1. *Almost periodic transformation groups*, Transactions of the American Mathematical Society, vol. 42 (1937), pp. 322–332.
2. *Pointwise periodic homeomorphisms*, American Journal of Mathematics, vol. 59 (1937), pp. 118–120.
3. *Measure preserving homeomorphisms at fixed points*, Bulletin of the American Mathematical Society, vol. 51 (1945), pp. 949–953.

MORSE, MARSTON

1. *A one-to-one representation of geodesics on a surface of negative curvature*, American Journal of Mathematics, vol. 43 (1921), pp. 33–51.
2. *Recurrent geodesics on a surface of negative curvature*, Transactions of the American Mathematical Society, vol. 22 (1921), pp. 84–100.
3. (With G. A. Hedlund) *Symbolic dynamics*, American Journal of Mathematics, vol. 60 (1938), pp. 815–866.
4. (With G. A. Hedlund) *Symbolic dynamics II. Sturmian trajectories*, American Journal of Mathematics, vol. 62 (1940), pp. 1–42.

NIEMYTZKI, V. V.

1. (With V. V. Stepanoff) *Qualitative Theory of Differential Equations*, Moscow-Leningrad, 1st ed. 1947, 2nd ed. 1949, (Russian).

2. *Topological problems of the theory of dynamical systems*, Uspehi Matematicheskikh Nauk (N. S.), vol. 4 (1949), pp. 91–153, (Russian); American Mathematical Society, Translation Number 103.
- VON NEUMANN, J.
1. *Almost periodic functions in a group*. I, Transactions of the American Mathematical Society, vol. 36 (1934), pp. 445–492.
- OXTOBY, J. C.
1. *Note on transitive transformations*, Proceedings of the National Academy of Sciences of the United States, vol. 23 (1937), pp. 443–446.
 2. (With S. M. Ulam) *Measure preserving homeomorphisms and metric transitivity*, Annals of Mathematics, vol. 42 (1941), pp. 874–920.
 3. *Ergodic sets*, Bulletin of the American Mathematical Society, vol. 58 (1952), pp. 116–136.
- RINOW, W.
1. See H. Hopf [2].
- SALENIUS, TAUNO
1. *Über dreidimensionale geschlossene Räume konstanter negativer Krümmung*, Den 11te Skandinaviske Matematikerkongress, Trondheim, 1949, pp. 107–112.
- SAMELSON, HANS
1. See Halmos [1].
- SCHWARTZMAN, S.
1. *On transformation groups*, Dissertation, Yale University, 1952.
- SEIDEL, W.
1. (With J. L. Walsh) *On approximation by euclidean and non-euclidean translations of an analytic function*, Bulletin of the American Mathematical Society, vol. 47 (1941), pp. 916–920.
- SMITH, P. A.
1. *Periodic and nearly periodic transformations*, Lectures in Topology, Ann Arbor, 1941, pp. 159–190.
- STEPANOFF, V. V.
1. (With A. Tychonoff) *Über die Räume der fastperiodischen Funktionen*, Recueil Mathématique, Nouvelle Série, vol. 41 (1934), pp. 166–178.
 2. See Niemytzki [2].
- STONE, A. H.
1. See Erdős [1].
- TYCHONOFF, A.
1. See Stepanoff [1].
- ULAM, S. M.
1. See Oxtoby [2].
- UTZ, W. R.
1. *Unstable homeomorphisms*, Proceedings of the American Mathematical Society, vol. 1 (1950), pp. 769–774.
- VAUGHAN, H. E.
1. See Halmos [3].
- WALLMAN, H.
1. See Hurewicz [1].
- WALSH, J. L.
1. See W. Seidel [1].
- WEIL, ANDRÉ
1. *L'intégration dans les groupes topologiques et ses applications*, Paris, 1940.
- WHYBURN, G. T.
1. *Analytic Topology*, American Mathematical Society Colloquium Publications, vol. 28, New York, 1942.

WILLIAMS, C. W.

1. *Recurrence and incompressibility*, Proceedings of the American Mathematical Society, vol. 2 (1951), pp. 798-806.

WINTNER, A.

1. See Hartman [1].

ZIPPIN, LEO

1. *Transformation groups*, Lectures in Topology, Ann Arbor, 1941, pp. 191-221.
Mathematics in U. S. S. R. for thirty years, 1917-1947, Moscow-Leningrad, 1948, pp. 508-517.

A limited number of copies of the *Bibliography for Dynamical Topology*, Fifth Edition, September 1972, compiled by Walter H. Gottschalk, Wesleyan University, is available (at cost) from the Department of Mathematics, Wesleyan University, Middletown, Connecticut 06457. This Bibliography contains 178 pages and lists 2475 numbered items.

INDEX

- Admissible orbit pairs, 10.38
- Admissible subsets of T , 3.12
- Almost periodic, 3.38
- Asymptotic, 10.21
- Asymptotic geodesics, 13.08
- Averaging process, 4.75
- Axiom
 - continuity, 1.01
 - homomorphism, 1.01
 - identity, 1.01
- Axis of a hyperbolic transformation, 13.05
- Baire subset, 9.11
- Base, 3.29
- Bisequence, 12.03
- Bisequence space, 12.08
- Block, 12.03
- Block, n -, 12.04
- Bounded, 14.02
- Bounded set of mappings, 11.11
- Cantor-manifold, 2.16
- Center, 7.17
- Centralizer of A in G , 9.31
- Compactification of a topological group, 4.47
- Compactive, 8.11
- Composition, 4.74
- Contained in, 12.04
- Cylinder homeomorphism, 14.03
- Decomposition, 1.37
- Discrete, 3.15
- Discrete in M , 13.41
- Dual, 12.27
- Dyadic tree, 14.15
- Envelope
 - x -, 2.08
- Equicontinuous set of mappings, 11.07
- Expansive, 10.31
- Extension
 - left S -, 1.43
 - right S -, 1.43
 - Q_x -, 12.29
- Extensive in T , 3.37
- Fixed point, 3.06, 10.33
 - positive, 13.05
 - negative, 13.05
- Flow, 1.30
 - geodesic, 13.10
 - geodesic partition, 13.25
 - horocycle, 13.12
 - horocycle partition, 13.31
- Fundamental region, 13.45
- Generative topological group, 6.01
- Group
 - motion preserving, 9.28
 - orbit preserving, 9.30
 - phase, 1.01
 - transition, 1.08
- Homeomorphism
 - motion preserving, 9.26
 - orbit preserving, 9.29
 - periodic, 10.33
 - pointwise periodic, 10.33
- Homeomorphism group, 1.14
 - discrete, 1.14
 - topological, 1.12
 - total, 1.14
- Horocycle, 13.07
- Hyperbolic
 - area, 13.01
 - circle, 13.06
 - distance, 13.05
 - length, 13.01
 - line, 13.03
 - plane, 13.01
 - ray, 13.03
- Indexed, left, 12.46
- Indexed, right, 12.46
- Intrinsic property, 1.03
- Invariant set, 1.22
- Isochronous, 3.38
 - weakly, 3.38
- Isometry with axis, 13.05
- Isomorphism
 - topological, 1.02
 - uniform, 1.02
- Length (of a block), 12.04
- Limit-entire, 13.26
- Limit-partial, 13.26
- Limit point
 - P -, 6.33
 - α -, 10.15
 - ω -, 10.15

- Limit set, 13.13
 - P -, 6.33
 - α -, 10.15
 - ω -, 10.15
- Locally bounded set of mappings, 11.30
- Mean, 4.81
 - α -, 4.78
- Minimal
 - orbit-closure, 2.11
 - S -, 2.11
 - totally, 2.27
- Mixing, regionally, 9.02
- Mobile, 13.22
- Motion-projection, 1.08
- Motion-space, 1.08
- Motion
 - x -, 1.08
- Nonseparated, 10.02, 10.38
- Normalizer, 9.31
- Orbit, 1.26, 10.12
- Orbit-closure, 1.26
 - minimal, 2.11
- Paddle motion, 13.05
- Partition, 1.27
- Period
 - of a homeomorphism, 10.33
 - of a point, 10.33
 - of T , 3.04
 - of T at x , 3.02
- Periodic, 3.06
 - almost, 3.38
 - at x , 3.06
 - pointwise, 3.06
 - regularly almost, 3.38
 - weakly almost, 3.38
- Periodic point, under homeomorphism, 10.33
- Permutation of a set, 11.15
- Topological group
 - monothetic, 4.49
 - solenoidal, 4.49
- Phase
 - group, 1.01
 - projection, 1.01
 - space, 1.01
- Point at infinity
 - of a horocycle, 13.07
 - of an h -ray, 13.03
- Points at infinity of an h -line, 13.03
- Projection
 - motion, 1.08
 - phase, 1.01
 - transition, 1.08
- Property
 - intrinsic topological, 1.03
 - intrinsic uniform, 1.03
- Ray
 - left, 12.02
 - right, 12.02
- Recurrent, 3.38, 10.18
- Recursive, 3.13
 - uniformly, 3.42, 3.44, 3.47, 3.50
 - weakly, 3.42, 3.44, 3.47, 3.50
- Regularly almost periodic, 3.38
- Replete in T , 3.37
- Representation, A -, 12.42
- Restriction, 1.32
- Reverse, 12.04
- Rotor, 13.32
- Saturation, 1.34
- Semigroup, 2.05
- Semiorbit, 10.12
- Separated, 10.02, 10.38
- Sequence
 - left, 12.03
 - right, 12.03
- Shift transformation, 12.11
- Similar, 12.04
- Space product, 1.49
- Star, \mathcal{A} -, 1.34
- Star-closed, 1.35
- Star-indexed, 2.35
- Star-open, 1.35
- Subblock, 12.04
- Subgroup restriction, 1.32
- Subsequence
 - left, 12.04
 - right, 12.04
- Subspace restriction, 1.32
- Symbol class, 12.01
- Symbolic flow, 12.11
- Syndetic, 2.02, 2.03
- Topology
 - bilateral compact-index, of a set of permutations, 11.34
 - bilateral compact-open, of a set of permutations, 11.44
 - bilateral space-index, of a set of permutations, 11.16
 - compact-index, of a set of permutations, 11.34
 - inverse compact-index, of a set of permutations, 11.34
 - inverse compact-open, of a set of permutations, 11.44

- inverse space-index, of a set of permutations, 11.16
- point-index, of a set of mappings, 11.24
- space-index, of a set of mappings, 11.01
- space-index, of a set of permutations, 11.16
- Total power, 4.74
- Total homeomorphism group, 1.14
- Totally minimal, 2.27
- Totally noncompactive, 8.12
- Trace, 2.40
- Transformation group, 1.01
 - bilateral, of T , 1.51
 - discrete, 1.14
 - effective, 1.09
 - equicontinuous, 1.52
 - equicontinuous at x , 1.52
 - functional, over (X, T, π) to Y , 1.68
 - φ -inverse partition, 1.42
 - left, of T , 1.51
 - left functional, over T to Y , 1.63
 - left, of T/S induced by T , 1.56
 - left, of S induced by T under φ , 1.58
 - left uniform functional, over T to Y , 1.62
 - partition, 1.39
 - Φ -orbit partition, 1.40
 - Φ -orbit-closure partition, 1.41
 - topological, 1.01
 - right, of T , 1.51
 - right functional, over T to Y , 1.63
 - right, of S induced by T under φ , 1.58
 - right, of $T \setminus S$ induced by T , 1.56
 - right uniform functional, over T to Y , 1.62
 - uniform functional, over (X, T, π) to Y , 1.66
 - uniformly continuous, 4.36
 - uniformly equicontinuous, 1.52
- Transformation subgroup, 1.32
- Transient, 13.28
- Transition
 - group, 1.08
 - projection, 1.08
 - t -, 1.08
- Transitive, 9.02
- Translate, n -, 12.04
- Uniform convergence at a point, 9.34
- Uniformity
 - bilateral compact-index, of a set of permutations, 11.34
 - bilateral space-index, of a set of permutations, 11.16
 - compact-index, of a set of mappings, 11.24
 - compact-index, of a set of permutations, 11.34
 - inverse compact-index, of a set of permutations, 11.34
 - inverse space-index, of a set of permutations, 11.16
 - partition, 2.34
 - point-index, of a set of mappings, 11.24
 - space-index, of a set of mappings, 11.01
 - space-index, of a set of permutations, 11.16
- Uniformly equicontinuous set of mappings, 11.08
- Union of a sequence of blocks, 12.05
- Unitangent, 13.10, 13.48
 - space, 13.10, 13.48
- Universally transitive, 9.02
- Weakly recursive, 3.13, 3.42, 3.44, 3.47, 3.50

ISBN 0-8218-1036-7



9 780821 810361

COLL/36

AMS *on the Web*
www.ams.org