

# Subject Index

- adjoint transfer factor, 58
- admissible, 549
- admissible invariant distribution, 389
- anisotropic, 517
- automorphic representation, vii
  
- Bernstein center, 388
- bilinear forms, 515
  
- central character of a parameter, 306
- compatible families
  - decomposable, 151
  - global, 151
  - local, 57
- Coxeter-Dynkin diagram, 517
- cuspidal function
  - global, 19
  - local, 59
- cuspidal lift, 352
  
- discrete part of the trace formula, 124
- dual group, 1
- duality operator, 389
  
- elliptic, 12
- elliptic element
  - in  $T(G)$ , 157
- elliptic endoscopic datum, 11, 133
- elliptic representation
  - of  $\Lambda_F$ , 7
- embedded eigenvalues, 187
- endoscopic datum, 11, 52, 130
- $\varepsilon$ -pair, 254
- $\varepsilon$ -parameter, 254
- exponents, 318
- extended pure inner twist, 542
  
- fundamental lemma, 53
  
- Galois cohomology, 514
- generalized Ramanujan conjecture, 25
  
- generic parameter
  - in  $\tilde{\mathcal{F}}$ , 256
- generic representation, 110
- global intertwining relation, 177
- globally  $(B, \chi)$ -generic, 483
  
- Hasse principle, 513, 524, 525
- Hecke type, 129
  - for  $\mathcal{F}$ , 152
  - for  $\tilde{\mathcal{E}}_{\text{ell}}(N)$ , 151
  
- inertia subgroup, 43
  - wild, 43
- infinitesimal character, 68
- inner form, 514
- inner twist, 514
- intertwining operator
  - normalized, 83, 96
  - unnormalized, 96
- invariant orbital integral, 52
- isobaric automorphic representation, 3, 22
- isomorphism, 514
- isomorphism of endoscopic data, 12
  
- $L$ -equivalence, 4
- $L$ -group, 1
- $L$ -homomorphism, 2
- $L$ -packet, 4
  - tempered, 42
- $\lambda$ -factor, 90
- Langlands group (global), ix, 3
- Levi embedding, 57
- Levi subset, 59
- local intertwining relation
  - for  $\tilde{G}$ , 108
  - for  $\tilde{G}(N)$ , 118
  - for  $G$ , 100
- locally  $(B, \chi)$ -generic, 483
- locally  $G$ -relevant, 526

- locally relevant, 131
- LSK conjecture, 53
- multiset, 42
- normalized character, 470
- normalized stable character, 471
- orthogonality relations, 354, 477
- parabolic subset, 59
  - standard, 390
  - pseudo- $z$ -embedding, 544
  - pseudocoefficient, 312, 368
  - pure inner twist, 540
- quadratic parameter, 396
- $R$ -group, 100, 154
- Rankin-Selberg convolution
  - global, 22
  - local, 17
- regular automorphic representation of
  - $GL(N)$ , 504
- regular Hecke family, 504
- root number
  - orthogonal, 96
  - symplectic, 96
- $S$ -packet ( $S$ -set, set over  $S$ ), 41
- self-dual representation
  - of  $\Lambda_F$ , 7
- self-intertwining operator, 96
- simple parameter, 29
- skew-symmetric square  $L$ -function, 49
- Speh character, 431
- spherical fundamental lemma, 143
- spherical parameter, 304
- spherical representation, 304
- stabilization, 134
- stable linear form, 53
- stable multiplicity formula, 171
- stable orbital integral, 53
- standard representation
  - of  $G(F)$ , 155
  - of  $\tilde{G}(N, F)$ , 66
  - of  $L_{F, \text{reg}}^*$ , 507
  - of  $\tilde{L}_{F, \text{reg}}^*$ , 511
  - of  $\mathcal{L}_\psi$ , 216
- strongly isomorphic, 537
- strongly regular, 52
- supercuspidal representation, 18
- symmetric inner twist, 552
- symmetric orbit  $\mathfrak{o}_\alpha$ , 225
- symmetric square  $L$ -function, 49
- $\tau$ -equivalent, 160
- tamely ramified parameter, 396
- Tate-Nakayama pairing, 541
- the discrete part of the trace formula, 124
- the packet  $\tilde{\Sigma}_\psi$ , 388
- the three exceptional cases, 333
- Tits indices, 515, 517
- transfer factor
  - global, 131
  - local, 52
- twisted cuspidal function, 317
- twisted endoscopic group, 11
- unnormalized intertwining operator, 82
- unramified group, 53
- virtual character, 430
- weighted fundamental lemma, 135
- Weil group, 3
- Weyl character formula, 477
- Whittaker datum, 55
- Whittaker functional, 62, 110
- Whittaker integral, 111
- Whittaker model, 110
- $(\mathbb{Z}/2\mathbb{Z})$ -symmetry, 482, 502
- $z$ -embedding, 542
- $z$ -extension, 542
- Zelevinsky conjecture, 390