

# Contents

<b>1</b>	<b>Introduction</b> . . . . .	<b>1</b>
<b>2</b>	<b>Some background on (locally) smooth operators</b> . . . . .	<b>15</b>
<b>3</b>	<b>A limiting absorption principle for interacting, massless Dirac operators</b>	<b>23</b>
<b>4</b>	<b>On the absence of eigenvalues for interacting, massless Dirac operators</b>	<b>37</b>
<b>5</b>	<b>The Green's functions of <math>H_0(m)</math> and <math>H_0</math></b> . . . . .	<b>41</b>
<b>6</b>	<b>Trace ideal properties of <math>F_1(\cdot)(H_0 - zI_{[L^2(\mathbb{R}^n)]^N})^{-1}F_2(\cdot)</math> and <math>\langle \cdot \rangle^{-\delta}(H_0 - zI_{[L^2(\mathbb{R}^n)]^N})^{-1}\langle \cdot \rangle^{-\delta}</math></b> . . . . .	<b>49</b>
<b>7</b>	<b>Powers of resolvents and trace ideals</b> . . . . .	<b>63</b>
<b>8</b>	<b>The spectral shift function: Abstract facts</b> . . . . .	<b>69</b>
<b>9</b>	<b>Representing <math>\xi(\cdot; S, S_0)</math> in terms of regularized Fredholm determinants</b>	<b>81</b>
<b>10</b>	<b>Analysis of <math>\text{Im}(F_{H, H_0}(\lambda + i0))</math>, <math>\lambda \in \mathbb{R}</math></b> . . . . .	<b>91</b>
<b>11</b>	<b>Analysis of <math>G_{H, H_0}</math></b> . . . . .	<b>119</b>
<b>12</b>	<b>Analysis of <math>\xi(\cdot; H, H_0)</math> and an application to the Witten index for a class of non-Fredholm operators</b> . . . . .	<b>145</b>
<b>A</b>	<b>Some remarks on block matrix operators</b> . . . . .	<b>157</b>
<b>B</b>	<b>Asymptotic results for Hankel functions</b> . . . . .	<b>163</b>
	B.1 Asymptotics of $H_\nu^{(1)}(\zeta)$ as $ \zeta  \rightarrow \infty$ . . . . .	164
	B.2 Asymptotics of $H_\nu^{(1)}(\zeta)$ as $ \zeta  \rightarrow 0$ . . . . .	170
<b>C</b>	<b>Expansions and estimates of the free Dirac Green's matrix</b> . . . . .	<b>175</b>
<b>D</b>	<b>A product formula for modified Fredholm determinants</b> . . . . .	<b>191</b>

<b>E Notational conventions</b> . . . . .	199
<b>References</b> . . . . .	203