
Preface

The main goal of this book is to introduce beginning graduate students to analytic number theory. In addition, large parts of it are suitable for advanced undergraduate students with a good grasp of analytic techniques.

Throughout, the emphasis has been put on exposing the main ideas rather than providing the most general results known. Any student wishing to do serious research in analytic number theory should broaden and deepen their knowledge by consulting some of the several excellent research-level books on the subject. Examples include: the books of Davenport [31] and of Montgomery-Vaughan [146] for classical multiplicative number theory; Tenenbaum's book [172] for probabilistic number theory and the saddle-point method; the book by Iwaniec-Kowalski [114] for the general theory of L -functions, of modular forms and of exponential sums; Montgomery's book [144] for the harmonic analytic aspects of analytic number theory; and the book of Friedlander-Iwaniec [59] for sieve methods.

Using the book

The book borrows the structure of Davenport's masterpiece *Multiplicative Number Theory* with several short- to medium-length chapters. Each chapter is accompanied by various exercises. Some of them aim to exemplify the concepts discussed, while others are used to guide the students to self-discover certain more advanced topics. A star next to an exercise indicates that its solution requires total mastery of the material.

The contents of the book are naturally divided into six parts as indicated in the table of contents. The first two parts study elementary and classical complex-analytic methods. They could thus serve as the manual for an

introductory graduate course to analytic number theory. The last three parts of the book are devoted to the theory of sieves: Part 4 presents the basic elements of the theory of the small sieve, whereas Part 5 explores the method of bilinear sums and develops the large sieve. These techniques are then combined in Part 6 to study the spacing distribution of prime numbers and prove some of the recent spectacular results about small and large gaps between primes. Finally, Part 3 studies multiplicative functions and the anatomy of integers, and serves as a bridge between the complex-analytic techniques and the more elementary theory of sieves. Topics from it could be presented either in the end of an introductory course to analytic number theory (Chapter 13 most appropriately), or in the beginning of a more advanced course on sieves (the most relevant material is contained in Chapters 14 and 15, as well as in Theorem 16.1).

Certain portions of the book can be used as a reference for an undergraduate course. More precisely, Chapters 1–8 can serve as the core of such a course, followed by a selection of topics from Chapters 14, 15, 17 and 21.

A short guide to the main theorems of the book. Below is a list of the main results proven and of their prerequisites.

Chebyshev's and Mertens' estimates are presented in Chapters 2 and 3, respectively. Their proofs rest on the material contained in Part 1.

The landmark *Prime Number Theorem* is proven in Chapter 8. Understanding it requires a good grasp of all preceding chapters.

The *Siegel-Walfisz theorem*, which is a uniform version of the Prime Number Theorem for arithmetic progressions, is presented in Chapter 12. Its proof builds on all of the material preceding it.

The *Landau-Selberg-Delange method* is a key tool in the study of multiplicative functions. It is presented in Chapter 13. Appreciating its proof requires a firm understanding of Chapters 1–8 for the main analytic tools, as well as of Chapter 12 for dealing with uniformity issues.

The foundations of *probabilistic number theory* are explained in Chapters 15 and 16, where the *Erdős-Kac theorem* and the *Sathe-Selberg theorem* are proven. The main prerequisites can be found in Part 1 and in Chapter 14. In addition, Chapter 13 is needed for the Sathe-Selberg theorem.

The *Fundamental Lemma of Sieve Theory* is proven in Chapter 19. Its proof uses ideas and techniques from Part 1 and Chapters 14–17.

Vinogradov's method, one of the foundations of modern analytic number theory, is presented in Chapter 23. It builds on the material of Chapters 1–12 and 19.

The *Hardy-Littlewood circle method* is presented in Chapter 24. It is used to detect additive patterns among the primes and, more specifically, to count ternary arithmetic progressions all of whose members are primes.

The *Bombieri-Vinogradov theorem*, often called the “Generalized Riemann Hypothesis on average”, is established in Chapter 26. Understanding its proof requires mastery of Vinogradov’s method (Chapter 23) and of the *large sieve* (Chapter 25).

Linnik’s theorem provides a very strong bound on the least prime in an arithmetic progression. It is proven in Chapter 27 and its prerequisites are Chapters 1–12, 17–20, 22–23 and 25.

The breakthrough of Zhang-Maynard-Tao about the existence of infinitely many *bounded gaps between primes* is presented in Chapter 28. Its proof requires a firm understanding of the Fundamental Lemma of Sieve Theory (Chapter 19), of Selberg’s sieve (Chapter 21) and of the Bombieri-Vinogradov theorem (Chapter 26).

The recent developments about *large gaps between primes* of Ford-Green-Konyagin-Tao and Maynard are presented in Chapter 29. Understanding them necessitates knowledge of the same concepts as the proof of the existence of bounded gaps between primes, with the addition of the results on smooth numbers presented in Chapters 14 and 16.

Maier discovered in 1985 that the distribution of prime numbers has certain unexpected irregularities. His results are presented in Chapter 30 and they assume knowledge of Linnik’s theorem (and of its prerequisites), as well as of Buchstab’s function (see Chapter 14 and, more precisely, Theorem 14.4).

Acknowledgments

Many people have helped me greatly in many different ways in writing this book.

I am indebted to Leo Goldmakher and James Maynard, with whom I discussed the contents of the book extensively at various stages of the writing process. In addition, an early version of the manuscript was used as a teaching reference by Wei Ho at the University of Michigan, and by Leo Goldmakher at Williams College. I am grateful to them and their students for the valuable feedback they provided.

I am obliged to Martin Čech, Tony Haddad, Youcef Mokrani, Alexis Leroux-Lapierre, Joëlle Matte, Kunjakanan Nath, Stelios Sachpazis, Simon St-Amant, Jeremie Turcotte and Peter Zenz, who patiently studied earlier versions of the book, catching various errors and providing many excellent comments.

I have had very useful mathematical conversations with Sandro Bettin, Brian Conrey, Chantal David, Ben Green, Adam Harper, Jean Lagacé and K. Soundararajan on certain topics of the book; I am grateful to them for their astute remarks. Furthermore, I would like to thank the anonymous reviewers for their suggestions that helped me improve the exposition of the ideas in the manuscript, especially those related to the bilinear methods presented in Part 5.

I am indebted to Kevin Ford and Andrew Granville, who taught me analytic number theory. Their influence is evident throughout the book.

A special thanks goes to Ina Mette, Marcia Almeida and Becky Rivard for guiding me through the publishing process. I would also like to thank Brian Bartling and Barbara Beeton for their assistance with several typesetting questions, as well as Alexis Leroux-Lapierre for his help with designing the figures that appear in the book.

Last but not least, I would like to thank my wife Jennifer Crisafulli for her love, support and companionship. This book could not have been written without her and I wholeheartedly dedicate it to her.

Funding. During the writing process, I was supported by the Natural Sciences and Engineering Research Council of Canada (Discovery Grant 2018-05699) and by the Fonds de recherche du Québec—Nature et technologies (projet de recherche en équipe—256442). Part of the book writing took place during my visit at the Mathematical Sciences Research Institute of Berkeley in the Spring of 2017 (funded by the National Science Foundation under Grant No. DMS-1440140), at the University of Oxford in the Spring of 2019 (funded by Ben Green’s Simons Investigator Grant 376201) and at the University of Genova in June 2019 (funded by the Istituto Nazionale di Alta Matematica “Francesco Severi”). I would like to thank my hosts for their support and hospitality.