

Subject Index

- abelian part, 187
- abelian scheme, 185
- abelian variety, 186
- absolute Galois group, 82
- absolutely irreducible, 85
- additive group, 175
- additive reduction, 17
- analytic expression
 - elliptic curve, 71
 - modular curve, 72
 - modular form, 73
- augmentation, 132

- Cartier divisor, 183
- Cartier dual, 177
- character group, 186
- closed condition, 182
- coarse moduli scheme, 39
- commutative group scheme, 175
- complete, 133
- complete intersection, 131
- completed group algebra, 144
- conductor, 17, 98
- connected component, 103
- constant group scheme, 175
- cyclic group scheme, 176
- cyclotomic character, 83

- deformation ring, 125
- degree
 - morphism of elliptic curves, 23
- discriminant, 14, 27
- elliptic curve, 26

- elliptic curve over a field, 13
- equivalent of liftings, 122
- étale covering, 178
- étale local, 179
- étale sheaf, 179
- étale sheafification, 179

- fine moduli scheme, 39
- finite G_K -module, 86
- finite étale commutative group scheme, 176
- finite flat commutative group scheme, 176
- finite morphism, 173
- Fitting ideal, 156
- flat covering, 178
- flat local, 179
- flat sheaf, 178
- free, 180
- Frobenius conjugacy class, 82
- Frobenius substitution, 82
- full Hecke algebra, 136
- functor
 - over \mathcal{O} , 159
 - over S , 37
- G -torsor, 179
- Galois representation
 - defined by the N -torsion points of E , 89
- $\Gamma_0(N)$ -structure, 41
- $\Gamma(N)$ -structure, 45
- geometric Frobenius, 23
- good, 3, 10

- $G_{\mathbf{Q}}$ -module, 96
- Galois representation, 98
- good reduction, 16
- Hecke algebra
 - over \mathcal{O} , 128
 - over \mathbf{Q} , 57
- Hecke module, 136
- Hecke operator, 57
- inertia group, 97, 180
- irreducible, 84
- j -invariant, 36
- ℓ -adic cyclotomic character, 83
- ℓ -adic representation, 83
 - associated with f , 92
 - determined by E , 89
- L -function
 - of a modular form, 64
 - of an elliptic curve, 22
- lifting, 122, 131
- local
 - over S , 179
- modular, 6
 - ℓ -adic representation, 92
 - elliptic curve, 65
 - mod ℓ representation, 93
 - of level N , 8
- mod ℓ representation, 83
- morphism of elliptic curves, 22
- multiplication-by- N morphism, 23, 28
- multiplicative, 186
- multiplicative group, 175
- multiplicative reduction, 17
- multiplier, 134
- Néron model, 107
- open subscheme defined by \mathcal{P} , 179
- ordinary, 98
- primary form, 5, 62
- primitive form, 64
 - over K , 70
- profinitely generated complete local
 - \mathcal{O} -algebra, 122
- q -expansion, 61
- q -expansion principle, 61
- quasi-finite morphism, 173
- quotient, 177
- reduced Hecke algebra
 - over \mathcal{O} , 128
 - over \mathbf{Q} , 68
- reducible, 84
- RTM -triple, 133
- Selmer group, 139
- semistable
 - elliptic curve, 17
 - Galois representation, 98
- semistable model, 31
- smooth model, 27
- smooth of relative dimension d , 171
- stable reduction, 16
- subfunctor, 160
- surjection
 - of RTM -triples, 133
- Tate module, 24, 89, 103, 186
- torus, 186
- torus part, 187
- type, 161
- type \mathcal{D}_{Σ} , 123
- unipotent, 186
- universal element, 39, 160
- unramified, 82