

CONTENTS

1. Introduction	1
1.1. Main assumptions	1
1.2. Main results	4
Acknowledgments	6
2. Lower bound	7
2.1. Main statement	7
2.2. Preliminary lemmas	7
2.3. Proof of Proposition 2.1	10
3. Quasimode construction – Type V1	17
3.1. Local coordinates and approximate operator	17
3.2. The formal construction	19
3.3. Quasimode and remainder	22
4. Quasimode construction – Type V2	23
5. V1 potentials: 1D operators	27
5.1. Motivation	27
5.2. Realization on the entire real line	28
5.3. Dirichlet realization in the half-line	31
5.4. Application to 2D separable operators	37
6. V1 potentials: 2D simplification	41
6.1. The operator \mathcal{B}_ε	41
6.2. Large $ \sigma $ simplification	42
6.3. Resolvent estimates for \mathcal{B}_ε	47
6.4. Curvature effects	53
6.5. A linear potential estimate	60
7. Simplified operators: V2 potentials	61
7.1. Definition and preliminary estimates	61
7.2. Large $ \sigma $ simplification	63
7.3. Second simplified operator	65
8. Upper bound	71

8.1. Goals and notation	71
8.2. Refined partition of unity	72
8.3. Localized resolvent estimates	74
8.4. Approximate resolvent error	78
8.5. Eigenvalue existence	89
A. Examples of potentials satisfying (1.4)	91
Bibliography	93