
Chapter 7

Homotopy and the Fundamental Group

The group G will be called the fundamental group of the manifold V .

HENRI POINCARÉ, 1895

The properties of a topological space that we have developed so far have depended on the choice of topology, the collection of open sets. Taking a different tack, we introduce a different structure, algebraic in nature, associated to a space together with a choice of base point (X, x_0) . This structure will allow us to bring to bear the power of algebraic arguments. The fundamental group was introduced by Poincaré in his investigations of the action of a group on a manifold [66].

The first step in defining the fundamental group is to study more deeply the relation of homotopy between continuous functions $f_0: X \rightarrow Y$ and $f_1: X \rightarrow Y$. Recall that f_0 is *homotopic* to f_1 , denoted $f_0 \simeq f_1$, if there is a continuous function (a *homotopy*)

$$H: X \times [0, 1] \rightarrow Y \text{ with } H(x, 0) = f_0(x) \text{ and } H(x, 1) = f_1(x).$$

The choice of notation anticipates an interpretation of the homotopy —if we write $H(x, t) = f_t(x)$, then a homotopy is a deformation of

the mapping f_0 into the mapping f_1 through the family of mappings f_t .

Theorem 7.1. *The relation $f \simeq g$ is an equivalence relation on the set $\text{Hom}(X, Y)$ of continuous mappings from X to Y .*

Proof. Let $f: X \rightarrow Y$ be a given mapping. The homotopy $H(x, t) = f(x)$ is a continuous mapping $H: X \times [0, 1] \rightarrow Y$ and so $f \simeq f$.

If $f_0 \simeq f_1$ and $H: X \times [0, 1] \rightarrow Y$ is a homotopy between f_0 and f_1 , then the mapping $H': X \times [0, 1] \rightarrow Y$ given by $H'(x, t) = H(x, 1 - t)$ is continuous and a homotopy between f_1 and f_0 , that is, $f_1 \simeq f_0$.

Finally, for $f_0 \simeq f_1$ and $f_1 \simeq f_2$, suppose that $H_1: X \times [0, 1] \rightarrow Y$ is a homotopy between f_0 and f_1 , and $H_2: X \times [0, 1] \rightarrow Y$ is a homotopy between f_1 and f_2 . Define the homotopy $H: X \times [0, 1] \rightarrow Y$ by

$$H(x, t) = \begin{cases} H_1(x, 2t), & \text{if } 0 \leq t \leq 1/2, \\ H_2(x, 2t - 1), & \text{if } 1/2 \leq t \leq 1. \end{cases}$$

Since $H_1(x, 1) = f_1(x) = H_2(x, 0)$, the piecewise definition of H gives a continuous function (Theorem 4.4). By definition, $H(x, 0) = f_0(x)$ and $H(x, 1) = f_2(x)$ and so $f_0 \simeq f_2$. \square

We denote the equivalence class under homotopy of a mapping $f: X \rightarrow Y$ by $[f]$ and the set of homotopy classes of maps between X and Y by $[X, Y]$. If $F: W \rightarrow X$ and $G: Y \rightarrow Z$ are continuous mappings, then the sets $[X, Y]$, $[W, X]$, and $[Y, Z]$ are related.

Proposition 7.2. *Continuous mappings $F: W \rightarrow X$ and $G: Y \rightarrow Z$ induce well-defined functions $F^*: [X, Y] \rightarrow [W, Y]$ and $G_*: [X, Y] \rightarrow [X, Z]$ by $F^*([h]) = [h \circ F]$ and $G_*([h]) = [G \circ h]$ for $[h] \in [X, Y]$.*

Proof. We need to show that if $h \simeq h'$, then $h \circ F \simeq h' \circ F$ and $G \circ h \simeq G \circ h'$. Fixing a homotopy $H: X \times [0, 1] \rightarrow Y$ with $H(x, 0) = h(x)$ and $H(x, 1) = h'(x)$, then the desired homotopies are $H_F(w, t) = H(F(w), t)$ and $H_G(x, t) = G(H(x, t))$. \square

To a space X we associate a space particularly rich in structure, the mapping space of paths in X , $\text{map}([0, 1], X)$. Recall that $\text{map}([0, 1], X)$ is the set of continuous mappings $\text{Hom}([0, 1], X)$ with

the compact-open topology. The space $\text{map}([0, 1], X)$ has the following properties.

(1) X embeds into $\text{map}([0, 1], X)$ by associating to each point $x \in X$ the *constant path* $c_x(t) = x$ for all $t \in [0, 1]$.

(2) Given a path $\lambda: [0, 1] \rightarrow X$, we can *reverse* the path by composing with $t \mapsto 1 - t$. Let $\lambda^{-1}(t) = \lambda(1 - t)$.

(3) Given a pair of paths $\lambda, \mu: [0, 1] \rightarrow X$ for which $\lambda(1) = \mu(0)$, we can *compose* paths by

$$\lambda * \mu(t) = \begin{cases} \lambda(2t), & \text{if } 0 \leq t \leq 1/2, \\ \mu(2t - 1), & \text{if } 1/2 \leq t \leq 1. \end{cases}$$

Thus, for certain pairs of paths λ and μ , we obtain a new path $\lambda * \mu \in \text{map}([0, 1], X)$.

Composition of paths is always defined when we restrict to a certain subspace of $\text{map}([0, 1], X)$.

Definition 7.3. Suppose X is a space and $x_0 \in X$ is a choice of base point in X . The **space of based loops** in X , denoted $\Omega(X, x_0)$, is the subspace of $\text{map}([0, 1], X)$,

$$\Omega(X, x_0) = \{\lambda \in \text{map}([0, 1], X) \mid \lambda(0) = \lambda(1) = x_0\}.$$


Composition of loops determines a binary operation $*$: $\Omega(X, x_0) \times \Omega(X, x_0) \rightarrow \Omega(X, x_0)$.

We restrict the notion of homotopy when applied to the space of based loops in X in order to stay in that space during the deformation.

Definition 7.4. Given two based loops λ and μ , a **loop homotopy** between them is a homotopy of paths $H: [0, 1] \times [0, 1] \rightarrow X$ with $H(t, 0) = \lambda(t)$, $H(t, 1) = \mu(t)$, and $H(0, s) = H(1, s) = x_0$. That is, for each $s \in [0, 1]$, the path $t \mapsto H(t, s)$ is a loop at x_0 .

The relation of loop homotopy on $\Omega(X, x_0)$ is an equivalence relation; the proof follows the proof of Theorem 7.1. We denote the set of equivalence classes under loop homotopy by $\pi_1(X, x_0) = [\Omega(X, x_0)]$, a notation for the first of a family of such sets, to be explained later. As it turns out, $\pi_1(X, x_0)$ enjoys some remarkable properties.

Theorem 7.5. *Composition of loops induces a group structure on $\pi_1(X, x_0)$ with identity element $[c_{x_0}(t)]$ and inverses given by $[\lambda]^{-1} = [\lambda^{-1}]$.*




Proof. We begin by showing that composition of loops induces a well-defined binary operation on the homotopy classes of loops. Given $[\lambda]$ and $[\mu]$, we then define $[\lambda]*[\mu] = [\lambda*\mu]$. Suppose that $[\lambda] = [\lambda']$ and $[\mu] = [\mu']$. We must show that $\lambda*\mu \simeq \lambda'*\mu'$. If $H: [0, 1] \times [0, 1] \rightarrow X$ is a loop homotopy between λ and λ' and $H': [0, 1] \times [0, 1] \rightarrow X$ a loop homotopy between μ and μ' , then form $H'': [0, 1] \times [0, 1] \rightarrow X$ defined by

$$H''(t, s) = \begin{cases} H(2t, s), & \text{if } 0 \leq t \leq 1/2, \\ H'(2t - 1, s), & \text{if } 1/2 \leq t \leq 1. \end{cases}$$


Since $H''(0, s) = H(0, s) = x_0$ and $H''(1, s) = H'(1, s) = x_0$, H'' is a loop homotopy. Also $H''(t, 0) = \lambda*\mu(t)$ and $H''(t, 1) = \lambda'*\mu'(t)$, and the binary operation is well defined on equivalence classes of loops.

We next show that $*$ is associative. Notice that $(\lambda * \mu) * \nu \neq \lambda * (\mu * \nu)$; we only get 1/4 of the interval for λ in the first product and 1/2 of the interval in the second product. We define the explicit homotopy after its picture, which makes the point more clearly:




$$H(t, s) = \begin{cases} \lambda(4t/(1+s)), & \text{if } 0 \leq t \leq (s+1)/4, \\ \mu(4t-1-s), & \text{if } (s+1)/4 \leq t \leq (s+2)/4, \\ \nu\left(1 - \frac{4(1-t)}{(2-s)}\right), & \text{if } (s+2)/4 \leq t \leq 1. \end{cases}$$

The class of the constant map, $e(t) = c_{x_0}(t) = x_0$, gives the identity for $\pi_1(X, x_0)$. To see this, we show, for all $\lambda \in \Omega(X, x_0)$, that $\lambda * e \simeq \lambda \simeq e * \lambda$ via loop homotopies. This is accomplished in the case $\lambda \simeq e * \lambda$ by the homotopy:



$$F(t, s) = \begin{cases} x_0, & \text{if } 0 \leq t \leq s/2, \\ \lambda((2t-s)/(2-s)), & \text{if } s/2 \leq t \leq 1. \end{cases}$$

The case $\lambda \simeq \lambda * e$ is similar. Finally, inverses are constructed by using the reverse loop $\lambda^{-1}(t) = \lambda(1-t)$. To show that $\lambda * \lambda^{-1} \simeq e$ consider the homotopy:



$$G(t, s) = \begin{cases} \lambda(2t), & \text{if } 0 \leq t \leq s/2, \\ \lambda(s), & \text{if } s/2 \leq t \leq 1 - (s/2), \\ \lambda(2-2t), & \text{if } 1 - (s/2) \leq t \leq 1. \end{cases}$$

The homotopy resembles the loop, moving out for a while, waiting a little, and then shrinking back along itself. The proof that $\lambda^{-1} * \lambda \simeq e$ is similar. \square

Definition 7.6. The group $\pi_1(X, x_0)$ is called the **fundamental group of X at the base point x_0** .

Suppose x_1 is another choice of basepoint for X . If X is path-connected, there is a path $\gamma: [0, 1] \rightarrow X$ with $\gamma(0) = x_0$ and $\gamma(1) = x_1$. This path induces a mapping $u_\gamma: \pi_1(X, x_0) \rightarrow \pi_1(X, x_1)$ by $[\lambda] \mapsto [\gamma^{-1} * \lambda * \gamma]$, that is, follow γ^{-1} from x_1 to x_0 , then follow λ around and back to x_0 , then follow γ back to x_1 , all giving a loop based at x_1 . Notice

$$\begin{aligned} u_\gamma([\lambda] * [\mu]) &= u_\gamma([\lambda * \mu]) \\ &= [\gamma^{-1} * \lambda * \mu * \gamma] \\ &= [\gamma^{-1} * \lambda * \gamma * \gamma^{-1} * \mu * \gamma] \\ &= [\gamma^{-1} * \lambda * \gamma] * [\gamma^{-1} * \mu * \gamma] = u_\gamma([\lambda]) * u_\gamma([\mu]). \end{aligned}$$

Thus u_γ is a homomorphism. The mapping $u_{\gamma^{-1}}: \pi_1(X, x_1) \rightarrow \pi_1(X, x_0)$ is an inverse, since $[\gamma * (\gamma^{-1} * \lambda * \gamma) * \gamma^{-1}] = [\lambda]$. Thus $\pi_1(X, x_0)$ is isomorphic to $\pi_1(X, x_1)$ whenever x_0 is joined to x_1 by a path. Though it is a bit of a lie, we write $\pi_1(X)$ for a space X that is path-connected since any choice of basepoint gives an isomorphic group. In this case, $\pi_1(X)$ denotes an isomorphism class of groups.

Following Proposition 7.2, a continuous function $f: X \rightarrow Y$ induces a mapping

$$f_*: \pi_1(X, x_0) \rightarrow \pi_1(Y, f(x_0)), \text{ given by } f_*([\lambda]) = [f \circ \lambda].$$

In fact, f_* is a homomorphism of groups:

$$\begin{aligned} f_*([\lambda] * [\mu]) &= f_*([\lambda * \mu]) = [f \circ (\lambda * \mu)] \\ &= [(f \circ \lambda) * (f \circ \mu)] = [f \circ \lambda] * [f \circ \mu] \\ &= f_*([\lambda]) * f_*([\mu]). \end{aligned}$$

Furthermore, when we have continuous mappings $f: X \rightarrow Y$ and $g: Y \rightarrow Z$, we obtain $f_*: \pi_1(X, x_0) \rightarrow \pi_1(Y, f(x_0))$ and $g_*: \pi_1(Y, f(x_0)) \rightarrow \pi_1(Z, g \circ f(x_0))$. Observe that

$$g_* \circ f_*([\lambda]) = g_*([f \circ \lambda]) = [g \circ f \circ \lambda] = (g \circ f)_*([\lambda]),$$

so we have $(g \circ f)_* = g_* \circ f_*$. It is evident that the identity mapping $\text{id}: X \rightarrow X$ induces the identity homomorphism of groups $\pi_1(X, x_0) \rightarrow \pi_1(X, x_0)$. We can summarize these observations by the (post-1945) remark that π_1 is a functor from pointed spaces and pointed maps to groups and group homomorphisms. Since we are focusing on classical notions in topology (pre-1935) and category theory was christened later, we will not use this language in what follows. For an introduction to this framework see [53].

The behavior of the induced homomorphisms under composition has the following consequence.

Corollary 7.7. *The fundamental group is a topological invariant of a space. That is, if $f: X \rightarrow Y$ is a homeomorphism, then the groups $\pi_1(X, x_0)$ and $\pi_1(Y, f(x_0))$ are isomorphic.*

Proof. Suppose $f: X \rightarrow Y$ has continuous inverse $g: Y \rightarrow X$. Then $g \circ f = \text{id}_X$ and $f \circ g = \text{id}_Y$. It follows that $g_* \circ f_* = \text{id}$ and $f_* \circ g_* = \text{id}$ on $\pi_1(X, x_0)$ and $\pi_1(Y, f(x_0))$, respectively. Thus f_* and g_* are group isomorphisms. \square

Before we do some calculations we derive a few more formal properties of the fundamental group. In particular, what conditions imply $\pi_1(X) = \{e\}$, and how does the fundamental group behave under the formation of subspaces, products, and quotients?

Definition 7.8. A subspace $A \subset X$ is a **retract** of X if there is a continuous function, the retraction, $r: X \rightarrow A$ for which $r(a) = a$ for all $a \in A$. The subset $A \subset X$ is a **deformation retraction** if A is a retract of X and the composition $i \circ r: X \rightarrow A \hookrightarrow X$ is homotopic to the identity on X via a homotopy that fixes A , that is, there is a homotopy $H: X \times [0, 1] \rightarrow X$ with


$$H(x, 0) = x, H(x, 1) = r(x), \text{ and } H(a, t) = a$$

for all $a \in A$ and all $t \in [0, 1]$.

Proposition 7.9. *If $A \subset X$ is a retract with retraction $r: X \rightarrow A$, then the inclusion $i: A \rightarrow X$ induces an injective homomorphism $i_*: \pi_1(A, a) \rightarrow \pi_1(X, a)$ and the retraction induces a surjective homomorphism $r_*: \pi_1(X, a) \rightarrow \pi_1(A, a)$.*

Proof. The composite $r \circ i: A \rightarrow X \rightarrow A$ is the identity mapping on A and so the composite $r_* \circ i_*: \pi_1(A, a) \rightarrow \pi_1(X, a) \rightarrow \pi_1(A, a)$ is the identity on $\pi_1(A, a)$. If $i_*([\lambda]) = i_*([\lambda'])$, then $[\lambda] = r_* i_*([\lambda]) = r_* i_*([\lambda']) = r_* i_*([\lambda']) = [\lambda']$, and so the homomorphism i_* is injective. If $[\lambda] \in \pi_1(A, a)$, then $r_*(i_*([\lambda])) = [\lambda]$ and so r_* is onto. \square

Examples. Represent the Möbius band M by glueing the left and right edges of $[0, 1] \times [0, 1]$ with a twist (Chapter 4). Let $A = \{[(t, \frac{1}{2})] \mid 0 \leq t \leq 1\} \subset M$ be the circle in the middle of the band. After the identification, A is homeomorphic to S^1 . Define the map $r: M \rightarrow A$ by projecting straight down or up to this line, that is, $[(t, s)] \mapsto [(t, \frac{1}{2})]$. It is easy to see that r is continuous and $r|_A = \text{id}_A$ so we have a retract. Thus the composite $r_* \circ i_*: \pi_1(S^1) \rightarrow \pi_1(M) \rightarrow \pi_1(S^1)$ is the identity on $\pi_1(S^1)$.



For any space X , the inclusion followed by projection

$$X \cong X \times \{0\} \hookrightarrow X \times [0, 1] \rightarrow X$$

is the identity and so X is a retract of $X \times [0, 1]$. In fact, X is a deformation retraction via the deformation $H: X \times [0, 1] \times [0, 1] \rightarrow X \times [0, 1]$ given by $H(x, t, s) = (x, ts)$: when $s = 1$, $H(x, t, 1) = (x, t)$ and for $s = 0$ we have $H(x, t, 0) = (x, 0)$.

Recall that a subset K of \mathbb{R}^n is **convex** if whenever \mathbf{x} and \mathbf{y} are in K , then for all $t \in [0, 1]$, $t\mathbf{x} + (1-t)\mathbf{y} \in K$. If $K \subset \mathbb{R}^n$ is convex, let $\mathbf{x}_0 \in K$. Then K is a deformation retraction of the one-point subset $\{\mathbf{x}_0\}$ by the homotopy $H(\mathbf{x}, t) = t\mathbf{x}_0 + (1-t)\mathbf{x}$. When $t = 0$ we have $H(\mathbf{x}, 0) = \mathbf{x}$ and when $t = 1$, $H(\mathbf{x}, 1) = \mathbf{x}_0$. The retraction $K \rightarrow \{\mathbf{x}_0\}$ is thus a deformation of the identity on K . Examples of convex subsets of \mathbb{R}^n include \mathbb{R}^n itself, any open ball $B(\mathbf{x}, \epsilon)$, and the boxes $[a_1, b_1] \times \cdots \times [a_n, b_n]$.

More generally, there is always the retract $\{x_0\} \hookrightarrow X \rightarrow \{x_0\}$, which leads to the trivial homomorphisms of groups $\{e\} \rightarrow \pi_1(X, x_0) \rightarrow \{e\}$. This retract is not always a deformation retract. We call a space **contractible** when it is a deformation retract of one of its points.

Deformation retracts give isomorphic fundamental groups.

Theorem 7.10. *If A is a deformation retract of X , then the inclusion $i: A \rightarrow X$ induces an isomorphism $i_*: \pi_1(A, a) \rightarrow \pi_1(X, a)$.*

Proof. From the definition of a deformation retract, the composite $i \circ r: X \rightarrow A \hookrightarrow X$ is homotopic to id_X via a homotopy fixing the points in A , that is, there is a homotopy $H: X \times [0, 1] \rightarrow X$ with $H(x, 0) = i \circ r(x)$, $H(x, 1) = x$, and $H(a, t) = a$ for all $t \in [0, 1]$. We show that $i_* \circ r_*([\lambda]) = [\lambda]$. In fact we show a little more:

Lemma 7.11. *If $f, g: (X, x_0) \rightarrow (Y, y_0)$ are continuous functions, homotopic through basepoint preserving maps, then $f_* = g_*: \pi_1(X, x_0) \rightarrow \pi_1(Y, y_0)$.*

Proof. Suppose there is a homotopy $G: X \times [0, 1] \rightarrow Y$ with $G(x, 0) = f(x)$, $G(x, 1) = g(x)$, and $G(x_0, t) = y_0$ for all $t \in [0, 1]$. Consider a loop based at x_0 , $\lambda: [0, 1] \rightarrow X$, and the compositions $f \circ \lambda$, $g \circ \lambda$, and $G \circ (\lambda \times \text{id}): [0, 1] \times [0, 1] \rightarrow Y$:

$$\begin{aligned} G(\lambda(s), 0) &= f \circ \lambda(s), \\ G(\lambda(s), 1) &= g \circ \lambda(s), \\ G(\lambda(0), t) &= G(\lambda(1), t) = y_0 \text{ for all } t \in [0, 1]. \end{aligned}$$

Thus $f_*[\lambda] = [f \circ \lambda] = [g \circ \lambda] = g_*[\lambda]$. Hence $f_* = g_*: \pi_1(X, x_0) \rightarrow \pi_1(Y, y_0)$. \square

A deformation retract gives a basepoint preserving homotopy between $i \circ r$ and id_X , so we have $\text{id} = i_* \circ r_*: \pi_1(X, a) \rightarrow \pi_1(X, a)$. By Proposition 7.9, we already know i_* is injective; i_* is surjective because for $[\lambda]$ any class in $\pi_1(X, a)$, one has $[\lambda] = i_*(r_*([\lambda]))$. \square

Examples. A convex subset of \mathbb{R}^n is a deformation retract of any point \mathbf{x}_0 in the set. It follows from $\pi_1(\{\mathbf{x}_0\}) = \{e\}$, that for any convex subset $K \subset \mathbb{R}^n$, $\pi_1(K, \mathbf{x}_0) = \{e\}$. Of course, this includes

$\pi_1(\mathbb{R}^n, \mathbf{0}) = \{e\}$. Next consider $\mathbb{R}^n - \{\mathbf{0}\}$. The $(n-1)$ -sphere $S^{n-1} \subset \mathbb{R}^n$ is a deformation retract of $\mathbb{R}^n - \{\mathbf{0}\}$ as follows: Let

$$F: (\mathbb{R}^n - \{\mathbf{0}\}) \times [0, 1] \rightarrow \mathbb{R}^n - \{\mathbf{0}\}$$

be given by

$$F(\mathbf{x}, t) = (1-t)\mathbf{x} + t\frac{\mathbf{x}}{\|\mathbf{x}\|}.$$

Here $F(\mathbf{x}, 0) = \mathbf{x}$ and $F(\mathbf{x}, 1) = \mathbf{x}/\|\mathbf{x}\| \in S^{n-1}$. By Theorem 7.10,

$$\pi_1(\mathbb{R}^n - \{\mathbf{0}\}, \mathbf{x}_0/\|\mathbf{x}_0\|) \cong \pi_1(S^{n-1}, \mathbf{x}_0/\|\mathbf{x}_0\|).$$


A space X is said to be **simply-connected** (or *1-connected*) if it is path-connected and $\pi_1(X) = \{e\}$. Any convex subset of \mathbb{R}^n or, more generally, any contractible space is simply-connected. Furthermore, simple connectivity is a topological property.

Theorem 7.12. *Suppose $X = U \cup V$, where U and V are open, simply-connected subspaces and $U \cap V$ is path-connected. Then X is simply-connected.*

Proof. Choose a point $x_0 \in U \cap V$ as basepoint. Let $\lambda: [0, 1] \rightarrow X$ be a loop based at x_0 . Since λ is continuous, $\{\lambda^{-1}(U), \lambda^{-1}(V)\}$ is an open cover of the compact space $[0, 1]$. The Lebesgue Lemma gives points $0 = t_0 < t_1 < t_2 < \cdots < t_n = 1$ with $\lambda([t_{i-1}, t_i]) \subset U$ or V . We can join x_0 to $\lambda(t_i)$ by a path γ_i . Define for $i \geq 1$,

$$\lambda_i(s) = \lambda((t_i - t_{i-1})s + t_{i-1}), \quad 0 \leq s \leq 1,$$

for the path along λ joining $\lambda(t_{i-1})$ to $\lambda(t_i)$.



Then $\lambda \simeq \lambda_1 * \lambda_2 * \cdots * \lambda_n$ and, furthermore,

$$\lambda \simeq (\lambda_1 * \gamma_1^{-1}) * (\gamma_1 * \lambda_2 * \gamma_2^{-1}) * (\gamma_2 * \lambda_3 * \gamma_3^{-1}) * \cdots * (\gamma_{n-1} * \lambda_n).$$

Each $\gamma_i * \lambda_{i+1} * \gamma_{i+1}^{-1}$ lies in U or V . Since U and V are simply-connected, each of these loops is homotopic to the constant map. Thus $\lambda \simeq c_{x_0}$. It follows that $\pi_1(X, x_0) \cong \{e\}$. \square

Corollary 7.13. $\pi_1(S^n) \cong \{e\}$ for $n \geq 2$.

Proof. We can decompose S^n as a union of $U = \{(r_0, r_1, \dots, r_n) \in S^n \mid r_n > -1/4\}$ and $V = \{(r_0, r_1, \dots, r_n) \in S^n \mid r_n < 1/4\}$. By stereographic projection from each pole, we can establish that U and V are homeomorphic to an open disk in \mathbb{R}^n , which is convex. The intersection $U \cap V$ is homeomorphic to $S^{n-1} \times (-1/4, 1/4)$, which is path-connected when $n \geq 2$. \square

Since $S^{n-1} \subset \mathbb{R}^n - \{\mathbf{0}\}$ is a deformation retract, we have proven:

Corollary 7.14. $\pi_1(\mathbb{R}^n - \{\mathbf{0}\}) \cong \{e\}$ for $n \geq 3$.

In Chapter 8 we will consider the case $\pi_1(S^1)$ in detail.

We next consider the fundamental group of a product $X \times Y$.

Theorem 7.15. *Let (X, x_0) and (Y, y_0) be pointed spaces. Then $\pi_1(X \times Y, (x_0, y_0))$ is isomorphic to $\pi_1(X, x_0) \times \pi_1(Y, y_0)$, the direct product of these two groups.*

Recall that if G and H are groups, the direct product $G \times H$ has underlying set the cartesian product of G and H and binary operation $(g_1, h_1) \cdot (g_2, h_2) = (g_1 g_2, h_1 h_2)$.

Proof. Recall from Chapter 4 that a mapping $\lambda: [0, 1] \rightarrow X \times Y$ is continuous if and only if $\text{pr}_1 \circ \lambda: [0, 1] \rightarrow X$ and $\text{pr}_2 \circ \lambda: [0, 1] \rightarrow Y$ are continuous. If λ is a loop at (x_0, y_0) , then $\text{pr}_1 \circ \lambda$ is a loop at x_0 and $\text{pr}_2 \circ \lambda$ is a loop at y_0 . We leave it to the reader to prove that

- 1) If $\lambda \simeq \lambda': [0, 1] \rightarrow X \times Y$, then $\text{pr}_i \circ \lambda \simeq \text{pr}_i \circ \lambda'$ for $i = 1, 2$.
- 2) If we take $\lambda * \lambda': [0, 1] \rightarrow X \times Y$, then $\text{pr}_i \circ (\lambda * \lambda') = (\text{pr}_i \circ \lambda) * (\text{pr}_i \circ \lambda')$.

These facts allow us to define a homomorphism

$$\text{pr}_{1*} \times \text{pr}_{2*} : \pi_1(X \times Y, (x_0, y_0)) \rightarrow \pi_1(X, x_0) \times \pi_1(Y, y_0)$$

by $\text{pr}_{1*} \times \text{pr}_{2*}([\lambda]) = ([\text{pr}_1 \circ \lambda], [\text{pr}_2 \circ \lambda])$. The inverse homomorphism is given by $([\lambda], [\mu]) \mapsto [(\lambda, \mu)(t)]$, where $(\lambda, \mu)(t) = (\lambda(t), \mu(t))$. Thus we have an isomorphism. \square

We can use such results to show that certain subspaces of a space are *not* deformation retracts. For example, if $\pi_1(X, x_0)$ is a nontrivial group, then $\pi_1(X \times X, (x_0, x_0))$ is not isomorphic to $\pi_1(X \times \{x_0\}, (x_0, x_0))$. Although $X \times \{x_0\}$ is a retract of $X \times X$ via

$$X \times \{x_0\} \hookrightarrow X \times X \rightarrow X \times \{x_0\},$$

it is not a deformation retract of $X \times X$.

Extra structure on a space can lead to more structure on the fundamental group. Recall (exercises of Chapter 4) that a topological group, (G, e) , is a Hausdorff topological space with basepoint $e \in G$ together with a continuous function (the group operation) $m: G \times G \rightarrow G$, satisfying $m(g, e) = m(e, g) = g$ for all $g \in G$, as well as another continuous function (the inverse) $\text{inv}: G \rightarrow G$ with $m(g, \text{inv}(g)) = e = m(\text{inv}(g), g)$ for all $g \in G$.

Theorem 7.15 allows us to define a new binary operation on $\pi_1(G, e)$, the composite of the isomorphism of the theorem with the homomorphism induced by m :

$$\mu_* : \pi_1(G, e) \times \pi_1(G, e) \rightarrow \pi_1(G \times G, (e, e)) \rightarrow \pi_1(G, e).$$

We denote the binary operation by $\mu_*([\lambda], [\nu]) = [\lambda \sharp \nu]$. On the level of loops, this mapping is given explicitly by $(\lambda, \mu) \mapsto \lambda \sharp \mu$, where $(\lambda \sharp \mu)(t) = m(\lambda(t), \mu(t))$. We next compare this binary operation with the usual multiplication of loops for the fundamental group.

Theorem 7.16. *If G is a topological group, then $\pi_1(G, e)$ is an abelian group.*


Proof. We first show that \sharp and the usual multiplication $*$ on $\pi_1(G, e)$ are actually the same binary operation! We argue as follows: Represent $\lambda * \mu(t)$ by $\lambda' \sharp \mu'(t)$, where

$$\lambda'(t) = \begin{cases} \lambda(2t), & 0 \leq t \leq \frac{1}{2}, \\ e, & \frac{1}{2} \leq t \leq 1, \end{cases} \quad \mu'(t) = \begin{cases} e, & 0 \leq t \leq \frac{1}{2}, \\ \mu(2t - 1), & \frac{1}{2} \leq t \leq 1. \end{cases}$$

Since $\lambda(1) = e = \mu(0)$ and $m(e, \mu'(t)) = \mu'(t)$, $m(\lambda'(t), e) = \lambda'(t)$, we see $\lambda * \mu(t) = m(\lambda'(t), \mu'(t))$. We next show that $\lambda * \mu$ is loop homotopic to $\lambda \sharp \mu$. Define two functions $h_1, h_2: [0, 1] \times [0, 1] \rightarrow [0, 1]$ by

$$h_1(t, s) = \begin{cases} 2t/(2 - s), & 0 \leq t \leq 1 - (s/2), \\ 1, & 1 - s/2 \leq t \leq 1, \end{cases}$$

$$h_2(t, s) = \begin{cases} 0, & 0 \leq t \leq s/2, \\ (2t - s)/(2 - s), & s/2 \leq t \leq 1. \end{cases}$$



Let $F(t, s) = m(\lambda(h_1(t, s)), \mu(h_2(t, s)))$. Since it is a composition of continuous functions, F is continuous. Notice

$$F(t, 0) = m(\lambda(h_1(t, 0)), \mu(h_2(t, 0))) = m(\lambda(t), \mu(t)) = \lambda \sharp \mu(t)$$

and


$$F(t, 1) = m(\lambda(h_1(t, 1)), \mu(h_2(t, 1))) = m(\lambda'(t), \mu'(t)) = \lambda * \mu(t).$$

Thus $\lambda * \mu$ is loop homotopic to $\lambda \sharp \mu$ and we get the same binary operation.

Given two loops λ and μ , consider the function

$$\mathcal{G}: [0, 1] \times [0, 1] \rightarrow G, \quad \mathcal{G}(t, s) = m(\lambda(t), \mu(s)).$$

The four corners are mapped to e and the diagonal from the lower left to the upper right is given by $\lambda \sharp \mu$. We will take some liberties



and argue with diagrams to construct a loop homotopy from $\lambda * \mu$ to $\mu * \lambda$.

Slice the square filled in by \mathcal{G} along the diagonal and paste in a rectangle that is simply a product of $\lambda \# \mu$ with an interval. Put the resulting hexagon into a square and fill in the remaining regions as the constant map at e , the identity element of G , in the trapezoidal regions and as λ or μ in the triangles where the path lies along the lines joining a vertex to the opposite side.

The diagram gives a homotopy from $\lambda * \mu$ to $\mu * \lambda$. It follows then that $[\lambda] * [\mu] = [\mu] * [\lambda]$ and so $\pi_1(G, e)$ is abelian. \square

Since S^1 is the topological group of unit length complex numbers, we have proved:

Corollary 7.17. $\pi_1(S^1, 1)$ is abelian.

Exercises

1. The unit sphere in \mathbb{R} is the set $S^0 = \{-1, 1\}$. Show that the set of homotopy classes of basepoint preserving mappings $[(S^0, -1), (X, x_0)]$ is the same set as $\pi_0(X)$, the set of path components of X .
2. Suppose that $f: X \rightarrow S^2$ is a continuous mapping that is *not* onto. Show that f is homotopic to a constant mapping.
3. If X is a space, recall that the *cone on X* is the quotient space $CX = X \times [0, 1] / X \times \{1\}$. Suppose $f: X \rightarrow Y$ is a continuous function and f is homotopic to a constant mapping

$c_y: X \rightarrow Y$ for some $y \in Y$. Show that there is an extension of f , $\hat{f}: CX \rightarrow Y$, so that $f = \hat{f} \circ i$, where $i: X \rightarrow CX$ is the inclusion, $i(x) = [(x, 0)]$.

4. Suppose that X is a path-connected space. When is it true that for any pair of points $p, q \in X$ all paths from p to q induce the same isomorphism between $\pi_1(X, p)$ and $\pi_1(X, q)$?
5. Prove that a disk minus two points is a deformation retract of a figure 8 (that is, $S^1 \vee S^1$).
6. A *starlike space* is a slightly weaker notion than a convex space—in a starlike space $X \subset \mathbb{R}^n$, there is a point $x_0 \in X$ so that for any other point $y \in X$ and any $t \in [0, 1]$ the point $tx_0 + (1 - t)y$ is in X . Give an example of a starlike space that is not convex. Show that a starlike space is a deformation retract of a point.
7. If $K = \alpha(S^1) \subset \mathbb{R}^3$ is a knot, that is, a homeomorphic image of a circle in \mathbb{R}^3 , then the complement of the knot $\mathbb{R}^3 - K$ has fundamental group $\pi_1(\mathbb{R}^3 - K)$. In fact, this group is an invariant of the knot in a sense that can be made precise. Give a plausibility argument that $\pi_1(\mathbb{R}^2 - K) \neq \{0\}$. See [69] for a thorough treatment of this important invariant of knots.

Chapter 8

Computations and Covering Spaces

... it is necessary, in order to affirm that a manifold is simply-connected, to study its fundamental group, ...

HENRI POINCARÉ, 1904

We have defined the fundamental group and showed that it is a topological invariant, that is, homeomorphic spaces have isomorphic fundamental groups. But we have yet to consider a space whose fundamental group is nontrivial. Two familiar spaces, S^1 and $\mathbb{R}P^2$, will provide examples.


The method of computation focuses on the properties of the mappings,

$$w: \mathbb{R} \rightarrow S^1 \quad \text{and} \quad p: S^2 \rightarrow \mathbb{R}P^2 \\ w(r) = \cos(2\pi r) + i \sin(2\pi r) = e^{2\pi i r} \quad \text{and} \quad p(\mathbf{x}) = [\pm \mathbf{x}].$$

These mappings share certain important properties.

Definition 8.1. Let X be a space. A **covering space** of X is a path-connected space \tilde{X} and a mapping $p: \tilde{X} \rightarrow X$ such that, for every $x \in X$, there is an open, path-connected subset U with $x \in U$ for which each path component of $p^{-1}(U)$ is homeomorphic to U by

restriction of the mapping p . Such open sets are called **elementary neighborhoods**.



For example, if $e^{i\theta} \in S^1$, then for $0 < \epsilon < \pi$, the open set $U = \{e^{i\alpha} \mid \theta - \epsilon < \alpha < \theta + \epsilon\}$ in S^1 has inverse image under w given by

$$w^{-1}(U) = \bigcup_{k \in \mathbb{Z}} \left(\frac{\theta}{2\pi} - \frac{\epsilon}{2\pi} + k, \frac{\theta}{2\pi} + \frac{\epsilon}{2\pi} + k \right).$$

Since $\epsilon/2\pi < 1/2$, the intervals in the union are all disjoint. Furthermore, w restricted to any one of these intervals has an inverse given by a branch of the logarithm. In the case of the quotient map $p: S^2 \rightarrow \mathbb{RP}^2$, for a connected open set $V \subset S^2$ satisfying $V \cap -V = \emptyset$, we have $p(V)$ open in \mathbb{RP}^2 and $p^{-1}(p(V)) = V \cup -V$. Since the components of $p^{-1}(p(V))$ are V and $-V$, it is an elementary neighborhood. For any $[\pm \mathbf{x}] \in \mathbb{RP}^2$, there is such an elementary neighborhood containing $[\pm \mathbf{x}]$ and so $p: S^2 \rightarrow \mathbb{RP}^2$ is a covering space.

Henceforth we will assume that all spaces are path-connected and locally path-connected to avoid pathological cases. The most useful property of covering spaces is the ability to lift paths in X to paths in \tilde{X} while preserving the homotopy relation.

Lemma 8.2. *Let $p: \tilde{X} \rightarrow X$ be a covering space and let $\tilde{x}_0 \in \tilde{X}$ with $p(\tilde{x}_0) = x_0 \in X$. If $\lambda: [0, 1] \rightarrow X$ is any path with $\lambda(0) = x_0$, then there exists a unique path $\hat{\lambda}: [0, 1] \rightarrow \tilde{X}$ with $\hat{\lambda}(0) = \tilde{x}_0$ and $p \circ \hat{\lambda} = \lambda$.*

Proof. Cover X by elementary neighborhoods. If $\lambda([0, 1]) \subset U$ for some elementary neighborhood, then $x_0 \in U$ and $\tilde{x}_0 \in p^{-1}(U)$. It follows that \tilde{x}_0 lies in some component C_0 of $p^{-1}(U)$ that is homeomorphic to U via $p|_{C_0}: C_0 \rightarrow U$. Let $(p|_{C_0})^{-1}: U \rightarrow C_0$ denote the inverse of this homeomorphism and let $\hat{\lambda} = (p|_{C_0})^{-1} \circ \lambda$. Then $\hat{\lambda}(0) = (p|_{C_0})^{-1}(x_0) = \tilde{x}_0$, since \tilde{x}_0 is the only point in \tilde{X} corresponding to x_0 in this component. Finally, $p \circ \hat{\lambda} = p \circ (p|_{C_0})^{-1} \circ \lambda = \lambda$.

If $\lambda([0, 1]) \not\subset U$, consider the collection $\{\lambda^{-1}(U') \subset [0, 1] \mid U', \text{ an elementary neighborhood}\}$. This is a cover of $[0, 1]$, which is a compact metric space, and so by Lebesgue's Lemma we can choose $0 = t_0 < t_1 < \cdots < t_{n-1} < t_n = 1$ with each $\lambda([t_{i-1}, t_i])$ a subset of some elementary neighborhood (take $t_i - t_{i-1} < \delta$, the Lebesgue number). Using the argument above, lift λ on $[0, t_1]$. Then take $\lambda(t_1)$ as x_0 and $\hat{\lambda}(t_1)$ as \tilde{x}_0 and lift λ to $[t_1, t_2]$. Continuing in this manner, we construct $\hat{\lambda}$ on $[0, 1]$ with $\hat{\lambda}(0) = \tilde{x}_0$ and $p \circ \hat{\lambda} = \lambda$.

To show that $\hat{\lambda}$ constructed in this manner is unique, we prove a more general result that implies uniqueness.

Lemma 8.3. *Let $p: \tilde{X} \rightarrow X$ be a covering space and Y a connected, locally connected space. Given two mappings $f_1, f_2: Y \rightarrow \tilde{X}$ with $p \circ f_1 = p \circ f_2$, then the set $\{y \in Y \mid f_1(y) = f_2(y)\}$ is either empty or all of Y .*

Proof. Consider the subset of Y given by $B = \{y \in Y \mid f_1(y) = f_2(y)\}$. We show that B is both open and closed. If $y \in \text{cls } B$, consider $x = p \circ f_1(y) = p \circ f_2(y)$ and U an elementary neighborhood containing x . The subset $(p \circ f_1)^{-1}(U) \cap (p \circ f_2)^{-1}(U)$ contains y . Because Y is locally connected, there is an open set W for which $y \in W \subset (p \circ f_1)^{-1}(U) \cap (p \circ f_2)^{-1}(U)$ with W connected. Then $f_1(W)$ and $f_2(W)$ are connected subsets of $p^{-1}(U) \subset \tilde{X}$. Since W is open and $y \in \text{cls } B$, there is a point $z \in W$ with $z \in B$. Thus $f_1(z) = f_2(z)$ and $f_1(W) \cap f_2(W) \neq \emptyset$; therefore, $f_1(W)$ and $f_2(W)$ must lie in the same component of $p^{-1}(U)$. Since $p \circ f_1(y) = p \circ f_2(y)$ and the component in which we find both $f_1(y)$ and $f_2(y)$ is homeomorphic to U by the restriction of p , we have $f_1(y) = f_2(y)$. Thus $y \in B$ and B is closed.

If we let $y \in B$, the argument above shows that the sets $f_1(W)$ and $f_2(W)$ lie in the same component C_0 of $p^{-1}(U)$. It follows that, for all $w \in W$,

$$f_1(w) = (p|_{C_0})^{-1} \circ p \circ f_1(w) = (p|_{C_0})^{-1} \circ p \circ f_2(w) = f_2(w)$$

and so W is contained in B . Thus B is open.

The only subsets of Y that are both open and closed are Y itself and \emptyset and so we have proved the lemma. \square

Using Lemma 8.3, two lifts of a path $\lambda: [0, 1] \rightarrow X$ which begin at the same point in \tilde{X} must be the same lift. This is the uniqueness part of Lemma 8.2. \square

Having lifted paths in X to paths in \tilde{X} , we next lift certain homotopies between paths.

Lemma 8.4. *Let $p: \tilde{X} \rightarrow X$ be a covering space and let $\eta_0, \eta_1: [0, 1] \rightarrow \tilde{X}$ be two paths in \tilde{X} with $\eta_0(0) = \eta_1(0) = \tilde{x}_0$. If $p \circ \eta_0(1) = x_1 = p \circ \eta_1(1)$ and $p \circ \eta_0 \simeq p \circ \eta_1$ via a homotopy that fixes the endpoints of the paths in X , then $\eta_1 \simeq \eta_2$ in \tilde{X} and, in particular, $\eta_0(1) = \eta_1(1)$.*

Proof. Let $H: [0, 1] \times [0, 1] \rightarrow X$ be a homotopy between $p \circ \eta_0$ and $p \circ \eta_1$. In this case, we have, for all $s, t \in [0, 1]$,

$$\begin{aligned} H(s, 0) = p \circ \eta_0(s) & \quad \text{and} \quad H(0, t) = p(\tilde{x}_0) \\ H(s, 1) = p \circ \eta_1(s) & \quad H(1, t) = p \circ \eta_0(1) = p \circ \eta_1(1). \end{aligned}$$

Since $[0, 1] \times [0, 1]$ is a compact metric space, when we cover it by the collection $\{H^{-1}(U) \mid U, \text{ an elementary neighborhood of } X\}$, we can apply Lebesgue's Lemma to get $\delta > 0$ for which any subset of $[0, 1] \times [0, 1]$ of diameter $< \delta$ lies in some $H^{-1}(U)$. If we subdivide the interval $[0, 1]$,

$$0 = s_0 < s_1 < \cdots < s_{m-1} < s_m = 1$$

and

$$0 = t_0 < t_1 < \cdots < t_{n-1} < t_n = 1,$$

so that $s_i - s_{i-1} < \delta/2$ and $t_j - t_{j-1} < \delta/2$, then H maps each subrectangle $[s_{i-1}, s_i] \times [t_{j-1}, t_j]$ into an elementary neighborhood for all i and j .

To construct the lifting $\hat{H}: [0, 1] \times [0, 1] \rightarrow \tilde{X}$ and show it is a homotopy between η_0 and η_1 , begin by lifting H on $[0, s_1] \times [0, t_1]$ to \tilde{X} by using $\hat{H} = (p|_{C_{11}})^{-1} \circ H$, where C_{11} is the component of $p^{-1}(U_{11})$ containing $\eta_0(0)$ and $H([0, s_1] \times [0, t_1]) \subset U_{11}$, an elementary neighborhood. Having done this, extend \hat{H} next to $[s_1, s_2] \times [0, t_1]$. Notice that \hat{H} has been defined on the line segment $\{s_1\} \times [0, t_1]$ which is connected and this determines the component of $p^{-1}(U_{21})$ for the elementary neighborhood U_{21} which contains $H([s_1, s_2] \times [0, t_1])$. Once the component, say C_{21} , is determined, extend \hat{H} by $\hat{H} = (p|_{C_{21}})^{-1} \circ H$. Continue in this manner until \hat{H} is defined on $[0, 1] \times [0, t_1]$. Next, extend to $[0, 1] \times [t_1, t_2]$ using the fact that the value of \hat{H} has been determined on each successive subrectangle along the left and bottom edges, as a connected subset. Continue along each row until \hat{H} is defined on $[0, 1] \times [0, 1]$. By Lemma 8.3, \hat{H} is unique fulfilling the condition $\hat{H}(0, 0) = \eta_0(0)$. Since $\eta_0(s)$ is also a lift of $H(s, 0)$, we have that $\hat{H}(s, 0) = \eta_0(s)$. The condition $H(0, t) = p \circ \eta_0(0)$ implies that $\hat{H}(0, t) = \eta_0(0)$, that is, the homotopy \hat{H} is constant on the subset $\{0\} \times [0, 1]$. Thus, the lift $\hat{H}(s, 1)$ of the path $p \circ \eta_1(s)$ in X begins at $\eta_0(0) = \eta_1(0)$, and $\eta_1(s)$ is also such a lift. By uniqueness, $\hat{H}(s, 1) = \eta_1(s)$. Finally, $H(1, t) = p \circ \eta_0(1) = p \circ \eta_1(1)$ for all $t \in [0, 1]$, $\hat{H}(1, t) = \eta_0(1)$, and we conclude that $\eta_0(1) = \eta_1(1)$ since $\hat{H}(1, t)$ is constant. \square

Uniqueness of liftings of homotopies provides considerable control over the fundamental group through a covering space, giving us a toehold for computation.

Corollary 8.5. *Suppose $p: \tilde{X} \rightarrow X$ is a covering space: (1) If $\eta: [0, 1] \rightarrow \tilde{X}$ is a loop at \tilde{x}_0 and $p \circ \eta$ is homotopic to the constant loop c_{x_0} for $x_0 = p(\tilde{x}_0)$, then $\eta \simeq c_{\tilde{x}_0}$. (2) The induced homomorphism $p_*: \pi_1(\tilde{X}, \tilde{x}_0) \rightarrow \pi_1(X, x_0)$ is injective. (3) For all $x \in X$, the subsets $p^{-1}(\{x\})$ of \tilde{X} have the same cardinality.*

Proof. (1) One lift of c_{x_0} is simply the constant path $c_{\tilde{x}_0}$. By Lemma 8.4, $p \circ \eta \simeq p \circ c_{\tilde{x}_0} = c_{x_0}$ implies $\eta \simeq c_{\tilde{x}_0}$.

(2) If $p_*([\lambda]) = p_*([\mu])$, then, because p_* is a homomorphism, $p_*([\lambda] * [\mu^{-1}]) = [c_{x_0}]$, that is, $p \circ (\lambda * \mu^{-1}) \simeq c_{x_0}$. By (1), $\lambda * \mu^{-1} \simeq c_{\tilde{x}_0}$ or $\lambda \simeq \mu$, that is, $[\lambda] = [\mu]$.

(3) Suppose x_0 and x_1 are in X and $\lambda: [0, 1] \rightarrow X$ is a path joining x_0 to x_1 . Suppose $y \in p^{-1}(\{x_0\})$. We define a mapping $\Lambda: p^{-1}(\{x_0\}) \rightarrow p^{-1}(\{x_1\})$ by lifting λ to $\lambda_y: [0, 1] \rightarrow \tilde{X}$ with $\lambda_y(0) = y$. Define $\Lambda(y) = \lambda_y(1)$. Since λ_y is uniquely determined by being a lift of $p \circ \lambda_y = \lambda$ with $\lambda_y(0) = y$, the function Λ is well defined. By Lemma 8.3, lifts of λ beginning at different elements in $p^{-1}(\{x_0\})$ must end at different points in $p^{-1}(\{x_1\})$ and so Λ is injective. Using lifts of λ^{-1} we deduce that Λ is surjective. (Notice that a different choice of λ might give a different one-one correspondence Λ .) \square

For $w: \mathbb{R} \rightarrow S^1$, $w(r) = e^{2\pi ir}$, we find that $w^{-1}(1 + 0i) = \mathbb{Z} \subset \mathbb{R}$ and so $w^{-1}(\{z\})$ is countably infinite for each $z \in S^1$. For $p: S^2 \rightarrow \mathbb{R}P^2$, $p^{-1}(\{\pm \mathbf{x}_0\})$ contains two elements, \mathbf{x}_0 and $-\mathbf{x}_0$. In general, if we lift a loop $\omega: [0, 1] \rightarrow X$ at x_0 in X , the proof of (3) of Corollary 8.5 obtains a mapping $\Omega: p^{-1}(\{x_0\}) \rightarrow p^{-1}(\{x_0\})$ by lifting the loop. By remark (1) of the corollary, if Ω is nontrivial, then the loop ω is not homotopic to the constant map. This observation is enough to prove the following.

Theorem 8.6. **A.** $\pi_1(S^1) \cong \mathbb{Z}$. **B.** $\pi_1(\mathbb{R}P^2) \cong \mathbb{Z}/2\mathbb{Z}$.

Proof of A. If $\beta: [0, 1] \rightarrow S^1$ is any loop at $1 \in S^1$, then the lift of β to $\hat{\beta}: [0, 1] \rightarrow \mathbb{R}$ satisfies $\hat{\beta}(1) \in \mathbb{Z}$. The properties of liftings determine a function $\Xi: \pi_1(S^1) \rightarrow \mathbb{Z}$ given by $[\beta] \mapsto \hat{\beta}(1)$.

Let $\alpha: [0, 1] \rightarrow S^1$ be given by $\alpha(t) = (\cos(2\pi t), \sin(2\pi t))$. Since $\alpha = w|_{[0,1]}$, we see that one lift of α to \mathbb{R} is just the identity and $\hat{\alpha}(1) = 1$. It follows that α is not homotopic to the constant map at 1 , c_1 . Next consider α^n for $n \in \mathbb{Z}$, given by $\alpha^n(t) = (\cos(2\pi nt), \sin(2\pi nt))$. By the same argument for α , $\hat{\alpha}^n(1) = n$ and so the mapping $\Xi: \pi_1(S^1) \rightarrow \mathbb{Z}$ is surjective. Since each $\alpha^n \not\cong c_1$ for $n \neq 0$, the subgroup generated by $[\alpha]$, isomorphic to \mathbb{Z} , is a subgroup of $\pi_1(S^1)$.

To finish the proof of **A**, we show that if β is any loop based at 1 in S^1 , then $\beta \simeq \alpha^n$ for some $n \in \mathbb{Z}$. Let $U_1 = \{(x, y) \in S^1 \mid y > -1/10\}$ and $U_2 = \{(x, y) \in S^1 \mid y < 1/10\}$. The pair $\beta^{-1}(U_1), \beta^{-1}(U_2)$ is an open cover of $[0, 1]$ and by Lebesgue's Lemma we can subdivide $[0, 1]$ as $0 = t_0 < t_1 < \dots < t_{m-1} < t_m = 1$ so that

$$\text{i) } \beta([t_i, t_{i+1}]) \subset U_1 \text{ or } \beta([t_i, t_{i+1}]) \subset U_2 \text{ for } 0 \leq i < m.$$

Form the union of consecutive subintervals when both are mapped to the same U_j , $j = 1$ or 2 . In detail, let $s_0 = 0$ and $s_1 = t_{i_1}$, where $\beta([0, t_{i_1}]) \subset U_{j_1}$ for j_1 one of 1 or 2 and $\beta([t_{i_1}, t_{i_1+1}]) \not\subset U_{j_1}$. Let $U_{j_2} \neq U_{j_1}$ and $\beta([t_{i_1}, t_{i_1+1}]) \subset U_{j_2}$. Then let $s_2 = t_{i_2}$, where $\beta([t_{i_1}, t_{i_2}]) \subset U_{j_2}$ but $\beta([t_{i_2}, t_{i_2+1}]) \not\subset U_{j_2}$. Continue in this manner to get

$$0 = s_0 < s_1 < \cdots < s_{k-1} < s_k = 1$$

so that

- ii) $\beta([s_{j-1}, s_j])$ and $\beta([s_j, s_{j+1}])$ are not both contained in the same U_k , for $k = 1, 2$.

Let $\beta_j: [0, 1] \rightarrow S^1$ denote the reparameterization of $\beta|_{[s_j, s_{j+1}]}$ so that $\beta \simeq \beta_0 * \beta_1 * \cdots * \beta_{k-1}$, and each β_j is a path in exactly one of U_1 or U_2 . Furthermore, $\beta(s_j) \in U_1 \cap U_2$, a subspace of two components, one of which contains $1 = e^{2\pi i 0}$ and the other $-1 = e^{\pi i}$. For $0 < j < m$ choose a path $\lambda_j: [0, 1] \rightarrow U_1 \cap U_2$ with $\lambda_j(0) = \beta(s_j) = \beta_{j-1}(s_j)$ and $\lambda_j(1) = 1$ or -1 , depending on the component. Define


$$\begin{aligned} \gamma_1 &= \beta_0 * \lambda_1, \\ \gamma_j &= \lambda_{j-1}^{-1} * \beta_{j-1} * \lambda_j \text{ for } 1 < j < k, \\ \gamma_k &= \lambda_{m-1}^{-1} * \beta_{k-1}. \end{aligned}$$

By cancelling $\lambda_j * \lambda_j^{-1}$, $\beta \simeq \gamma_1 * \gamma_2 * \cdots * \gamma_k$. Consider the paths γ_ℓ . If γ_ℓ is a closed path, it lies in U_1 or U_2 , which are simply-connected and so $\gamma_\ell \simeq c_1$ or $\gamma_\ell \simeq c_{-1}$. If γ_ℓ is not closed, then it crosses between the components of $U_1 \cap U_2$. It follows that $\gamma_\ell \simeq \eta_1^{\pm 1}$ or $\gamma_\ell \simeq \eta_2^{\pm 1}$, where $\eta_1(t) = (\cos(\pi t), \sin(\pi t))$, the path joining 1 to -1 in U_1 , and $\eta_2(t) = (\cos(\pi t + \pi), \sin(\pi t + \pi))$, the path joining -1 to 1 in U_2 . Making the cancellations of the type $\eta_1 \eta_1^{-1} \simeq c_1$ or $\eta_2 \eta_2^{-1} \simeq c_{-1}$, we are left with three possibilities:

$$\begin{aligned} \beta &\simeq c_1, \quad \beta \simeq \eta_1 * \eta_2 * \eta_1 * \eta_2 * \cdots * \eta_1 * \eta_2, \text{ or} \\ &\beta \simeq \eta_2^{-1} * \eta_1^{-1} * \eta_2^{-1} * \cdots * \eta_2^{-1} * \eta_1^{-1}, \end{aligned}$$

after cancelling out $c_{\pm 1}$. The ordering is determined by the fact that β begins and ends at 1 , and each γ_ℓ either joins 1 to -1 , joins -1 to 1 , or it simply stays put. After cancellation of the paths that stay put or products of paths that are homotopic to the constant path, we are

left with such a product in that order. Finally, $w|_{[0,1]} = \alpha \simeq \eta_1 * \eta_2$ and so $\beta \simeq \alpha^n$ for some $n \in \mathbb{Z}$. \square



Proof of B. Consider the model of the projective plane given by the *di-gon*, a disk with each point on the boundary identified with the point symmetric with respect to the origin. Let $x_0 \in \mathbb{RP}^2$ be the point $x_0 = [\pm(1, 0, 0)]$. Let $p: S^2 \rightarrow \mathbb{RP}^2$ denote the covering space $p(\mathbf{x}) = [\pm\mathbf{x}]$. Let the loop a in \mathbb{RP}^2 denote *half of the equator*, and lift a to S^2 . We get a path \hat{a} from $(1, 0, 0)$ to $(-1, 0, 0)$ along the equator of S^2 . By Corollary 8.5, $\hat{a} \not\sim c_{x_0}$. In the di-gon representation of \mathbb{RP}^2 , $a * a = a^2$ surrounds the disk, and so a^2 can be contracted to c_{x_0} by shrinking to the center of the disk and translating over to x_0 . It follows that $\pi_1(\mathbb{RP}^2)$ contains $\mathbb{Z}/2\mathbb{Z}$. To finish, we need show that any loop at x_0 is homotopic to a^n for some $n \in \mathbb{Z}$. Using the di-gon we see that away from the image of the path a^2 a path lies in the contractible interior of a disk. The disk can be used to push any loop onto a as often as it crosses between the copies of x_0 . Thus we see that any loop based at x_0 is homotopic to a^n for some $n \in \mathbb{Z}$ and so homotopic to a or c_{x_0} . This implies that

$$\pi_1(\mathbb{RP}^2) = \langle [a] \rangle / ([a]^2 = [c_{x_0}]) \cong \mathbb{Z}/2\mathbb{Z}.$$

This completes the proof of Theorem 8.6. \square

Covering spaces can be developed much further. We refer the reader to [55] or [51] for thorough treatments. Let's turn now to applications. We first return to the central question of the text:

Invariance of Dimension for $(2, n)$. For $n \neq 2$, \mathbb{R}^n and \mathbb{R}^2 are not homeomorphic.

Proof. We assume that $n \geq 2$ since the case of $n = 1$ is covered in Chapter 5. If $\mathbb{R}^n \cong \mathbb{R}^2$, then, by composing with a translation if needed, we can choose a homeomorphism $f: \mathbb{R}^n \rightarrow \mathbb{R}^2$ for which $f(\mathbf{0}) = (0, 0)$. Such a mapping induces a homeomorphism $\mathbb{R}^n - \{\mathbf{0}\} \cong \mathbb{R}^2 - \{(0, 0)\}$. Since S^{n-1} is a deformation retract of $\mathbb{R}^n - \{\mathbf{0}\}$, by Theorem 7.10, $\pi_1(\mathbb{R}^n - \{\mathbf{0}\}) \cong \pi_1(S^{n-1})$. For $n > 2$, Corollary 7.13 states that $\pi_1(S^{n-1}) \cong \{e\}$, while, for $n = 2$, $\pi_1(S^1) \cong \mathbb{Z}$. Since the fundamental group is a topological invariant, it must be the case that $n = 2$. \square

This argument is characteristic of the power of introducing algebraic structures as topological invariants of spaces. Our goal in later chapters is to generalize these ideas.

Recall the somewhat unexpected topological property introduced in the exercises of Chapter 2: A space X has the *fixed point property* (**FPP**) if any continuous mapping $f: X \rightarrow X$ has a fixed point, that is, there exists a point $x_0 \in X$ with $f(x_0) = x_0$. By the Intermediate Value Theorem we can prove that the interval $[0, 1]$ has the FPP: if $f: [0, 1] \rightarrow [0, 1]$ is continuous, then define $g(x) = f(x) - x: [0, 1] \rightarrow \mathbb{R}$. If $f(0) \neq 0$ and $f(1) \neq 1$, then $g(0) > 0$ and $g(1) < 0$ and so g must take the value 0 somewhere. If $g(x) = 0$, then $f(x) = x$.


What is the generalization of the space $[0, 1]$ to higher dimensions? Candidates include $[0, 1] \times [0, 1]$ in dimension 2 or maybe the **two-disk** $e^2 = \{\mathbf{x} \in \mathbb{R}^2 \mid \|\mathbf{x}\| \leq 1\} = \text{cls } B(\mathbf{0}, 1)$. The choice between these two candidates is irrelevant since the fixed point property is a topological property and they are homeomorphic. (Can you prove it?) We generalize the fixed point property for the interval $[0, 1]$ to the two-disk.

Theorem 8.7 (Brouwer's Theorem in dimension 2). *The two-disk $e^2 = \{\mathbf{x} \in \mathbb{R}^2 \mid \|\mathbf{x}\| \leq 1\} \subset \mathbb{R}^2$ has the fixed point property.*

Proof. Suppose $f: e^2 \rightarrow e^2$ is a continuous function without a fixed point. Then for each $\mathbf{x} \in e^2$, $f(\mathbf{x}) \neq \mathbf{x}$. Define $g: e^2 \rightarrow S^1$ by

$$g(\mathbf{x}) = \text{intersection of the ray from } f(\mathbf{x}) \text{ to } \mathbf{x} \text{ with } S^1.$$

To see that $g(\mathbf{x})$ is continuous on e^2 , we apply some vector geometry: write $Q = f(\mathbf{x})$, $Z = g(\mathbf{x})$. Let $O = (0, 0)$ and define



$X = (\mathbf{x} - Q)/\|\mathbf{x} - Q\|$. Then, $g(\mathbf{x}) = Z = Q + tX$ for some $t \geq 0$ for which $Q + tX \in S^1$, that is, $(Q + tX) \cdot (Q + tX) = 1$. This condition can be rewritten to solve for t , namely,

$$(Q + tX) \cdot (Q + tX) = t^2(X \cdot X) + 2t(Q \cdot X) + Q \cdot Q = 1.$$

The quadratic formula gives

$$\begin{aligned} t_{\mathbf{x}} &= -Q \cdot X + \sqrt{(Q \cdot X)^2 + 1 - Q \cdot Q} \\ &= -f(\mathbf{x}) \cdot \frac{\mathbf{x} - f(\mathbf{x})}{\|\mathbf{x} - f(\mathbf{x})\|} + \sqrt{\left(f(\mathbf{x}) \cdot \frac{\mathbf{x} - f(\mathbf{x})}{\|\mathbf{x} - f(\mathbf{x})\|}\right)^2 + 1 - f(\mathbf{x}) \cdot f(\mathbf{x})}. \end{aligned}$$

Note that this choice of signs gives $t_{\mathbf{x}} \geq 0$, and $t_{\mathbf{x}} = 0$ implies $f(\mathbf{x}) = \mathbf{x}$. Since we have assumed that this doesn't happen, $t_{\mathbf{x}} > 0$. Furthermore, $t_{\mathbf{x}}$ is a continuous function of \mathbf{x} . We can write $g(\mathbf{x})$ explicitly as

$$g(\mathbf{x}) = f(\mathbf{x}) + t_{\mathbf{x}} \frac{\mathbf{x} - f(\mathbf{x})}{\|\mathbf{x} - f(\mathbf{x})\|}$$

and so $g(\mathbf{x})$ is continuous.

By the definition of the mapping g , if $\mathbf{x} \in S^1 \subset e^2$, then $g(\mathbf{x}) = \mathbf{x}$. We have constructed a continuous mapping $g: e^2 \rightarrow S^1$ for which $g \circ i = \text{id}_{S^1}$, that is, the identity mapping on S^1 can be factored:

$$\text{id}_{S^1}: S^1 \xrightarrow{i} e^2 \xrightarrow{g} S^1.$$

This composite leads to a composite of group homomorphisms and fundamental groups:

$$\text{id}: \pi_1(S^1) \xrightarrow{i_*} \pi_1(e^2) \xrightarrow{g_*} \pi_1(S^1).$$

However, $\pi_1(e^2) = \{[c_1]\}$ and so $g_* \circ i_*([c_1]) = [c_1] \neq [\alpha]$ and $g_* \circ i_* \neq \text{id}$, a contradiction. Therefore, a continuous function $f: e^2 \rightarrow e^2$ without fixed points is not possible. \square

Corollary 8.8. S^1 is not a retract of e^2 .

More powerful tools will be developed in later chapters to prove a generalization of Theorem 8.7 and its corollary. Brouwer proved this general result around 1911 [11].

We next apply the fact that $\pi_1(\mathbb{R}P^2) \cong \mathbb{Z}/2\mathbb{Z}$. Recall that $\mathbb{R}P^2$ is the space of lines through the origin in \mathbb{R}^3 . The lower dimensional analogue is the space $\mathbb{R}P^1$ consisting of lines through the origin in \mathbb{R}^2 . We can identify a line with the angle it makes with the x -axis. To obtain every line through the origin, we only need angles $0 \leq \theta \leq \pi$, where the x -axis is identified with the angles 0 and π . Hence $\mathbb{R}P^1 \cong [0, \pi]/(0 \sim \pi) \cong S^1$. Thus $\pi_1(\mathbb{R}P^1) \cong \mathbb{Z}$. The analogue of the covering map $p: S^2 \rightarrow \mathbb{R}P^2$ in this case is $\bar{p}: S^1 \rightarrow \mathbb{R}P^1$ given by $e^{2\pi i\theta} \mapsto [\pm e^{2\pi i\theta}]$. In fact, $\bar{p}_*: \pi_1(S^1) \rightarrow \pi_1(\mathbb{R}P^1)$ is described as a homomorphism $\mathbb{Z} \rightarrow \mathbb{Z}$ given by multiplication by two, because the generator $[\alpha]$ wraps around $\mathbb{R}P^1$ twice.

In Chapter 5 we proved that a continuous mapping $f: S^1 \rightarrow \mathbb{R}$ must send some point and its negative to the same value, that is, there is always a point $x_0 \in S^1$ with $f(x_0) = f(-x_0)$. We can generalize that result to S^2 .

Theorem 8.9. *If $f: S^2 \rightarrow \mathbb{R}^2$ is a continuous function, then there exists a point $\mathbf{x} \in S^2$ with $f(\mathbf{x}) = f(-\mathbf{x})$.*

We proceed by proving an associated result.

Proposition 8.10 (The Borsuk-Ulam Theorem for $n = 2$). *There does not exist a continuous function $f: S^2 \rightarrow S^1$ that satisfies $f(-\mathbf{x}) = -f(\mathbf{x})$ for all $\mathbf{x} \in S^2$.*

Proof of the Borsuk-Ulam Theorem. Assume such a function exists. The condition satisfied by f can be written $f(\pm\mathbf{x}) = \pm f(\mathbf{x})$.

It follows that f induces $\hat{f}: \mathbb{R}P^2 \rightarrow \mathbb{R}P^1$ and \hat{f} fits into a diagram:

$$\begin{array}{ccc} S^2 & \xrightarrow{f} & S^1 \\ \downarrow p & & \downarrow \bar{p} \\ \mathbb{R}P^2 & \xrightarrow{\hat{f}} & \mathbb{R}P^1 \end{array}$$

for which $\bar{p} \circ f = \hat{f} \circ p$. Consider the induced homomorphism $\hat{f}_*: \pi_1(\mathbb{R}P^2) \rightarrow \pi_1(\mathbb{R}P^1)$. By Theorem 8.6, \hat{f}_* is a homomorphism $\mathbb{Z}/2\mathbb{Z} \rightarrow \mathbb{Z}$. However, any such homomorphism must be the trivial homomorphism. (Do you know why?) Let $\lambda: [0, 1] \rightarrow S^2$ denote a path from the North Pole to the South Pole along a meridian of constant longitude. It follows that $[p \circ \lambda] = [\alpha]$, a generator for $\mathbb{Z}/2\mathbb{Z} \cong \pi_1(\mathbb{R}P^2)$. Since the North and South Poles are antipodal, these points are identified in $\mathbb{R}P^1$ after passage through f and \bar{p} . Hence $[\bar{p} \circ f \circ \lambda]$ is nontrivial in $\pi_1(\mathbb{R}P^1)$. But $[\bar{p} \circ f \circ \lambda] = [\hat{f} \circ p \circ \lambda] = \hat{f}_*([p \circ \lambda]) = 0$, a contradiction. \square

Corollary 8.11. *If $f: S^2 \rightarrow \mathbb{R}^2$ is a continuous function such that $f(-\mathbf{x}) = -f(\mathbf{x})$ for all $\mathbf{x} \in S^2$, then $f(\mathbf{x}) = (0, 0)$ for some $\mathbf{x} \in S^2$.*

Proof. If not, then $g(\mathbf{x}) = f(\mathbf{x})/\|f(\mathbf{x})\|$ would be a continuous function $g: S^2 \rightarrow S^1$ with $g(-\mathbf{x}) = -g(\mathbf{x})$ for all $\mathbf{x} \in S^2$. \square

Proof of Theorem 8.9. Suppose for every $\mathbf{x} \in S^2$ that $f(\mathbf{x}) \neq f(-\mathbf{x})$. Then define $g(\mathbf{x}) = f(\mathbf{x}) - f(-\mathbf{x})$. Notice that g is continuous, $g(-\mathbf{x}) = -g(\mathbf{x})$, and $g(\mathbf{x}) \neq \mathbf{0}$ for all $\mathbf{x} \in S^2$, a contradiction. \square

Corollary 8.12. *No subset of \mathbb{R}^2 is homeomorphic to S^2 .*

The corollary tells us that there is no cartographic map homeomorphic to the entire sphere.

Finally, we derive an unexpected corollary of our analysis of the fundamental group of the circle, namely, the Fundamental Theorem of Algebra. This topological proof gives a complete proof avoiding the difficulties in the approach of Gauss in Chapter 5 based on connectedness.

The Fundamental Theorem of Algebra. *If $p(z) = z^n + a_{n-1}z^{n-1} + \cdots + a_1z + a_0$ is a polynomial with complex coefficients, then there is a complex number z_0 with $p(z_0) = 0$.*

Proof. Recall that $\mathbb{C} \cong \mathbb{R}^2$ and the n th power mapping $h: z \mapsto z^n$ induces a mapping $h: S^1 \rightarrow S^1$ which can be written as $e^{i\theta} \mapsto e^{in\theta}$. Lifting this mapping to the covering space $w: \mathbb{R} \rightarrow S^1$, it represents $n \in \mathbb{Z} \cong \pi_1(S^1)$ via the identification of $\pi_1(S^1)$ with \mathbb{Z} given by $[\beta] \mapsto \hat{\beta}(1)$.

Viewed as a mapping, $h: S^1 \rightarrow S^1$, h induces the homomorphism $h_*: \pi_1(S^1) \rightarrow \pi_1(S^1)$. The law of exponents implies that

$$h_*(\theta \mapsto e^{\pi im\theta}) = (\theta \mapsto (e^{\pi im\theta})^n = e^{\pi inm\theta}),$$

that is, h_* is multiplication by n .

We first consider a special case of the theorem—suppose

$$|a_{n-1}| + |a_{n-2}| + \cdots + |a_0| < 1.$$

Suppose $p(z)$ has no root in $e^2 = \{z \in \mathbb{C} \mid |z| \leq 1\}$. Define the mapping $\hat{p}: e^2 \rightarrow \mathbb{R}^2 - \{\mathbf{0}\}$ by $\hat{p}(z) = p(z)$. Restricting to $S^1 = \partial e^2$ we get $\hat{p}|: S^1 \rightarrow \mathbb{R}^2 - \{\mathbf{0}\}$. Since $\hat{p}|$ can be extended to e^2 , it follows (exercise) that $\hat{p}|$ is homotopic to a constant map. However, consider the mapping

$$F(z, t) = z^n + t(a_{n-1}z^{n-1} + \cdots + a_0),$$

which gives a homotopy between $F(z, 0) = z^n$ and $F(z, 1) = p(z)$. If $F(z, t)$ never vanishes on S^1 , the homotopy implies $\hat{p}| \simeq z^n$. To establish this condition, for $|z| = 1$ we estimate

$$\begin{aligned} |F(z, t)| &\geq |z^n| - |t(a_{n-1}z^{n-1} + \cdots + a_0)| \\ &\geq 1 - t(|a_{n-1}z^{n-1}| + \cdots + |a_0|) \\ &= 1 - t(|a_{n-1}| + \cdots + |a_0|) > 0. \end{aligned}$$

As a class in $\pi_1(S^1)$, $[(z \mapsto z^n)]$ is not homotopic to the constant map while $\hat{p}|$ is, so we get a contradiction.

To reduce the general case to this special case, let $t \in \mathbb{R}$, $t \neq 0$, and let $u = tz$. So

$$\begin{aligned} p(u) &= u^n + a_{n-1}u^{n-1} + \cdots + a_1u + a_0 \\ &= (tz)^n + a_{n-1}(tz)^{n-1} + \cdots + a_1tz + a_0. \end{aligned}$$

If $p(u) = 0$, then

$$z^n + \frac{a_{n-1}}{t}z^{n-1} + \cdots + \frac{a_1}{t^{n-1}}z + \frac{a_0}{t^n} = 0.$$

So given a zero for $p(u)$ we get a zero for $\tilde{p}_t(z)$ with $\tilde{p}_t(z) = z^n + \frac{a_{n-1}}{t}z^{n-1} + \cdots + \frac{a_0}{t^n}$ and vice versa. Taking t large enough we can guarantee

$$\left| \frac{a_{n-1}}{t} \right| + \cdots + \left| \frac{a_1}{t^{n-1}} \right| + \left| \frac{a_0}{t^n} \right| < 1$$

and we can apply the special case. \square

In Chapter 7 we proved that a subspace A of a space X , which is a deformation retract of X , shares the same fundamental group as X . Furthermore, if X and Y are homeomorphic spaces, they share the same fundamental group. We generalize these conditions to identify an important relation between spaces.

Definition 8.13. Two spaces are **homotopy equivalent**, denoted $X \simeq Y$, if there are mappings $f: X \rightarrow Y$ and $g: Y \rightarrow X$ with $g \circ f \simeq \text{id}_X$ and $f \circ g \simeq \text{id}_Y$.

If $A \subset X$ is a deformation retract, then there is a mapping $r: X \rightarrow A$ for which $\text{id}_A = r \circ i: A \rightarrow A$ and $\text{id}_X \simeq i \circ r: X \rightarrow X$. Thus A is homotopy equivalent to X and homotopy equivalence generalizes the relation of deformation retraction. Contractible spaces are homotopy equivalent to a one-point space so homotopy equivalence is a weaker notion than homeomorphism.

Proposition 8.14. *In a set of topological spaces, homotopy equivalence is an equivalence relation.*

Proof. It suffices to check transitivity since the other properties are clear. Suppose $X \simeq Y$ and $Y \simeq Z$ via mappings $f: X \rightarrow Y$, $g: Y \rightarrow$

X , $t: Y \rightarrow Z$, and $u: Z \rightarrow Y$. Consider $t \circ f: X \rightarrow Z$ and $g \circ u: Z \rightarrow X$. Then

$$(g \circ u) \circ (t \circ f) \simeq g \circ (u \circ t) \circ f \simeq g \circ \text{id}_Y \circ f = g \circ f \simeq \text{id}_X \quad \text{and}$$

$$(t \circ f) \circ (g \circ u) \simeq t \circ (f \circ g) \circ u \simeq t \circ \text{id}_X \circ u = t \circ u \simeq \text{id}_Z.$$

Fixing a universe, that is, a set in which all relevant spaces are elements, the equivalence class of a space X is called its **homotopy type**. The effectiveness of the fundamental group to distinguish spaces is limited by homotopy equivalence.

Proposition 8.15. *If X and Y are homotopy-equivalent spaces via mappings $f: X \rightarrow Y$ and $g: Y \rightarrow X$, then the induced mappings $f_*: \pi_1(X, x_0) \rightarrow \pi_1(Y, f(x_0))$ and $g_*: \pi_1(Y, y_0) \rightarrow \pi_1(X, g(y_0))$ are isomorphisms.*


Proof. Let $H: X \times [0, 1] \rightarrow X$ be a homotopy between $g \circ f$ and id_X . Let $\gamma: [0, 1] \rightarrow X$ be the path $\gamma(t) = H(x_0, t)$, so that $\gamma(0) = g \circ f(x_0)$ and $\gamma(1) = x_0$. We can write the induced homomorphisms:

$$\pi_1(X, x_0) \xrightarrow{f_*} \pi_1(Y, f(x_0)) \xrightarrow{g_*} \pi_1(X, g \circ f(x_0)) \xrightarrow{u_\gamma} \pi_1(X, x_0).$$

We claim that this composite is the identity homomorphism. Consider $[\lambda] \in \pi_1(X, x_0)$. The result of the composite on this element is

$$[\lambda] \mapsto [f \circ \lambda] \mapsto [g \circ f \circ \lambda] \mapsto [\gamma^{-1} * (g \circ f \circ \lambda) * \gamma].$$

Apply the homotopy H to get a homotopy from $g \circ f \circ \lambda$ to λ by $H(\lambda(t), s)$. We use this homotopy to construct one from $\gamma^{-1} * (g \circ f \circ \lambda) * \gamma$ to λ by reparameterizing according to the diagram:



In the triangles, we have taken γ and opened it into a triangle with the pictured curves given by isobars (constant paths). It follows from the homotopy that $[\gamma^{-1} * (g \circ f \circ \lambda) * \gamma] = [\lambda]$. This implies that f_* is injective and g_* surjective. To finish the proof consider the composite

$$\begin{aligned} \pi_1(Y, f(x_0)) &\xrightarrow{g_*} \pi_1(X, g \circ f(x_0)) \\ &\xrightarrow{f_*} \pi_1(Y, f \circ g \circ f(x_0)) \xrightarrow{u_\eta} \pi_1(Y, f(x_0)), \end{aligned}$$

where $\eta: [0, 1] \rightarrow Y$ is the path $\eta(t) = \bar{H}(f(x_0), t)$ in the homotopy \bar{H} between $f \circ g$ and id_Y . The same argument applies *mutatis mutandis* to show that f_* is surjective and g_* is injective and hence both homomorphisms are isomorphisms. \square

Homotopy equivalence is cruder than homeomorphism but includes it as a special case. To give an idea of how crude homotopy equivalence is, notice that, for all n , \mathbb{R}^n is homotopy equivalent to a point. The letters of the alphabet as subspaces of \mathbb{R}^2 show other failures to distinguish between different topological spaces.

$$A \simeq D \simeq S^1, B \simeq S^1 \vee S^1, C \simeq E \simeq F \simeq *, \dots$$

Proposition 8.15 shows that the fundamental group is a **homotopy invariant**, that is, if $X \simeq Y$, then $\pi_1(X) \cong \pi_1(Y)$. Thinking of the fundamental group as a filter that distinguishes spaces, it can only hope to catch homotopy inequivalent spaces. In later chapters we will consider other homotopy invariants. Poincaré [66] introduced the fundamental group to distinguish certain manifolds that were indistinguishable via other combinatorial invariants.

Exercises

1. Suppose that $f: S^1 \rightarrow S^1$ has an extension $\hat{f}: e^2 \rightarrow S^1$, that is, the mapping \hat{f} satisfies $\hat{f} \circ i = f$, where $i: S^1 \rightarrow e^2$ is the inclusion. Show that f is **null-homotopic**, that is, f is homotopic to the constant mapping.
2. Though we will not prove it, one of the useful theorems for computing the fundamental groups of spaces is the **Seifert-vanKampen Theorem**. A special case of this theorem

is the following: *If a path-connected space X is a union $X = U \cup V$ with V simply-connected and $x_0 \in U \cap V$, then the inclusion $i: U \rightarrow X$ induces a surjection $i_*: \pi_1(U, x_0) \rightarrow \pi_1(X, x_0)$ with kernel given by the smallest subgroup of $\pi_1(U, x_0)$ containing $j_*(\pi_1(U \cap V, x_0))$, where $j: U \cap V \hookrightarrow U$ denotes the inclusion.* Use the descriptions of $\mathbb{R}P^2$ of previous chapters and this theorem to make another computation of $\pi_1(\mathbb{R}P^2)$.

3. Suppose that X is simply-connected and $p: \tilde{X} \rightarrow X$ is a covering space of X . Show that p is a homeomorphism.
4. Let $\Omega(X, x_0)$ denote the based loop space of X given by $\Omega(X, x_0) = \{\lambda: [0, 1] \rightarrow X \mid \lambda \text{ is continuous and } \lambda(0) = \lambda(1) = x_0\}$.

This subspace of $\text{map}(I, X)$ is topologized with the compact-open topology. Show that

- i) $\pi_0(\Omega(X, x_0))$, the collection of path-components of $\Omega(X, x_0)$, is in one-one correspondence with $\pi_1(X, x_0)$.
 - ii) Show that the loop multiplication $m: \Omega(X, x_0) \times \Omega(X, x_0) \rightarrow \Omega(X, x_0)$ given by $m(\lambda, \mu) = \lambda * \mu$ is a continuous multiplication on $\Omega(X, x_0)$.
5. We know from Theorem 7.15 and Theorem 8.6 that the fundamental group of the torus $S^1 \times S^1$ is $\mathbb{Z} \times \mathbb{Z}$. Use the argument for the computation of $\pi_1(\mathbb{R}P^2) \cong \mathbb{Z}/2\mathbb{Z}$ to prove $\pi_1(S^1 \times S^1) \cong \mathbb{Z} \times \mathbb{Z}$ by viewing the torus as a quotient of $[0, 1] \times [0, 1]$.
 6. Let's make a space—take two distinct 2-spheres, S^2 , and join them by a line segment—a space resembling dumbbells, but with a very thin connector. Denote this space by X and show that it is simply-connected.