

# Index

- $A$ , **158**
- $a$ , superscript, **30**
- acceptable subterminal pair, **7**
- agreeable subgroup, **25**
- almost strongly  $p$ -embedded subgroup, **4**
  - types of, **4**
- Alperin, J., **314**
- Alt*, **1**
- anchored component, **453**
- Aschbacher, M., **x**, **12**
  - classical involution theorem, **29**
- $B^*$ , **157**
- $B_0$ , **175**
- Background Results, **xi**
- Baer, R., **332**
- Bender, H., **xi**
- Bennett, C., **12**
- Blok, R., **xi**, **10**, **12**
- breather, **303**
- brush, **393**
- $C^o$ ,  $C(V)^o$ , **15**
- Chev*, **1**
- $C_I$ ,  $C_{I'}$ ,  $C_I^o$ , **16**
- classical involution, **89**, **294**, **299**
- compatible, **14**
- $\mathcal{C}_p$ , **5**
- CT-pairs and CT-systems, **376–383**
- CT-system, *see also* weak CT-system
- Curtis-Tits theorem, **9**
- $C(V)$ , **15**
- $D_e$ , **29**, **155**
- Devillers, A., **12**
- Dickson, L. E., **xi**, **323**
- $D_n^+(q)$ ,  $D_n^-(q)$ , **16**
- $d_p$ , **6**
  - $d_p(G)$ , **6**
  - $d_p(K)$ , **6**
- Dunlap, J., **12**
- $E_6^+(q)$ ,  $E_6^-(q)$ , **16**
- $\epsilon_q$ 
  - when  $p$  is odd, **16**, **155**
  - when  $p = 2$ , **30**
- $E^q(X)$ , **94**
- even type, **303**
  - restricted, **8**
- extendible, **13**
- $\mathcal{F}$ ,  $\mathcal{F}(K)$ , **6**
- $\Phi_m$  ( $m$ th cyclotomic polynomial), **7**
- $f$ ,  $f(K)$ , **6**
- Finkelstein, L., **x**, **15**
- $G_0$ , **29**
- $G_0$ , **2**
- $G_0 \in \mathcal{C}hev$ , construction of, **29–290**
- $\mathcal{G}_2^i$ -groups, **5**
- $\gamma(G)$ , **4**
- Generic Case, **4**
- Gilman, R. H., **x**, **15**, **29**
- Gilman-Griess theory, **14**
- Gorenstein, D., **x**
- $\mathcal{G}_p$ , **5**
- $\mathcal{G}_p^i$ , **5**
- $\mathcal{G}_p^i$ -groups, **6**
- Gramlich, R., **xi**, **12**
- Griess, R. L., Jr., **x**, **15**, **29**
- $G^*$ -trick, **253**
- Harris, M. E., **x**
- $H_D$ , **18**
- Hoffman, C., **xi**, **10**, **12**
- Holt, D., **318**
- Horn, M., **12**
- $hs$ , superscript, **30**
- ignorable, **7**
- $\mathcal{IL}_p^o(G)$ , **4**
- $\mathcal{I}_p^o(G)$ , **4**
- $\mathcal{J}_p^*(G)$ , **7**
- $\mathcal{J}_p(G)$ , **6**
- $J_r$ ,  $J_{\bar{r}}$ , **171**
- $\kappa$ , **15**
- $\mathcal{K}^{(7)*}$ , **1**
- $\mathcal{K}^{(7)+}$ , **3**

- Köhl, R., xi, 12  
 $\mathcal{K}$ -proper, **1**  
 $\mathcal{K}_p$ , **5**  
 $K$ -preuniqueness subgroup, **4**  
  
 $L_2^q(X)$ , **94**  
 long root  $p$ -element, **474**  
 $\mathcal{L}_p^o(G)$ , **4**  
 $L^+(q)$ ,  $L^-(q)$ , 15  
  
 Mal'cev, A. I., 337, 405  
 McLaughlin, J., xi  
 milestone, 303  
 $\widehat{m}_p(K)$ , **5**  
 Mühlherr, B., 12  
  
 $N$ ,  $N(x, K, y, L)$ , x, **2**, 29  
 Natural, Mr., *see also* Griess, R. L., Jr.  
 neighborhood, **2**  
 Nickel, W., 12  
  
 $\mathcal{P}$ , **18**  
 $P$ -pairs and  $P$ -systems, 376–383  
 pair, **4**  
     acceptable subterminal, **7**  
      $p$ -component preuniqueness subgroup, 4  
      $p$ -generic, 5  
     Phan system, *see also* weak  $P$ -system  
     Phan theorem, 13, 14  
     Phan theory, 10–14  
     Phan, K.-W., 12  
     Phan-able, **11**  
     PP-pair, **377**  
     primitive prime divisor, **147**, 386  
      $p$ -Thin Configuration, 1  
  
 $\mathcal{Q}(G)$ ,  $\mathcal{Q}_p(G)$ , **156**  
 $\mathcal{Q}^*(G)$ , **156**  
  
 $\mathcal{R}$ , **173**  
 ranking functions  
      $\mathcal{F}$ , **6**  
      $\mathcal{F}$ ,  $\mathcal{F}(K)$ , **6**  
      $\tau$ ,  $\tau(K)$ , **6**  
      $f$ ,  $f(K)$ , **6**  
 reflection module, **326**  
 $\rho(K, I)$ , **436**  
  
 Serežkin, V. N., 322  
 Shpectorov, S., xi, 10, 12  
 Solomon 2-fusion systems, 79, 393  
 Solomon, R., 15  
 splitting  $K \in \mathcal{C}hev(r)$ , **8**  
 $Spor$ , **1**  
 standard CT-pair, **9**  
 standard Phan pair, standard  $P$ -pair, **10**  
 Suzuki, M., 332  
  
 $\tau$ ,  $\tau(K)$ , **6**  
 Theorem  $\mathcal{C}_7^*$ , **1**  
     Stage 4b+, 3  
     Stage 5+, 291  
 Thompson, J. G., x, xi, 46, 246  
 torally embedded, **339**  
 $\mathcal{T}_p$ , 5  
 triality-free, **475**  
  
 $u$ , superscript, **30**  
 uniqueness subgroup  
     almost strongly  $p$ -embedded subgroup  
         strongly  $p$ -embedded type, 4  
         strongly closed type, 4–5  
     strong  $p$ -uniqueness subgroup, 4  
     strong  $p$ -uniqueness subgroup of  
         component type, 4  
  
 $V_I$ ,  $V_{I'}$ , **16**  
 $V_{\kappa, p}$ , **16**  
  
 $W$ , **158**  
 $W^*$ , **159**  
 $W_1$ , **173**  
 Wagner, A., 322  
 Walter, J. H., x  
 weak CT-system, weak Curtis-Tits system,  
     **9**  
 weak CTP-system, **14**  
 weak  $P$ -system  
     extendible, **13**  
     weak Phan system, weak  $P$ -system, **13**  
 $W_H$ ,  $W_K$ ,  $W_L$ ,  $W_{L_u}$ , **158**  
 Wielandt, H., 359  
 Witzel, S., 12  
 $W(\mathcal{L})$ , **159**  
 Wong, W., 15  
 Wong, W. J., x  
 $W \equiv W(\mathcal{L})$ , 159  
 $W \equiv_{\nabla} W(\mathcal{L})$ ,  $W \equiv W(\mathcal{L})$ , **322**  
  
 Zalesskiĭ, A. E., 322