

# Index

- 2-local  $p$ -rank, **2**  
 $\downarrow_p$ , **24**  
 $\uparrow_p$ , **24**
- almost strongly  $p$ -embedded subgroup, **35**  
*Alt*, **2**  
 Aschbacher, M., 9, 10, 31, 95, 237, 245,  
 250, 253, 266, 272, 274  
 A-terminal  $p$ -component, **94**
- $\mathcal{B}_3^1$ , **312**  
 $\mathcal{B}_3$ -group, **311**  
 Background Results, 9  
 balanced signalizer functor, 11  
 barely in  $\mathcal{C}_p$ , **145**  
 Bender, H., 3  
 Bender-Thompson lemma, 13  
 bicharacteristic group, 6  
 $B$ -narrow component, **146**  
 $\mathcal{B}^p(G)$ , **34**  
 $\mathcal{B}_{*,c}^p(G)$ , **89, 104**  
 $\mathcal{B}_*^p(G)$ , **34, 104**  
 $\mathcal{B}_{*,o}^p(G)$ , **89, 104**  
 broad, **90**  
 broad triple, **6, 146**  
 $B$ -terminal  $p$ -component, **94**  
 $Bt\mathcal{K}_{\varepsilon xc}^3(G)$ , **5, 90**  
 $Bt\mathcal{K}^p(G)$ , **5, 90, 144**  
 $B$ -wide component, **146**
- $\mathcal{C}_2, \mathcal{C}_2$ -group, **33**  
 $\mathcal{C}_2^o$ , **2**  
 $C(b, J)$ , **8**  
*Chev*, *Chev*( $r$ ), **2**  
 complete  $(S)U_3(2)$ -component, **126**  
 component preuniqueness hypothesis, **95**  
 constrained  $\{z, b\}$ -neighborhood, **8, 209**  
 constrained neighborhood of  $U_4(2)$ -type,  
**209**  
 constrained neighborhood of sporadic type,  
**209**  
 constrained neighborhood, 210  
 core of a constrained neighborhood, **209**  
 core of a neighborhood, **8**
- Corollary  $\mathcal{C}_6$ , 310  
 Corollary  $\mathcal{C}_6^*$ , 312  
 $\mathcal{C}_3, \mathcal{C}_3$ -group, **33**  
 $\mathcal{C}_p, \mathcal{C}_p$ -group,  $p$  odd, **33**  
 $\mathcal{C}_p^o$ , **145**  
 $\mathcal{C}_p^*$ , **145**  
 Curtis-Tits theorem, 9  
 $C(x, K)$ , **19**
- $d$ , **310**  
 degenerate neighborhood, **90**  
 degenerate nonconstrained neighborhood, **6**  
 $d_{\mathcal{T}, p}$ , **310**
- $\mathcal{E}_1(P)$ , **11**  
 $\mathcal{E}^p(X), \mathcal{E}_m^p(X)$ , **11**  
 even type, **2**  
 exceptional situation, **313**
- faithful symplectic pair, **89, 143**  
 field triple, **44**  
 Finkelstein, L., 9, 10, 253  
 Fischer, B., 31  
 flat  $\mathcal{T}\mathcal{G}_3$ -group, **312**  
 frame, **441**
- Gilman ordering, refinement of, 313  
 Gilman, R., 95, 313  
 Gilman-maximal, **25**  
 $\Gamma_K$ , 250  
 $\Gamma_{P,2}^{o1}(G)$ , **93**  
 $\mathcal{G}_p$ -depth, 312  
 $\Gamma'_{P,k}(X)$ , **11**  
 $\Gamma_{P,k}(X)$ , **11**  
 Gramlich, R., 9  
 Griess, R. L., 10, 31
- $H_b^*$ , **327**  
 Horn, M., 9
- $\mathcal{I}_3^*(K)$ , **121**  
 $\mathcal{I}_p^o$ -rigid pair, **311**  
 $\mathcal{I}_p^o$ -terminal  $\mathcal{T}_p$ -pair,  $\mathcal{T}_p^i$ -pair, **311**  
 $\mathcal{I}_p^o$ -rigid, **25**  
 $\mathcal{I}_p^o$ -terminal, **25**

- $\mathcal{I}_p^o(X)$ , **2**  
 $\mathcal{I}_p(X)$ , **2**  
*J*-*p* central element, **196**  
 $\mathcal{K}$ , **2**  
 $\mathcal{KbI}\mathcal{J}^p(G)$ , **90**  
 $\mathcal{KbI}^p(G)$ , **90**, 146  
 Klinger, K., 3  
 Koehl, R., 9  
 $\mathcal{K}_p$ , **2**  
 $L_2(p^p)$  field triple, **37**, 44  
 $L_3^{cs}(H)$ , **126**  
 $\mathcal{L}\mathcal{C}_p$ -uniqueness subgroup, **35**  
 locally unbalancing triple, **13**, 36  
 $\mathcal{L}_p^o(G)$ , **2**  
 $m_{2,p}(G)$ , **2**, **34**  
 Mason, G., 3  
 mate, **5**, **90**, 148  
 maximal symplectic triple, **219**  
 Meierfrankenfeld, U., 10, 31  
 $\widehat{m}_p(K)$ , **16**, **312**  
 $\mathcal{V}_G^*(D; 2)$ , **77**  
 neighborhood of  $U_4(2)$ -type, **8**  
 neighbors in a neighborhood, **8**  
 $\mathcal{V}_G^*(B; 2)$ , **5**  
 Nickel, W., 9  
 nonconstrained  $\{t, b\}$  neighborhood, **6**, **90**  
 nonconstrained neighborhood, **90**, 210  
 nondegenerate constrained neighborhood,  
**8**, **209**  
 nondegenerate neighborhood, **90**, 160  
 nondegenerate nonconstrained  
 neighborhood, **6**  
 $\mathcal{V}_X(B; 2)$ , **34**  
 $N(x, K)$ , **311**  
 one-step rigid, **25**  
 ordered frame, **441**  
 outer well-generated, 100  
*p*-component preuniqueness subgroup, **35**  
 Phan, K.-W., 9  
 Powell, M., 95  
 preuniqueness subgroup, 35  
*p*-saturated, **374**  
*p*-Thin Configuration, 314  
*p*-uniqueness subgroup, **35**  
 $q(K)$ , for  $K \in \mathcal{C}hev$ , **165**  
 regular, **90**  
 regular triple, **5**, 146  
 regularly embedded, **90**  
 regularly embedded triple, 146  
 restricted even type, 2  
 $\sigma_0(G)$ , **3**, 309  
 saturated, **374**  
 Segev, Y., 10, 31  
 $\sigma(G)$ , **2**, 309  
 Smith, S. D., 10  
 Solomon, R., 9, 10  
*Spor*, **2**  
 sporadic type neighborhood, **8**  
 $S^p(X)$ , **34**, **89**  
 strong *p*-uniqueness subgroup, 3, 34, **35**  
 strongly *p*-embedded subgroup, 34, **35**  
 strongly balanced, **39**  
 strongly locally balanced component, 14, 15  
 symplectic pair, **89**, 143  
 symplectic pair (trivial, faithful), **4**  
 $\{t, b\}$ -neighborhood, **160**  
 $\mathcal{T}\mathcal{G}_3$ -group, **311**  
 $\mathcal{T}\mathcal{G}_p(G)$ , **313**  
 Theorem  $\mathcal{C}_5$ , **1**  
 Theorem  $\mathcal{C}_5$ : Stage 1, **4**, **34**  
 Theorem  $\mathcal{C}_5$ : Stage 2, **6**  
 Theorem  $\mathcal{C}_5$ : Stage 3, **9**  
 Theorem  $\mathcal{C}_5$ : Stage 4, **9**  
 Theorem  $\mathcal{C}_6$ , 310  
 Theorem  $\mathcal{C}_6^*$ , 312  
 Theorem  $\mathcal{C}_7$ , proof of, 314  
 Theorems  $\mathcal{C}_7, \mathcal{C}_7^*$ , 312  
 $\Theta_1(a)$ , 11  
 $\Theta_1(G; A)$ , **11**  
 Thompson, J. G., 3  
 Thwaites, G., 95  
 $\mathcal{T}\mathcal{J}'_p(G)$ , **313**  
 $\mathcal{T}\mathcal{J}^*_p(G)$ , **313**  
 $\mathcal{T}\mathcal{J}_p(G)$ , **313**  
 $\mathcal{T}_p$ , **309**  
 $\mathcal{T}_p^i$ ,  $1 \leq i \leq 5$ , **310**  
 trivial symplectic pair, **89**, 143  
 unambiguously, **347**  
 unbalancing triple, **13**, 36  
 $\mathbf{V}_1, \mathbf{V}_2, \mathbf{V}_3, \mathbf{V}_4, \mathbf{V}_5, \mathbf{V}_6, \mathbf{V}_K$ , **1**  
 Walter, J. H., 3  
 weak Curtis-Tits system, 273  
 weak Phan system, 275  
 width, 220  
 width of an extraspecial or symplectic type  
 group, **20**  
 Wong, W., 9, 10  
 $w(S)$ , **20**  
 $Z_4$ -semirigid, **24**  
 $Z_4$ -semirigid in  $J$ , **24**  
 $\{z, b\}$ -neighborhood, **209**