ARC INDEX OF PRETZEL KNOTS OF TYPE $(-p, q, r)$

HWA JEONG LEE AND GYO TAEK JIN

(Communicated by Daniel Ruberman)

ABSTRACT. We computed the arc index for some of the pretzel knots $K = P(-p, q, r)$ with $p, q, r \geq 2$, $r \geq q$ and at most one of p, q, r is even. If $q = 2$, then the arc index $\alpha(K)$ equals the minimal crossing number $c(K)$. If $p \geq 3$ and $q = 3$, then $\alpha(K) = c(K) - 1$. If $p \geq 5$ and $q = 4$, then $\alpha(K) = c(K) - 2$.

1. ARC PRESENTATION

Let D be a diagram of a knot or a link L . Suppose that there is a simple closed curve C meeting D in k distinct points which divide D into k arcs $\alpha_1, \alpha_2, \dots, \alpha_k$ with the following properties:

- (1) Each α_i has no self-crossing.
- (2) If α_i crosses over α_j at a crossing, then $i > j$ and it crosses over α_j at any other crossings with α_j .
- (3) For each i , there exists an embedded disk d_i such that $\partial d_i = C$ and $\alpha_i \subset d_i$.
- (4) $d_i \cap d_j = C$, for distinct i and j .

Then the pair (D, C) is called an *arc presentation* of L with k arcs, and C is called the *axis* of the arc presentation. Figure 1 shows an arc presentation of the trefoil knot. The thick round curve is the axis. It is known that every knot or link has an arc presentation [3, 4]. For a given knot or link L , the minimal number of arcs in all arc presentations of L is called the *arc index* of L , denoted by $\alpha(L)$.

FIGURE 1. An arc presentation of the right-handed trefoil knot

By removing a point P from C away from L , we may identify $C \setminus P$ with the z -axis and each $d_i \setminus P$ with a vertical half plane along the z -axis. This shows that an arc presentation is equivalent to an open-book presentation.

Received by the editors April 3, 2012 and, in revised form, September 12, 2012, August 24, 2013 and December 20, 2013.

2010 *Mathematics Subject Classification.* Primary 57M27; Secondary 57M25.

The first author was supported in part by the National Research Foundation of Korea Grant funded by the Korean Government (NRF-2010-0024630).

The second author was supported in part by the National Research Foundation of Korea Grant funded by the Korean Government (NRF-2011-0027989).

FIGURE 2. An open-book presentation of the right-handed trefoil knot

Given a link L , let $c(L)$ denote the minimal crossing number of L .

Theorem 1.1 (Jin-Park). *A prime link L is nonalternating if and only if*

$$\alpha(L) \leq c(L).$$

2. KAUFFMAN POLYNOMIAL

The *Kauffman polynomial* $F_L(a, z)$ of an oriented knot or link L is defined by

$$F_L(a, z) = a^{-w(D)} \Lambda_D(a, z)$$

where D is a diagram of L , $w(D)$ the writhe of D and $\Lambda_D(a, z)$ the polynomial determined by the rules K1, K2 and K3.

(K1) $\Lambda_O(a, z) = 1$ where O is the trivial knot diagram.

(K2) For any four diagrams D_+ , D_- , D_0 and D_∞ which are identical outside a small disk in which they differ as shown below,

we have the relation

$$\Lambda_{D_+}(a, z) + \Lambda_{D_-}(a, z) = z(\Lambda_{D_0}(a, z) + \Lambda_{D_\infty}(a, z)).$$

(K3) For any three diagrams D_+ , D and D_- which are identical outside a small disk in which they differ as shown below,

we have the relation

$$a \Lambda_{D_+}(a, z) = \Lambda_D(a, z) = a^{-1} \Lambda_{D_-}(a, z).$$

For a connected sum and a split union of two diagrams, Λ satisfies the following properties:

(K4) If D is a connected sum of D_1 and D_2 , then

$$\Lambda_D(a, z) = \Lambda_{D_1}(a, z) \Lambda_{D_2}(a, z).$$

(K5) If D is the split union of D_1 and D_2 , then

$$\Lambda_D(a, z) = (z^{-1}a - 1 + z^{-1}a^{-1}) \Lambda_{D_1}(a, z) \Lambda_{D_2}(a, z).$$

The Laurent degree in the variable a of the Kauffman polynomial $F_L(a, z)$ is denoted by $\text{spread}_a(F_L)$ and defined by the formula

$$\text{spread}_a(F_L) = \max\text{-deg}_a(F_L) - \min\text{-deg}_a(F_L).$$

Notice that $\text{spread}_a(F_L) = \text{spread}_a(\Lambda_D)$ for any diagram D of L . The following theorem gives an important lower bound for the arc index.

Theorem 2.1 (Morton-Beltrami). *Let L be a link. Then*

$$\alpha(L) \geq \text{spread}_a(F_L) + 2.$$

If L is nonsplit and alternating, then the equality holds so that $\alpha(L) = c(L) + 2$. This is shown by Bae and Park [1] using arc presentations in the form of wheel diagrams.

3. PRETZEL KNOTS

Given a sequence of integers $p_1, p_2, \dots, p_n$, we connect two disjoint disks by n bands with p_i half twists, $i = 1, 2, \dots, n$, so that the boundary of the resulting surface is a link as shown in Figure 3. This link is called the *pretzel link* of type $(p_1, p_2, \dots, p_n)$ and denoted by $P(p_1, p_2, \dots, p_n)$.

FIGURE 3. Pretzel links $P(p_1, p_2, \dots, p_n)$ and $P(-p, q, r)$

In the case $n = 3$, the pretzel links satisfy the following properties:

Proposition 3.1. *Let p, q , and r be nonzero integers.*

- (1) *The link type of $P(p, q, r)$ is independent of the order of p, q, r .*
- (2) *$P(p, q, r)$ is a knot if and only if at most one of p, q, r is an even number.*

In this work, we compute the arc index for the pretzel knots $K = P(-p, q, r)$ with $p, q, r \geq 2$. By Proposition 3.1(1), we may assume that $r \geq q$. By Theorem 3.2, we know that $P(-p, q, r)$ is a minimal crossing diagram of K , i.e., $c(K) = p + q + r$.

Theorem 3.2 (Lickorish-Thistlethwaite). *If a link L admits a reduced Montesinos diagram having n crossings, then L cannot be projected with fewer than n crossings.*

This work was motivated by Theorem 3.3 which is a special case of Theorem 1.1.

Theorem 3.3 (Beltrami-Cromwell). *If $K = P(-p, q, r)$ is a knot with $p, q, r \geq 2$, then*

$$\alpha(K) \leq c(K) = p + q + r.$$

By computing $\text{spread}_a(F_K)$ and finding arc presentations of $K = P(-p, q, r)$ with the minimum number of arcs for various values of p, q and r , we obtained sharper results.

4. MAIN RESULTS

Theorem 4.1. *If $K = P(-2, q, r)$ is a knot with $3 \leq q \leq r$, then*

$$\alpha(K) \leq c(K) - 1.$$

Theorem 4.2. *If $K = P(-p, 2, r)$ is a knot with $p \geq 3$, $r \geq 3$, then*

$$\alpha(K) = c(K).$$

Theorem 4.3. *If $K = P(-p, 3, r)$ is a knot with $p \geq 3$, $r \geq 3$, then*

$$\alpha(K) = c(K) - 1.$$

Theorem 4.4. *If $K = P(-p, 4, r)$ is a knot with $p \geq 5$, $r \geq 5$, then*

$$\alpha(K) = c(K) - 2.$$

Theorem 4.5. *If $K = P(-3, 4, r)$ is a knot with $r \geq 7$, then*

$$c(K) - 4 \leq \alpha(K) \leq c(K) - 2.$$

5. ARC PRESENTATIONS OF $P(-p, q, r)$

Proposition 5.1. *If $K = P(-2, q, r)$ is a knot with $3 \leq q \leq r$, then K has an arc presentation with $q + r + 1$ arcs.*

FIGURE 4. An arc presentation of $P(-2, q, r)$

Proof. Figure 4 shows a pretzel diagram of $P(-2, q, r)$ and its arc presentation with $q + r + 1$ arcs. The thick curve is the axis of the arc presentation which cuts the knot at 1 place in the leftmost box, $q - 1$ places in the second, 2 places in the third, and $r - 1$ places in the fourth. The $q + r + 1$ arcs of the knot satisfy the four properties of an arc presentation. $\square$

Proposition 5.2. *If $K = P(-p, q, r)$ is a knot with $p \geq 3$ and $2 \leq q \leq 3 \leq r$, then K has an arc presentation with $p + r + 2$ arcs.*

Proof. For each of $q = 2, 3$, Figure 5 shows a pretzel diagram of $P(-p, q, r)$ and its arc presentation with $p + r + 2$ arcs. The thick curve is the axis of the arc presentation which cuts the knot at $p - 1$ places in the leftmost box, 4 places in the second, and $r - 1$ places in the third. The $p + r + 2$ arcs of the knot satisfy the four properties of an arc presentation. $\square$

FIGURE 5. Arc presentations of $P(-p, q, r)$ for $q = 2, 3$

Proposition 5.3. *If $K = P(-p, q, r)$ is a knot with $p \geq 3$ and $4 \leq q \leq r$, then K has an arc presentation with $p + q + r - 2$ arcs.*

Proof. In Figure 6, the diagram (a) shows a pretzel diagram of $P(-p, q, r)$ with $p \geq 3$ and $4 \leq q \leq r$. The diagram (b) is obtained from (a) by two applications of the Reidemeister move of type 3. The diagram (c) shows an arc presentation with $p + q + r - 1$ arcs. The diagram (d) is obtained from (c) by isotoping the arc labeled x over the axis so that there are only $p + q + r - 2$ arcs. Each of the seven boxes, from left to right, contains 1, $p - 3$, 2, 3, $q - 3$, 1, and $r - 3$ arcs, respectively. $\square$

FIGURE 6. Arc presentations of $P(-p, q, r)$ with $p \geq 3$ and $4 \leq q \leq r$

6. THE KAUFFMAN POLYNOMIAL OF THE PRETZEL KNOTS $P(-p, q, r)$

For any link diagram D , the polynomial Λ_D is of the form

$$\Lambda_D(a, z) = \sum_{i=m}^n f_i(z) a^i$$

where m, n are integers with $m \leq n$, and $f_i(z)$'s are polynomials in z with integer coefficients such that $f_m(z) \neq 0$ and $f_n(z) \neq 0$. To simplify our computation of $\text{spread}_a(\Lambda_D)$ we use the notation

$$f_m(z) = \langle k_m z^{h_m} \rangle,$$

$$f_n(z) = \langle k_n z^{h_n} \rangle,$$

$$\sum_{i=m}^n f_i(z) a^i = [\langle k_n z^{h_n} \rangle a^n, \langle k_m z^{h_m} \rangle a^m], \quad (m < n)$$

where $k_m z^{h_m}$ and $k_n z^{h_n}$ are the highest degree terms in $f_m(z)$ and $f_n(z)$, respectively. For example, we write

$$z(z^2 - 1)a^{-1} + z^2 a^{-2} - 2za^{-3} = [\langle z^3 \rangle a^{-1}, \langle -2z \rangle a^{-3}].$$

We also use the notation $\Lambda_{(p_1, p_2, \dots, p_n)}$ for Λ_D when $D = P(p_1, p_2, \dots, p_n)$.

FIGURE 7. Links $P(-m, 0)$, $P(0, n)$, and $P(-m, n)$

Let the polynomial $\Phi_i(z)$ be defined by $\Phi_0(z) = 1$, $\Phi_1(z) = z$ and

$$\Phi_{i+1}(z) = z\Phi_i(z) - \Phi_{i-1}(z).^1$$

Lemma 6.1. *Let m, n be nonnegative integers. Then*

$$\Lambda_{(-m, n)} = \begin{cases} [\langle z \rangle a^{k-1}, \langle z^{k-1} \rangle a^{-1}] & \text{if } k = m - n > 1, \\ a^{-1} & \text{if } k = m - n = 1, \\ z^{-1}a - 1 + z^{-1}a^{-1} & \text{if } k = n - m = 0, \\ a & \text{if } k = n - m = 1, \\ [\langle z^{k-1} \rangle a, \langle z \rangle a^{-(k-1)}] & \text{if } k = n - m > 1. \end{cases}$$

More precisely,

$$\Lambda_{(-m, n)} = \begin{cases} [za^{m-1}, z^{-1}\Phi_m(z)a^{-1}] & \text{if } m \geq 3, \quad n = 0, \\ [z^{-1}\Phi_n(z)a, za^{-(n-1)}] & \text{if } m = 0, \quad n \geq 3. \end{cases}$$

¹ $\Phi_i(2z)$'s are the Chebyshev's polynomials of the second type.

Proof. Using the skein relations K1, K2, and K3, we have the following

$$\Lambda_{(-1,0)} = a^{-1}, \quad \Lambda_{(0,1)} = a$$

$$\Lambda_{(0,0)} = z^{-1}a - 1 + z^{-1}a^{-1}.$$

Let D_v be a vertical integer tangle which has $|v|$ times half-twists in the positive or negative direction according to the sign of v for an integer v . D_{+1} , D_{-1} , D_0 , and D_∞ are exemplified in (K2). In the sublemma below, we use the following simplified notation:

$$\Lambda_v = \Lambda_{D_v}, \quad \Lambda_\infty = \Lambda_{D_\infty}.$$

Sublemma. *For $m \geq 2$ and $n \geq 2$, we have*

$$(6.1) \quad \Lambda_{-m} = -\Phi_{m-2}(z) \Lambda_0 + \Phi_{m-1}(z) \Lambda_{-1} + \left(\sum_{i=0}^{m-2} z \Phi_i(z) a^{m-1-i} \right) \Lambda_\infty,$$

$$(6.2) \quad \Lambda_n = -\Phi_{n-2}(z) \Lambda_0 + \Phi_{n-1}(z) \Lambda_{+1} + \left(\sum_{j=0}^{n-2} z \Phi_j(z) a^{-(n-1-j)} \right) \Lambda_\infty.$$

Proof of Sublemma. As shown below, equation (6.1) holds when $m = 2, 3$.

$$\begin{aligned} \Lambda_{-2} &= -\Lambda_0 + z\Lambda_{-1} + za\Lambda_\infty \\ &= -\Phi_0(z) \Lambda_0 + \Phi_1(z) \Lambda_{-1} + z\Phi_0(z)a \Lambda_\infty \\ \Lambda_{-3} &= -\Lambda_{-1} + z\Lambda_{-2} + za^2\Lambda_\infty \\ &= -\Phi_0(z) \Lambda_{-1} + z(-\Phi_0(z) \Lambda_0 + \Phi_1(z) \Lambda_{-1} + z\Phi_0(z)a \Lambda_\infty) + za^2\Lambda_\infty \\ &= -\Phi_1(z) \Lambda_0 + \Phi_2(z) \Lambda_{-1} + \left(\sum_{i=0}^1 z \Phi_i(z) a^{2-i} \right) \Lambda_\infty. \end{aligned}$$

Suppose that equation (6.1) holds for $2 \leq m \leq k-1$ for some $k \geq 4$. Then

$$\begin{aligned} \Lambda_{-k} &= -\Lambda_{-(k-2)} + z\Lambda_{-(k-1)} + za^{k-1}\Lambda_\infty \\ &= - \left(-\Phi_{k-4}(z) \Lambda_0 + \Phi_{k-3}(z) \Lambda_{-1} + \left(\sum_{i=0}^{k-4} z \Phi_i(z) a^{k-3-i} \right) \Lambda_\infty \right) \\ &\quad + z \left(-\Phi_{k-3}(z) \Lambda_0 + \Phi_{k-2}(z) \Lambda_{-1} + \left(\sum_{i=0}^{k-3} z \Phi_i(z) a^{k-2-i} \right) \Lambda_\infty \right) + za^{k-1}\Lambda_\infty \\ &= -\Phi_{k-2}(z) \Lambda_0 + \Phi_{k-1}(z) \Lambda_{-1} + \left(\sum_{i=0}^{k-2} z \Phi_i(z) a^{k-1-i} \right) \Lambda_\infty. \end{aligned}$$

This proves that equation (6.1) holds for $m \geq 2$. In a similar manner, we can prove that equation (6.2) holds for $n \geq 2$. □

□

For $m \geq 2$ and $n \geq 2$, using equations (6.1) and (6.2) on m and n respectively, we obtain

$$\begin{aligned}
\Lambda_{(-m,0)} &= -\Phi_{m-2}(z)\Lambda_{(0,0)} + \Phi_{m-1}(z)\Lambda_{(-1,0)} + \sum_{i=0}^{m-2} z\Phi_i(z)a^{m-1-i} \\
&= \begin{cases} [(z - z^{-1})a, (z - z^{-1})a^{-1}] & \text{if } m = 2, \\ [za^{m-1}, \{\Phi_{m-1}(z) - z^{-1}\Phi_{m-2}(z)\}a^{-1}] & \text{if } m > 2 \end{cases} \\
&= \begin{cases} [z^{-1}\Phi_2(z)a, z^{-1}\Phi_2(z)a^{-1}] & \text{if } m = 2, \\ [za^{m-1}, z^{-1}\Phi_m(z)a^{-1}] & \text{if } m > 2. \end{cases} \\
\Lambda_{(0,n)} &= -\Phi_{n-2}(z)\Lambda_{(0,0)} + \Phi_{n-1}(z)\Lambda_{(0,1)} + \sum_{j=0}^{n-2} z\Phi_j(z)a^{-(n-1-j)} \\
&= \begin{cases} [(z - z^{-1})a, (z - z^{-1})a^{-1}] & \text{if } n = 2, \\ [\{\Phi_{n-1}(z) - z^{-1}\Phi_{n-2}(z)\}a, za^{-(n-1)}] & \text{if } n > 2 \end{cases} \\
&= \begin{cases} [z^{-1}\Phi_2(z)a, z^{-1}\Phi_2(z)a^{-1}] & \text{if } n = 2, \\ [z^{-1}\Phi_n(z)a, za^{-(n-1)}] & \text{if } n > 2. \end{cases}
\end{aligned}$$

Since $\Lambda_D(a, z)$ is an invariant under regular isotopy of diagrams, we have

$$\Lambda_{(-m,n)} = \begin{cases} \Lambda_{(-k,0)} & \text{if } k = m - n \geq 1, \\ \Lambda_{(0,0)} & \text{if } k = m - n = 0, \\ \Lambda_{(0,k)} & \text{if } k = n - m \geq 1. \end{cases}$$

This completes the proof. $\square$

Lemma 6.2.

$$\begin{aligned}
\Lambda_{(-p,0,r)} &= \begin{cases} 1 & \text{if } p = r = 1, \\ [\langle z \rangle a^p, \langle z^{p-1} \rangle a^0] & \text{if } p > 1, \ r = 1, \\ [\langle z^r \rangle a^p, \langle z^p \rangle a^{-r}] & \text{if } p > 1, \ r > 1. \end{cases} \\
\Lambda_{(-p,1,r)} &= \begin{cases} a^2 & \text{if } p = 0, \ r = 1, \\ [\langle z^{r-1} \rangle a^2, \langle z \rangle a^{-(r-2)}] & \text{if } p = 0, \ r > 1, \\ a^{-r} & \text{if } p = 1, \ r \geq 1, \\ [\langle z^{r+1} \rangle a^p, \langle z^{p-1} \rangle a^{-r}] & \text{if } p > 1, \ r \geq 1. \end{cases}
\end{aligned}$$

More precisely, for $p \geq 3$ and $r \geq 3$

$$\Lambda_{(-p,0,r)} = [\Phi_r(z)a^p, \Phi_p(z)a^{-r}]$$

and for $r \geq 2$

$$\Lambda_{(-p,1,r)} = \begin{cases} [z^{-1}\Phi_{r+2}(z)a^2, \Phi_1(z)a^{-r}] & \text{if } p = 2, \\ [\Phi_{r+1}(z)a^p, \Phi_{p-1}(z)a^{-r}] & \text{if } p > 2. \end{cases}$$

Proof. Since $P(-p,0,r) = P(-p,0) \sharp P(0,r)$, we have $\Lambda_{(-p,0,r)} = \Lambda_{(-p,0)} \Lambda_{(0,r)}$. Therefore the formula about $\Lambda_{(-p,0,r)}$ follows from Lemma 6.1.

Now we consider the formula about $\Lambda_{(-p,1,r)}$. Three cases with $p = 0$ or $p = 1$ follow from K1, K2, K3, and Lemma 6.1. The other case is derived by equation (6.1). For $p \geq 2$, we have

$$\begin{aligned}
 \Lambda_{(-p,1,r)} &= -\Phi_{p-2}(z)\Lambda_{(0,1,r)} + \Phi_{p-1}(z)\Lambda_{(-1,1,r)} + \sum_{i=0}^{p-2} z\Phi_i(z)a^{p-1-i}\Lambda_{(1,r)} \\
 &= -\Phi_{p-2}(z)a\Lambda_{(0,r)} + \Phi_{p-1}(z)a^{-r} + \sum_{i=0}^{p-2} z\Phi_i(z)a^{p-1-i}\Lambda_{(0,r+1)} \\
 &= \begin{cases} [\{-z^{-1}\Phi_r(z) + \Phi_{r+1}(z)\}a^2, \Phi_1(z)a^{-r}] & \text{if } p = 2, \\ [\Phi_{r+1}(z)a^p, \Phi_{p-1}(z)a^{-r}] & \text{if } p > 2 \end{cases} \\
 &= \begin{cases} [z^{-1}\Phi_{r+2}(z)a^2, \Phi_{p-1}(z)a^{-r}] & \text{if } p = 2, \\ [\Phi_{r+1}(z)a^p, \Phi_{p-1}(z)a^{-r}] & \text{if } p > 2. \end{cases}
 \end{aligned}$$

This completes the proof. $\square$

Proposition 6.3. $\text{spread}_a(\Lambda_{(-p,2,r)}(a, z)) = p + r$ for $p \geq 3$, and $r \geq 3$.

Proof. For $p \geq 3$, we show that

$$(6.3) \quad \Lambda_{(-p,2,r)} = \begin{cases} [\langle z^3 \rangle a^p, \langle z^{p-1} \rangle a^{-2}] & (\text{if } r = 1), \\ [\langle z^4 \rangle a^p, \langle 3z^{p-2} \rangle a^{-2}] & (\text{if } r = 2), \\ [\langle z^{r+2} \rangle a^p, \langle z^{p-2} \rangle a^{-r}] & (\text{if } r \geq 3). \end{cases}$$

Using K1, K2, K3 and Lemmas 6.1 and 6.2, we obtain

$$\begin{aligned}
 \Lambda_{(-p,2,1)} &= -\Lambda_{(-p,0,1)} + z\Lambda_{(-p,1,1)} + za^{-1}\Lambda_{(-p,1)} \\
 &= -[\langle z \rangle a^p, \langle z^{p-1} \rangle a^0] + z[\langle z^2 \rangle a^p, \langle z^{p-1} \rangle a^{-1}] + za^{-1}[\langle z \rangle a^{p-2}, \langle z^{p-2} \rangle a^{-1}] \\
 &= [\langle z^3 \rangle a^p, \langle z^{p-1} \rangle a^{-2}], \\
 \Lambda_{(-p,2,2)} &= -\Lambda_{(-p,0,2)} + z\Lambda_{(-p,1,2)} + za^{-1}\Lambda_{(-p,2)} \\
 &= -[za^{p-1}, z^{-1}\Phi_p(z)a^{-1}][z^{-1}\Phi_2(z)a, z^{-1}\Phi_2(z)a^{-1}] \\
 &\quad + z[\Phi_3(z)a^p, \Phi_{p-1}(z)a^{-2}] \\
 &\quad + za^{-1} \begin{cases} a^{-1} & (\text{if } p = 3), \\ [z^{-1}\Phi_2(z)a, z^{-1}\Phi_2(z)a^{-1}] & (\text{if } p = 4), \\ [za^{p-3}, z^{-1}\Phi_{p-2}(z)a^{-1}] & (\text{if } p \geq 5) \end{cases} \\
 &= -[\Phi_2(z)a^p, z^{-2}\Phi_2(z)\Phi_p(z)a^{-2}] + z[\Phi_3(z)a^p, \Phi_{p-1}(z)a^{-2}] \\
 &\quad + \begin{cases} za^{-2} & (\text{if } p = 3), \\ za^{-1}[z^{-1}\Phi_2(z)a, z^{-1}\Phi_2(z)a^{-1}] & (\text{if } p = 4), \\ za^{-1}[za^{p-3}, z^{-1}\Phi_{p-2}(z)a^{-1}] & (\text{if } p \geq 5) \end{cases} \\
 &= [\Phi_4(z)a^p, \langle 3z^{p-2} \rangle a^{-2}],
 \end{aligned}$$

which prove the first two cases of (6.3). Now we show the third case of (6.3) by an induction on r . For $r = 3$, we have

$$\begin{aligned}\Lambda_{(-p,2,3)} &= -\Lambda_{(-p,2,1)} + z\Lambda_{(-p,2,2)} + za^{-2}\Lambda_{(-p,2)} \\ &= -[\langle z^3 \rangle a^p, \langle z^{p-1} \rangle a^{-2}] + z[\langle z^4 \rangle a^p, \langle 3z^{p-2} \rangle a^{-2}] \\ &\quad + \begin{cases} za^{-3} & (\text{if } p = 3), \\ [\langle z^2 \rangle a^{p-5}, \langle z^{p-2} \rangle a^{-3}] & (\text{if } p \geq 4) \end{cases} \\ &= [\langle z^5 \rangle a^p, \langle z^{p-2} \rangle a^{-3}],\end{aligned}$$

and for $r \geq 4$, inductively, we have

$$\begin{aligned}\Lambda_{(-p,2,r)} &= -\Lambda_{(-p,2,r-2)} + z\Lambda_{(-p,2,r-1)} + za^{-(r-1)}\Lambda_{(-p,2)} \\ &= -[\langle z^r \rangle a^p, \langle * \rangle a^{-(r-2)}] + z[\langle z^{r+1} \rangle a^p, \langle z^{p-2} \rangle a^{-(r-1)}] \\ &\quad + \begin{cases} za^{-r} & (\text{if } p = 3), \\ [\langle z^2 \rangle a^{p-r-2}, \langle z^{p-2} \rangle a^{-r}] & (\text{if } p \geq 4) \end{cases} \\ &= [\langle z^{r+2} \rangle a^p, \langle z^{p-2} \rangle a^{-r}]\end{aligned}$$

where $\langle * \rangle a^{-(r-2)}$ indicates that the lowest a -degree of $\Lambda_{(-p,2,r-2)}$ is not smaller than $-(r-2)$.

This completes the proof. $\square$

Proposition 6.4. $\text{spread}_a(\Lambda_{(-p,3,r)}(a, z)) = p + r$ for $p \geq 3$ and $r \geq 3$.

Proof. We show that

$$\Lambda_{(-p,3,r)} = \begin{cases} [\langle z^6 \rangle a^p, \langle 2z^{p-3} \rangle a^{-3}] & (\text{if } r = 3), \\ [\langle z^{r+3} \rangle a^p, \langle z^{p-3} \rangle a^{-r}] & (\text{if } r \geq 4). \end{cases}$$

Using equation (6.2) and Lemmas 6.1 and 6.2, we obtain

$$\begin{aligned}\Lambda_{(-p,3,3)} &= -\Phi_1(z)\Lambda_{(-p,0,3)} + \Phi_2(z)\Lambda_{(-p,1,3)} + (za^{-2} + z^2a^{-1})\Lambda_{(-p,3)} \\ &= -\Phi_1(z)[za^{p-1}, z^{-1}\Phi_p(z)a^{-1}][z^{-1}\Phi_3(z)a, za^{-2}] \\ &\quad + \Phi_2(z)[\Phi_4a^p, \Phi_{p-1}(z)a^{-3}] \\ &\quad + (za^{-2} + z^2a^{-1}) \begin{cases} (z^{-1}a - 1 + z^{-1}a^{-1}) & (\text{if } p = 3), \\ a^{-1} & (\text{if } p = 4), \\ [\langle z \rangle a^{p-4}, \langle z^{p-4} \rangle a^{-1}] & (\text{if } p \geq 5) \end{cases} \\ &= [\langle z^6 \rangle a^p, \Phi_{p-3}(z)a^{-3}] + \begin{cases} [z, a^{-3}] & (\text{if } p = 3), \\ [z^2a^{-2}, za^{-3}] & (\text{if } p = 4), \\ [\langle z^3 \rangle a^{p-5}, \langle z^{p-3} \rangle a^{-3}] & (\text{if } p \geq 5) \end{cases} \\ &= [\langle z^6 \rangle a^p, \langle 2z^{p-3} \rangle a^{-3}].\end{aligned}$$

Using K1, K2, K3 and the results above, we obtain

$$\begin{aligned}
 \Lambda_{(-p,3,4)} &= -\Lambda_{(-p,2,3)} + z\Lambda_{(-p,3,3)} + za^{-3}\Lambda_{(-p,3)} \quad (\because \Lambda_{(-p,3,2)} = \Lambda_{(-p,2,3)}) \\
 &= -[\langle z^5 \rangle a^p, \langle z^{p-2} \rangle a^{-2}] + z[\langle z^6 \rangle a^p, \langle 2z^{p-3} \rangle a^{-3}] \\
 &\quad + za^{-3} \begin{cases} (z^{-1}a - 1 + z^{-1}a^{-1}) & (\text{if } p = 3), \\ a^{-1} & (\text{if } p = 4), \\ [\langle z \rangle a^{p-4}, \langle z^{p-4} \rangle a^{-1}] & (\text{if } p \geq 5) \end{cases} \\
 &= [\langle z^7 \rangle a^p, \langle z^{p-3} \rangle a^{-4}],
 \end{aligned}$$

and, for $r \geq 5$, inductively, we have

$$\begin{aligned}
 \Lambda_{(-p,3,r)} &= -\Lambda_{(-p,3,r-2)} + z\Lambda_{(-p,3,r-1)} + za^{-(r-1)}\Lambda_{(-p,3)} \\
 &= -[\langle z^{r+1} \rangle a^p, \langle * \rangle a^{-(r-2)}] + z[\langle z^{r+2} \rangle a^p, \langle z^{p-3} \rangle a^{-(r-1)}] \\
 &\quad + za^{-(r-1)} \begin{cases} (z^{-1}a - 1 + z^{-1}a^{-1}) & (\text{if } p = 3), \\ a^{-1} & (\text{if } p = 4), \\ [\langle z \rangle a^{p-4}, \langle z^{p-4} \rangle a^{-1}] & (\text{if } p \geq 5) \end{cases} \\
 &= [\langle z^{r+3} \rangle a^p, \langle z^{p-3} \rangle a^{-r}].
 \end{aligned}$$

This completes the proof. $\square$

Proposition 6.5. $\text{spread}_a(\Lambda_{(-3,4,r)}(a, z)) = r + 1$ for $r \geq 7$.

Proof.

$$\Lambda_{(-3,4,0)} = \Lambda_{(-3,0)}\Lambda_{(0,4)} = [\Phi_4(z)a^3, \Phi_3(z)a^{-4}].$$

Using equation (6.1) and K1, K3 and Lemma 6.1, we obtain

$$\begin{aligned}
 \Lambda_{(-3,4,1)} &= -\Phi_1(z)\Lambda_{(0,4,1)} + \Phi_2(z)\Lambda_{(-1,4,1)} + \left(\sum_{i=0}^1 z\Phi_i(z)a^{2-i}\right)\Lambda_{(4,1)} \\
 &= -za\Lambda_{(0,4)} + \Phi_2(z)a^{-4} + (za^2 + z^2a)\Lambda_{(0,5)} \\
 &= -za[\langle z^{-1}\Phi_4(z)a, za^{-3} \rangle] + \Phi_2(z)a^{-4} + (za^2 + z^2a)[z^{-1}\Phi_5(z)a, za^{-4}] \\
 &= [\Phi_5(z)a^3, \Phi_2(z)a^{-4}].
 \end{aligned}$$

Using equation (6.2) and the above two formulas, for $r \geq 7$ we obtain

$$\begin{aligned}
 \Lambda_{(-3,4,r)} &= -\Phi_{r-2}(z)\Lambda_{(-3,4,0)} + \Phi_{r-1}(z)\Lambda_{(-3,4,1)} + \sum_{i=0}^{r-2} z\Phi_i(z)a^{-(r-1-i)}\Lambda_{(-3,4)} \\
 &= -\Phi_{r-2}(z)[\Phi_4(z)a^3, \Phi_3(z)a^{-4}] \\
 &\quad + \Phi_{r-1}(z)[\Phi_5(z)a^3, \Phi_2(z)a^{-4}] + \sum_{i=0}^{r-2} z\Phi_i(z)a^{-(r-2-i)} \\
 &= [\langle z^{r+4} \rangle a^3, za^{-(r-2)}].
 \end{aligned}$$

This completes the proof. $\square$

Proposition 6.6. $\text{spread}_a(\Lambda_{(-p,4,r)}(a, z)) = p + r$ for $p \geq 5$ and $r \geq 5$.

Proof. Using equation (6.2) and Lemmas 6.1 and 6.2, we obtain

$$\begin{aligned}
\Lambda_{(-p,4,r)} &= -\Phi_2(z)\Lambda_{(-p,0,r)} + \Phi_3(z)\Lambda_{(-p,1,r)} + \sum_{i=0}^2 z\Phi_i(z)a^{-(3-i)}\Lambda_{(-p,r)} \\
&= -(z^2 - 1) [\Phi_r(z)a^p, \Phi_p(z)a^{-r}] \\
&\quad + (z^3 - 2z) [\Phi_{r+1}(z)a^p, \Phi_{p-1}(z)a^{-r}] \\
&\quad + \{za^{-3} + z^2a^{-2} + (z^3 - z)a^{-1}\} \begin{cases} [\langle z \rangle a^{k-1}, \langle z^{k-1} \rangle a^{-1}] & \text{if } k = p - r > 1, \\ a^{-1} & \text{if } k = p - r = 1, \\ z^{-1}a - 1 + z^{-1}a^{-1} & \text{if } k = p - r = 0, \\ a & \text{if } k = r - p = 1, \\ [\langle z^{k-1} \rangle a, \langle z \rangle a^{-(k-1)}] & \text{if } k = r - p > 1 \end{cases} \\
&= [\langle z^{r+4} \rangle a^p, \{-(z^2 - 1)\Phi_p(z) + (z^3 - 2z)\Phi_{p-1}(z)\}a^{-r}] \\
&= [\langle z^{r+4} \rangle a^p, \Phi_{p-4}(z)a^{-r}].
\end{aligned}$$

This completes the proof. $\square$

7. PROOFS OF MAIN RESULTS AND COMMENTS

Theorem 4.1 is proved by Proposition 5.1. Table 1 shows that the upper bound ‘ $c(K) - 1$ ’ for the arc index in Theorem 4.1 is best possible. It also shows that the lower bound ‘ $\text{spread}_a(F_K) + 2$ ’ in Theorem 2.1 is best possible.

TABLE 1. Examples of Theorem 4.1

Pretzel knot K	DT Name ²	$\text{spread}_a(F_K) + 2$	arc index	$c(K) - 1$
$P(-2, 3, 3)$	$8n3$	6	7	7
$P(-2, 3, 5)$	$10n21$	6	8	9
$P(-2, 3, 7)$	$12n242$	9	9	11
$P(-2, 5, 5)$	$12n725$	10	10	11

The proof of Theorem 4.2 is a combination of Propositions 5.2 and 6.3. The proof of Theorem 4.3 is a combination of Propositions 5.2 and 6.4. The proof of Theorem 4.4 is a combination of Propositions 5.3 and 6.6. The proof of Theorem 4.5 is a combination of Propositions 5.3 and 6.5. Table 2 shows that the upper bound ‘ $c(K) - 2$ ’ is best possible but the lower bound ‘ $c(K) - 4$ ’ may not be best possible.

TABLE 2. Examples of Theorem 4.5

Pretzel knot K	DT Name	$c(K) - 4$	arc index	$c(K) - 2$
$P(-3, 4, 5)$	$12n475$	8	10	10
$P(-3, 4, 7)$	$14n12205$	10	11	12

²The Dowker-Thistlethwaite name. See [10].

ACKNOWLEDGMENT

The authors would like to thank the referee for valuable comments and suggestions.

REFERENCES

- [1] Yongju Bae and Chan-Young Park, *An upper bound of arc index of links*, Math. Proc. Cambridge Philos. Soc. **129** (2000), no. 3, 491–500, DOI 10.1017/S0305004100004576. MR1780500 (2002f:57009)
- [2] Elisabetta Beltrami and Peter R. Cromwell, *Minimal arc-presentations of some nonalternating knots*, Topology Appl. **81** (1997), no. 2, 137–145, DOI 10.1016/S0166-8641(97)00024-2. MR1481138 (98h:57012)
- [3] H. Brunn, *Über verknötete Kurven*, Mathematiker-Kongresses Zurich 1897, Leipzig (1898) 256–259.
- [4] Peter R. Cromwell, *Embedding knots and links in an open book. I. Basic properties*, Topology Appl. **64** (1995), no. 1, 37–58, DOI 10.1016/0166-8641(94)00087-J. MR1339757 (96g:57006)
- [5] Peter R. Cromwell and Ian J. Nutt, *Embedding knots and links in an open book. II. Bounds on arc index*, Math. Proc. Cambridge Philos. Soc. **119** (1996), no. 2, 309–319, DOI 10.1017/S0305004100074181. MR1357047 (96m:57011)
- [6] Gyo Taek Jin and Wang Keun Park, *Prime knots with arc index up to 11 and an upper bound of arc index for non-alternating knots*, J. Knot Theory Ramifications **19** (2010), no. 12, 1655–1672, DOI 10.1142/S0218216510008595. MR2755494 (2012d:57006)
- [7] W. B. R. Lickorish and M. B. Thistlethwaite, *Some links with nontrivial polynomials and their crossing-numbers*, Comment. Math. Helv. **63** (1988), no. 4, 527–539, DOI 10.1007/BF02566777. MR966948 (90a:57010)
- [8] Hugh R. Morton and Elisabetta Beltrami, *Arc index and the Kauffman polynomial*, Math. Proc. Cambridge Philos. Soc. **123** (1998), no. 1, 41–48, DOI 10.1017/S0305004197002090. MR1474862 (98i:57015)
- [9] Knotscape, <http://www.math.utk.edu/~morwen/knotscape.html>
- [10] Table of Knot Invariants, <http://www.indiana.edu/~knotinfo/>

DEPARTMENT OF MATHEMATICS, CHUNG-ANG UNIVERSITY, 221 HEUKSEOK-DONG, DONGJAK-GU, SEOUL 156-756, KOREA

E-mail address: hjwith@cau.ac.kr

Current address: Department of Mathematical Sciences, KAIST, 291 Daehak-ro, Yuseong-gu, Daejeon 305-701, Korea

E-mail address: hjwith@kaist.ac.kr

DEPARTMENT OF MATHEMATICAL SCIENCES, KAIST, 291 DAEHAK-RO, YUSEONG-GU, DAEJEON 305-701, KOREA

E-mail address: trefoil@kaist.ac.kr