

## PROPER HOLOMORPHIC EMBEDDINGS WITH SMALL LIMIT SETS

FRANC FORSTNERIČ

(Communicated by Filippo Bracci)

ABSTRACT. Let  $X$  be a Stein manifold of dimension  $n \geq 1$ . Given a continuous positive increasing function  $h$  on  $\mathbb{R}_+ = [0, \infty)$  with  $\lim_{t \rightarrow \infty} h(t) = \infty$ , we construct a proper holomorphic embedding  $f = (z, w) : X \hookrightarrow \mathbb{C}^{n+1} \times \mathbb{C}^n$  satisfying  $|w(x)| < h(|z(x)|)$  for all  $x \in X$ . In particular,  $f$  may be chosen such that its limit set at infinity is a linearly embedded copy of  $\mathbb{C}\mathbb{P}^n$  in  $\mathbb{C}\mathbb{P}^{2n}$ .

### 1. THE MAIN RESULT

A theorem of Remmert [23], Narasimhan [21], and Bishop [4] states that every Stein manifold  $X$  of dimension  $n \geq 1$  admits a proper holomorphic map to  $\mathbb{C}^{n+1}$ , a proper holomorphic immersion to  $\mathbb{C}^{2n}$ , and a proper holomorphic embedding in  $\mathbb{C}^{2n+1}$ . (See also [18, Chap. VII.C].) We are interested in the question how much space proper holomorphic embeddings or immersions  $X \rightarrow \mathbb{C}^N$  need, and how small can their limit sets at infinity be.

By Remmert [22], the image  $A = f(X) \subset \mathbb{C}^N$  of a proper holomorphic map  $f : X \rightarrow \mathbb{C}^N$  is a closed complex subvariety of pure dimension  $n = \dim X$ . Such an  $A$  is algebraic if and only if it is contained, after a unitary change of coordinates on  $\mathbb{C}^N$ , in a domain of the form

$$D = \{(z, w) \in \mathbb{C}^n \times \mathbb{C}^p = \mathbb{C}^N : |w| < C(1 + |z|)\}$$

for some  $C > 0$  (see Chirka [5, Theorem 2, p. 77]). Equivalently, if  $H = \mathbb{C}\mathbb{P}^N \setminus \mathbb{C}^n \cong \mathbb{C}\mathbb{P}^{N-1}$  denotes the hyperplane at infinity and  $A_\infty = \overline{A} \cap H$ , where  $\overline{A}$  is the topological closure of  $A$  in  $\mathbb{C}\mathbb{P}^N$ , then  $A$  is algebraic if and only if there is a linear subspace  $L \cong \mathbb{C}\mathbb{P}^{N-n-1}$  of  $H \cong \mathbb{C}\mathbb{P}^{N-1}$  such that  $L \cap A_\infty = \emptyset$ . If this holds then  $\overline{A}$  and  $A_\infty$  are algebraic subvarieties of pure dimension  $n$  and  $n-1$ , respectively. If  $X$  is not algebraic then the image of any proper holomorphic immersion  $f : X \rightarrow \mathbb{C}^N$  is not algebraic either, so its limit set  $f(X)_\infty \subset \mathbb{C}\mathbb{P}^{N-1}$  has a nonempty intersection with every linear subspace  $\mathbb{C}\mathbb{P}^{N-n-1} \cong L \subset \mathbb{C}\mathbb{P}^{N-1}$ .

We construct proper holomorphic embeddings with images in small Hartogs domains.

**Theorem 1.1.** *Let  $X$  be a Stein manifold of dimension  $n \geq 1$ . Given a continuous increasing function  $h : [0, \infty) \rightarrow (0, \infty)$  with  $\lim_{t \rightarrow \infty} h(t) = \infty$  there exist a*

---

Received by the editors October 30, 2023, and, in revised form, January 18, 2024.

2020 *Mathematics Subject Classification.* Primary 32H02; Secondary 32E10, 32Q56.

*Key words and phrases.* Stein manifold, proper holomorphic embedding.

The author was supported by the European Union (ERC Advanced grant HPDR, 101053085) and grants P1-0291, J1-3005, N1-0237 from ARIS, Republic of Slovenia.

proper holomorphic embedding  $(z, w) : X \hookrightarrow \mathbb{C}^{n+1} \times \mathbb{C}^n$  and a proper holomorphic immersion  $(z, w) : X \rightarrow \mathbb{C}^{n+1} \times \mathbb{C}^{n-1}$  satisfying

$$(1.1) \quad |w(x)| < h(|z(x)|) \quad \text{for all } x \in X.$$

Furthermore, given a compact  $\mathcal{O}(X)$ -convex set  $K$  in  $X$ , an open neighbourhood  $U \subset X$  of  $K$ , and a holomorphic map  $f_0 = (z_0, w_0) : U \rightarrow \mathbb{C}^{n+1} \times \mathbb{C}^p$  satisfying (1.1) for all  $x \in K$ , we can approximate  $f_0$  uniformly on  $K$  by a proper holomorphic embedding  $f = (z, w) : X \rightarrow \mathbb{C}^{n+1} \times \mathbb{C}^p$  if  $p \geq n$ , resp. immersion if  $p = n - 1$ , satisfying (1.1).

The function  $h$  in Theorem 1.1 can be chosen to grow arbitrarily slowly, and hence the image  $f(X)$  may be arbitrarily close to the subspace  $\mathbb{C}^{n+1} \times \{0\}^p$  in the Fubini–Study metric on  $\mathbb{C}\mathbb{P}^{n+1+p}$ . Choosing  $h$  such that  $\lim_{t \rightarrow \infty} h(t)/t = 0$  gives Corollary 1.2.

**Corollary 1.2.** *Every Stein manifold  $X$  of dimension  $n \geq 1$  admits a proper holomorphic embedding  $f : X \hookrightarrow \mathbb{C}^{2n+1}$  whose limit set  $f(X)_\infty = \overline{f(X)} \cap H$  is a linearly embedded copy of  $\mathbb{C}\mathbb{P}^n$  in  $H = \mathbb{C}\mathbb{P}^{2n+1} \setminus \mathbb{C}^{2n+1} \cong \mathbb{C}\mathbb{P}^{2n}$ . In particular, every open Riemann surface  $X$  admits a proper holomorphic embedding in  $\mathbb{C}^3$  whose limit set is a projective line  $\mathbb{C}\mathbb{P}^1 \subset \mathbb{C}\mathbb{P}^2$ . The analogous result holds for proper holomorphic immersions  $X \rightarrow \mathbb{C}^{2n}$ .*

By the preceding discussion, the limit set  $f(X)_\infty$  intersects every projective subspace  $L \subset \mathbb{C}\mathbb{P}^{N-1}$  of dimension  $N - n - 1$ , unless  $f(X)$  is algebraic. Therefore, the nonalgebraic embeddings given by Corollary 1.2 have the smallest possible limit sets.

Given a nonalgebraic complex subvariety  $X$  of  $\mathbb{C}^N$ , its closure  $\overline{X} \subset \mathbb{C}\mathbb{P}^N$  and the limit set  $X_\infty \subset \mathbb{C}\mathbb{P}^{N-1}$  need not be analytic subvarieties, and for any pair of integers  $1 \leq n < N$  there are  $n$ -dimensional closed complex submanifolds  $X \subset \mathbb{C}^N$  with  $X_\infty = \mathbb{C}\mathbb{P}^{N-1}$ . (This always holds if  $N = n + 1$  and  $X$  is nonalgebraic.) Indeed, if  $X$  is a closed complex subvariety of  $\mathbb{C}^N$  ( $N > 1$ ) then for any closed discrete set  $B = \{b_j\}_{j \in \mathbb{N}} \subset \mathbb{C}^N$  there exist a domain  $\Omega \subset \mathbb{C}^N$  containing  $X$  and a biholomorphic map  $\Phi : \Omega \rightarrow \mathbb{C}^N$  such that  $B \subset \Phi(X)$  (see [13, Theorem 6.1] or [14, Theorem 4.17.1 (i)]). Note that  $X' = \Phi(X)$  is a closed complex subvariety of  $\mathbb{C}^N$ . Choosing  $B$  such that its closure in  $\mathbb{C}\mathbb{P}^N$  contains the hyperplane at infinity implies  $X'_\infty = \mathbb{C}\mathbb{P}^{N-1}$ . A characterization of the closed subsets of  $\mathbb{C}\mathbb{P}^{N-1}$  which are limit sets of closed complex subvarieties of  $\mathbb{C}^N$  of a given dimension does not seem to be known.

The corollary is especially interesting in dimension  $n = 1$ . A long-standing open question (the Forster conjecture [11], also called the Bell–Narasimhan conjecture [2, 3]) asks whether every open Riemann surface,  $X$ , admits a proper holomorphic embedding in  $\mathbb{C}^2$ . Recent surveys of this subject can be found in [14, Secs. 9.10–9.11] and the preprint [1] by Alarcón and López, where the authors constructed a proper harmonic embedding of any open Riemann surface in  $\mathbb{C} \times \mathbb{R}^2 \cong \mathbb{C}^2$  with a holomorphic first coordinate function. Note that if  $X \rightarrow \mathbb{C}^2$  is a proper holomorphic map with nonalgebraic image then  $f(X)_\infty = \mathbb{C}\mathbb{P}^1$ . (There are algebraic open Riemann surfaces which do not embed as smooth proper affine curves in  $\mathbb{C}^2$ .) Corollary 1.2 gives proper holomorphic embeddings  $f : X \hookrightarrow \mathbb{C}^3$  whose images are arbitrarily close to the subspace  $\mathbb{C}^2 \times \{0\}$  in the Fubini–Study metric on  $\mathbb{C}\mathbb{P}^3$ , and  $f(X)_\infty = \mathbb{C}\mathbb{P}^1$ .

It was recently shown by Drinovec Drnovšek and Forstnerič [8, Theorem 1.3] that, under a mild condition on an unbounded closed convex set  $E \subset \mathbb{C}^N$ , proper holomorphic embeddings  $f : X \hookrightarrow \mathbb{C}^N$  from any Stein manifold  $X$  with  $2 \dim X < N$  such that  $f(X) \subset \Omega = \mathbb{C}^N \setminus E$  are dense in the space  $\mathcal{O}(X, \Omega)$  of all holomorphic maps  $X \rightarrow \Omega$ . A similar result holds for immersions if  $2 \dim X \leq N$ . Their proof relies on the fact, proved by Forstnerič and Wold [17], that such a domain  $\Omega$  is an Oka domain. (See [14, Definition 5.4.1 and Theorem 5.4.4] for the definition and the main results concerning Oka manifolds.) Note that the domains in Theorem 1.1 are much smaller than those in [8, Theorem 1.3] when the codimension is at least 2. On the other hand, Theorem 1.1 does not pertain to proper maps in codimension 1 (the case  $p = 0$ ). We do not know whether a Hartogs domain of the form

$$(1.2) \quad \Omega = \{(z, w) \in \mathbb{C}^{n+1} \times \mathbb{C}^p : |w| < h(|z|)\}, \quad n \geq 1, p \geq 1,$$

which appears in Theorem 1.1, is an Oka domain, except if  $p = 1$ , the function  $h > 0$  grows at least linearly at infinity, and  $\log h(|z|)$  is plurisubharmonic on  $\mathbb{C}^{n+1}$  (see Forstnerič and Kusakabe [15, Proposition 3.1]). Our proof does not require that  $\Omega$  be an Oka domain.

We mention that a Stein manifold of dimension  $n \geq 2$  admits a proper holomorphic embedding  $X \hookrightarrow \mathbb{C}^N$  with  $N = \lfloor \frac{3n}{2} \rfloor + 1$  and a proper holomorphic immersion with  $N = \lfloor \frac{3n+1}{2} \rfloor$ ; see Eliashberg and Gromov [10], Schürmann [24], and [14, Theorem 9.3.1]. The proofs are very delicate and depend on Oka theory. We do not know whether one can expect a similar control of the range of the embedding in these dimensions.

## 2. PROOF OF THEOREM 1.1

Our proof of Theorem 1.1 relies on the following technical result, which is a special case of [9, Theorem 1.1] by Drinovec Drnovšek and Forstnerič. (See also [16, Theorem 6], which is based on the same result.) Similar results were obtained earlier by Dor [6, 7].

**Theorem 2.1.** *Assume that  $X$  is a Stein manifold of dimension  $n \geq 1$ ,  $D$  is a relatively compact, smoothly bounded, strongly pseudoconvex domain in  $X$ ,  $K$  is a compact set contained in  $D$ ,  $t_0$  is a real number,  $\sigma : \mathbb{C}^{n+1} \rightarrow \mathbb{R}$  is a strongly plurisubharmonic exhaustion function which has no critical points in the set  $\{\sigma \geq t_0\}$ , and  $g_0 : \overline{D} \rightarrow \mathbb{C}^{n+1}$  is a continuous map that is holomorphic in  $D$  and satisfies  $g_0(\overline{D} \setminus K) \subset \{\sigma > t_0\}$ . Given numbers  $t_1 > t_0$  and  $\epsilon > 0$ , there is a holomorphic map  $g : \overline{D} \rightarrow \mathbb{C}^{n+1}$  satisfying the following conditions:*

- (a)  $g(bD) \subset \{\sigma > t_1\}$ .
- (b)  $\sigma(g(x)) > \sigma(g_0(x)) - \epsilon$  for all  $x \in \overline{D}$ .
- (c)  $|g(x) - g_0(x)| < \epsilon$  for all  $x \in K$ .

Note that if  $\epsilon > 0$  is small enough then condition (b) implies

$$g(\overline{D} \setminus K) \subset \{\sigma > t_0\}.$$

The analogous result holds much more generally, and we only stated the case that will be used here. For condition (b), see [9, Lemma 5.3], which is the main inductive step in [9, proof of Theorem 1.1]. We remark that a map from a compact set in a complex manifold is said to be holomorphic if it is holomorphic in an open neighbourhood of the said set.

*Proof of Theorem 1.1.* We shall construct proper holomorphic embeddings  $X \hookrightarrow \mathbb{C}^N$  with  $N \geq 2n + 1$  satisfying (1.1); the same arguments will yield immersions when  $N = 2n$ .

Let  $\Omega \subset \mathbb{C}^N$  be a domain of the form (1.2) with coordinates  $(z, w) \in \mathbb{C}^{n+1} \times \mathbb{C}^p$  where  $p \geq n$ ,  $N = n + 1 + p$ , and the function  $h : [0, \infty) \rightarrow (0, \infty)$  is as in the theorem. We shall use Theorem 2.1 with the exhaustion function  $\sigma(z) = |z|$  on  $\mathbb{C}^{n+1}$ ; the nonsmooth point at the origin will not matter. We denote by  $\mathbb{B}$  the open unit ball in  $\mathbb{C}^{n+1}$ .

Since the set  $K \subset X$  is compact and  $\mathcal{O}(X)$ -convex, there exist a smooth strongly plurisubharmonic Morse exhaustion function  $\rho : X \rightarrow \mathbb{R}_+$  and a sequence  $0 < c_0 < c_1 < \dots$  with  $\lim_{i \rightarrow \infty} c_i = +\infty$  such that every  $c_i$  is a regular value of  $\rho$  and, setting

$$D_i = \{x \in X : \rho(x) < c_i\} \quad \text{for } i = 0, 1, 2, \dots,$$

we have that  $K \subset D_0 \subset \bar{D}_0 \subset U$ , where  $U \subset X$  is a neighbourhood of  $K$  as in the theorem (see [19, Theorem 5.1.6, p. 117]). We may assume that the given holomorphic map  $f_0 = (z_0, w_0) : U \rightarrow \Omega$  satisfies condition (1.1) for all  $x \in \bar{D}_0$  and  $z_0(x) \neq 0$  for all  $x \in bD_0$ . (We shall use the subscript in  $z_i$  and  $w_i$  as an index in the induction process; a notation for the components of these maps will not be needed.) Pick a number  $t_0 \in \mathbb{R}$  with

$$0 < t_0 < \min_{x \in bD_0} |z_0(x)|.$$

Choose a number  $t_{-1}$  with  $0 < t_{-1} < t_0$  and close enough to  $t_0$  such that the sublevel set  $D_{-1} = \{\rho < t_{-1}\}$  satisfies  $K \subset D_{-1} \subset \bar{D}_{-1} \subset D_0$  and

$$z_0(\overline{D_0 \setminus D_{-1}}) \subset \mathbb{C}^{n+1} \setminus t_0 \bar{\mathbb{B}}.$$

Note that the set  $\bar{D}_i$  is  $\mathcal{O}(X)$ -convex for every  $i = -1, 0, 1, \dots$

By the Oka–Weil theorem, we can approximate the map  $w_0 : U \rightarrow \mathbb{C}^p$  uniformly on  $\bar{D}_0$  by a holomorphic map  $w_1 : X \rightarrow \mathbb{C}^p$  such that  $(z_0, w_1)(\bar{D}_0) \subset \Omega$ . We shall now construct a holomorphic map  $z_1 : \bar{D}_1 \rightarrow \mathbb{C}^{n+1}$  such that the holomorphic map  $f_1 = (z_1, w_1) : \bar{D}_1 \rightarrow \Omega$  enjoys suitable properties to be explained. This will be the first step of an induction procedure.

Pick a number  $t_1 \geq t_0 + 1$  so big that

$$(2.1) \quad h(t_1) > \max\{|w_1(z)| : z \in \bar{D}_1\}.$$

(Such a number exists since  $\lim_{t \rightarrow \infty} h(t) = +\infty$ .) Fix  $\epsilon > 0$  whose precise value will be determined later. Let  $\tilde{z}_0 : \bar{D}_0 \rightarrow \mathbb{C}^{n+1}$  be a holomorphic map given by Theorem 2.1 (with  $\tilde{z}_0 = g$  in the notation of that theorem, applied to the map  $g_0 = z_0$ , the compact set  $\bar{D}_{-1} \subset D_0$ , and the numbers  $\epsilon$  and  $t_0 < t_1$ ). Condition (b) in Theorem 2.1 gives

$$|\tilde{z}_0(x)| > |z_0(x)| - \epsilon \quad \text{for all } x \in \bar{D}_0.$$

Since the function  $h$  in (1.2) is continuous, it follows that if  $\epsilon > 0$  is small enough then the map  $(\tilde{z}_0, w_1) : \bar{D}_0 \rightarrow \mathbb{C}^N$  has range in  $\Omega$ , and we have that

$$(2.2) \quad \tilde{z}_0(bD_0) \subset \mathbb{C}^{n+1} \setminus t_1 \bar{\mathbb{B}}, \quad \tilde{z}_0(\overline{D_0 \setminus D_{-1}}) \subset \mathbb{C}^{n+1} \setminus t_0 \bar{\mathbb{B}}, \quad |\tilde{z}_0 - z_0| < \epsilon \quad \text{on } \bar{D}_{-1}.$$

We now use the fact that  $\mathbb{C}^{n+1} \setminus t_1 \bar{\mathbb{B}}$  is an Oka domain (see Kusakabe [20, Corollary 1.3]). Hence, the main result of Oka theory gives a holomorphic map  $z_1 : \bar{D}_1 \rightarrow \mathbb{C}^{n+1}$  satisfying

$$(2.3) \quad z_1(\overline{D_1 \setminus D_0}) \subset \mathbb{C}^{n+1} \setminus t_1 \bar{\mathbb{B}} \quad \text{and} \quad |z_1 - \tilde{z}_0| < \epsilon \quad \text{on } \bar{D}_0.$$

(See [12, Theorem 1.3] for a precise statement of a more general result. In the special case at hand, the existence of a map  $z_1$  satisfying (2.3) was proved by a more involved argument in the paper [16] by Forstnerič and Ritter, predating Kusakabe's work [20].) If the number  $\epsilon > 0$  is chosen small enough, it follows from (2.1)–(2.3) and the definition of  $\Omega$  (1.2) that

$$(2.4) \quad z_1(\overline{D_0 \setminus D_{-1}}) \subset \mathbb{C}^{n+1} \setminus t_0\mathbb{B} \quad \text{and} \quad (z_1, w_1)(\overline{D_1}) \subset \Omega.$$

Since the dimension of the target space  $\mathbb{C}^N$  is at least  $2 \dim X + 1$ , we may assume after a small perturbation that the map  $f_1 = (z_1, w_1) : \overline{D_1} \hookrightarrow \Omega$  is an embedding satisfying the above conditions (see [14, Corollary 8.9.3]). Assuming as we may that all approximations are close enough, we also have that  $|f_1 - f_0| < \epsilon_0$  on  $\overline{D_{-1}}$  for a given  $\epsilon_0 > 0$ .

Continuing inductively, we obtain an increasing sequence  $t_0 < t_1 < t_2 < \dots$  with  $\lim_{i \rightarrow \infty} t_i = \infty$ , a decreasing sequence  $\epsilon_0 > \epsilon_1 > \epsilon_2 > \dots > 0$  with  $\lim_{i \rightarrow \infty} \epsilon_i = 0$ , and a sequence of holomorphic embeddings  $f_i = (z_i, w_i) : \overline{D_i} \hookrightarrow \mathbb{C}^{2n+1}$  satisfying the following conditions for  $i = 1, 2, \dots$

- (i)  $f_i(\overline{D_i}) \subset \Omega$ .
- (ii)  $z_i(\overline{D_i \setminus D_{i-1}}) \subset \mathbb{C}^{n+1} \setminus t_i\overline{\mathbb{B}}$ .
- (iii)  $z_i(\overline{D_{i-1} \setminus D_{i-2}}) \subset \mathbb{C}^{n+1} \setminus t_{i-1}\overline{\mathbb{B}}$ .
- (iv)  $|f_i - f_{i-1}| < \epsilon_{i-1}$  on  $\overline{D_{i-2}}$ .
- (v)  $t_i \geq t_{i-1} + 1$ .
- (vi)  $0 < \epsilon_i < \epsilon_{i-1}/2$ .
- (vii) Every holomorphic map  $f : \overline{D_i} \rightarrow \mathbb{C}^N$  with  $|f - f_i| < 2\epsilon_i$  on  $\overline{D_{i-1}}$  is an embedding on  $\overline{D_{i-2}}$  and satisfies  $f(\overline{D_{i-1}}) \subset \Omega$ .

Note that conditions (i) and (ii) also holds for  $i = 0$  by the assumptions on  $f_0$ , and conditions (i)–(v) hold for  $i = 1$  by the construction of the map  $f_1$ .

The inductive step is similar to the one from  $i = 0$  to  $i = 1$ , which was explained above. Assume inductively that conditions (i)–(v) hold for some  $i \in \{1, 2, \dots\}$ . Pick a number  $\epsilon_i$  satisfying conditions (vi) and (vii). Also, fix a number  $\epsilon > 0$  whose precise value will be determined during this induction step. By the Oka–Weil theorem, there is a holomorphic map  $w_{i+1} : X \rightarrow \mathbb{C}^p$  with  $|w_{i+1} - w_i| < \epsilon$  on  $\overline{D_i}$ . Choose a number  $t_{i+1} \geq t_i + 1$  so big that

$$(2.5) \quad h(t_{i+1}) > \max\{|w_{i+1}(x)| : x \in \overline{D_{i+1}}\}.$$

If  $\epsilon > 0$  is chosen small enough then Theorem 2.1, applied to the map  $g_0 = z_i : \overline{D_i} \rightarrow \mathbb{C}^{n+1}$ , the compact set  $\overline{D_{i-1}} \subset D_i$ , and the numbers  $t_i < t_{i+1}$  furnishes a holomorphic map  $\tilde{z}_i : \overline{D_i} \rightarrow \mathbb{C}^{n+1}$  such that the map  $(\tilde{z}_i, w_{i+1}) : \overline{D_0} \rightarrow \mathbb{C}^N$  has range in  $\Omega$  and the following conditions hold:

$$\tilde{z}_i(bD_i) \subset \mathbb{C}^{n+1} \setminus t_{i+1}\overline{\mathbb{B}}, \quad \tilde{z}_i(\overline{D_i \setminus D_{i-1}}) \subset \mathbb{C}^{n+1} \setminus t_i\overline{\mathbb{B}}, \quad |\tilde{z}_i - z_i| < \epsilon \quad \text{on } \overline{D_{i-1}}.$$

(For  $i = 0$  these are conditions (2.2).) Since  $\mathbb{C}^{n+1} \setminus t_{i+1}\overline{\mathbb{B}}$  is an Oka domain (see [20, Corollary 1.3]), there is a holomorphic map  $z_{i+1} : \overline{D_{i+1}} \rightarrow \mathbb{C}^{n+1}$  satisfying

$$z_{i+1}(\overline{D_{i+1} \setminus D_i}) \subset \mathbb{C}^{n+1} \setminus t_{i+1}\overline{\mathbb{B}} \quad \text{and} \quad |z_{i+1} - \tilde{z}_i| < \epsilon \quad \text{on } \overline{D_i}.$$

(This is an analogue of condition (2.3).) Finally, we perturb the holomorphic map

$$f_{i+1} = (z_{i+1}, w_{i+1}) : \overline{D_{i+1}} \rightarrow \mathbb{C}^N$$

slightly to make it an embedding. If all approximations are close enough then  $f_{i+1}$  satisfies conditions (i)–(iv), and (v) holds by the choice of  $t_{i+1}$ . This completes the induction step.

Conditions (iv) and (vi) imply that the sequence  $f_i$  converges to the limit map

$$f = (z, w) = \lim_{i \rightarrow \infty} f_i : X \rightarrow \mathbb{C}^N$$

satisfying  $|f - f_i| < 2\epsilon_i$  on  $\bar{D}_{i-1}$  for every  $i = 0, 1, \dots$  (In particular,  $|f - f_0| < 2\epsilon_0$  on  $K$ .) Conditions (i), (vi), and (vii) then imply that  $f$  is a holomorphic embedding with  $f(X) \subset \Omega$ . Finally, conditions (ii)–(vi) imply that the map  $z : X \rightarrow \mathbb{C}^{n+1}$  is proper, and hence  $f$  is proper as map to  $\mathbb{C}^N$ .  $\square$

#### REFERENCES

- [1] A. Alarcon and F. J. Lopez, *Proper harmonic embeddings of open Riemann surfaces into  $\mathbb{R}^4$* , <https://arxiv.org/abs/2206.03566>, 2022.
- [2] S. R. Bell, J.-L. Brylinski, A. T. Huckleberry, R. Narasimhan, C. Okonek, G. Schumacher, A. Van de Ven, and S. Zucker, *Complex manifolds*, Springer-Verlag, Berlin, 1998. Corrected reprint of the 1990 translation [*Several complex variables. VI*, Encyclopaedia, Math. Sci., 69, Springer, Berlin, 1990; MR1095088 (91i:32001)], DOI 10.1007/978-3-642-61299-2. MR1608852
- [3] Steven R. Bell and Raghavan Narasimhan, *Proper holomorphic mappings of complex spaces*, *Several complex variables, VI*, Encyclopaedia Math. Sci., vol. 69, Springer, Berlin, 1990, pp. 1–38. MR1095089
- [4] Errett Bishop, *Mappings of partially analytic spaces*, *Amer. J. Math.* **83** (1961), 209–242, DOI 10.2307/2372953. MR123732
- [5] E. M. Chirka, *Complex analytic sets*, *Mathematics and its Applications (Soviet Series)*, vol. 46, Kluwer Academic Publishers Group, Dordrecht, 1989. Translated from the Russian by R. A. M. Hoksbergen, DOI 10.1007/978-94-009-2366-9. MR1111477
- [6] Avner Dor, *Approximation by proper holomorphic maps into convex domains*, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* (4) **20** (1993), no. 1, 147–162. MR1216002
- [7] Avner Dor, *Immersion and embeddings in domains of holomorphy*, *Trans. Amer. Math. Soc.* **347** (1995), no. 8, 2813–2849, DOI 10.2307/2154757. MR1282885
- [8] Barbara Drinovec Drnovšek and Franc Forstnerič, *Proper holomorphic maps in Euclidean spaces avoiding unbounded convex sets*, *J. Geom. Anal.* **33** (2023), no. 6, Paper No. 170, 22, DOI 10.1007/s12220-023-01222-z. MR4567571
- [9] Barbara Drinovec Drnovšek and Franc Forstnerič, *Strongly pseudoconvex domains as subvarieties of complex manifolds*, *Amer. J. Math.* **132** (2010), no. 2, 331–360, DOI 10.1353/ajm.0.0106. MR2654777
- [10] Yakov Eliashberg and Mikhael Gromov, *Embeddings of Stein manifolds of dimension  $n$  into the affine space of dimension  $3n/2 + 1$* , *Ann. of Math.* (2) **136** (1992), no. 1, 123–135, DOI 10.2307/2946547. MR1173927
- [11] Otto Forster, *Plongements des variétés de Stein (French)*, *Comment. Math. Helv.* **45** (1970), 170–184, DOI 10.1007/BF02567324. MR269880
- [12] Franc Forstnerič, *Recent developments on Oka manifolds*, *Indag. Math. (N.S.)* **34** (2023), no. 2, 367–417, DOI 10.1016/j.indag.2023.01.005. MR4547869
- [13] Franc Forstnerič, *Interpolation by holomorphic automorphisms and embeddings in  $\mathbb{C}^n$* , *J. Geom. Anal.* **9** (1999), no. 1, 93–117, DOI 10.1007/BF02923090. MR1760722
- [14] Franc Forstnerič, *Stein manifolds and holomorphic mappings*, 2nd ed., *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, vol. 56, Springer, Cham, 2017. The homotopy principle in complex analysis, DOI 10.1007/978-3-319-61058-0. MR3700709
- [15] F. Forstnerič and Y. Kusakabe, *Oka tubes in holomorphic line bundles*, [arXiv:2310.14871](https://arxiv.org/abs/2310.14871), 2023.
- [16] Franc Forstnerič and Tyson Ritter, *Oka properties of ball complements*, *Math. Z.* **277** (2014), no. 1-2, 325–338, DOI 10.1007/s00209-013-1258-2. MR3205776

- [17] Franc Forstnerič and Erlend Fornæss Wold, *Oka domains in Euclidean spaces*, Int. Math. Res. Not. IMRN **3** (2024), 1801–1824, DOI 10.1093/imrn/rnac347. MR4702264
- [18] Robert C. Gunning and Hugo Rossi, *Analytic functions of several complex variables*, AMS Chelsea Publishing, Providence, RI, 2009. Reprint of the 1965 original, DOI 10.1090/chel/368. MR2568219
- [19] Lars Hörmander, *An introduction to complex analysis in several variables*, 3rd ed., North-Holland Mathematical Library, vol. 7, North-Holland Publishing Co., Amsterdam, 1990. MR1045639
- [20] Yuta Kusakabe, *Oka properties of complements of holomorphically convex sets*, Ann. Math. (2), **199** (2024), no. 2, 899–917, DOI 10.4007/annals.2024.199.2.7.
- [21] Raghavan Narasimhan, *Imbedding of holomorphically complete complex spaces*, Amer. J. Math. **82** (1960), 917–934, DOI 10.2307/2372949. MR148942
- [22] Reinhold Remmert, *Projektionen analytischer Mengen* (German), Math. Ann. **130** (1956), 410–441, DOI 10.1007/BF01343236. MR86353
- [23] Reinhold Remmert, *Sur les espaces analytiques holomorphiquement séparables et holomorphiquement convexes* (French), C. R. Acad. Sci. Paris **243** (1956), 118–121. MR79808
- [24] J. Schürmann, *Embeddings of Stein spaces into affine spaces of minimal dimension*, Math. Ann. **307** (1997), no. 3, 381–399, DOI 10.1007/s002080050040. MR1437045

FACULTY OF MATHEMATICS AND PHYSICS, UNIVERSITY OF LJUBLJANA, JADRANSKA 19, SI–1000 LJUBLJANA, SLOVENIA; and INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS, JADRANSKA 19, SI–1000 LJUBLJANA, SLOVENIA

*Email address:* `franc.forstneric@fmf.uni-lj.si`