

COMMENTS ON CANONICAL LINES

R. B. RASMUSEN AND B. L. HAGEN

1. **Introduction.** In this paper we propose to find the equations of the canonical edges of Green using a conjugate net as the parametric net of an analytic surface; to give a new interpretation to the line called by Davis the associate line of collineation; and, finally, to make a generalization of the canonical quadric of Davis.

2. **Analytic basis.** Let the projective homogeneous coordinates x^1, \dots, x^4 of a point P_x on a surface S in ordinary space be analytic functions of two independent variables u, v . In the notation¹ of Lane if the parametric curves on S form a conjugate net, the coordinates x of the point P_x and the coordinates y of a point which is the harmonic conjugate of the point P_x with respect to the foci of the axis of the point P_x , satisfy a system of equations of the form

$$(1) \quad \begin{aligned} x_{uu} &= px + \alpha x_u + Ly, \\ x_{uv} &= cx + ax_u + bx_v, \\ x_{vv} &= qx + \delta x_v + Ny, \end{aligned} \quad LN \neq 0.$$

The ray-points of the net at the point P_x are given by the formulas

$$x_{-1} = x_u - bx, \quad x_1 = x_v - ax.$$

Some of the invariants of the net are

$$(2) \quad \begin{aligned} H &= c + ab - a_u, & K &= c + ab - b_v, \\ \mathfrak{S} &= c + ab + b_v - \delta_u, & \mathfrak{R} &= c + ab + a_u - \alpha_v, \\ 8\mathfrak{B}' &= 4a - 2\delta + (\log r)_v, & r &= N/L, \\ 8\mathfrak{C}' &= 4b - 2\alpha - (\log r)_u. \end{aligned}$$

If the covariant tetrahedron, x, x_1, x_{-1}, y is used as the local tetrahedron of reference, a power series expansion² for one nonhomogeneous coordinate z of a point on the surface in terms of the other two coordinates x, y is

$$(3) \quad \begin{aligned} z &= \frac{1}{2}(Lx^2 + Ny^2) + \frac{4}{3}(L\mathfrak{C}'x^3 + N\mathfrak{B}'y^3) + c_0x^4 \\ &\quad + 4c_1x^3y + 4c_3xy^3 + c_4y^4 + \dots, \end{aligned}$$

¹ Lane, *Conjugate nets and the lines of curvature*, American Journal of Mathematics, vol. 53 (1931), p. 574.

² Lane, *A canonical power series expansion for a surface*, Transactions of this Society, vol. 37 (1935), p. 481.

where the coefficients c_0, c_1, c_3, c_4 are given by

$$c_0 = \frac{1}{3}L\mathfrak{C}'[12\mathfrak{C}' + (\log \mathfrak{C}'r^{1/2})_u], \quad 4c_1 = \frac{1}{6}L(H - \mathfrak{S}),$$

$$c_4 = \frac{1}{3}N\mathfrak{B}'[12\mathfrak{B}' + (\log \mathfrak{B}'r^{-1/2})_v], \quad 4c_3 = \frac{1}{6}N(K - \mathfrak{R}),$$

and the coefficient of x^2y^2 is zero.

We will assume that the surface S is non-ruled,³ that is, that $\mathfrak{B}'^2 + r\mathfrak{C}'^2 \neq 0$.

3. The second canonical edge of Green. Segre has shown⁴ that the second⁵ edge of Green is the line of inflexion of the nodal cubic in which the tangent plane at a point P_x of an analytic surface S intersects the four-parameter family of nondegenerate algebraic cubic surfaces having fourth order contact with S at P_x .

The equation in nonhomogeneous coordinates of a general algebraic cubic surface through the point P_x is

$$(4) \quad \begin{aligned} & a_1x + a_2y + a_3z + b_1x^2 + b_2y^2 + b_3z^2 + b_4xy + b_5xz \\ & + b_6yz + d_1x^3 + d_2y^3 + d_3z^3 + d_4x^2y + d_5x^2z + d_6xy^2 \\ & + d_7y^2z + d_8xz^2 + d_9yz^2 + d_{10}xyz = 0. \end{aligned}$$

If this surface has fourth order contact with the surface S at the point P_x , its coefficients satisfy the following conditions:

$$(5) \quad \begin{aligned} & a_1 = a_2 = b_4 = 0, \quad a_3L + 2b_1 = 0, \quad a_3N + 2b_2 = 0, \\ & 8L\mathfrak{C}'a_3 + 3Lb_5 + d_1 = 0, \quad Lb_6 + 2d_4 = 0, \\ & 8N\mathfrak{B}'a_3 + 3Nb_6 + d_2 = 0, \quad Nb_5 + 2d_6 = 0, \\ & 12c_0a_3 + 3L^2b_3 + 16L\mathfrak{C}'b_5 + 6Ld_5 = 0, \\ & 12c_4a_3 + 3N^2b_3 + 16N\mathfrak{B}'b_6 + 6Nd_7 = 0, \\ & 12c_1a_3 + 8L\mathfrak{C}'b_6 + 3Ld_{10} = 0, \\ & 12c_3a_3 + 8N\mathfrak{B}'b_5 + 3Nd_{10} = 0, \\ & LNb_3 + Nd_5 + Ld_7 = 0. \end{aligned}$$

The tangent plane, $z=0$, intersects this cubic surface in the cubic curve

³ Green, *Projective differential geometry of one-parameter families of curves and conjugate nets on a curved surface*, First Memoir, American Journal of Mathematics, vol. 37 (1915), p. 239.

⁴ B. Segre, *La cubique indicatrice de l'élément linéaire projectif d'une surface*, Comptes Rendus de l'Académie des Sciences, Paris, vol. 184, p. 729.

⁵ This classification of the edges of Green is the reverse of that introduced by Green.

$$(6) \quad z = b_1x^2 + b_2y^2 + d_1x^3 + d_2y^3 + d_4x^2y + d_6xy^2 = 0,$$

where the coefficients $b_1, b_2, d_1, d_2, d_4, d_6$ can be expressed in terms of the invariants given by equations (2) and a_3 .

Following the usual method of solving a cubic with its Hessian and discarding the solution giving the double point tangents, we obtain the following equation of the line of inflexions of the cubic:

$$(7) \quad z = a_3 + (b_5 + 2\mathfrak{C}'a_3)x + (b_6 + 2\mathfrak{B}'a_3)y = 0,$$

where

$$(8) \quad \begin{aligned} b_5 &= \frac{-a_3}{8(N\mathfrak{C}'^2 + L\mathfrak{B}'^2)} \left\{ -L\mathfrak{B}'[(H - \mathfrak{S}) - (K - \mathfrak{R})] \right. \\ &\quad \left. + 2\mathfrak{C}'[12(N\mathfrak{C}'^2 + L\mathfrak{B}'^2) + N\mathfrak{C}'(\log \mathfrak{C}'r^{1/2})_u \right. \\ &\quad \left. + L\mathfrak{B}'(\log \mathfrak{B}'r^{-1/2})_v] \right\}, \\ b_6 &= \frac{-a_3}{8(N\mathfrak{C}'^2 + L\mathfrak{B}'^2)} \left\{ N\mathfrak{C}'[(H - \mathfrak{S}) - (K - \mathfrak{R})] \right. \\ &\quad \left. + 2\mathfrak{B}'[12(N\mathfrak{C}'^2 + L\mathfrak{B}'^2) + N\mathfrak{C}'(\log \mathfrak{C}'r^{1/2})_u \right. \\ &\quad \left. + L\mathfrak{B}'(\log \mathfrak{B}'r^{-1/2})_v] \right\}. \end{aligned}$$

Any line of the second canonical pencil⁶ of Davis may have its equation in the form

$$z = 1 + k(\mathfrak{C}'x + \mathfrak{B}'y) = 0,$$

where k is a constant. The flex-ray (7) is a line of this pencil in case

$$2\mathfrak{C}' + b_5/a_3 = k\mathfrak{C}', \quad 2\mathfrak{B}' + b_6/a_3 = k\mathfrak{B}'.$$

Eliminating k from these equations, we obtain

$$(\mathfrak{B}'^2 + r\mathfrak{C}'^2)[(H - \mathfrak{S}) - (K - \mathfrak{R})] = 0.$$

Therefore, *necessary and sufficient conditions for the second edge of Green to be a line of the second canonical pencil of Davis are*

$$(9) \quad \begin{aligned} H - \mathfrak{S} &= K - \mathfrak{R}, \\ [r\mathfrak{C}'(\log \mathfrak{C}'r^{1/2})_u + \mathfrak{B}'(\log \mathfrak{B}'r^{-1/2})_v]/(\mathfrak{B}'^2 + r\mathfrak{C}'^2) &= \text{const.} \end{aligned}$$

A line of the first canonical pencil⁷ of Davis joins the points $(1, 0, 0, 0)$ and $(0, -k\mathfrak{C}'/L, -k\mathfrak{B}'/N, 1)$. It can be verified that the

⁶ W. M. Davis, *Contributions to the Theory of Conjugate Nets*, Chicago Doctor's Thesis, 1933, p. 18.

⁷ Davis, *ibid.*, p. 20.

first edge of Green is the line joining the point $(1, 0, 0, 0)$ to the point $[0, -(b_5/a_3+4\mathfrak{C}')/L, -(b_5/a_3+4\mathfrak{B}')/N, 1]$. The first edge of Green is a line of the first canonical pencil of Davis if conditions (9) hold. This is a special case of the following theorem.⁸

If one line of an arbitrarily selected pair of lines which are reciprocal with respect to a quadric of Darboux is a line of the second canonical pencil of Davis, its reciprocal is a line of the first canonical pencil of Davis, and conversely.

4. The associate line of collineation. The cubic surface

$$z = \frac{1}{2}(Lx^2 + Ny^2) + \frac{4}{3}(L\mathfrak{C}'x^3 + N\mathfrak{B}'y^3)$$

obtained by truncating the power series (3) can be interpreted⁹ geometrically as follows: first, it has third order contact with the analytic surface S given by equations (3) at the point $(1, 0, 0, 0)$; second, it has a unode of second order at the point $(0, 0, 0, 1)$. This cubic surface intersects the tangent plane $z=0$ in the nodal cubic

$$z = 3(x^2 + ry^2) + 8(\mathfrak{C}'x^3 + r\mathfrak{B}'y^3) = 0$$

which has for its line of inflexion $z = 1+2(\mathfrak{C}'x+r\mathfrak{B}'y)=0$. This is the *associate line of collineation*,¹⁰ and is a line of the second canonical pencil of Davis.

5. A generalization of the canonical quadric of Davis. Davis has shown¹¹ that at each point P_x of a conjugate net on an analytic surface S there exists a unique quadric surface which has the following properties: the quadric has second order contact with the surface sustaining the net; the axis and ray of the net at P_x are reciprocal polars with respect to the quadric; and the quadric passes through the point P_y which is the harmonic conjugate of P_x with respect to the focal points of the axis. We shall generalize this definition by showing that at each point P_x of a conjugate net there exists a two-parameter family of quadric surfaces having the following properties: *the quadrics have second order contact with the surface sustaining the net and any line of the second canonical pencil of Davis has as its polar reciprocal with respect to any quadric of this family the line of the first canonical pencil defined by the same value of k .*

⁸ We are indebted to P. O. Bell for this theorem in its general form.

⁹ Lane, *Projective Differential Geometry of Curves and Surfaces*, The University of Chicago Press, 1932, p. 296.

¹⁰ Davis, *ibid.*, p. 25.

¹¹ Davis, *ibid.*, p. 11.

The equation¹² of all quadric surfaces having second order contact with the analytic surface S , given by equation (3), is

$$(10) \quad Lx^2 + Ny^2 + z(-2 + k_2x + k_3y + k_4z) = 0,$$

where k_2, k_3, k_4 are arbitrary constants when P_x is fixed, and functions of u, v when P_x varies over S .

If we demand that the polar reciprocals of any line of the first canonical pencil of Davis with respect to a quadric given by equation (10) be the line of the second canonical pencil of David defined by the same value of k , we obtain $k_2 = -4k\mathfrak{C}'$, $k_3 = -4k\mathfrak{B}'$. Substituting these values in (10), we have

$$(11) \quad Lx^2 + Ny^2 + z(-2 - 4k\mathfrak{C}'x - 4k\mathfrak{B}'y + k_4z) = 0.$$

If $k=0$, and the quadric passes through the point P_y we have the canonical quadric of Davis; also if $k=-1$, we have the quadrics of Darboux.

It is interesting to note that for $k=-1$ the line in the second canonical pencil of Davis is the line of inflexions of the ray-point cubic.¹³

The second canonical tangent¹⁴ of Davis is the line joining the canonical point $(0, \mathfrak{B}', -\mathfrak{C}', 0)$ to the point $P(1, 0, 0, 0)$. The coordinates of any point on this line are

$$(12) \quad (\lambda, \mu\mathfrak{B}', -\mu\mathfrak{C}', 0),$$

where λ and μ are arbitrary constants. The polar plane of this point with respect to the quadrics whose equation is given by (11) is

$$(13) \quad \mu(L\mathfrak{B}'x - N\mathfrak{C}'y) - \lambda z = 0,$$

which is independent of k .

For $\lambda=0$, the point given by (12) is the canonical point of Davis and the plane (13) is the canonical plane of Davis.

Consequently, we have the theorem: *the polar plane of a point on the second canonical tangent of Davis with respect to a quadric of the family defined by equation (11) is a member of a pencil whose axis is the first canonical tangent of Davis. This plane is dependent on the selection of the point but is independent of the selection of the quadric of the family (11).*

WOODROW WILSON JUNIOR COLLEGE AND
PENNSYLVANIA STATE COLLEGE

¹² Davis, *ibid.*, p. 11.

¹³ Lane, *Conjugate nets and the lines of curvature*, American Journal of Mathematics, vol. 53 (1931), p. 577.

¹⁴ Davis, *ibid.*, pp. 17-18.