DELIGNE’S CATEGORY $\underline{\text{Rep}}(GL_\delta)$ AND REPRESENTATIONS OF GENERAL LINEAR SUPERGROUPS

JONATHAN COMES AND BENJAMIN WILSON

ABSTRACT. We classify indecomposable summands of mixed tensor powers of the natural representation for the general linear supergroup up to isomorphism. We also give a formula for the characters of these summands in terms of composite supersymmetric Schur polynomials, and give a method for decomposing their tensor products. Along the way, we describe indecomposable objects in $\underline{\text{Rep}}(GL_\delta)$ and explain how to decompose their tensor products.

CONTENTS

1. Introduction	568
2. Category-theoretic preliminaries	571
3. The category $\underline{\text{Rep}}(GL_\delta)$	575
4. Indecomposable objects in $\underline{\text{Rep}}(GL_\delta)$	579
5. Connection to representations of the general linear group	587
6. The lifting map	590
7. Decomposing tensor products in $\underline{\text{Rep}}(GL_\delta)$	596
8. Representations of the general linear supergroup	598
Acknowledgements	608
References	608

1. INTRODUCTION

1.1. Classical Schur-Weyl duality concerns the commuting actions of the symmetric group Σ_r and the general linear group GL_d on the tensor power $V^{\otimes r}$ of the natural representation V of GL_d . It enables the labelling of the isomorphism classes of indecomposable summands of the tensor power by partitions $\lambda \vdash r$ with height $l(\lambda) \leq d$ (this is *Weyl’s Strip Theorem*), and the description of the characters of these summands in terms of Schur polynomials. Schur-Weyl duality for the general linear supergroup $GL(m|n)$, established by Sergeev [25] and by Berele and Regev [4], provides similar insights into structure of the tensor power $V^{\otimes r}$ of the natural representation V of $GL(m|n)$. In this case, the isomorphism classes of the indecomposable summands of $V^{\otimes r}$ are parametrized by those partitions $\lambda \vdash r$ that are $(m|n)$ -hook, that is, partitions whose Young diagram can be covered by an m -wide, n -high hook, and the characters of the summands are described by the so-called supersymmetric Schur polynomials.

Received by the editors September 5, 2011 and, in revised form, July 31, 2012.
 2010 *Mathematics Subject Classification*. Primary 17B10, 18D10.

©2012 American Mathematical Society
 Reverts to public domain 28 years from publication

An analogue of Schur-Weyl duality for the *mixed tensor powers*

$$T(r, s) = T_{V, V^*}(r, s) = V^{\otimes r} \otimes (V^*)^{\otimes s}$$

of GL_d was developed by [2, 26, 18, 27]. Here, the walled Brauer algebras $B_{r,s}(\delta)$, with $\delta = d$, replace the group algebra of the symmetric group as the generic centralizer of the GL_d -action. It is known, in particular, that the indecomposable summands are labelled up to isomorphism by certain bipartitions (i.e., pairs of partitions). Schur-Weyl duality for mixed tensor powers also holds for the general linear supergroup $GL(m|n)$, where $\delta = m - n$ is the super dimension of the natural representation V . However, in this case, many fundamental questions remain unresolved. In this paper, we classify for the first time the indecomposable summands of the mixed tensor powers for the general linear supergroups up to isomorphism, and derive a character formula for the indecomposable summands in terms of composite supersymmetric Schur polynomials. In addition, we describe a method for the decomposition of tensor products of these indecomposable summands.

1.2. We work over a field $\mathbb{K}$ of characteristic zero throughout, identify finite dimensional representations of $GL(m|n)$ with integral representations of the Lie superalgebra $\mathfrak{gl}(m|n)$, and write $\text{Rep}(GL_d)$ and $\text{Rep}(\mathfrak{gl}(m|n))$ for the categories of finite-dimensional representations of GL_d and $\mathfrak{gl}(m|n)$, respectively. Fundamental to our approach is the tensor category $\underline{\text{Rep}}(GL_\delta)$, defined by Deligne [14, 15], that permits the simultaneous study of the mixed tensor powers for the general linear groups and the general linear supergroups. This category, which we refer to as *Deligne's category*, is constructed as the additive and Karoubi envelope of a “skeleton” tensor category $\underline{\text{Rep}}_0(GL_\delta)$ (cf. §3.5). Up to isomorphism, the objects $w_{r,s}$ of $\underline{\text{Rep}}_0(GL_\delta)$ are parametrized by pairs (r, s) of non-negative integers, representing the potencies of a mixed tensor power, and the morphism spaces are spanned by walled Brauer diagrams of the appropriate sizes. The structure of Deligne's category depends upon the parameter $\delta \in \mathbb{K}$ so that, in particular, the endomorphism algebras are the walled Brauer algebras $B_{r,s}(\delta)$. The universal property of Deligne's category guarantees that for any rigid δ -dimensional object V in a tensor category $\mathcal{T}$ satisfying hypotheses familiar from classical Schur-Weyl duality (cf. §4.7), there exists a full tensor functor $F : \underline{\text{Rep}}(GL_\delta) \rightarrow \mathcal{T}$ such that for any $r, s \geq 0$, $F(w_{r,s}) = T(r, s)$ is the corresponding mixed tensor power. In particular, for any $d > 0$ and for any $m, n \geq 0$ there exist full tensor functors

$$F_d : \underline{\text{Rep}}(GL_d) \rightarrow \text{Rep}(GL_d), \quad F_{m|n} : \underline{\text{Rep}}(GL_{m-n}) \rightarrow \text{Rep}(\mathfrak{gl}(m|n))$$

defined by the natural representations of GL_d and of $GL(m|n)$, respectively. Recent results in the representation theory of the walled Brauer algebra also play crucial roles. For any bipartition λ , let $\lambda^\bullet, \lambda^\circ$ be the partitions defined by $\lambda = (\lambda^\bullet, \lambda^\circ)$, let $l(\lambda) = l(\lambda^\bullet) + l(\lambda^\circ)$, and write $\lambda \vdash (|\lambda^\bullet|, |\lambda^\circ|)$. It was shown in [12] that the walled Brauer algebras $B_{r,s}(\delta)$ are cellular, with standard modules parametrized by the set of bipartitions

$$\Lambda_{r,s} = \{ \lambda \mid \lambda \vdash (r - i, s - i), \ 0 \leq i \leq \min(r, s) \}.$$

A recursive formula for the decomposition numbers for the walled Brauer algebras was described in [11], using cap diagrams introduced by Brundan and Stroppel [5]–[9]. The classification of the simple modules for the walled Brauer algebras up to isomorphism obtained in [12] enables the parametrization of the indecomposable objects of Deligne's category by bipartitions (cf. Theorem 4.6.2). Writing $L(\lambda)$ for

the indecomposable object corresponding to the bipartition λ , we have the following description for the decomposition of tensor products of indecomposable objects of $\text{Rep}(GL_\delta)$ for generic δ (see Theorem 7.1.1 for a precise statement).

Theorem 1.2.1. *For generic values of $\delta \in \mathbb{K}$,*

$$L(\lambda) \otimes L(\mu) \cong \bigoplus_{\nu} L(\nu)^{\oplus \Gamma_{\lambda, \mu}^{\nu}}$$

where the sum is over bipartitions ν and the coefficients $\Gamma_{\lambda, \mu}^{\nu}$ are given in terms of Littlewood-Richardson coefficients (see (16)).

This theorem is derived using Koike's Theorem [18], which gives the decomposition of a tensor product of irreducible rational representations for the general linear group. In the case of general linear supergroups, it is compatible with the result of Sergeev [25]. The calculation of the decomposition numbers for the walled Brauer algebras [11] and the cap diagrams of Brundan and Stroppel permit the definition of a “lifting isomorphism” in the spirit of Comes and Ostrik [10]. The lifting isomorphism relates the additive Grothendieck rings of $\text{Rep}(GL_\delta)$ in the singular and generic cases, thus enabling the decomposition of any tensor product of indecomposable objects of $\text{Rep}(GL_\delta)$ for any value of δ .

The known decomposition of the mixed tensor powers for GL_d [2, 26, 18, 27] finds the following expression in terms of the functor F_d :

Theorem 1.2.2 (see Theorem 5.2.2). *For any $d > 0$ and any bipartition λ , $F_d(L(\lambda))$ is an indecomposable object of $\text{Rep}(GL_d)$ and is non-zero if and only if $l(\lambda) \leq d$. Moreover, any non-zero indecomposable summand of a mixed tensor power $T(r, s)$ in $\text{Rep}(GL_d)$ is isomorphic to $F_d(L(\lambda))$ for precisely one bipartition $\lambda \in \Lambda_{r, s}$ with $l(\lambda) \leq d$.*

The Young diagram of a bipartition λ is obtained by superimposing the Young diagrams for the partitions $\lambda^\bullet$ and λ° so that their top and left edges coincide, and then rotating the Young diagram for λ° 180-degrees about its upper-left corner (cf. §4.1). A bipartition is $(m|n)$ -cross¹ if its Young diagram can be covered with an m -high, n -wide cross (cf. §8.7). The decomposition of the mixed tensor powers for $\mathfrak{gl}(m|n)$ can be described in terms of the functor $F_{m|n}$ as follows.

Theorem 1.2.3. *For any $m, n \geq 0$ and any bipartition λ , $F_{m|n}(L(\lambda))$ is an indecomposable object of $\text{Rep}(\mathfrak{gl}(m|n))$ and is non-zero if and only if λ is $(m|n)$ -cross. Moreover, any non-zero indecomposable summand of a mixed tensor power $T(r, s)$ in $\text{Rep}(\mathfrak{gl}(m|n))$ is isomorphic to $F_{m|n}(L(\lambda))$ for precisely one $(m|n)$ -cross bipartition $\lambda \in \Lambda_{r, s}$.*

The same result was recently obtained via a different approach by Brundan and Stroppel [5]. Their approach yields additional information about the modules $F_{m|n}(L(\lambda))$; for instance, the irreducible socles and heads are computed explicitly.

In the case when $s = 0$, a bipartition $\lambda \in \Lambda_{r, 0}$ is $(m|n)$ -cross if and only if $\lambda^\bullet$ is $(m|n)$ -hook. Thus, in this case, Theorem 1.2.3 gives the decomposition of the covariant tensor power $V^{\otimes r}$ familiar from the work of Sergeev [25] and of Berele and Regev [4] (similarly, in the contravariant case, i.e., when $r = 0$). On the other hand, if $n = 0$, then a bipartition λ is $(m|n)$ -cross if and only if $l(\lambda) \leq m$. Thus Theorem 1.2.2 can be seen as the special case of Theorem 1.2.3 where $n = 0$.

¹In [22], the term $\mathfrak{gl}(m|n)$ -standard is used.

For any bipartition μ , let $\mathbf{s}_\mu$ denote the corresponding composite supersymmetric Schur polynomial (see e.g. [22]). Then we have the following formula for the characters of the indecomposable summands of the mixed tensor powers for $GL(m|n)$.

Theorem 1.2.4 (see Theorem 8.5.2). *Let $m, n \geq 0$ and $\delta = m - n$. Then for any bipartition λ ,*

$$\text{ch } F_{m|n}(L(\lambda)) = \sum_{\mu} D_{\lambda, \mu}(\delta) \mathbf{s}_\mu,$$

where the $D_{\lambda, \mu}$ are the decomposition numbers for the walled Brauer algebras.

When $|\lambda^\circ| = 0$, the decomposition number $D_{\lambda, \mu}$ is 1 if $\lambda = \mu$ and 0 otherwise, and $\mathbf{s}_\lambda$ is the (non-composite) supersymmetric Schur polynomial associated to $\lambda^\bullet$. Thus, if $\lambda = (\lambda^\bullet, \emptyset)$ and $\lambda^\bullet$ is $(m|n)$ -hook, then the character formula of Theorem 1.2.4 reduces to that of Sergeev [25] and Berele and Regev [4].

1.3. The paper is organized as follows. We begin in §2 with a review of the category-theoretic notions necessary for the definition of Deligne's category and derivation of its universal property in §3. The indecomposable objects of Deligne's category are then classified in §4 using the cellular structure of the walled Brauer algebras as described in [12]. In §5, the representation theory of the general linear group is briefly recalled, and Koike's Theorem on the decomposition of tensor products of indecomposable representations is reviewed. The lifting isomorphism is defined in §6, relating the additive Grothendieck rings of Deligne's category in the singular and generic cases, and it is shown that the defining coefficients are the decomposition numbers of the walled Brauer algebras. The analogue of Koike's Theorem in Deligne's category for generic values of δ is presented in §7, and it is shown that the lifting isomorphism enables the decomposition of tensor products of indecomposables in all cases. In §8, composite supersymmetric Schur polynomials are introduced and the results of §6 and §7 are employed to derive the character formula for the indecomposable summands of the mixed tensor powers of the general linear supergroup. Next, the character formula is used to prove the classification of these indecomposable summands in terms of $(m|n)$ -cross bipartitions, as described by Theorem 1.2.3. Finally, we illustrate how to decompose tensor products of these indecomposable summands with an explicit example.

2. CATEGORY-THEORETIC PRELIMINARIES

This section presents definitions and some elementary results from representation theory expressed in a category-theoretic framework. Let $\mathbb{K}$ denote a field of characteristic zero. A category is said to be $\mathbb{K}$ -linear if the Hom sets are equipped with the structure of vector spaces over the field $\mathbb{K}$ in such a way that composition of morphisms is bilinear.

2.1. Monoidal categories. For any category $\mathcal{C}$, let $\sigma_{\mathcal{C}} : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C}$ denote the functor $(X, Y) \mapsto (Y, X)$. A *monoidal category* is a tuple $(\mathcal{C}, \otimes, \mathbf{1}, c)$ where $\mathcal{C}$ is a category,

$$T := - \otimes - : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$$

is a bifunctor and $\mathbf{1}$ is a distinguished object (the *unit for T*), satisfying

$$T \circ (T \times \text{id}_{\mathcal{C}}) = T \circ (\text{id}_{\mathcal{C}} \times T), \quad \mathbf{1} \otimes - = \text{id}_{\mathcal{C}} = - \otimes \mathbf{1},$$

and $c : T \Rightarrow T \circ \sigma_{\mathcal{C}}$ is a natural isomorphism (the *symmetric braiding*) satisfying

$$\begin{aligned} (\mathrm{id}_Y \otimes c_{X,Z}) \circ (c_{X,Y} \otimes \mathrm{id}_Z) &= c_{X,Y \otimes Z}, & c_{Y,X} \circ c_{X,Y} &= \mathrm{id}_{X \otimes Y}, \\ (c_{X,Z} \otimes \mathrm{id}_Y) \circ (\mathrm{id}_X \otimes c_{Y,Z}) &= c_{X \otimes Y,Z}, \end{aligned}$$

for all objects X, Y, Z of $\mathcal{C}$. When no confusion arises, we may write $\mathcal{C}$ for both the underlying category and monoidal category $(\mathcal{C}, \otimes_{\mathcal{C}}, \mathbb{1}_{\mathcal{C}}, c_{\mathcal{C}})$ and omit the subscript $\mathcal{C}$ when it is implicit. A monoidal category, in our sense, is elsewhere called a strict, symmetric monoidal category.

A *monoidal functor* is a tuple (F, η, α) where $F : \mathcal{C} \rightarrow \mathcal{D}$ is a functor, $\mathcal{C}, \mathcal{D}$ are monoidal categories, $\eta : F \circ T_{\mathcal{C}} \Rightarrow T_{\mathcal{D}} \circ (F \times F)$ is a natural transformation and $\alpha : F(\mathbb{1}_{\mathcal{C}}) \rightarrow \mathbb{1}_{\mathcal{D}}$ is an isomorphism such that

$$\begin{aligned} \eta_{Y,X} \circ F(c_{X,Y}) &= c_{FX,FY} \circ \eta_{X,Y}, & \mathrm{id}_{FX} \otimes (\alpha \circ \eta_{X,\mathbb{1}}) &= \mathrm{id}_{FX}, \\ (\eta_{X,Y} \otimes \mathrm{id}_{FZ}) \circ \eta_{X \otimes Y,Z} &= (\mathrm{id}_{FX} \otimes \eta_{Y,Z}) \circ \eta_{X,Y \otimes Z}. \end{aligned}$$

The monoidal functor is *strict* if $F(X \otimes Y) = FX \otimes FY$ and $\eta_{X,Y} = \mathrm{id}_{FX \otimes FY}$ for all objects X, Y of $\mathcal{C}$, $F\mathbb{1} = \mathbb{1}$ and $\alpha = \mathrm{id}_{\mathbb{1}}$. A monoidal functor, in our sense, is elsewhere called a non-strict symmetric monoidal functor.

Let $\mathcal{C}, \mathcal{D}$ be monoidal categories and let

$$F = (F, \eta, \alpha), F' = (F', \eta', \alpha') : \mathcal{C} \rightarrow \mathcal{D}$$

be monoidal functors. A *monoidal natural transformation* $\epsilon : F \Rightarrow F'$ is a natural transformation of the underlying functors $F \Rightarrow F'$ such that $\epsilon_{\mathbb{1}} = (\alpha')^{-1} \circ \alpha$ and

$$\eta'_{X,Y} \circ \epsilon_{X \otimes Y} = \epsilon_X \otimes \epsilon_Y \circ \eta_{X,Y},$$

for all objects X, Y .

We use monoidal categories in place of their familiar, non-strict, counterparts when convenient. This is without loss of generality, by MacLane's Coherence Theorem (see e.g. [20]). Thus $\mathrm{Vect}_{\mathbb{K}}$ denotes the monoidal category of finite-dimensional vector spaces and linear maps over $\mathbb{K}$, with $\mathbb{1}$ a one-dimensional vector space and the usual symmetric braiding, while $\mathrm{SVect}_{\mathbb{K}}$ denotes the category of all finite-dimensional superspaces and their morphisms over $\mathbb{K}$, with $\mathbb{1}$ a one-dimensional even superspace and symmetric braiding given by the rule of signs. Given a pure element u of a superspace, write $\bar{u}$ for its parity.

2.2. Tensor categories. Let $\mathcal{C}$ be a monoidal category and X an object of $\mathcal{C}$. A *dual* of X is a tuple $(X^*, \mathrm{ev}_X, \mathrm{coev}_X)$ where X^* is an object of $\mathcal{C}$ and $\mathrm{ev}_X, \mathrm{coev}_X$ are morphisms

$$\mathrm{ev}_X : X^* \otimes X \rightarrow \mathbb{1}, \quad \mathrm{coev}_X : \mathbb{1} \rightarrow X \otimes X^*,$$

of $\mathcal{C}$ such that

$$(\mathrm{id}_X \otimes \mathrm{ev}_X) \circ (\mathrm{coev}_X \otimes \mathrm{id}_X) = \mathrm{id}_X, \quad (\mathrm{ev}_X \otimes \mathrm{id}_{X^*}) \circ (\mathrm{id}_{X^*} \otimes \mathrm{coev}_X) = \mathrm{id}_{X^*}.$$

The category $\mathcal{C}$ is *rigid* if every object has a dual. A *tensor category* is a rigid $\mathbb{K}$ -linear monoidal category such that $\mathrm{End} \mathbb{1} = \mathbb{K}$ and $- \otimes -$ is a bilinear bifunctor. A (strict) *tensor functor* is a (strict) monoidal functor that is preadditive.

Let $\varphi : X \rightarrow Y$ be a $\mathcal{C}$ -morphism. Duals $(X^*, \mathrm{ev}_X, \mathrm{coev}_X)$ and $(Y^*, \mathrm{ev}_Y, \mathrm{coev}_Y)$ for X and Y define a *dual morphism* $\varphi^* : Y^* \rightarrow X^*$ by

$$\varphi^* = \mathrm{ev}_Y \otimes \mathrm{id}_{X^*} \circ \mathrm{id}_{Y^*} \otimes \varphi \otimes \mathrm{id}_{X^*} \circ \mathrm{id}_{Y^*} \otimes \mathrm{coev}_X,$$

and one has that

$$(1) \quad \text{ev}_X \circ (\varphi^* \otimes \text{id}_X) = \text{ev}_Y \circ (\text{id}_{Y^*} \otimes \varphi), \quad (\text{id}_X \otimes \varphi^*) \circ \text{coev}_X = (\varphi \otimes \text{id}_{Y^*}) \circ \text{coev}_Y.$$

Now let $\mathcal{D}$ be another monoidal category, and $F = (F, \eta, \alpha) : \mathcal{C} \rightarrow \mathcal{D}$ a monoidal functor. Then the functor F and the dual for X in $\mathcal{C}$ define a dual

$$(2) \quad (F(X^*), \text{ev}_{FX}, \text{coev}_{FX})$$

for FX in $\mathcal{D}$, where

$$\text{ev}_{FX} = \alpha \circ F(\text{ev}_X) \circ \eta_{X^*, X}^{-1}, \quad \text{coev}_{FX} = \eta_{X, X^*} \circ F(\text{coev}_X) \circ \alpha^{-1}.$$

We equip $\text{Vect}_{\mathbb{K}}$ and $\text{SVect}_{\mathbb{K}}$ with the structure of tensor category in the usual way.

Proposition 2.2.1. *Let $\mathcal{C}$ be a rigid monoidal category, $\mathcal{D}$ a monoidal category and*

$$\epsilon : (F, \eta, \alpha) \Rightarrow (F', \eta', \alpha')$$

a monoidal natural transformation of monoidal functors $\mathcal{C} \rightarrow \mathcal{D}$. Then ϵ is a natural isomorphism, and for all objects X in $\mathcal{C}$,

$$\epsilon_{X^*} = ((\epsilon_X)^*)^{-1} = (\epsilon_X^{-1})^*$$

with respect to the duals for FX and $F'X$ defined by (2).

2.3. Categorical dimension. Suppose that $\mathcal{C}$ is a rigid monoidal category. For any object X of $\mathcal{C}$ and $\varphi \in \text{End}_{\mathcal{C}} X$, define the *categorical trace* $\text{tr } \varphi$ by

$$\text{tr } \varphi = \text{ev}_X \circ c_{X, X^*} \circ (\varphi \otimes \text{id}_{X^*}) \circ \text{coev}_X \in \text{End}_{\mathcal{C}} \mathbb{1},$$

and define the *categorical dimension* of X by $\dim X = \text{tr}(\text{id}_X)$. The categorical trace and dimension do not depend on the choice of dual for X , and are preserved by any monoidal functor. When $\mathcal{C} = \text{Vect}_{\mathbb{K}}$ is the category of finite-dimensional vector spaces, categorical trace and dimension coincide with their elementary counterparts. When $\mathcal{C} = \text{SVect}_{\mathbb{K}}$ is the category of super vector spaces, the categorical dimension of an object coincides with the superdimension.

2.4. Functor categories. For categories $\mathcal{C}, \mathcal{D}$, define the following categories whose objects are functors $\mathcal{C} \rightarrow \mathcal{D}$ of the specified type and whose morphisms are natural transformations of the specified type:

category	$\mathcal{C}, \mathcal{D}$	functors	nat. trans.
$\text{Hom}(\mathcal{C}, \mathcal{D})$	any	any	any
$\text{Hom}^{\otimes}(\mathcal{C}, \mathcal{D})$	monoidal	monoidal	monoidal
$\text{Hom}_+(\mathcal{C}, \mathcal{D})$	preadditive	preadditive	any
$\text{Hom}_+^{\otimes}(\mathcal{C}, \mathcal{D})$	tensor	tensor	monoidal
$\text{Hom}_+^{\otimes\text{-str}}(\mathcal{C}, \mathcal{D})$	tensor	strict tensor	monoidal

2.5. The additive envelope. Let $\mathcal{C}$ be a preadditive category. An *additive envelope* of $\mathcal{C}$ is a pair $(\mathcal{C}^{\text{add}}, \iota)$ where $\mathcal{C}^{\text{add}}$ is an additive category and $\iota : \mathcal{C} \rightarrow \mathcal{C}^{\text{add}}$ is a fully-faithful preadditive functor such that for any additive category $\mathcal{D}$, the “restriction functor”

$$\text{Hom}_+(\mathcal{C}^{\text{add}}, \mathcal{D}) \xrightarrow{\sim} \text{Hom}_+(\mathcal{C}, \mathcal{D}), \quad F \mapsto F \circ \iota, \quad (\eta : F \Rightarrow F') \mapsto \eta_{\iota},$$

is an equivalence of categories². Thus an additive envelope is unique up to equivalence of categories, when it exists, and the category $\mathcal{C}$ may be identified with a full

²preadditive functors necessarily preserve biproducts, when they exist.

subcategory of $\mathcal{C}^{\text{add}}$ via the functor ι . Additive envelopes exist for all preadditive categories. If $\mathcal{C}$ is a tensor category, then $\mathcal{C}^{\text{add}}$ can be endowed with the structure of a tensor category such that ι is a tensor functor and for any additive tensor category $\mathcal{D}$,

$$(3) \quad \text{Hom}_+^{\otimes}(\mathcal{C}^{\text{add}}, \mathcal{D}) \xrightarrow{\sim} \text{Hom}_+^{\otimes}(\mathcal{C}, \mathcal{D}), \quad F \mapsto F \circ \iota, \quad (\eta : F \Rightarrow F') \mapsto \eta_{\iota},$$

is an equivalence of categories.

2.6. Splittings of idempotents & the Karoubi envelope. Let $\mathcal{C}$ be a category, X an object of $\mathcal{C}$ and $e^2 = e \in \text{End}_{\mathcal{C}}(X)$; one says that e is an *idempotent* of $\mathcal{C}$. Idempotents $e, e' \in \text{End}_{\mathcal{C}}(X)$ are *orthogonal* if $ee' = e'e = 0$. An idempotent is *primitive* if it is non-zero and cannot be written as the sum of two non-zero orthogonal idempotents. A *splitting* of e is a tuple $(\text{im } e, \iota_e, \pi_e)$ where $\text{im } e$ is an object of $\mathcal{C}$ and

$$\iota_e : \text{im } e \rightarrow X, \quad \pi_e : X \rightarrow \text{im } e,$$

are morphisms of $\mathcal{C}$ such that

- (1) $e = \iota_e \circ \pi_e$,
- (2) $\text{id}_{\text{im } e} = \pi_e \circ \iota_e$.

One says that the idempotent e *splits*, and calls $\text{im } e$ the *image* of e . A category is said to be *Karoubi* if every idempotent of the category splits. A *Karoubi envelope* of a category $\mathcal{C}$ is a tuple $(\mathcal{C}^{\text{kar}}, \iota)$ where $\mathcal{C}^{\text{kar}}$ is Karoubi and $\iota : \mathcal{C} \rightarrow \mathcal{C}^{\text{kar}}$ is a fully-faithful functor such that for any Karoubi category $\mathcal{D}$, the “restriction functor”

$$\text{Hom}(\mathcal{C}^{\text{kar}}, \mathcal{D}) \xrightarrow{\sim} \text{Hom}(\mathcal{C}, \mathcal{D}), \quad F \mapsto F \circ \iota, \quad (\eta : F \Rightarrow F') \mapsto \eta_{\iota},$$

is an equivalence of categories. Thus a Karoubi envelope is unique up to equivalence of categories and the category $\mathcal{C}$ may be identified with a full subcategory of $\mathcal{C}^{\text{kar}}$ via the functor ι . Karoubi envelopes exist for all categories, and if $\mathcal{C}$ is a tensor category, then $\mathcal{C}^{\text{kar}}$ can be endowed with the structure of a tensor category such that ι is a tensor functor and for any Karoubi tensor category $\mathcal{D}$

$$(4) \quad \text{Hom}_+^{\otimes}(\mathcal{C}^{\text{kar}}, \mathcal{D}) \xrightarrow{\sim} \text{Hom}_+^{\otimes}(\mathcal{C}, \mathcal{D}), \quad F \mapsto F \circ \iota, \quad (\eta : F \Rightarrow F') \mapsto \eta_{\iota},$$

is an equivalence of categories. Moreover, if $\mathcal{C}$ is an additive tensor category, then so is $\mathcal{C}^{\text{kar}}$.

2.7. Krull-Schmidt categories. A $\mathbb{K}$ -linear *Krull-Schmidt* category is a category that is $\mathbb{K}$ -linear, additive and Karoubi, with finite-dimensional Hom-spaces. Thus, if $\mathcal{C}$ is any $\mathbb{K}$ -linear category with finite-dimensional Hom-spaces, then the Karoubi envelope of the additive envelope $(\mathcal{C}^{\text{add}})^{\text{kar}}$ is an example of a $\mathbb{K}$ -linear Krull-Schmidt category. As indicated by the following proposition, objects in Krull-Schmidt categories possess essentially unique biproduct decompositions into indecomposable summands, as in the familiar case of finitely-generated modules over a finite-dimensional algebra.

Proposition 2.7.1. *Let $\mathcal{C}$ be a $\mathbb{K}$ -linear category considered as a full subcategory of its Karoubi envelope $\mathcal{C}^{\text{kar}}$, let A be an object of $\mathcal{C}$ and let $e, e', e'' \in \text{End}_{\mathcal{C}} A$ be idempotents with splittings chosen in $\mathcal{C}^{\text{kar}}$. Then:*

- (1) *$\text{im } e$ is indecomposable if and only if e is a primitive idempotent, and up to isomorphism, all indecomposables are so obtained.*
- (2) *if $e = e' + e''$ and e', e'' are orthogonal, then $\text{im } e \cong \text{im } e' \oplus \text{im } e''$.*

(3) Suppose further that $\text{End}_{\mathcal{C}} A$ is finite dimensional. Then $\text{im } e \cong \text{im } e'$ if and only if e and e' are conjugate in $\text{End}_{\mathcal{C}} A$.

Recall that a ring R is *semiperfect* if R/J is semisimple and idempotents of R/J lift to R , where $J = J(R)$ denotes the radical. In particular, any finite-dimensional algebra is semiperfect (see e.g. [1]).

Proposition 2.7.2. *Let $\mathcal{C}$ be a preadditive category and X an object of $\mathcal{C}$. Then*

- (1) *If $\text{End } X$ is local, then X is indecomposable.*
- (2) *Suppose further that $\mathcal{C}$ is Karoubi and $\text{End } X$ is semiperfect. Then if X is indecomposable, then $\text{End } X$ is local.*

Corollary 2.7.3. *Let $\mathcal{C}$ be a $\mathbb{K}$ -linear Krull-Schmidt category and let X be an object of $\mathcal{C}$. Then X is indecomposable if and only if $\text{End } X$ is local.*

Proposition 2.7.4. *Let $\mathcal{C}$ be a $\mathbb{K}$ -linear Krull-Schmidt category, $\mathcal{D}$ a preadditive category and $F : \mathcal{C} \rightarrow \mathcal{D}$ a full preadditive functor. Then FX is indecomposable object of $\mathcal{D}$ if X is an indecomposable object of $\mathcal{C}$. Moreover, if X, Y are indecomposable objects of $\mathcal{C}$ such that FX, FY are non-zero isomorphic objects of $\mathcal{D}$, then $X \cong Y$.*

Proof. The first part follows from Corollary 2.7.3 since homomorphic images of local rings are local. To prove the second part, suppose that X, Y are indecomposable objects of $\mathcal{C}$ and that $FX \cong FY$ are non-zero in $\mathcal{D}$. As F is full, there exist morphisms $\varphi : X \rightarrow Y$, $\psi : Y \rightarrow X$ such that $F(\psi \circ \varphi) = \text{id}_{FX}$. Thus $\alpha = \psi \circ \varphi$ is not nilpotent. Hence $\alpha \notin J$, where J is the radical of the finite-dimensional algebra $\text{End } X$. As $\text{End } X$ is local, J is the unique maximal left ideal, so it follows that $(\text{End } X)\alpha = \text{End } X$. Let $\beta \in \text{End } X$ be such that $\beta \circ \alpha = \beta \circ \psi \circ \varphi = \text{id}_X$. Then $\varphi \circ \beta \circ \psi$ is a non-zero idempotent in the local algebra $\text{End } Y$, hence is equal to id_Y . Thus $X \cong Y$. $\square$

2.8. The additive Grothendieck ring $R_{\mathcal{C}}$. Let $\mathcal{C}$ denote a $\mathbb{K}$ -linear Krull-Schmidt category, let $\mathbb{Z}[\mathcal{C}]$ denote the free $\mathbb{Z}$ -module generated by the isomorphism classes of the objects of $\mathcal{C}$, and let $(\cdot, \cdot)_{\mathcal{C}}$ denote the bilinear form on $\mathbb{Z}[\mathcal{C}]$ defined by bilinear extension of the rule

$$([U], [V])_{\mathcal{C}} = \dim_{\mathbb{K}} \text{Hom}_{\mathcal{C}}(U, V).$$

Write $R_{\mathcal{C}}$ for the quotient of $\mathbb{Z}[\mathcal{C}]$ by the relations

$$[A \oplus B] - [A] - [B]$$

for all objects A, B in $\mathcal{C}$, and endow it with the induced bilinear form. We call $R_{\mathcal{C}}$ the *additive Grothendieck group* of $\mathcal{C}$. If $\mathcal{C}$ is a $\mathbb{K}$ -linear Krull-Schmidt tensor category, then $R_{\mathcal{C}}$ is a commutative ring, called the *additive Grothendieck ring* of $\mathcal{C}$, via $[A][B] = [A \otimes B]$ for all objects A, B of $\mathcal{C}$.

3. THE CATEGORY $\underline{\text{Rep}}(GL_\delta)$

In this section we define Deligne's tensor category $\underline{\text{Rep}}(GL_\delta)$ and prove that it satisfies a certain universal property (see Proposition 3.5.1). To define $\underline{\text{Rep}}(GL_\delta)$ we first diagrammatically define a smaller "skeleton category". We then take the additive envelope (§2.5) followed by the Karoubi envelope (§2.6) of this skeleton category to get $\underline{\text{Rep}}(GL_\delta)$. To start, we introduce the diagrams we will use to define the skeleton category.

3.1. Words and diagrams. Suppose w and w' are finite (possibly empty) words in two letters denoted $\bullet$ (black letter) and $\circ$ (white letter). A (w, w') -*diagram* is a graph which satisfies the following conditions:

- (i) The vertices are positioned in two (possibly empty) horizontal rows.
- (ii) Each vertex is drawn as either $\bullet$ or $\circ$ so that the bottom (resp. top) row of vertices is the word w (resp. w').
- (iii) Each vertex is adjacent to exactly one edge.
- (iv) An edge is adjacent to both a black and a white vertex if and only if the vertices adjacent to that edge are either both in the top row or both in the bottom row.

An edge in a (w, w') -diagram is called a *propagating edge* if it is adjacent to a vertex in the top row and a vertex in the bottom row.

Example 3.1.1. (1) Let $\mathbf{1}$ denote the empty word. The empty graph is the unique $(\mathbf{1}, \mathbf{1})$ -diagram. On the other hand, there are two $(\bullet\bullet\circ\circ, \mathbf{1})$ -diagrams:

(2) There are six $(\bullet\circ\circ\bullet, \bullet\circ)$ -diagrams:

Each of the top four $(\bullet\circ\circ\bullet, \bullet\circ)$ -diagrams have two propagating edges, whereas the bottom two have no propagating edges.

Remark 3.1.2. Suppose w (resp. w') is a word with r (resp. r') black letters and s (resp. s') white letters. It is easy to show that a (w, w') -diagram exists if and only if $r + s' = r' + s$, in which case the number of (w, w') -diagrams is $(r + s')!$.

Suppose w, w' , and w'' are finite words in the letters $\bullet$ and $\circ$. Given a (w, w') -diagram X and a (w', w'') -diagram Y , we let $Y \star X$ denote the graph obtained by stacking Y atop X so that the top row of vertices of X are identified with the bottom row of vertices of Y . Next, we let $Y \cdot X$ denote the (w, w'') -diagram obtained by restricting $Y \star X$ to its top and bottom rows of vertices. Finally, let $\ell(X, Y)$ denote the number of cycles in $Y \star X$ (i.e. the number of connected components of $Y \star X$ minus the number of connected components of $Y \cdot X$). For example, if

and $\ell(X, Y) = 2$.

3.2. The skeleton category. Fix $\delta \in \mathbb{K}$. Using the setup from 3.1 we can now define the skeleton category $\underline{\text{Rep}}_0(GL_\delta)$.

Definition 3.2.1. The category $\underline{\text{Rep}}_0(GL_\delta)$ has:

Objects: finite words in the letters $\bullet$ and $\circ$.

Morphisms: $\text{Hom}(w, w')$ is the $\mathbb{K}$ -vector space on basis $\{(w, w')\text{-diagrams}\}$.

Composition: $\text{Hom}(w', w'') \times \text{Hom}(w, w') \rightarrow \text{Hom}(w, w'')$ sending $(f, g) \mapsto fg$ is the $\mathbb{K}$ -bilinear map satisfying $YX = \delta^{\ell(X, Y)} Y \cdot X$ whenever X is a (w, w') -diagram and Y is a (w', w'') -diagram.

To show $\underline{\text{Rep}}_0(GL_\delta)$ is indeed a category, it is easy to check that composition in $\underline{\text{Rep}}_0(GL_\delta)$ is associative. Also, if w is a finite word in $\bullet$ and $\circ$, then the (w, w) -diagram with each vertex in the bottom row adjacent to the vertex directly above it is the identity morphism in $\text{End}(w)$. For example,

$$\text{id}_{\bullet\circ\circ\bullet\bullet\circ\circ} = \begin{array}{cccccc} \bullet & \circ & \circ & \bullet & \bullet & \circ & \circ \\ | & | & | & | & | & | & | \\ \bullet & \circ & \circ & \bullet & \bullet & \circ & \circ \end{array}.$$

3.3. Tensor category structure of $\underline{\text{Rep}}_0(GL_\delta)$. We will now equip $\underline{\text{Rep}}_0(GL_\delta)$ with the structure of a tensor category in the sense of §2.2.

Definition 3.3.1. The bifunctor $- \otimes - : \underline{\text{Rep}}_0(GL_\delta) \times \underline{\text{Rep}}_0(GL_\delta) \rightarrow \underline{\text{Rep}}_0(GL_\delta)$ is defined as follows:

On objects: Set $w_1 \otimes w_2 = w_1 w_2$ (concatenation of words) for any objects w_1 and w_2 in $\underline{\text{Rep}}_0(GL_\delta)$.

On morphisms: Assume w_i and w'_i are finite words in $\bullet$ and $\circ$, and X_i is a (w_i, w'_i) -diagram for $i = 1, 2$. Let $X_1 \otimes X_2$ denote the $(w_1 w_2, w'_1 w'_2)$ -diagram obtained by placing X_1 directly to the left of X_2 . Now extend $\mathbb{K}$ -linearly in both arguments to define tensor products of arbitrary morphisms in $\underline{\text{Rep}}_0(GL_\delta)$.

Let $c_{w_1, w_2} : w_1 \otimes w_2 \rightarrow w_2 \otimes w_1$ be the $(w_1 w_2, w_2 w_1)$ -diagram such that the vertex in the bottom row corresponding to the i th letter in w_1 (resp. w_2) is adjacent to the vertex in the top row corresponding to the i th letter in w_1 (resp. w_2). For example,

$$c_{\circ\circ\bullet, \bullet\bullet\circ\circ} = \begin{array}{cccccc} \circ & \bullet & \bullet & \circ & \circ & \bullet \\ & \diagdown & \diagup & \diagdown & \diagup & \\ & \circ & \circ & \bullet & \bullet & \circ \\ & \diagup & \diagdown & \diagup & \diagdown & \\ \circ & \bullet & \bullet & \circ & \circ & \bullet \end{array}.$$

It is easy to see that Definition 3.3.1 gives $\underline{\text{Rep}}_0(GL_\delta)$ the structure of a monoidal category with unit object $\mathbf{1}$ (the empty word) and symmetric braiding c .

Next, we will show that $\underline{\text{Rep}}_0(GL_\delta)$ is rigid. To do so, given a finite word w in $\bullet$ and $\circ$ let w^* denote the word obtained from w by replacing all black letters with white letters and vice versa. Now define the morphism $\text{ev}_w : w^* \otimes w \rightarrow \mathbf{1}$ (resp. $\text{coev}_w : \mathbf{1} \rightarrow w \otimes w^*$) to be the $(w^* w, \mathbf{1})$ -diagram (resp. $(\mathbf{1}, w w^*)$ -diagram) such that the i th letter in w^* is adjacent to the i th letter in w for all i . For example,

$$\text{ev}_{\bullet\circ\bullet} = \begin{array}{ccccc} & & \bullet & \circ & \bullet \\ & \diagdown & \diagup & \diagdown & \diagup \\ & \circ & \circ & \bullet & \bullet \\ & \diagup & \diagdown & \diagup & \diagdown \\ \circ & \bullet & \bullet & \circ & \bullet \end{array} \quad \text{and} \quad \text{coev}_{\bullet\circ\bullet} = \begin{array}{ccccc} \bullet & \circ & \bullet & \circ & \bullet \\ & \diagdown & \diagup & \diagdown & \diagup \\ & \circ & \circ & \bullet & \bullet \\ & \diagup & \diagdown & \diagup & \diagdown \\ \circ & \bullet & \bullet & \circ & \bullet \end{array}.$$

It is easy to check that ev_w and coev_w make w^* a dual to w . Since $\text{End}(\mathbf{1}) = \mathbb{K}$ and $- \otimes -$ is bilinear, it follows that $\underline{\text{Rep}}_0(GL_\delta)$ is a tensor category.

3.4. Universal property of $\underline{\text{Rep}}_0(GL_\delta)$. Following §2.3 we can compute the categorical dimension of any object in $\underline{\text{Rep}}_0(GL_\delta)$. In particular,

$$\dim(\bullet) = \text{ev}_\bullet \circ c_{\bullet, \circ} \circ \text{coev}_\bullet = \delta$$

where the last equality follows from the fact that

$$\text{ev}_\bullet \star c_{\bullet, \circ} \star \text{coev}_\bullet = \text{diagram}$$

$\underline{\text{Rep}}_0(GL_\delta)$ is characterized as the universal tensor category generated by an object of dimension δ and its dual [14]. More precisely, it possesses the following universal property:

Proposition 3.4.1. *Given a tensor category $\mathcal{T}$, let $\mathcal{T}_\delta$ denote the category of δ -dimensional objects in $\mathcal{T}$ and their isomorphisms. Then the following functor induces an equivalence of categories:*

$$\begin{aligned} \Theta : \text{Hom}_+^{\otimes\text{-str}}(\underline{\text{Rep}}_0(GL_\delta), \mathcal{T}) &\rightarrow \mathcal{T}_\delta, \\ F &\mapsto F(\bullet), \\ (\eta : F \Rightarrow F') &\mapsto \eta_\bullet. \end{aligned}$$

Proof. Since tensor functors preserve categorical dimension, $\Theta(F) = F(\bullet)$ is an object of the category $\mathcal{T}_\delta$, and by Proposition 2.2.1, $\Theta(\eta) = \eta_\bullet$ is an isomorphism of $F(\bullet)$, hence a morphism of $\mathcal{T}_\delta$. Let X be an object of $\mathcal{T}_\delta$ and let $(X^*, \text{ev}_X, \text{coev}_X)$ be a dual for X in $\mathcal{T}$. Let w, w' be finite words in $\bullet$ and $\circ$. By the coherence theorem for tensor categories (see for instance [23] and references therein), the image of any (w, w') -diagram under a strict tensor functor is completely determined by the image of

$$(5) \quad \text{id}_\bullet, \text{id}_\circ, \text{ev}_\bullet, \text{coev}_\bullet.$$

Since

$$\text{ev}_X \circ c_{X, X^*} \circ \text{coev}_X = \dim X = \delta,$$

there exists a unique, well-defined, strict tensor functor $F_X : \underline{\text{Rep}}_0(GL_\delta) \rightarrow \mathcal{T}$ with

$$\begin{aligned} F_X : \quad \bullet &\mapsto X, & \circ &\mapsto X^*, \\ \text{ev}_\bullet &\mapsto \text{ev}_X, & \text{coev}_\bullet &\mapsto \text{coev}_X. \end{aligned}$$

Thus Θ is essentially surjective.

We now show that Θ is full. Let X, Y be objects of $\mathcal{T}_\delta$ and write F_X, F_Y for the functors defined X, Y as above. Suppose now that $\varphi : X \rightarrow Y$ is a morphism in $\mathcal{T}_\delta$; so φ is invertible. Define a family of isomorphisms

$$\epsilon = (\epsilon_w : F_X(w) \Rightarrow F_Y(w))_w$$

indexed by finite words w , by

$$\epsilon_\bullet = \varphi, \quad \epsilon_\circ = (\varphi^{-1})^*$$

and $\epsilon_{w \otimes w'} = \epsilon_w \otimes \epsilon_{w'}$ for all finite words w, w' . It remains to show that ϵ is a natural transformation, that is, for all morphisms $\sigma : w \rightarrow w'$ in $\underline{\text{Rep}}_0(GL_\delta)$, that $F_Y(\sigma) \circ \epsilon_w = \epsilon_{w'} \circ F_X(\sigma)$. For finite words w, w' , we have

$$\begin{aligned} F_Y(c_{w, w'}) \circ \epsilon_{ww'} &= c_{F_Y w, F_Y w'} \circ \epsilon_{ww'} \\ &= \epsilon_{w' w} \circ c_{F_X w, F_X w'} \\ &= \epsilon_{w' w} \circ F_X(c_{w, w'}). \end{aligned}$$

Thus, in verifying naturality, the words may be reordered. As ϵ is monoidal by construction, it suffices to check naturality in the cases where σ is one of the four morphisms (5). The cases of the identity morphisms are trivial. In the cases of the latter two morphisms, the naturality relations are precisely

$$\text{ev}_X = \text{ev}_Y \circ (\varphi^{-1})^* \otimes \varphi, \quad \varphi \otimes (\varphi^{-1})^* \circ \text{coev}_X = \text{coev}_Y,$$

which follow immediately from (1). Thus Θ is full.

Finally, suppose that ϵ is a morphism of $\mathcal{H}om_+^{\otimes\text{-str}}(\underline{\text{Rep}}_0(GL_\delta), \mathcal{T})$. Since ϵ is monoidal, it is determined by $\epsilon_\bullet$ and $\epsilon_\circ$. By Proposition 2.2.1, $\epsilon_\circ = (\epsilon_\bullet^{-1})^*$, and so ϵ is determined by $\epsilon_\bullet$ alone. Thus Θ is faithful. $\square$

3.5. Definition of $\underline{\text{Rep}}(GL_\delta)$. Let $\underline{\text{Rep}}(GL_\delta) = (\underline{\text{Rep}}_0(GL_\delta)^{\text{add}})^{\text{kar}}$ denote the Karoubi envelope of the additive envelope of $\underline{\text{Rep}}_0(GL_\delta)$, as per §2.5 and §2.6. Thus $\underline{\text{Rep}}_0(GL_\delta)$ may be identified with a full subcategory of $\underline{\text{Rep}}(GL_\delta)$, and the tensor category structure of $\underline{\text{Rep}}(GL_\delta)$ extends that of $\underline{\text{Rep}}_0(GL_\delta)$. For every idempotent e of $\underline{\text{Rep}}_0(GL_\delta)$, fix a splitting $(\text{im } e, \iota_e, \pi_e)$ of e in $\underline{\text{Rep}}(GL_\delta)$. As a notational convenience, whenever $e \in \text{End } X$ and $f \in \text{End } Y$ are idempotents of objects X, Y of $\underline{\text{Rep}}_0(GL_\delta)$, identify

$$\text{Hom}_{\underline{\text{Rep}}(GL_\delta)}(\text{im } e, \text{im } f) = f \text{Hom}_{\underline{\text{Rep}}_0(GL_\delta)}(X, Y)e \subset \text{Hom}_{\underline{\text{Rep}}_0(GL_\delta)}(X, Y)$$

via the morphisms $\iota_e, \iota_f, \pi_e, \pi_f$. For any tensor category $\mathcal{T}$, let $\mathcal{H}om'(\underline{\text{Rep}}(GL_\delta), \mathcal{T})$ denote the full subcategory of $\mathcal{H}om_+^{\otimes}(\underline{\text{Rep}}(GL_\delta), \mathcal{T})$ whose objects are those functors whose restriction $\underline{\text{Rep}}_0(GL_\delta) \rightarrow \mathcal{T}$ yields a strict tensor functor. The category $\underline{\text{Rep}}(GL_\delta)$ has the following universal property (see [14, Proposition 10.3]).

Proposition 3.5.1. *Suppose that $\mathcal{T}$ is a tensor category and let $\mathcal{T}_\delta$ be as in Proposition 3.4.1. Then the following functor induces an equivalence of categories:*

$$\begin{aligned} \mathcal{H}om'(\underline{\text{Rep}}(GL_\delta), \mathcal{T}) &\rightarrow \mathcal{T}_\delta, \\ F &\mapsto F(\bullet), \\ (\eta : F \Rightarrow F') &\mapsto \eta_\bullet. \end{aligned}$$

Proof. The universal properties of the additive envelope (3) and Karoubi envelope (4), yield

$$\mathcal{H}om_+^{\otimes}(\underline{\text{Rep}}(GL_\delta), \mathcal{T}) \cong \mathcal{H}om_+^{\otimes}(\underline{\text{Rep}}_0(GL_\delta), \mathcal{T})$$

via the restriction functors there described. As $\mathcal{H}om_+^{\otimes\text{-str}}(\underline{\text{Rep}}_0(GL_\delta), \mathcal{T})$ is a full subcategory of the latter, the result follows from Proposition 3.4.1. $\square$

4. INDECOMPOSABLE OBJECTS IN $\underline{\text{Rep}}(GL_\delta)$

The main goal of this section is to classify isomorphism classes of indecomposable objects in $\underline{\text{Rep}}(GL_\delta)$. By Proposition 2.7.1(1), these indecomposable objects correspond to primitive idempotents in endomorphism algebras of $\underline{\text{Rep}}_0(GL_\delta)$. In light of this, we first describe the classification of conjugacy classes of such primitive idempotents.

To start, let us fix some notation. For non-negative integers r and s , let $w_{r,s}$ denote the word with r black letters followed by s white letters:

$$w_{r,s} = \underbrace{\bullet \cdots \bullet}_r \underbrace{\circ \cdots \circ}_s.$$

Using the symmetric braiding it is easy to see that every object in $\underline{\text{Rep}}_0(GL_\delta)$ is naturally isomorphic to $w_{r,s}$ for some $r, s \geq 0$. Hence, we will only consider

endomorphisms of the $w_{r,s}$'s. We will write $\mathbb{K}B_{r,s}(\delta)$ (or just $B_{r,s}$) for the endomorphism algebra $\text{End}(w_{r,s})$. The algebras $B_{r,s}$ are the so-called *walled Brauer algebras* (compare with [2], [18], [27]). It is well known that conjugacy classes of primitive idempotents in an algebra A are in bijective correspondence with isomorphism classes of simple A -modules³, which in turn are in bijective correspondence with isomorphism classes of projective indecomposable A -modules, hereafter referred to as PIMs (see for example [3]). In this correspondence a primitive idempotent $e \in A$ corresponds to the PIM eA . The isomorphism classes of simples in the walled Brauer algebras are classified in [12]. To explain their classification, we first need to recall some properties of (bi)partitions and their relation to symmetric groups.

4.1. (Bi)partitions. A *partition* is a tuple of non-negative integers $\alpha = (\alpha_1, \alpha_2, \dots)$ whose *parts* (i.e. α_i 's) are such that $\alpha_i \geq \alpha_{i+1}$ for all $i > 0$, and $\alpha_i = 0$ for all but finitely many i . We write $|\alpha| = \sum_{i>0} \alpha_i$ for the *size* of α and we write $\alpha \vdash |\alpha|$. We define the *length* of α , written $l(\alpha)$, to be the smallest positive integer with $\alpha_{l(\alpha)+1} = 0$. We will sometimes write $(\dots 2^{a_2} 1^{a_1})$ for the partition with a_i parts equal to i . It will be convenient for us to identify a partition α with its *Young diagram* which consists of $l(\alpha)$ left-aligned rows of boxes, with α_i boxes in the i th row (reading from top to bottom). For example,

$$(5, 2, 2, 2, 1, 1, 0, \dots) = (5, 2^3, 1^2) =$$

Next, we let α^t denote the *transpose* of α , i.e., α_i^t is the number of boxes in the i th column of α . For example, $(5, 2^3, 1^2)^t = (6, 4, 1^3)$. Finally, we let $\mathcal{P}$ denote the set of all partitions.

Elements of $\mathcal{P} \times \mathcal{P}$ are called *bipartitions*. Given a bipartition λ , we let $\lambda^\bullet$ and λ° denote the partitions such that $\lambda = (\lambda^\bullet, \lambda^\circ)$. We write $|\lambda| = (|\lambda^\bullet|, |\lambda^\circ|)$ for the *size* of λ and we write $\lambda \vdash |\lambda|$. Moreover, we write $l(\lambda) := l(\lambda^\bullet) + l(\lambda^\circ)$ for the *length* of λ . We define a partial order on sizes of bipartitions by declaring that $(a, b) \leq (c, d)$ whenever $a \leq c$ and $b \leq d$. In particular, we write $|\mu| < |\lambda|$ to mean $|\mu| \leq |\lambda|$ and $\mu \neq \lambda$. Additionally, we set $\lambda^* = (\lambda^\bullet, \lambda^\circ)^* = (\lambda^\circ, \lambda^\bullet)$. We also have a bipartition-version of a Young diagram which we get as follows: first place the Young diagram of λ° atop the Young diagram of $\lambda^\bullet$ so that the upper left corners are overlapping, then rotate the Young diagram of λ° 180 degrees about its upper left corner. For example, the diagram associated to the bipartition $((4, 3, 1), (2^2, 1^2))$ is

³It will be convenient for us to work with right modules. However, the categories of right and left $B_{r,s}$ -modules are equivalent via the anti-automorphism on $B_{r,s}$ given by reading diagrams up rather than down the page.

4.2. Symmetric groups. For a non-negative integer r , let Σ_r denote the symmetric group on r -elements, and let $\mathbb{K}\Sigma_r$ denote the corresponding group algebra⁴. It is well known that the primitive idempotents in $\mathbb{K}\Sigma_r$ (up to conjugation) are in bijective correspondence with partitions of size r (see for example [16]). Given $\alpha \vdash r$, let $z_\alpha \in \mathbb{K}\Sigma_r$ denote the corresponding primitive idempotent. For example, $z_{(r)} = \frac{1}{r!} \sum_{\sigma \in \Sigma_r} \sigma$ so that the partition (r) corresponds to the trivial $\mathbb{K}\Sigma_r$ -module.

We now connect the theory of symmetric groups with that of the walled Brauer algebras. Regardless of δ , the walled Brauer algebras $B_{r,0}$ and $B_{0,r}$ are isomorphic to the group algebra $\mathbb{K}\Sigma_r$. These isomorphisms are given by

$$\begin{array}{ccccc} B_{r,0} & \xleftarrow{\sim} & \mathbb{K}\Sigma_r & \xrightarrow{\sim} & B_{0,r} \\ \sigma^\bullet & \longleftarrow & \sigma & \mapsto & \sigma^\circ \end{array}$$

where, given $\sigma \in \Sigma_r$, $\sigma^\bullet$ (resp. σ°) is the $(w_{r,0}, w_{r,0})$ -diagram (resp. $(w_{0,r}, w_{0,r})$ -diagram) whose i th bottom vertex is adjacent to its $\sigma(i)$ th top vertex (reading left to right) for $1 \leq i \leq r$. For example, if $\sigma \in \Sigma_5$ is the 3-cycle $2 \mapsto 3 \mapsto 5 \mapsto 2$, then

$$\sigma^\bullet = \begin{array}{c} \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ | \quad \diagup \quad \diagdown \quad \diagup \quad \diagdown \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \quad \text{and} \quad \sigma^\circ = \begin{array}{c} \circ \quad \circ \quad \circ \quad \circ \quad \circ \\ | \quad \diagdown \quad \diagup \quad \diagdown \quad \diagup \\ \circ \quad \circ \quad \circ \quad \circ \quad \circ \end{array}.$$

More generally, given non-negative integers r and s , we have an inclusion of algebras $\mathbb{K}[\Sigma_r \times \Sigma_s] \hookrightarrow B_{r,s}$ given by $(\sigma, \tau) \mapsto \sigma^\bullet \otimes \tau^\circ$ for all $\sigma \in \Sigma_r, \tau \in \Sigma_s$ (here $\mathbb{K}[\Sigma_r \times \Sigma_s]$ denotes the group algebra of the direct product $\Sigma_r \times \Sigma_s$). Using this embedding we can consider $\mathbb{K}[\Sigma_r \times \Sigma_s]$ as a subalgebra of $B_{r,s}$, and we will do so for the rest of the paper.

Now, let $J \subset B_{r,s}$ denote the $\mathbb{K}$ -span of all $(w_{r,s}, w_{r,s})$ -diagrams with less than $r + s$ propagating edges. One can show that J is a two-sided ideal in $B_{r,s}$ with $B_{r,s}/J \cong \mathbb{K}[\Sigma_r \times \Sigma_s]$ (see [12, Proposition 2.3 and (2)]). Hence, we have a surjection of algebras $\pi : B_{r,s} \twoheadrightarrow \mathbb{K}[\Sigma_r \times \Sigma_s]$. It is straightforward to show

$$(6) \quad \pi(a) = a \quad \text{for all } a \in \mathbb{K}[\Sigma_r \times \Sigma_s] \subset B_{r,s}.$$

4.3. Definition of the idempotent e_λ . Given a bipartition $\lambda \vdash (r, s)$, we set $z_\lambda := z_{\lambda^\bullet}^\bullet \otimes z_{\lambda^\circ}^\circ \in \mathbb{K}[\Sigma_r \times \Sigma_s] \subset B_{r,s}$. Note that z_λ is an idempotent defined up to conjugation. The assignment $\lambda \mapsto z_\lambda$ induces a bijection between bipartitions of size (r, s) and primitive idempotents in $\mathbb{K}[\Sigma_r \times \Sigma_s]$ (up to conjugation). It is important to notice that while z_λ is a primitive idempotent in $\mathbb{K}[\Sigma_r \times \Sigma_s]$, it will generally not be primitive in the (usually) larger algebra $B_{r,s}$. Let $z_\lambda = e_1 + \cdots + e_k$ be a decomposition of z_λ into mutually orthogonal primitive idempotents in $B_{r,s}$. Then $\pi(e_1), \dots, \pi(e_k)$ are mutually orthogonal idempotents in $\mathbb{K}[\Sigma_r \times \Sigma_s]$ whose sum, by (6), is z_λ . As z_λ is primitive in $\mathbb{K}[\Sigma_r \times \Sigma_s]$, there is a unique $i \in \{1, \dots, k\}$ such that $\pi(e_i) \neq 0$. Set $e_\lambda = e_i$. Again, note that $e_\lambda \in B_{r,s}$ is a primitive idempotent defined up to conjugation.

Example 4.3.1. (1) Let $\emptyset$ denote the empty partition $(0, 0, \dots)$. Then $z_\emptyset$ is the identity element of $\mathbb{K}\Sigma_0 = B_{0,0}$. Hence, if $\lambda \vdash (r, 0)$, then $z_\lambda = z_{\lambda^\bullet}^\bullet \otimes \text{id}_1 = z_{\lambda^\bullet}^\bullet$. Moreover, since $B_{r,0} = \mathbb{K}\Sigma_r$, we also have $e_\lambda = z_{\lambda^\bullet}^\bullet$. Similarly, if $\lambda \vdash (0, s)$, then $e_\lambda = z_{\lambda^\circ}^\circ$. As a special case, $e_{(\emptyset, \emptyset)} = \text{id}_1$ (the empty graph).

(2) Consider the bipartition $(\square, \square)$. $z_\square^\bullet = \text{id}_\bullet$ and $z_\square^\circ = \text{id}_\circ$ which implies $z_{(\square, \square)} = \text{id}_{\bullet\circ}$. If $\delta \neq 0$, then $\text{id}_{\bullet\circ}$ decomposes as $e_1 + e_2$ in $B_{1,1}$ where e_1 and e_2 are

⁴By convention, Σ_0 denotes the trivial group of size $0! = 1$, hence we identify $\mathbb{K}\Sigma_0 = B_{0,0} = \mathbb{K}$.

the following orthogonal primitive idempotents:

$$e_1 = \begin{array}{c} \bullet \\ | \\ \bullet \end{array} \begin{array}{c} \circ \\ | \\ \circ \end{array} - \frac{1}{\delta} \begin{array}{c} \bullet \quad \circ \\ \diagdown \quad \diagup \\ \bullet \quad \circ \end{array} \quad \text{and} \quad e_2 = \frac{1}{\delta} \begin{array}{c} \bullet \quad \circ \\ \diagdown \quad \diagup \\ \bullet \quad \circ \end{array}.$$

In this case $\pi(e_1) = \text{id}_{\bullet\circ}$ and $\pi(e_2) = 0$, hence $e_{(\square, \square)} = e_1$. If $\delta = 0$, then $\text{id}_{\bullet\circ}$ is primitive in $B_{1,1}$, and hence is equal to $e_{(\square, \square)}$.

We close this subsection with the following useful proposition. For a proof, we refer the reader to the proof of a completely analogous statement for $\underline{\text{Rep}}(S_t)$ found [10, Proposition 3.8].

Proposition 4.3.2. *The idempotents e_λ are absolutely primitive. In other words, if $\mathbb{K} \subset \mathbb{K}'$ is a field extension, then $e_\lambda \in \mathbb{K}B_{r,s}(\delta)$ is primitive when viewed as an idempotent in $\mathbb{K}'B_{r,s}(\delta)$.*

4.4. Definition of the idempotent $e_\lambda^{(i)}$. Next, we explain how to construct new idempotents from the e_λ 's. Consider the following morphisms:

$$\begin{aligned} \psi_{r,s} &= (\text{id}_\bullet)^{\otimes r} \otimes \text{coev}_\bullet \otimes (\text{id}_\circ)^{\otimes s}, \\ \hat{\psi}_{r,s} &= (\text{id}_\bullet)^{\otimes r} \otimes \text{ev}_\circ \otimes (\text{id}_\circ)^{\otimes s}, \\ \phi_{r,s} &= (\text{id}_\bullet)^{\otimes r} \otimes ((\text{ev}_\circ \otimes \text{id}_\circ)(\text{id}_\bullet \otimes c_{\circ,\circ})) \otimes (\text{id}_\circ)^{\otimes s-1} \quad (s > 0), \\ \hat{\phi}_{r,s} &= (\text{id}_\bullet)^{\otimes r-1} \otimes ((\text{id}_\bullet \otimes \text{ev}_\circ)(c_{\bullet,\bullet} \otimes \text{id}_\circ)) \otimes (\text{id}_\circ)^{\otimes s} \quad (r > 0). \end{aligned}$$

For example,

$$\begin{aligned} \psi_{2,3} &= \begin{array}{c} \bullet \quad \bullet \quad \bullet \quad \circ \quad \circ \quad \circ \\ \diagdown \quad \diagdown \quad \diagup \quad \diagup \quad \diagup \quad \diagup \\ \bullet \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \end{array}, & \hat{\psi}_{2,3} &= \begin{array}{c} \bullet \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \\ \diagdown \quad \diagdown \quad \diagup \quad \diagup \quad \diagup \quad \diagup \\ \bullet \quad \bullet \quad \bullet \quad \circ \quad \circ \quad \circ \end{array}, \\ \phi_{2,3} &= \begin{array}{c} \bullet \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \\ \diagdown \quad \diagdown \quad \diagup \quad \diagup \quad \diagup \quad \diagup \\ \bullet \quad \bullet \quad \bullet \quad \circ \quad \circ \quad \circ \end{array}, & \hat{\phi}_{2,3} &= \begin{array}{c} \bullet \quad \bullet \quad \circ \quad \circ \quad \circ \quad \circ \\ \diagdown \quad \diagdown \quad \diagup \quad \diagup \quad \diagup \quad \diagup \\ \bullet \quad \bullet \quad \bullet \quad \circ \quad \circ \quad \circ \end{array}. \end{aligned}$$

The following identities easily follow from the definitions above:

$$(7) \quad \phi_{r,s} \psi_{r,s} = \text{id}_{w_{r,s}}, \quad \hat{\phi}_{r,s} \psi_{r,s} = \text{id}_{w_{r,s}}, \quad \hat{\psi}_{r,s} \psi_{r,s} = \delta \text{id}_{w_{r,s}}.$$

Now, given a bipartition $\lambda \vdash (r, s)$ we set $e_\lambda^{(0)} = e_\lambda$ and define $e_\lambda^{(i)} \in B_{r+i, s+i}$ for $i > 0$ recursively by

$$e_\lambda^{(i)} = \begin{cases} \psi_{r+i-1, s+i-1} e_\lambda^{(i-1)} \phi_{r+i-1, s+i-1} & \text{if } s > 0, \\ \psi_{r+i-1, s+i-1} e_\lambda^{(i-1)} \hat{\phi}_{r+i-1, s+i-1} & \text{if } s = 0 \text{ and } r > 0, \\ \frac{1}{\delta} \psi_{i-1, i-1} e_\lambda^{(i-1)} \hat{\psi}_{i-1, i-1} & \text{if } \lambda = (\emptyset, \emptyset) \text{ and } \delta \neq 0. \end{cases}$$

Notice that $e_\lambda^{(i)}$ is undefined when $i > 0$, $\lambda = (\emptyset, \emptyset)$ and $\delta = 0$. However, $e_\lambda^{(i)}$ is defined (up to conjugation) in all other cases.

Example 4.4.1. (1) In Example 4.3.1(1) we found that $e_{(\emptyset, \emptyset)}$ is the empty graph. Hence for $\delta \neq 0$ we have

$$e_{(\emptyset, \emptyset)}^{(1)} = \frac{1}{\delta} \begin{array}{c} \bullet \quad \circ \\ \diagdown \quad \diagup \\ \bullet \quad \circ \end{array}, \quad e_{(\emptyset, \emptyset)}^{(2)} = \frac{1}{\delta^2} \begin{array}{c} \bullet \quad \bullet \quad \circ \quad \circ \\ \diagdown \quad \diagdown \quad \diagup \quad \diagup \\ \bullet \quad \bullet \quad \circ \quad \circ \end{array}, \quad e_{(\emptyset, \emptyset)}^{(3)} = \frac{1}{\delta^3} \begin{array}{c} \bullet \quad \bullet \quad \bullet \quad \circ \quad \circ \quad \circ \\ \diagdown \quad \diagdown \quad \diagdown \quad \diagup \quad \diagup \quad \diagup \\ \bullet \quad \bullet \quad \bullet \quad \circ \quad \circ \quad \circ \end{array}, \dots$$

(2) By Example 4.3.1(1), for any $\delta \in \mathbb{K}$ we have

$$e_{((2), \emptyset)} = \frac{1}{2} \left[\begin{array}{c} \bullet \quad \bullet \\ | \quad | \\ \bullet \quad \bullet \end{array} + \begin{array}{c} \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \end{array} \right].$$

$$\text{Hence } e_{((2), \emptyset)}^{(1)} = \frac{1}{2} \left[\begin{array}{c} \bullet \quad \bullet \quad \circ \\ | \quad | \quad \diagup \\ \bullet \quad \bullet \quad \circ \end{array} + \begin{array}{c} \bullet \quad \bullet \quad \circ \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \quad \circ \end{array} \right]$$

$$\text{and } e_{((2), \emptyset)}^{(2)} = \frac{1}{2} \left[\begin{array}{c} \bullet \quad \bullet \quad \bullet \quad \circ \\ | \quad | \quad | \quad \diagup \\ \bullet \quad \bullet \quad \bullet \quad \circ \end{array} + \begin{array}{c} \bullet \quad \bullet \quad \bullet \quad \circ \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \quad \bullet \quad \circ \end{array} \right].$$

(3) If $\delta = 0$, then it follows from Example 4.3.1(2) that

$$e_{(\square, \square)} = \begin{array}{c} \bullet \quad \circ \\ | \quad | \\ \bullet \quad \circ \end{array}, \quad e_{(\square, \square)}^{(1)} = \begin{array}{c} \bullet \quad \bullet \quad \circ \\ | \quad | \quad \diagup \\ \bullet \quad \bullet \quad \circ \end{array}, \quad e_{(\square, \square)}^{(2)} = \begin{array}{c} \bullet \quad \bullet \quad \bullet \quad \circ \\ | \quad | \quad | \quad \diagup \\ \bullet \quad \bullet \quad \bullet \quad \circ \end{array}, \dots$$

On the other hand, if $\delta \neq 0$, then by Example 4.3.1(2),

$$e_{(\square, \square)}^{(1)} = \begin{array}{c} \bullet \quad \bullet \quad \circ \\ | \quad | \quad \diagup \\ \bullet \quad \bullet \quad \circ \end{array} - \frac{1}{\delta} \begin{array}{c} \bullet \quad \bullet \quad \circ \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \quad \circ \end{array},$$

$$e_{(\square, \square)}^{(2)} = \begin{array}{c} \bullet \quad \bullet \quad \bullet \quad \circ \\ | \quad | \quad | \quad \diagup \\ \bullet \quad \bullet \quad \bullet \quad \circ \end{array} - \frac{1}{\delta} \begin{array}{c} \bullet \quad \bullet \quad \bullet \quad \circ \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \quad \bullet \quad \circ \end{array}.$$

It follows from (7) that $e_\lambda^{(i)}$ is an idempotent whenever it is defined. Hence, the image of $e_\lambda^{(i)}$ is an object in $\underline{\text{Rep}}(GL_\delta)$.

Proposition 4.4.2. *Given a bipartition λ , the objects $\text{im } e_\lambda$ and $\text{im } e_\lambda^{(i)}$ are isomorphic in $\underline{\text{Rep}}(GL_\delta)$ whenever $e_\lambda^{(i)}$ is defined.*

Proof. It suffices to show $\text{im } e_\lambda^{(i-1)} \cong \text{im } e_\lambda^{(i)}$ whenever $i > 0$. Using (7) it is easy to check that the following table lists mutually inverse morphisms between $\text{im } e_\lambda^{(i-1)}$ and $\text{im } e_\lambda^{(i)}$ in all desired cases:

$\text{im } e_\lambda^{(i-1)} \rightarrow \text{im } e_\lambda^{(i)}$	$\text{im } e_\lambda^{(i)} \rightarrow \text{im } e_\lambda^{(i-1)}$	case:
$\psi_{r+i-1, s+i-1} e_\lambda^{(i-1)}$	$e_\lambda^{(i-1)} \phi_{r+i-1, s+i-1}$	$s > 0$
$\psi_{r+i-1, s+i-1} e_\lambda^{(i-1)}$	$e_\lambda^{(i-1)} \hat{\phi}_{r+i-1, s+i-1}$	$s = 0, r > 0$
$\psi_{i-1, i-1} e_\lambda^{(i-1)}$	$\frac{1}{\delta} e_\lambda^{(i-1)} \hat{\psi}_{i-1, i-1}$	$\lambda = (\emptyset, \emptyset), \delta \neq 0$

□

Corollary 4.4.3. *$e_\lambda^{(i)}$ is absolutely primitive whenever it is defined.*

Proof. By Proposition 4.3.2, e_λ is absolutely primitive. Thus, by Proposition 2.7.1(1), $\text{im } e_\lambda \cong \text{im } e_\lambda^{(i)}$ is absolutely indecomposable, which implies $e_\lambda^{(i)}$ is absolutely primitive. □

4.5. Classification of primitive idempotents in walled Brauer algebras.

We are now in position to state the classification of conjugacy classes of primitive idempotents in walled Brauer algebras. The next theorem is merely a translation of the classification of simple modules for walled Brauer algebras [12, Theorem 2.7] to the language of primitive idempotents (see for instance [3]).

Theorem 4.5.1. (1) If $r \neq s$ or $\delta \neq 0$, then $\{e_\lambda^{(i)} \mid \lambda \vdash (r-i, s-i), 0 \leq i \leq \min(r, s)\}$ is a complete set of pairwise non-conjugate primitive idempotents in $B_{r,s}$.

(2) If $\delta = 0$ and $r > 0$, then $\{e_\lambda^{(i)} \mid \lambda \vdash (r-i, r-i), 0 \leq i < r\}$ is a complete set of pairwise non-conjugate primitive idempotents in $B_{r,r}$.

Corollary 4.5.2. Primitive idempotents in walled Brauer algebras are absolutely primitive.

Proof. This follows from Theorem 4.5.1 along with Corollary 4.4.3. $\square$

4.6. Classification of indecomposable objects in $\underline{\text{Rep}}(GL_\delta)$. Given a bipartition λ , let $L(\lambda)$ denote the image of e_λ in $\underline{\text{Rep}}(GL_\delta)$. Since the primitive idempotent e_λ is only defined up to conjugation, $L(\lambda)$ is an indecomposable object in $\underline{\text{Rep}}(GL_\delta)$ which is defined up to isomorphism. The following proposition concerning $L(\lambda)$ will be used to prove our upcoming classification of indecomposable objects. Since the result will be used later in the paper, we record it separately here:

Proposition 4.6.1. If λ and μ are bipartitions with $\text{Hom}(L(\lambda), L(\mu)) \neq 0$, then $|\lambda^\bullet| + |\mu^\circ| = |\lambda^\circ| + |\mu^\bullet|$.

Proof. $\text{Hom}(L(\lambda), L(\mu)) = \text{Hom}(\text{im } e_\lambda, \text{im } e_\mu) \subset \text{Hom}(w_{|\lambda^\bullet|, |\lambda^\circ|}, w_{|\mu^\bullet|, |\mu^\circ|})$, hence the proposition follows from Remark 3.1.2. $\square$

Now we are ready to classify indecomposable objects in $\underline{\text{Rep}}(GL_\delta)$.

Theorem 4.6.2. The assignment $\lambda \mapsto L(\lambda)$ induces a bijection

$$\left\{ \begin{array}{c} \text{bipartitions of} \\ \text{arbitrary size} \end{array} \right\} \xrightarrow{\text{bij.}} \left\{ \begin{array}{c} \text{non-zero indecomposable objects in} \\ \underline{\text{Rep}}(GL_\delta), \text{ up to isomorphism} \end{array} \right\}$$

Proof. By Proposition 2.7.1(1) every indecomposable object in $\underline{\text{Rep}}(GL_\delta)$ is isomorphic to the image of a primitive idempotent endomorphism in $\underline{\text{Rep}}_0(GL_\delta)$. Since every object in $\underline{\text{Rep}}_0(GL_\delta)$ is isomorphic to $w_{r,s}$ for some $r, s \geq 0$, it follows that each indecomposable object in $\underline{\text{Rep}}(GL_\delta)$ is isomorphic to the image of a primitive idempotent in $B_{r,s}$ for some $r, s \geq 0$. Hence, by Theorem 4.5.1 and Proposition 4.4.2 the assignment $\lambda \mapsto L(\lambda)$ is surjective.

Now suppose $\lambda \vdash (r, s)$ and $\mu \vdash (r', s')$ are two bipartitions with $L(\lambda) \cong L(\mu)$. For convenience, assume $r \geq r'$ so that $(r, s) = (r' + i, s' + i)$ for some integer $i \geq 0$ (Proposition 4.6.1). If $\mu = (\emptyset, \emptyset)$, then the existence of an isomorphism $L((\emptyset, \emptyset)) \rightarrow L(\lambda)$ implies that the composition map

$$(8) \quad \text{Hom}(w_{r,s}, \mathbf{1}) \times \text{Hom}(\mathbf{1}, w_{r,s}) \rightarrow B_{0,0}$$

is non-zero. If $\delta = 0$, then the map (8) is necessarily zero unless $r = s = 0$. Hence, if $\mu = (\emptyset, \emptyset)$ and $\delta = 0$, then $\lambda = (\emptyset, \emptyset)$ too. Now assume $\mu \neq (\emptyset, \emptyset)$ or $\delta \neq 0$ so that $e_\mu^{(i)}$ is defined. By Proposition 4.4.2, $\text{im } e_\mu^{(i)} \cong \text{im } e_\mu \cong \text{im } e_\lambda$, which implies $e_\mu^{(i)}$ and e_λ are conjugate idempotents in $B_{r,s}$ (see Proposition 2.7.1(3)). By Theorem 4.5.1, $\lambda = \mu$. Thus the assignment $\lambda \mapsto L(\lambda)$ is injective. $\square$

Remark 4.6.3. Instead of relying on [12], one can prove Theorem 4.6.2 with straightforward modifications of the proof of [10, Theorem 3.7].

We end this subsection with a couple propositions concerning $L(\lambda)$ which will be useful later.

Proposition 4.6.4. *Given a bipartition $\lambda \vdash (r, s)$,*

$$L((\lambda^\bullet, \emptyset)) \otimes L((\emptyset, \lambda^\circ)) = \text{im } z_\lambda = L(\lambda) \oplus L(\mu^{(1)}) \oplus \cdots \oplus L(\mu^{(k)})$$

for some bipartitions $\mu^{(1)}, \dots, \mu^{(k)}$ which have the property $\mu^{(j)} \vdash (r - i_j, s - i_j)$ with $0 < i_j \leq \min(r, s)$ for all $j = 1, \dots, k$.

Proof. First, using Example 4.3.1(1) we have $e_{(\lambda^\bullet, \emptyset)} \otimes e_{(\emptyset, \lambda^\circ)} = z_{\lambda^\bullet}^\bullet \otimes z_{\lambda^\circ}^\circ = z_\lambda$ which implies the left equality. For the right equality, notice that by the definition of e_λ we can write $z_\lambda = e_\lambda + e_1 + \cdots + e_k$ where $e_\lambda, e_1, \dots, e_k$ are mutually orthogonal primitive idempotents in $B_{r,s}$. Moreover, $\pi(e_j) = 0$ for all $j = 1, \dots, k$. By Theorem 4.5.1, there exists a bipartition $\mu^{(j)} \vdash (r - i_j, s - i_j)$ for some $0 \leq i_j \leq \min(r, s)$ such that e_j is conjugate to $e_{\mu^{(j)}}^{(i_j)}$ for all $j = 1, \dots, k$. It follows that $\pi(e_{\mu^{(j)}}^{(i_j)}) = 0$, which implies $i_j \neq 0$ for each $j = 1, \dots, k$. Finally, by Propositions 2.7.1(2) and 4.4.2 we are done. $\square$

The following example illustrates Proposition 4.6.4.

Example 4.6.5. (1) Assume $\delta \neq 0$. Then by Examples 4.3.1(2) and 4.4.1(1) we have the following orthogonal decomposition of $z_{(\square, \square)} = \text{id}_{\bullet \circ}$ into primitive idempotents: $z_{(\square, \square)} = e_{(\square, \square)} + e_{(\emptyset, \emptyset)}^{(1)}$. Hence $\bullet \circ = L((\square, \square)) \oplus L((\emptyset, \emptyset))$ in $\underline{\text{Rep}}(GL_\delta)$.

(2) When $\delta = 0$, $z_{(\square, \square)} = e_{(\square, \square)}$, by Example 4.3.1(2). Hence $\bullet \circ = L((\square, \square))$ in $\underline{\text{Rep}}(GL_0)$.

Proposition 4.6.6. $L(\lambda)^* = L(\lambda^*)$.

Proof. Given $\sigma \in \Sigma_r$, it is easy to check that the dual morphisms $(\sigma^\bullet)^* = (\sigma^\circ)^{-1}$ and $(\sigma^\circ)^* = (\sigma^\bullet)^{-1}$. Given a partition $\alpha \vdash r$, the idempotent $z_\alpha \in \mathbb{K}\Sigma_r$ is invariant under the operation $\mathbb{K}\Sigma_r \rightarrow \mathbb{K}\Sigma_r, \sigma \mapsto \sigma^{-1}$. Hence, $z_\lambda^* = (z_\lambda^\bullet \otimes z_{\lambda^\circ}^\circ)^* = z_{\lambda^\circ}^\circ \otimes z_{\lambda^\bullet}^\bullet$. Hence, up to isomorphism we have

$$(\text{im } z_\lambda)^* = \text{im}(z_\lambda^*) = \text{im}(z_{\lambda^\circ}^\circ) \otimes \text{im}(z_{\lambda^\bullet}^\bullet) = \text{im}(z_{\lambda^\circ}^\circ) \otimes \text{im}(z_{\lambda^\bullet}^\bullet) = \text{im}(z_{\lambda^*}).$$

The result now follows from Proposition 4.6.4 after inducting on $|\lambda|$. $\square$

4.7. Indecomposable summands of mixed tensor powers. Let V denote a δ -dimensional object of a tensor category $\mathcal{T}$, let V^* denote a dual for V in $\mathcal{T}$, and write

$$T(r, s) = T_{V, V^*}(r, s) = V^{\otimes r} \otimes V^{*\otimes s}$$

for the *mixed tensor power* of V . The following theorem gives a useful criterion for the fullness of the functor

$$F : \underline{\text{Rep}}(GL_\delta) \rightarrow \mathcal{T}, \quad F(\bullet) \mapsto V$$

defined by Proposition 3.5.1, and moreover, shows that any indecomposable summand of a $T(r, s)$ is isomorphic to the image under F of an indecomposable object from $\underline{\text{Rep}}(GL_\delta)$. This will be later applied in §5 and §8, where V will denote the natural representations of the general linear group and the general linear supergroup, respectively.

Theorem 4.7.1. *Suppose that the $\mathbb{K}$ -algebra maps $\mathbb{K}\Sigma_p \rightarrow \text{End}_{\mathcal{T}}(V^{\otimes p})$ defined by the symmetric braiding of $\mathcal{T}$ are surjective, and $\text{Hom}_{\mathcal{T}}(T(r, s), T(r', s')) = 0$ whenever $r + s' \neq r' + s$. Then:*

(1) *F is full.*

(2) *$\Lambda = \{F(L(\lambda)) \mid \lambda \text{ is a bipartition, } F(L(\lambda)) \neq 0\}$ is a complete set of indecomposable summands of the mixed tensor powers $T(r, s)$. Moreover, the members of Λ are pairwise non-isomorphic.*

Proof. To prove the first part, it suffices to show the restriction of F to $\underline{\text{Rep}}_0(GL_\delta)$ is full. Since every object in $\underline{\text{Rep}}_0(GL_\delta)$ is isomorphic (by braidings) to a word of the form $w_{r,s}$ for some r, s , and the restriction of F to $\underline{\text{Rep}}_0(GL_\delta)$ is strict (hence it preserves braidings) it suffices to show

$$(9) \quad F : \text{Hom}_{\underline{\text{Rep}}_0(GL_\delta)}(w_{r,s}, w_{r',s'}) \rightarrow \text{Hom}_{\mathcal{T}}(T(r, s), T(r', s'))$$

is surjective for every r, r', s, s' . By hypothesis, we can assume $r + s' = r' + s$. Consider the diagram

$$(10) \quad \begin{array}{ccc} \text{Hom}_{\underline{\text{Rep}}_0(GL_\delta)}(w_{r,s}, w_{r',s'}) & \longrightarrow & \text{End}_{\underline{\text{Rep}}_0(GL_\delta)}(\bullet^{\otimes r+s'}) \\ \downarrow F & & \downarrow F \\ \text{Hom}_{\mathcal{T}}(T(r, s), T(r', s')) & \longrightarrow & \text{End}_{\mathcal{T}}(V^{\otimes r+s'}) \end{array}$$

where the horizontal maps are given by

$$(11) \quad f \mapsto (\text{id}_{\bullet}^{\otimes r'} \otimes \text{ev}_{w_{s',0}} \otimes \text{id}_{\bullet}^{\otimes s})(f \otimes c_{w_{s,0}, w_{s',0}})(\text{id}_{\bullet}^{\otimes r} \otimes \text{coev}_{w_{0,s}} \otimes \text{id}_{\bullet}^{\otimes s'})$$

and

$$(12) \quad f \mapsto (\text{id}_V^{\otimes r'} \otimes \text{ev}_{T(s',0)} \otimes \text{id}_V^{\otimes s})(f \otimes c_{T(s,0), T(s',0)})(\text{id}_V^{\otimes r} \otimes \text{coev}_{T(0,s)} \otimes \text{id}_V^{\otimes s'}).$$

Since the restriction of F to $\underline{\text{Rep}}_0(GL_\delta)$ is a strict tensor functor, the diagram (10) commutes. Moreover, the maps (11) and (12) are $\mathbb{K}$ -vector space isomorphisms with inverses

$$f \mapsto (\text{id}_{\bullet}^{\otimes r'} \otimes \text{ev}_{w_{0,s}} \otimes \text{id}_{\circ}^{\otimes s'})(f \otimes c_{w_{0,s'}, w_{0,s}})(\text{id}_{\bullet}^{\otimes r} \otimes \text{coev}_{w_{s',0}} \otimes \text{id}_{\circ}^{\otimes s})$$

and

$$f \mapsto (\text{id}_V^{\otimes r'} \otimes \text{ev}_{T(0,s)} \otimes \text{id}_{V^*}^{\otimes s'})(f \otimes c_{T(0,s'), T(0,s)})(\text{id}_V^{\otimes r} \otimes \text{coev}_{T(s',0)} \otimes \text{id}_{V^*}^{\otimes s}),$$

respectively (this is easily verified using diagram calculus for tensor categories [23]). Since the rightmost vertical map is surjective, by hypothesis, we are done.

We now prove the second part. A summand W of $T(r, s)$ is the image of an idempotent in $\text{End}_{\mathcal{T}}(T(r, s))$. By part (1), such an idempotent has a pre-image $e \in B_{r,s}$ under F . Write $\text{im } e = L(\lambda^{(1)}) \oplus \cdots \oplus L(\lambda^{(k)})$ in $\underline{\text{Rep}}(GL_\delta)$ for some bipartitions $\lambda^{(i)}$ (see Theorem 4.6.2). Then $W = F(L(\lambda^{(1)})) \oplus \cdots \oplus F(L(\lambda^{(k)}))$ in $\mathcal{T}$, and by Proposition 2.7.4 the $F(L(\lambda^{(i)}))$ are indecomposable. Thus if W is indecomposable, then $W = F(L(\lambda^{(i)}))$ for some i . $\square$

4.8. Generic semisimplicity of $\underline{\text{Rep}}(GL_\delta)$. We close this section with the following theorem which tells us exactly when the category $\underline{\text{Rep}}(GL_\delta)$ is semisimple⁵ (compare with [14, Théorème 10.4]).

Theorem 4.8.1. *$\underline{\text{Rep}}(GL_\delta)$ is semisimple if and only if δ is not an integer.*

Proof. The semisimplicity of the walled Brauer algebras is completely determined in [12, Theorem 6.3]. In particular, if $\delta \in \mathbb{Z}$, then $B_{r,s}(\delta)$ is not semisimple for some r and s . In this case, using Theorem 4.5.1, there exist distinct bipartitions $\lambda \vdash (r-i, s-i)$ and $\mu \vdash (r-j, s-j)$ for some $i, j \geq 0$ with $e_\mu^{(j)} B_{r,s} e_\lambda^{(i)} \neq 0$. Since $e_\mu^{(j)} B_{r,s} e_\lambda^{(i)} = \text{Hom}(\text{im } e_\lambda^{(i)}, \text{im } e_\mu^{(j)}) = \text{Hom}(L(\lambda), L(\mu))$, we are done in case $\delta \in \mathbb{Z}$.

Now assume $\delta \notin \mathbb{Z}$, and let λ and μ be bipartitions with $\text{Hom}(L(\lambda), L(\mu)) \neq 0$. If $\lambda \vdash (r, s)$, then $\mu \vdash (r-i, s-i)$ for some i (Proposition 4.6.1). We proceed under the assumption $i \geq 0$, the case $i \leq 0$ being dual, by Proposition 4.6.6. Then

$$(13) \quad 0 \neq \text{Hom}(L(\lambda), L(\mu)) = e_\mu^{(i)} B_{r,s} e_\lambda.$$

Since $\delta \notin \mathbb{Z}$, $B_{r,s}$ is a semisimple algebra, hence (13) can only be true if e_λ and $e_\mu^{(i)}$ are conjugate, which implies $\lambda = \mu$ (Theorem 4.5.1). Moreover, the semisimplicity of $B_{r,s}$ implies $\text{End}(L(\lambda)) = e_\lambda B_{r,s} e_\lambda$ is indeed a division algebra. $\square$

5. CONNECTION TO REPRESENTATIONS OF THE GENERAL LINEAR GROUP

Fix a non-negative integer d and consider the category $\text{Rep}(GL_d)$ of finite dimensional representations of the general linear group GL_d over $\mathbb{K}$. In this short section we describe how the categories $\underline{\text{Rep}}(GL_\delta)$ and $\text{Rep}(GL_d)$ are related. One can view this section as a preview of §8 where we show how $\underline{\text{Rep}}(GL_\delta)$ is related to representations of general linear supergroups. However, we will see in §7 that formulas for decomposing tensor products in $\underline{\text{Rep}}(GL_\delta)$ can be obtained by interpolating decomposition formulas in $\text{Rep}(GL_d)$; hence this preview of §8 is also important to the structure of the paper.

5.1. The category $\text{Rep}(GL_d)$. Let $GL_d = GL(d, \mathbb{K})$ denote the general linear group, that is, the group of invertible $d \times d$ -matrices with entries from $\mathbb{K}$, and write V for the d -dimensional natural module. Let $\text{Rep}(GL_d)$ denote the category of finite-dimensional GL_d -modules. With the tensor product of GL_d -modules defined in the usual way, and $\mathbb{1}$ denoting the trivial one-dimensional module, $\text{Rep}(GL_d)$ becomes a monoidal category. Note that, in particular, we consider the monoidal category $\text{Rep}(GL_d)$ to be strict.

For any object U of $\text{Rep}(GL_d)$, let $(U^*, \text{ev}_U, \text{coev}_U)$ be the dual for U in $\text{Vect}_{\mathbb{K}}$. Recall that U^* is a GL_d -module via

$$(14) \quad (x \cdot \phi)(u) = \phi(x^{-1} \cdot u), \quad (x \in GL_d, u \in U, \phi \in U^*).$$

Then $\text{ev}_U, \text{coev}_U$ are GL_d -module maps, and so $(U^*, \text{ev}_U, \text{coev}_U)$ is a dual for U in $\text{Rep}(GL_d)$. Thus $\text{Rep}(GL_d)$ is a tensor category. It is well known that $\text{Rep}(GL_d)$ is semisimple (see, e.g. [16]). In particular, an object is simple if and only if it is indecomposable.

⁵Recall that a $\mathbb{K}$ -linear Krull Schmidt category $\mathcal{C}$ is semisimple if and only if $\text{End}_{\mathcal{C}}(L)$ is a finite $\mathbb{K}$ -dimensional division algebra for all indecomposable objects L and $\text{Hom}_{\mathcal{C}}(L, L') = 0$ for all non-isomorphic indecomposable objects L, L' .

Let ε_i denote the function that takes any matrix to the its (i, i) entry for each $i = 1, \dots, d$, and let Γ denote the set of weights. For $k \in \mathbb{Z}$, letting

$$\Gamma_k = \{\gamma = \sum_{i=1}^d \gamma_i \varepsilon_i \mid \gamma_i \in \mathbb{Z}, \gamma_1 \geq \dots \geq \gamma_d, \sum_i \gamma_i = k\},$$

we have $\Gamma = \bigsqcup_{k \in \mathbb{Z}} \Gamma_k$. Write γ_i for the coefficients of $\gamma \in \Gamma$ with respect to the ε_i . For $\gamma \in \Gamma$, let $V(\gamma)$ denote the finite-dimensional highest-weight GL_d -module of highest-weight γ . The highest-weight modules classify the indecomposable objects of $\text{Rep}(GL_d)$ up to isomorphism. That is, any indecomposable object of $\text{Rep}(GL_d)$ is isomorphic to V_γ for some $\gamma \in \Gamma$, and if $V_\gamma \cong V_{\gamma'}$ for $\gamma, \gamma' \in \Gamma$, then $\gamma = \gamma'$. The weights Γ are in bijection with bipartitions λ with $l(\lambda) \leq d$, via

$$(15) \quad wt(\lambda) = \sum_{i>0} \lambda_i^\bullet \varepsilon_i - \sum_{j>0} \lambda_j^\circ \varepsilon_{d-j+1}.$$

Writing V_λ for $V_{wt(\lambda)}$, the isomorphism classes of indecomposable objects in $\text{Rep}(GL_d)$ are thus parameterized by such bipartitions.

For $k \in \mathbb{Z}$, define a partial order $\leq$ on Γ_k by declaring

$$\gamma \leq \gamma' \iff \gamma_1 + \dots + \gamma_k \leq \gamma'_1 + \dots + \gamma'_k, \quad k = 1, \dots, d.$$

If $\gamma \in \Gamma$, then there exists k such that all weights of the highest-weight module $V(\gamma)$ belong to Γ_k (k is the rational degree of the module). Thus the set of the weights of $V(\gamma)$ are partially ordered. If γ' is a weight of $V(\gamma)$, then $\gamma' \leq \gamma$.

Proposition 5.1.1. *Let λ be a bipartition with $l(\lambda) \leq d$. Then $(V_\lambda)^* = V_{\lambda^*}$.*

Proof. Duality defines an endofunctor of $\text{Rep}(GL_d)$. Thus $(V_\lambda)^*$ is indecomposable, so $(V_\lambda)^* \cong V_\mu$ for some bipartition μ with $l(\mu) \leq d$. It follows immediately from (14) that the dual of a weight space of V_λ is a weight space of $(V_\lambda)^*$ with the weight negated. Negation inverts the partial order on weights, so $wt(\mu)$ is the lowest weight of V_λ .

Recall that the symmetric group Σ_d embeds in GL_d as the permutation matrices. Thus Σ_d acts on V_λ and hence on its set of weights. If σ denotes the longest element of Σ_d , and γ is a weight, then $(\gamma^\sigma)_i = \gamma_{d-i+1}$ for $i = 1, \dots, d$, and furthermore, $wt(\nu^*) = wt(\nu)^\sigma$ for any bipartition ν . Finally, σ is an anti-involution of the poset of weights, and so $wt(\lambda)^\sigma = wt(\mu)$ is the lowest weight of V_λ , and so $\mu = \lambda^*$. $\square$

Theorem 5.1.2 (Compare with [18, Theorem 2.4]). *Fix bipartitions $\lambda \vdash (r, s)$ and $\mu \vdash (r', s')$ such that $l(\lambda), l(\mu) \leq d$. For each bipartition ν with $l(\nu) \leq d$ let $\Gamma_{\lambda, \mu}^\nu$ be such that*

$$V_\lambda \otimes V_\mu = \bigoplus_{\nu} V_\nu^{\oplus \Gamma_{\lambda, \mu}^\nu}.$$

Then $\Gamma_{\lambda, \mu}^\nu = 0$ unless the $|\nu| \leq (r + r', s + s')$. Moreover, if $l(\lambda) + l(\mu) \leq d$, then

$$(16) \quad \Gamma_{\lambda, \mu}^\nu = \sum_{\alpha, \beta, \gamma, \theta \in \mathcal{P}} \left(\sum_{\kappa \in \mathcal{P}} LR_{\kappa, \alpha}^{\lambda^\bullet} LR_{\kappa, \beta}^{\mu^\circ} \right) \left(\sum_{\gamma \in \mathcal{P}} LR_{\gamma, \eta}^{\lambda^\circ} LR_{\gamma, \theta}^{\mu^\bullet} \right) LR_{\alpha, \theta}^{\nu^\bullet} LR_{\beta, \eta}^{\nu^\circ}$$

where $LR_{\beta, \gamma}^\alpha$'s are the Littlewood Richardson coefficients.

5.2. The functor $F_d : \underline{\text{Rep}}(GL_d) \rightarrow \text{Rep}(GL_d)$. Write $F_d : \underline{\text{Rep}}(GL_d) \rightarrow \text{Rep}(GL_d)$ for the tensor functor which sends $\bullet \mapsto V$ defined by Proposition 3.5.1. By classical Schur-Weyl duality (see [28]), the $\mathbb{K}$ -algebra map $\mathbb{K}\Sigma_p \rightarrow \text{End}_{\text{Rep}(GL_d)}(V^{\otimes p})$ defined by the symmetric braiding is surjective, for any $p \geq 0$. Write $T(r, s)$ for the mixed tensor powers of V as per section §4.7. For any $\zeta \in \mathbb{K}$, the central element $\zeta \cdot \text{id} \in GL_d$ acts on $T(r, s)$ by the scalar ζ^{r-s} , so $\text{Hom}_{\text{Rep}(GL_d)}(T(r, s), T(r', s')) = 0$ unless $r + s' = r' + s$. Thus, by Theorem 4.7.1, the functor F_d is full (this is the so-called First Fundamental Theorem of invariant theory) and any indecomposable summand of a mixed tensor powers $T(r, s)$ is isomorphic to $F_d(L(\lambda))$ for some bipartition λ .

Given a bipartition λ , we write $W(\lambda) = F_d(L(\lambda))$. In this section we will completely describe $W(\lambda)$. We start by assuming one of $\lambda^\bullet, \lambda^\circ$ is $\emptyset$.

Proposition 5.2.1. *Assume $\lambda \vdash (r, s)$ with $rs = 0$. If $l(\lambda) \leq d$, then $W(\lambda) = V_\lambda$. If $l(\lambda) > d$, then $W(\lambda) = 0$.*

Proof. First, Σ_r acts on $V^{\otimes r}$ by permuting tensors. Since strict tensor functors preserve symmetric braidings, this action coincides with $F_d : \mathbb{K}\Sigma_r \rightarrow \text{End}(V^{\otimes r})$. Hence if $r = 0$, then, by Example 4.3.1(1), $W(\lambda)$ is the image of the idempotent $z_{\lambda^\bullet} \in \mathbb{K}\Sigma_r$ acting on $V^{\otimes r}$. This is precisely Weyl's construction of V_λ (see for example [16]). The case $s = 0$ follows from the case $r = 0$ using Propositions 4.6.6 and 5.1.1. $\square$

In fact, Proposition 5.2.1 holds without the assumption that $rs = 0$.

Theorem 5.2.2. *Suppose λ is an arbitrary bipartition. Then*

$$W(\lambda) = \begin{cases} V_\lambda & \text{if } l(\lambda) \leq d, \\ 0 & \text{if } l(\lambda) > d. \end{cases}$$

Proof. We induct on the size of λ . The base case $\lambda = (\emptyset, \emptyset)$ is clear, so assume $\lambda \vdash (r, s)$ with $rs \neq 0$. By Example 4.3.1(1) we have $z_\lambda = e_{(\lambda^\bullet, \emptyset)} \otimes e_{(\emptyset, \lambda^\circ)}$ from which it follows that

$$(17) \quad F_d(\text{im } z_\lambda) = W((\lambda^\bullet, \emptyset)) \otimes W((\emptyset, \lambda^\circ)).$$

Also, by Proposition 4.6.4,

$$(18) \quad F_d(\text{im } z_\lambda) = W(\lambda) \oplus W(\mu^{(1)}) \oplus \cdots \oplus W(\mu^{(k)})$$

where $\mu^{(1)}, \dots, \mu^{(k)}$ are bipartitions whose sizes are strictly smaller than (r, s) .

Assume $l(\lambda) \leq d$. Then by (17) and Proposition 5.2.1, $F_d(\text{im } z_\lambda)$ has a highest weight vector with weight $wt(\lambda)$. On the other hand, since $l(\lambda) \leq d$ there is no cancelation in (15). Hence, by induction, $W(\mu^{(j)})$ does not have a highest weight vector of weight $wt(\lambda)$ for any $j = 1, \dots, k$. Thus, by (18), $W(\lambda)$ has a highest weight vector with weight $wt(\lambda)$. Finally, by Proposition 2.7.4, $W(\lambda)$ is simple.

Now assume $l(\lambda) > d$. If either $l(\lambda^\bullet)$ or $l(\lambda^\circ)$ is greater than d we are done by Proposition 5.2.1, hence we can assume $l(\lambda^\bullet), l(\lambda^\circ) \leq d$. By Theorem 5.1.2, the highest weights in (17) are all of the form $wt(\nu)$ with $|\nu| \leq (r, s)$. Thus, by induction, $F_d(z_\lambda)$ decomposes as a direct sum of $W(\nu)$'s with $|\nu| \leq (r, s)$. Hence, by (18) and Proposition 2.7.4, $W(\lambda) = 0$. $\square$

6. THE LIFTING MAP

As before, we fix $\delta \in \mathbb{K}$. Let t be an indeterminate. In this section we construct a ring isomorphism (called the *lifting map*) between the Grothendieck rings of $\underline{\text{Rep}}(GL_\delta)$ and $\underline{\text{Rep}}(GL_t)$. The definition of the lifting map is not an explicit one; however, we show in §6.4 that values of the lifting map can be computed using combinatorics of certain diagrams introduced by Brundan and Stroppel. We will see later (§7) that the lifting map will play a crucial role in our ability to decompose tensor products in $\underline{\text{Rep}}(GL_\delta)$. We begin by fixing some notation for the Grothendieck rings mentioned above.

6.1. The rings R_δ and R_t . Write $\mathbb{K}(t)$, $\mathbb{K}[[t - \delta]]$, and $\mathbb{K}((t - \delta))$ for the field of fractions in t , ring of power series in $t - \delta$, and the field of Laurent series⁶ in $t - \delta$, respectively. Then we have the following categories with their corresponding additive Grothendieck rings (see §2.8):

category		additive Grothendieck ring	bilinear form
$\underline{\text{Rep}}(GL_\delta)$	over $\mathbb{K}$	R_δ	$(-, -)_\delta$
$\underline{\text{Rep}}(GL_t)$	over $\mathbb{K}(t)$	R_t	$(-, -)_t$
$\underline{\text{Rep}}(GL_t)$	over $\mathbb{K}((t - \delta))$	$R_{t,\delta}$	$(-, -)_{t,\delta}$

By Theorem 4.6.2 we can identify the elements of R_δ , R_t , and $R_{t,\delta}$ with formal $\mathbb{Z}$ -linear combinations of bipartitions, and we will do so for the rest of the paper. In particular, the rings R_δ , R_t , and $R_{t,\delta}$ are clearly isomorphic as abelian groups. However, the multiplication in these rings depends on the parameter:

Example 6.1.1. It is always true that $e_{(\square, \emptyset)} = \text{id}_\bullet$ and $e_{(\emptyset, \square)} = \text{id}_\circ$, which implies $L((\square, \emptyset)) = \bullet$ and $L((\emptyset, \square)) = \circ$. Hence $L((\square, \emptyset)) \otimes L((\emptyset, \square)) = \bullet \circ$ always. Thus, it follows from example 4.3.1(2) that

$$(\square, \emptyset)(\emptyset, \square) = (\square, \square) \in R_0, \quad \text{whereas} \quad (\square, \emptyset)(\emptyset, \square) = (\square, \square) + (\emptyset, \emptyset) \in R_t.$$

The next proposition shows that although the multiplication of bipartitions in R_δ and R_t may differ, the rings R_t and $R_{t,\delta}$ can be identified regardless of δ .

Proposition 6.1.2. (1) *The $\mathbb{Z}$ -linear map $R_t \rightarrow R_{t,\delta}$ with $\lambda \mapsto \lambda$ for each bipartition λ is a ring isomorphism.*

$$(2) \quad (\lambda, \mu)_t = (\lambda, \mu)_{t,\delta} = \begin{cases} 1 & \text{if } \lambda = \mu, \\ 0 & \text{if } \lambda \neq \mu. \end{cases}$$

Proof. (1) Suppose

$$(19) \quad e_\lambda \otimes e_\mu = e_1 + \cdots + e_k$$

is a decomposition of $e_\lambda \otimes e_\mu$ into mutually orthogonal primitive idempotents over $\mathbb{K}(t)$. Then $\lambda\mu = \sum_\nu a_\nu \nu \in R_t$ where a_ν is the number of summands in (19) correspond to the bipartition ν . By Corollary 4.5.2, viewing (19) over the larger field $\mathbb{K}((t - \delta)) \supset \mathbb{K}(t)$ still gives an orthogonal decomposition of $e_\lambda \otimes e_\mu$ into primitive idempotents, hence $\lambda\mu = \sum_\nu a_\nu \nu \in R_{t,\delta}$ also.

(2) By Corollary 4.5.2, we can work over the algebraic closure of $\mathbb{K}(t)$ (resp. $\mathbb{K}((t - \delta))$) to compute $(\lambda, \mu)_t$ (resp. $(\lambda, \mu)_{t,\delta}$). The result now follows from the fact that $\underline{\text{Rep}}(GL_t)$ is semisimple over any field containing the indeterminate t (see Theorem 4.8.1). $\square$

⁶In other words, $\mathbb{K}((t - \delta))$ is the field of fractions of $\mathbb{K}[[t - \delta]]$.

With Proposition 6.1.2 in mind, for the rest of the paper we will identify $R_{t,\delta}$ with R_t for every δ and write R_t for both.

6.2. The ring map $\text{lift}_\delta : R_\delta \rightarrow R_t$. Fix a bipartition $\lambda \vdash (r, s)$ and consider the idempotent $e_\lambda \in \mathbb{K}B_{r,s}(\delta)$. We can lift e_λ to an idempotent $\tilde{e} \in \mathbb{K}((t - \delta))B_{r,s}(t)$, i.e., $\tilde{e}$ is of the form $\tilde{e} = \sum_X a_X X$ with $a_X \in \mathbb{K}[[t - \delta]]$ for all $(w_{r,s}, w_{r,s})$ -diagrams X , and $\tilde{e}|_{t=\delta} = e$ (see [10, Theorem A.2]). Now, given another bipartition μ , let $D_{\lambda,\mu} = D_{\lambda,\mu}(\delta)$ denote the number of times $L(\mu)$ occurs in a decomposition of $\text{im}(\tilde{e})$ into a direct sum of indecomposables in $\underline{\text{Rep}}(GL_t)$ over $\mathbb{K}((t - \delta))$. One can show that $D_{\lambda,\mu}$ does not depend on the choice of representative for e_λ or on the choice of $\tilde{e}$ (compare with [10, Theorem 3.9]). Now, let $\text{lift}_\delta : R_\delta \rightarrow R_t$ be the $\mathbb{Z}$ -linear map defined on bipartitions by

$$\text{lift}_\delta(\lambda) = \sum_{\mu} D_{\lambda,\mu} \mu.$$

Example 6.2.1. If $\lambda \vdash (r, 0)$, then, by Example 4.3.1(1), $e_\lambda = z_{\lambda^\bullet}^\bullet \in \mathbb{K}B_{r,0}(\delta)$. Since $z_{\lambda^\bullet}^\bullet$ does not depend on δ , it can be lifted to $z_{\lambda^\bullet}^\bullet = e_\lambda \in \mathbb{K}((t))B_{r,0}(t)$. Hence, $\text{lift}_\delta(\lambda) = \lambda$ for all $\lambda \vdash (r, 0)$, $\delta \in \mathbb{K}$. Similarly, $\text{lift}_\delta(\lambda) = \lambda$ whenever $\lambda \vdash (0, s)$.

Example 6.2.2. (1) Assume $\delta = 0$. By Example 4.3.1(2), $e_{(\square, \square)} = \text{id}_{\bullet \circ} \in \mathbb{K}B_{1,1}(0)$ which lifts to $\text{id}_{\bullet \circ} \in \mathbb{K}((t))B_{1,1}(t)$. By Example 4.6.5(1), $\bullet \circ = L((\square, \square)) \oplus L((\emptyset, \emptyset))$ in $\underline{\text{Rep}}(GL_t)$. Thus $\text{lift}_0((\square, \square)) = (\square, \square) + (\emptyset, \emptyset)$.

(2) Assume $\delta \neq 0$. An explicit expression for $e_{(\square, \square)} \in \mathbb{K}B_{1,1}(\delta)$ is given in Example 4.3.1(2). Since $\frac{1}{t} = \sum_{n=0}^{\infty} \frac{(-1)^n}{\delta^{n+1}} (t - \delta)^n \in \mathbb{K}[[t - \delta]]$, a lift of $e_{(\square, \square)}$ is obtained by replacing δ with t in that expression. Hence, $\text{lift}_\delta((\square, \square)) = (\square, \square)$.

The following theorem lists properties of lift_δ which are very useful for this paper.

Theorem 6.2.3. (1) $\text{lift}_\delta : R_\delta \rightarrow R_t$ is a ring isomorphism for every $\delta \in \mathbb{K}$.

(2) $D_{\lambda,\lambda} = 1$ for all λ . Moreover, $D_{\lambda,\mu} = 0$ unless $\mu = \lambda$ or $\mu \vdash (|\lambda^\bullet| - i, |\lambda^\circ| - i)$ for some $i > 0$.

(3) Fix a bipartition λ . $\text{lift}_\delta(\lambda) = \lambda$ for all but finitely many $\delta \in \mathbb{K}$.

(4) $(\text{lift}_\delta(x), \text{lift}_\delta(y))_t = (x, y)_\delta$ for all $x, y \in R_\delta$.

For a proof of Theorem 6.2.3 we refer the reader to [10, Proposition 3.12] where the analogous statements are proved for $\underline{\text{Rep}}(S_t)$.

An important consequence of Theorem 6.2.3 is the following:

Corollary 6.2.4. $(\lambda, \mu)_\delta = \sum_{\nu} D_{\lambda,\nu} D_{\mu,\nu}$ for all bipartitions λ and μ .

Proof.

$$\begin{aligned} (\lambda, \mu)_\delta &= (\sum_{\nu} D_{\lambda,\nu} \nu, \sum_{\nu'} D_{\mu,\nu'} \nu')_t \quad (\text{Theorem 6.2.3(4)}) \\ &= \sum_{\nu, \nu'} D_{\lambda,\nu} D_{\mu,\nu'} (\nu, \nu')_t \\ &= \sum_{\nu} D_{\lambda,\nu} D_{\mu,\nu} \quad (\text{Proposition 6.1.2(2)}). \end{aligned}$$

□

6.3. The diagrams of Brundan and Stroppel. We will soon show that $\text{lift}_\delta(\lambda)$ can be computed explicitly using certain diagrams introduced by Brundan and Stroppel [5]–[9]. In this subsection we introduce these diagrams and give a few examples.

As usual, we fix $\delta \in \mathbb{K}$. Given a bipartition λ , set

$$\begin{aligned} I_\wedge(\lambda) &= \{\lambda_1^\bullet, \lambda_2^\bullet - 1, \lambda_3^\bullet - 2, \dots\}, \\ I_\vee(\lambda, \delta) &= \{1 - \delta - \lambda_1^\circ, 2 - \delta - \lambda_2^\circ, 3 - \delta - \lambda_3^\circ, \dots\}. \end{aligned}$$

Now, let $x_\lambda = x_\lambda(\delta)$ be the diagram obtained by labeling the integer vertices on the number line according to the following rule: label the i th vertex by

$$\begin{cases} \bigcirc & \text{if } i \notin I_\wedge(\lambda) \cup I_\vee(\lambda, \delta), \\ \wedge & \text{if } i \in I_\wedge(\lambda) \setminus I_\vee(\lambda, \delta), \\ \vee & \text{if } i \in I_\vee(\lambda, \delta) \setminus I_\wedge(\lambda), \\ \times & \text{if } i \in I_\wedge(\lambda) \cap I_\vee(\lambda, \delta). \end{cases}$$

For example,

$$\begin{aligned} x_{(\emptyset, \emptyset)}(0) &= \cdots \quad \begin{array}{cccccccccccccccc} & \wedge & \wedge & \wedge & \wedge & \wedge & \wedge & \vee & \vee & \vee & \vee & \vee & \vee & \vee & \vee & \cdots \\ & -5 & -4 & -3 & -2 & -1 & 0 & 1 & 2 & 3 & 4 & 5 & 6 & & & \end{array}, \\ x_{((3,1),(2))}(-1) &= \cdots \quad \begin{array}{cccccccccccccccc} & \wedge & \wedge & \wedge & \wedge & \bigcirc & \times & \bigcirc & \bigcirc & \times & \vee & \vee & \vee & \vee & \cdots \\ & -5 & -4 & -3 & -2 & -1 & 0 & 1 & 2 & 3 & 4 & 5 & 6 & & \end{array}, \\ x_{((2^4),(2^5))}(3) &= \cdots \quad \begin{array}{cccccccccccccccc} & \wedge & \times & \vee & \vee & \times & \times & \wedge & \wedge & \vee & \vee & \vee & \vee & \vee & \cdots \\ & -5 & -4 & -3 & -2 & -1 & 0 & 1 & 2 & 3 & 4 & 5 & 6 & & \end{array}, \\ x_{((2^2,1),(5,2))}(\sqrt{2}) &= \cdots \quad \begin{array}{cccccccccccccccc} & \wedge & \wedge & \wedge & \bigcirc & \wedge & \bigcirc & \wedge & \wedge & \bigcirc & \bigcirc & \bigcirc & \bigcirc & \cdots \\ & -5 & -4 & -3 & -2 & -1 & 0 & 1 & 2 & 3 & 4 & 5 & 6 & \end{array}. \end{aligned}$$

Remark 6.3.1. Notice that the integer i in x_λ is labelled $\wedge$ for $i \leq 0$. Moreover, if $\delta \in \mathbb{Z}$ (resp. $\delta \notin \mathbb{Z}$), then i is labelled by $\vee$ (resp. $\bigcirc$) for $i \geq 0$. In fact, it is not difficult to show that when $\delta \in \mathbb{Z}$ there is a bijection between the set of all bipartitions and the set of all diagrams with (1) i labelled $\wedge$ for $i \leq 0$; (2) i labelled $\vee$ for $i \geq 0$; and (3) the number of $\times$'s minus the number of $\bigcirc$'s equal to δ .

Next, we construct the *cap diagram* $c_\lambda = c_\lambda(\delta)$ in the following recursive manner:

Step 0: Start with x_λ .

Step n : Draw a cap connecting vertices i and j on the number line whenever (i) $i < j$; (ii) i is labelled by $\vee$ and j is labelled by $\wedge$ in x_λ ; and (iii) each integer between i and j in x_λ is either labelled by $\bigcirc$, labelled by $\times$, or already part of a cap from an earlier step.

It follows from Remark 6.3.1 that no new caps will be added after a finite number of steps, leaving us with the cap diagram c_λ .

Example 6.3.2. If $\delta = 1$ and $\lambda = ((5^2, 4^2, 3^2), (5^3, 4, 3, 2))$, then

$$c_\lambda = \cdots \quad \begin{array}{cccccccccccccccc} & \wedge & \wedge & \vee & \vee & \vee & \vee & \times & \bigcirc & \times & \wedge & \vee & \vee & \vee & \vee & \cdots \\ & -7 & -6 & -5 & -4 & -3 & -2 & -1 & 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 \end{array}$$

Given integers $i < j$, we say (i, j) is a $\vee\wedge$ -pair in x_λ if there is a cap from j to i in c_λ . For instance, in Example 6.3.2 there are four $\vee\wedge$ -pairs: $(-5, 5)$, $(-4, 2)$, $(-3, -2)$, and $(3, 4)$. Next, given bipartitions μ and λ , we say that μ is *linked* to λ if there exists an integer $k \geq 0$ and bipartitions $\nu^{(n)}$ for $0 \leq n \leq k$ such that

(i) $\nu^{(0)} = \lambda$, (ii) $\nu^{(k)} = \mu$, and (iii) $x_{\nu^{(n)}}$ is obtained from $x_{\nu^{(n-1)}}$ by swapping the labels of some $\vee \wedge$ -pair in x_λ whenever $0 < n \leq k$. Finally, set

$$D'_{\lambda,\mu} = D'_{\lambda,\mu}(\delta) = \begin{cases} 1 & \text{if } \mu \text{ is linked to } \lambda, \\ 0 & \text{otherwise.} \end{cases}$$

Remark 6.3.3. It is shown in [11] that $D'_{\lambda,\mu}$ give decomposition numbers for walled Brauer algebras. This is easy to see when $\delta \notin \mathbb{Z}$. Indeed, when $\delta \notin \mathbb{Z}$ there are no $\vee$ labels on x_λ , so there are no $\vee \wedge$ -pairs; hence $D'_{\lambda,\mu} \neq 0$ if and only if $\lambda = \mu$.

Example 6.3.4. Fix $\delta = -1$. In this example we will compute the numbers $D'_{\lambda,\mu}$ where $\lambda = ((3, 2), (3, 1))$ and μ is arbitrary.

Now we swap labels on $\vee \wedge$ -pairs in x_λ to determine which bipartitions μ are linked to λ . The following table lists our results:

$\vee \wedge$ -pairs swapped	x_μ	μ
none		$((3, 2), (3, 1))$
$(-1, 1)$		$((3), (1^2))$
$(2, 3)$		$((2^2), (3))$
both		$((2), (1))$

Hence $D'_{\lambda,\mu} = 1$ when μ is one of the four bipartitions listed in the table above, and $D'_{\lambda,\mu} = 0$ for all other μ .

Example 6.3.5. In this example we compute $D'_{(\square,\square),\mu}(\delta)$ for all bipartitions μ and all $\delta \in \mathbb{K}$. Since $I_\wedge((\square, \square)) = \{1, -1, -2, \dots\}$, the diagram $x_{(\square,\square)}(\delta)$ has a $\vee \wedge$ -pair if and only if $0 \in I_\vee((\square, \square), \delta)$ and $1 \notin I_\vee((\square, \square), \delta)$, which occurs if and only if $\delta = 0$ since $I_\vee((\square, \square), \delta) = \{-\delta, 2 - \delta, 3 - \delta, \dots\}$. Hence, when $\delta \neq 0$ we have

$$D'_{(\square,\square),\mu}(\delta) = \begin{cases} 1 & \text{if } \mu = (\square, \square), \\ 0 & \text{otherwise.} \end{cases}$$

On the other hand,

Swapping the $\vee \wedge$ -pair $(0, 1)$ in $x_{(\square,\square)}(0)$ gives $x_{(\emptyset,\emptyset)}(0)$. Hence

$$D'_{(\square,\square),\mu}(0) = \begin{cases} 1 & \text{if } \mu = (\square, \square) \text{ or } \mu = (\emptyset, \emptyset), \\ 0 & \text{otherwise.} \end{cases}$$

In particular, $D'_{(\square, \square), \mu} = D_{(\square, \square), \mu}$ for all μ regardless of δ (see Example 6.2.2). In §6.4 we will show $D'_{\lambda, \mu} = D_{\lambda, \mu}$ always (see Corollary 6.4.2).

The following proposition describes how swapping the labels on $\vee \wedge$ -pairs affects the size of the corresponding bipartitions.

Proposition 6.3.6. *Suppose λ and μ are bipartitions and (i, j) is a $\vee \wedge$ -pair in x_λ . If x_μ is obtained from x_λ by swapping the labels of i and j , then*

$$|\mu| = (|\lambda^\bullet| + i - j, |\lambda^\circ| + i - j).$$

Proof. Set $a_k = \lambda_k^\bullet - k + 1$ and $b_k = k - \delta - \lambda_k^\circ$ for each $k > 0$ so that

$$\begin{aligned} I_\wedge(\lambda) &= \{a_1, a_2, a_3, \dots\}, \\ I_\vee(\lambda, \delta) &= \{b_1, b_2, b_3, \dots\}. \end{aligned}$$

Then $|\lambda^\bullet| = \sum_{k>0} (a_k + k - 1)$ and $|\lambda^\circ| = \sum_{k>0} (k - \delta - b_k)$. Now let $L, M \in \mathbb{Z}$ be such that $a_M = j$ and $b_L = i$. Then swapping the labels of i and j in x_λ results in x_μ with

$$\begin{aligned} I_\wedge(\mu) &= \{a_1, \dots, a_{M-1}, a_{M+1}, \dots, a_N, b_L, a_{N+1}, a_{N+2}, \dots\}, \\ I_\vee(\mu, \delta) &= \{b_1, \dots, b_{L-1}, b_{L+1}, \dots, b_{N'}, a_M, b_{N'+1}, b_{N'+2}, \dots\} \end{aligned}$$

for some N, N' . Hence

$$\begin{aligned} |\mu^\bullet| &= \sum_{\substack{0 < k < M \\ \text{or } k > N}} (a_k + k - 1) + \sum_{M < k \leq N} (a_k + (k - 1) - 1) + (b_L + N - 1) \\ &= \sum_{k>0} (a_k + k - 1) - (a_M + M - 1) - (N - M) + (b_L + N - 1) \\ &= |\lambda^\bullet| + i - j. \end{aligned}$$

Similarly,

$$\begin{aligned} |\mu^\circ| &= \sum_{\substack{0 < k < L \\ \text{or } k > N'}} (k - \delta - b_k) + \sum_{L < k \leq N'} (k - 1 - \delta - b_k) + (N' - \delta - a_M) \\ &= \sum_{k>0} (k - \delta - b_k) - (L - \delta - b_L) - (N' - L) + (N' - \delta - a_M) \\ &= |\lambda^\circ| + i - j. \end{aligned} \quad \square$$

The next corollary, an immediate consequence of Proposition 6.3.6, will be useful later.

Corollary 6.3.7. $D'_{\lambda, \lambda} = 1$ for all λ . Moreover, $D'_{\lambda, \mu} = 0$ unless $\mu = \lambda$ or $|\mu| = (|\lambda^\bullet| - i, |\lambda^\circ| - i)$ for some $i > 0$.

6.4. Computing $\text{lift}_\delta(\lambda)$. The following theorem will allow us to explicitly compute values of the lifting map using the combinatorics developed in §6.3 (see Corollary 6.4.2). Our proof of this theorem relies heavily on the results in [11].

Theorem 6.4.1. $\sum_\nu D'_{\lambda, \nu} D'_{\mu, \nu} = (\lambda, \mu)_\delta$ for all bipartitions λ, μ and all $\delta \in \mathbb{K}$.

Proof. It follows from Corollary 6.2.4 that the statement of the theorem is symmetric in λ and μ , hence we may assume $|\mu| \not\geq |\lambda|$. Suppose $\lambda \vdash (r, s)$ and $\mu \vdash (r', s')$. $(\lambda, \mu)_\delta$ is zero unless $r + s' = r' + s$ (Proposition 4.6.1). Hence, by Corollary 6.3.7, it suffices to consider the case $\mu \vdash (r - i, s - i)$ for some $i \geq 0$.

Assume $\delta \neq 0$. Then $B_{r,s}$ is a quasi-hereditary (hence cellular) algebra (see [12, Corollary 2.8]) with decomposition numbers given by $D'_{\lambda,\nu}$ (see [11, Theorem 4.10]⁷). In particular, this implies that the $B_{r,s}$ -module homomorphisms between the projective modules $e_\lambda B_{r,s}$ and $e_\mu^{(i)} B_{r,s}$ satisfy the following:

$$(20) \quad \sum_{\nu} D'_{\lambda,\nu} D'_{\mu,\nu} = \dim_{\mathbb{K}} \text{Hom}_{B_{r,s}}(e_\lambda B_{r,s}, e_\mu^{(i)} B_{r,s})$$

(see, for instance [17, Theorem 3.7(iii)]). Since

$$(21) \quad \text{Hom}_{B_{r,s}}(e_\lambda B_{r,s}, e_\mu^{(i)} B_{r,s}) = e_\mu^{(i)} B_{r,s} e_\lambda = \text{Hom}_{\underline{\text{Rep}}(GL_\delta)}(L(\lambda), L(\mu)),$$

it follows that $(\lambda, \mu)_\delta$ agrees with (20).

If $\delta = 0$, the algebra $B_{r,s}$ is no longer quasi-hereditary, but it is still cellular (see [12, Theorem 2.7]). The decomposition numbers are still given by $D'_{\lambda,\nu}$, however, there is no PIM labelled by $(\emptyset, \emptyset)$ in this case, hence we must require $\lambda \neq (\emptyset, \emptyset)$. Hence, if neither λ nor μ is $(\emptyset, \emptyset)$, (20) and (21) still hold and we are done as before. Since we are assuming $|\mu| \not\geq |\lambda|$, to complete the proof of the theorem we only need to prove the case $\mu = (\emptyset, \emptyset)$, $\lambda \vdash (r, r)$ and $\delta = 0$. Since $D'_{(\emptyset, \emptyset), \nu} = 0$ whenever $\nu \neq (\emptyset, \emptyset)$ and $D'_{(\emptyset, \emptyset), (\emptyset, \emptyset)} = 1$, we have $\sum_{\nu} D'_{\lambda,\nu} D'_{(\emptyset, \emptyset), \nu} = D'_{\lambda, (\emptyset, \emptyset)}$. The decomposition number $D'_{\lambda, (\emptyset, \emptyset)}$ is the composition multiplicity of the simple $B_{r,r}$ -module labelled by λ in $\text{Hom}(w_{r,r}, \mathbf{1})$, the standard $B_{r,r}$ -module labelled by $(\emptyset, \emptyset)$. Hence,

$$(22) \quad D'_{\lambda, (\emptyset, \emptyset)} = \dim_{\mathbb{K}} \text{Hom}_{B_{r,r}}(e_\lambda B_{r,r}, \text{Hom}(w_{r,r}, \mathbf{1})).$$

Since

$$\text{Hom}_{B_{r,r}}(e_\lambda B_{r,r}, \text{Hom}(w_{r,r}, \mathbf{1})) = \text{Hom}(w_{r,r}, \mathbf{1}) e_\lambda = \text{Hom}_{\underline{\text{Rep}}(GL_0)}(L(\lambda), L((\emptyset, \emptyset))),$$

$(\lambda, (\emptyset, \emptyset))_0$ agrees with (22). $\square$

Corollary 6.4.2. $D_{\lambda,\mu}(\delta) = D'_{\lambda,\mu}(\delta)$ for all bipartitions λ, μ and all $\delta \in \mathbb{K}$.

Proof. First, put the following partial order on pairs of bipartitions: $(\lambda, \mu) > (\lambda', \mu')$ means either $|\lambda| > |\lambda'|$, or $\lambda = \lambda'$ and $|\mu| > |\mu'|$. We prove the corollary by inducting on this partial order. First notice that $D_{(\emptyset, \emptyset), (\emptyset, \emptyset)} = 1 = D'_{(\emptyset, \emptyset), (\emptyset, \emptyset)}$. Now assume $(\lambda, \mu) \neq ((\emptyset, \emptyset), (\emptyset, \emptyset))$. By Theorem 6.2.3(2) and Corollary 6.3.7 we may assume $|\lambda| > |\mu|$. Thus

$$\begin{aligned} D_{\lambda,\mu} &= (\lambda, \mu)_\delta - \sum_{\substack{\nu \\ |\nu| < |\mu|}} D_{\lambda,\nu} D_{\mu,\nu} \quad (\text{Corollary 6.2.4 and Theorem 6.2.3(2)}) \\ &= (\lambda, \mu)_\delta - \sum_{\substack{\nu \\ |\nu| < |\mu|}} D'_{\lambda,\nu} D'_{\mu,\nu} \quad (\text{Induction}) \\ &= D'_{\lambda,\mu} \quad (\text{Theorem 6.4.1 and Corollary 6.3.7}). \quad \square \end{aligned}$$

Example 6.4.3. Using Corollary 6.4.2 and Example 6.3.4 we have

$$\text{lift}_{-1}(((3, 2), (3, 1))) = ((3, 2), (3, 1)) + ((3), (1^2)) + ((2^2), (3)) + ((2), (1)).$$

⁷The results in [11] are only proved for $\mathbb{K} = \mathbb{C}$, however, using Corollary 4.4.3 it can be shown that their results hold over arbitrary fields of characteristic zero.

7. DECOMPOSING TENSOR PRODUCTS IN $\underline{\text{Rep}}(GL_\delta)$

In this section we give a generic decomposition formula for decomposing tensor products of indecomposable objects in $\underline{\text{Rep}}(GL_t)$. We then show how this generic decomposition formula along with the lifting map from the previous section can be used to decompose arbitrary tensor products in $\underline{\text{Rep}}(GL_\delta)$. Throughout this section we will work in the Grothendieck rings R_δ and R_t (see §6.1).

7.1. The generic case. The following theorem explains how to decompose the tensor product of two indecomposable objects in $\underline{\text{Rep}}(GL_t)$.

Theorem 7.1.1. *Given bipartitions λ, μ , and ν , let $\Gamma_{\lambda, \mu}^\nu$ be as in (16). Then $\lambda\mu = \sum_\nu \Gamma_{\lambda, \mu}^\nu \nu$ in R_t .*

Proof. Fix bipartitions λ and μ and let $\nu^{(1)}, \dots, \nu^{(k)}$ be bipartitions such that

$$(23) \quad \lambda\mu = \nu^{(1)} + \dots + \nu^{(k)}$$

in R_t . By Theorem 6.2.3(3) there exists a positive integer d which simultaneously satisfies: (i) $d \geq l(\lambda) + l(\mu)$; (ii) $d \geq l(\nu^{(i)})$ for each $i = 1, \dots, k$; and (iii) lift_d fixes $\lambda, \mu, \nu^{(1)}, \dots, \nu^{(k)}$. Now, lift_d is a ring isomorphism (Theorem 6.2.3(1)), hence (23) holds in R_d by assumption (iii). Since F_d is a tensor functor, by Theorem 5.2.2 along with assumptions (i) and (ii) we have

$$V_\lambda \otimes V_\mu = V_{\nu^{(1)}} \oplus \dots \oplus V_{\nu^{(k)}}$$

in $\text{Rep}(GL_d)$. The result now follows from Theorem 5.1.2 and assumption (i). $\square$

The following corollary lists special cases of Theorem 7.1.1, which are easy to prove using basic properties of Littlewood Richardson coefficients:

Corollary 7.1.2. *The following equations hold in R_t .*

$$(24) \quad (\lambda^\bullet, \emptyset)(\mu^\bullet, \emptyset) = \sum_{\alpha \in \mathcal{P}} LR_{\lambda^\bullet, \mu^\bullet}^\alpha(\alpha, \emptyset),$$

$$(25) \quad (\emptyset, \lambda^\circ)(\emptyset, \mu^\circ) = \sum_{\alpha \in \mathcal{P}} LR_{\lambda^\circ, \mu^\circ}^\alpha(\emptyset, \alpha),$$

$$(26) \quad (\lambda^\bullet, \emptyset)(\emptyset, \mu^\circ) = \sum_\nu \sum_{\kappa \in \mathcal{P}} LR_{\kappa, \nu}^{\lambda^\bullet} LR_{\kappa, \nu}^{\mu^\circ} \nu,$$

$$(27) \quad \lambda(\square, \emptyset) = \sum_{\lambda^{\bullet+}} (\lambda^{\bullet+}, \lambda^\circ) + \sum_{\lambda^{\circ-}} (\lambda^\bullet, \lambda^{\circ-}),$$

$$(28) \quad \lambda(\emptyset, \square) = \sum_{\lambda^{\circ+}} (\lambda^\bullet, \lambda^{\circ+}) + \sum_{\lambda^{\bullet-}} (\lambda^{\bullet-}, \lambda^\circ),$$

where the sums in (27) are taken over all partitions $\lambda^{\bullet+}$ (resp. $\lambda^{\circ-}$) obtained from the Young diagram $\lambda^\bullet$ (resp. λ°) by adding one box (resp. removing one box). Similarly for (28).

Example 7.1.3. In this example we compute $((2), \emptyset)(\square, \square) \in R_t$. By Example 6.1.1 (or using (26), (27), or (28)) we have $(\square, \emptyset)(\emptyset, \square) = (\square, \square) + (\emptyset, \emptyset) \in R_t$. Hence, using (27) and (28) we have the following in R_t :

$$\begin{aligned} ((2), \emptyset)(\square, \square) &= ((2), \emptyset)((\square, \emptyset)(\emptyset, \square) - (\emptyset, \emptyset)) \\ &= ((2), \emptyset)(\square, \emptyset)(\emptyset, \square) - ((2), \emptyset) \\ &= (((2), 1), \emptyset) + ((3), \emptyset)(\emptyset, \square) - ((2), \emptyset) \\ &= ((2), 1), \square + ((3), \square) + ((1^2), \emptyset) + ((2), \emptyset). \end{aligned}$$

7.2. Decomposing arbitrary tensor products. To compute the product of two bipartitions $\lambda, \mu \in R_\delta$ for arbitrary $\delta \in \mathbb{K}$ (i.e., to decompose tensor products in $\text{Rep}(GL_\delta)$) we

- (1) determine the coefficients in $\text{lift}_\delta(\lambda\mu) = \sum_{\nu, \nu'} D_{\lambda, \nu} D_{\mu, \nu'} \nu\nu'$ using Corollary 6.4.2,
- (2) use the results in §7.1 to expand $\sum_{\nu, \nu'} D_{\lambda, \nu} D_{\mu, \nu'} \nu\nu' = \nu^{(1)} + \dots + \nu^{(k)} \in R_t$,
- (3) determine $\text{lift}_\delta^{-1}(\nu^{(1)} + \dots + \nu^{(k)}) = \lambda\mu$, which by Theorem 6.2.3(2) consists of a sum of a subset of the bipartitions $\nu^{(1)}, \dots, \nu^{(k)}$.

The following examples illustrate the process described above:

Example 7.2.1. Consider $((2^2), (3, 1))(\square, \emptyset) \in R_{-1}$. Since

$$c_{((2^2), (3, 1))}(-1) = \cdots \quad \begin{array}{ccccccccccc} & & & & \curvearrowright & & & & & & \\ \wedge & \wedge & & \circ & \wedge & \times & \circ & \vee & \vee & & \\ -3 & -2 & -1 & 0 & 1 & 2 & 3 & 4 & 5 & & \end{array} \cdots$$

and

$$x_{((2), (1^2))}(-1) = \cdots \quad \begin{array}{ccccccccccc} & & & \vee & \times & & \vee & \vee & & & \\ \wedge & \wedge & \wedge & \circ & \wedge & \times & \circ & \vee & \vee & & \\ -3 & -2 & -1 & 0 & 1 & 2 & 3 & 4 & 5 & & \end{array} \cdots$$

by Corollary 6.4.2 we have $\text{lift}_{-1}(((2^2), (3, 1))) = ((2^2), (3, 1)) + ((2), (1^2))$. Moreover, by Example 6.2.1, $\text{lift}_{-1}((\square, \emptyset)) = (\square, \emptyset)$. Since lift_{-1} is a ring map (Theorem 6.2.3(1)) it follows that

$$\text{lift}_{-1}(((2^2), (3, 1))(\square, \emptyset)) = ((2^2), (3, 1))(\square, \emptyset) + ((2), (1^2))(\square, \emptyset),$$

which, by (27), is equal to

$$((3, 2), (3, 1)) + ((2^2, 1), (3, 1)) + ((2^2), (3)) + ((2^2), (2, 1)) + ((2, 1), (1^2)) + ((3), (1^2)) + ((2), \square).$$

Now, by Example 6.3.4 along with Corollary 6.4.2

$$\text{lift}_{-1}(((3, 2), (3, 1))) = ((3, 2), (3, 1)) + ((3), (1^2)) + ((2^2), (3)) + ((2), \square).$$

Similarly, one can show $\text{lift}_{-1}(((2^2, 1), (3, 1))) = ((2^2, 1), (3, 1))$ and

$$\text{lift}_{-1}(((2^2), (2, 1))) = ((2^2), (2, 1)) + ((2, 1), (1^2)).$$

Hence,

$$\text{lift}_{-1}(((2^2), (3, 1))(\square, \emptyset)) = \text{lift}_{-1}(((3, 2), (3, 1)) + ((2^2, 1), (3, 1)) + ((2^2), (2, 1))).$$

It follows from Theorem 6.2.3(1) that

$$((2^2), (3, 1))(\square, \emptyset) = ((3, 2), (3, 1)) + ((2^2, 1), (3, 1)) + ((2^2), (2, 1)) \in R_{-1}.$$

Example 7.2.2. In this example we compute $((2), \emptyset)(\square, \square) \in R_\delta$ for arbitrary $\delta \in \mathbb{K}$. First, since lift_δ is a ring map (Theorem 6.2.3(1)), by Examples 6.2.1 and 6.2.2 we have

$$\text{lift}_\delta(((2), \emptyset)(\square, \square)) = \begin{cases} ((2), \emptyset)(\square, \square) + ((2), \emptyset) & \text{if } \delta = 0, \\ ((2), \emptyset)(\square, \square) & \text{if } \delta \neq 0. \end{cases}$$

Hence, by Example 7.1.3 we have

$$\text{lift}_\delta(((2), \emptyset)(\square, \square)) = \begin{cases} ((2, 1), \square) + ((3), \square) + ((1^2), \emptyset) + 2((2), \emptyset) & \text{if } \delta = 0, \\ ((2, 1), \square) + ((3), \square) + ((1^2), \emptyset) + ((2), \emptyset) & \text{if } \delta \neq 0. \end{cases}$$

Now, $\text{lift}_\delta(((1^2), \emptyset)) = ((1^2), \emptyset)$ and $\text{lift}_\delta(((2), \emptyset)) = ((2), \emptyset)$ for all δ (Example 6.2.1). Moreover, using Corollary 6.4.2, we compute

$$\text{lift}_\delta(((2, 1), \square)) = \begin{cases} ((2, 1), \square) + ((1^2), \emptyset) & \text{if } \delta = -1, \\ ((2, 1), \square) + ((2), \emptyset) & \text{if } \delta = 1, \\ ((2, 1), \square) & \text{if } \delta \neq \pm 1. \end{cases}$$

$$\text{lift}_\delta(((3), \square)) = \begin{cases} ((3), \square) + ((2), \emptyset) & \text{if } \delta = -2, \\ ((3), \square) & \text{if } \delta \neq -2. \end{cases}$$

Since lift_δ is an isomorphism for all δ , we have the following in R_δ :

$$((2), \emptyset)(\square, \square) = \begin{cases} ((2, 1), \square) + ((3), \square) + ((1^2), \emptyset) + 2((2), \emptyset) & \text{if } \delta = 0, \\ ((2, 1), \square) + ((3), \square) + ((2), \emptyset) & \text{if } \delta = -1, \\ ((2, 1), \square) + ((3), \square) + ((1^2), \emptyset) & \text{if } \delta \in \{1, -2\}, \\ ((2, 1), \square) + ((3), \square) + ((1^2), \emptyset) + ((2), \emptyset) & \text{otherwise.} \end{cases}$$

8. REPRESENTATIONS OF THE GENERAL LINEAR SUPERGROUP

Fix $m, n \geq 0$ and consider the algebraic supergroup $GL(m|n)$ over $\mathbb{K}$. In this paper we will only deal with finite dimensional representations of $GL(m|n)$, which can be identified with integrable representations of the corresponding Lie superalgebra $\mathfrak{gl}(m|n)$. We prefer to exploit this identification and work with $\mathfrak{gl}(m|n)$ rather than $GL(m|n)$, and we will do so for the rest of the paper. We begin by fixing notation and conventions for representations of $\mathfrak{gl}(m|n)$.

8.1. The category $\text{Rep}(\mathfrak{gl}(m|n))$. Let $V = V_{\bar{0}} \oplus V_{\bar{1}}$ denote a superspace over $\mathbb{K}$ with $\dim_{\mathbb{K}} V_{\bar{0}} = m$, $\dim_{\mathbb{K}} V_{\bar{1}} = n$. Write $\mathfrak{gl}(m|n)$ for the associated *general linear Lie superalgebra*, that is, for the Lie superalgebra of endomorphisms of the superspace V , considered as $(m+n) \times (m+n)$ matrices. Then V is called the *natural module* for $\mathfrak{gl}(m|n)$.

Let $\text{Rep}(\mathfrak{gl}(m|n))$ denote the category of finite-dimensional $\mathfrak{gl}(m|n)$ -modules. Given $\mathfrak{gl}(m|n)$ -modules U, U' , the tensor product of superspaces $U \otimes U'$ is again a $\mathfrak{gl}(m|n)$ -module with action

$$(29) \quad x \cdot (u \otimes u') = (x \cdot u) \otimes u' + (-1)^{\bar{x}\bar{u}} u \otimes (x \cdot u'),$$

for $x \in \mathfrak{gl}(m|n)$, $u \in U$, $u' \in U'$. Thus, $\text{Rep}(\mathfrak{gl}(m|n))$ is a monoidal category, where the one-dimensional purely-even module $\mathbb{1}$ carries the trivial action. Note that, in particular, we consider the tensor product of $\text{Rep}(\mathfrak{gl}(m|n))$ to be strict.

If U is a $\mathfrak{gl}(m|n)$ -module, then the dual superspace U^* is again a $\mathfrak{gl}(m|n)$ -module with action

$$(30) \quad (x \cdot \phi)(u) = -(-1)^{\bar{x}\bar{\phi}} \phi(x \cdot u),$$

for $x \in \mathfrak{gl}(m|n)$, $\phi \in U^*$, $u \in U$. The maps ev_U , coev_U are maps of $\mathfrak{gl}(m|n)$ -modules, and so $\text{Rep}(\mathfrak{gl}(m|n))$ is a tensor category.

It is well known that the category $\text{Rep}(\mathfrak{gl}(m|n))$ is not semisimple when $n > 0$ or $m > 0$ (see e.g. [24]).

8.2. Characters. Let $\mathfrak{h} \subset \mathfrak{gl}(m|n)$ denote the subalgebra of diagonal matrices. For any object U of $\text{Rep}(\mathfrak{gl}(m|n))$, let

$$\text{ch } U = \sum_{\mu \in \mathfrak{h}^*} \dim_{\mathbb{K}} U_{\mu} e^{\mu},$$

denote the *character* of U , where $\dim_{\mathbb{K}} U_{\mu}$ denotes the dimension of the μ -weight space of U as a vector space over $\mathbb{K}$, and e^{μ} denotes the formal exponential. Addition and multiplication of characters of objects in $\text{Rep}(\mathfrak{gl}(m|n))$ are defined componentwise and by convolution, respectively. Since $\mathfrak{h}$ is purely even, it follows that from the definition of the biproduct and equation (29) that

$$\text{ch}(U \oplus U') = \text{ch } U + \text{ch } U', \quad \text{ch}(U \otimes U') = \text{ch } U \cdot \text{ch } U'$$

for any objects U, U' of $\text{Rep}(\mathfrak{gl}(m|n))$.

For any $r, s \geq 0$, write $T(r, s) = V^{\otimes r} \otimes (V^*)^{\otimes s}$, as per §4.7. Let $\{\varepsilon_i\}_{1 \leq i \leq m+n}$ denote the diagonal coordinate functions of $\mathfrak{gl}(m|n)$, so that ε_i takes any matrix to its (i, i) -entry. Then for any $r, s \geq 0$, all weights of $T(r, s)$ are integral linear combinations of the ε_i (hence also of any submodule of $T(r, s)$). Let

$$x_i = e^{\varepsilon_i}, \quad 1 \leq i \leq m, \quad y_j = e^{\varepsilon_{m+j}}, \quad 1 \leq j \leq n.$$

If U is a submodule of $T(r, s)$, then $\text{ch } U = \text{ch } U(x|y)$ is a Laurent polynomial in the variables $x = \{x_i\}$ and $y = \{y_j\}$. Write $\bar{x} = \{x_i^{-1}\}_{1 \leq i \leq m}$ and $\bar{y} = \{y_i^{-1}\}_{1 \leq i \leq n}$. Then it follows from equation (30) that

$$\text{ch}(U^*)(x|y) = \text{ch } U(\bar{x}|\bar{y}).$$

Example 8.2.1. One has that $\text{ch } \mathbb{1} = 1$, while

$$\text{ch } V = x_1 + \cdots + x_m + y_1 + \cdots + y_n, \quad \text{ch } V^* = x_1^{-1} + \cdots + x_m^{-1} + y_1^{-1} + \cdots + y_n^{-1}.$$

8.3. The functor $F_{m|n} : \underline{\text{Rep}}(GL_{m-n}) \rightarrow \text{Rep}(\mathfrak{gl}(m|n))$. Let

$$F_{m|n} : \underline{\text{Rep}}(GL_{m-n}) \rightarrow \text{Rep}(\mathfrak{gl}(m|n))$$

denote the tensor functor which sends $\bullet \mapsto V$ defined by Proposition 3.5.1. For any $\zeta \in \mathbb{K}$, the central element $\zeta \cdot \text{id}_{m+n} \in \mathfrak{gl}(m|n)$ acts on the mixed tensor power $T(r, s)$ by the scalar ζ^{r-s} , and so $\text{Hom}_{\text{Rep}(\mathfrak{gl}(m|n))}(T(r, s), T(r', s')) = 0$ unless $r+s' = r'+s$. Moreover, it is well known that the $\mathbb{K}$ -algebra map $\mathbb{K}\Sigma_p \rightarrow \text{End}_{\text{Rep}(\mathfrak{gl}(m|n))}(V^{\otimes p})$ defined by the symmetric braiding is surjective, for any $p \geq 0$ (see [4, Remark 4.15] or [25]). Thus, by Theorem 4.7.1, the functor $F_{m|n}$ is full and any indecomposable summand of mixed tensor powers $T(r, s)$ is isomorphic to $F_{m|n}(L(\lambda))$ for some bipartition λ . Generalizing our notation from §5, we set $W(\lambda) = F_{m|n}(L(\lambda))$. The rest of this section is devoted to describing $W(\lambda)$ for arbitrary λ . More precisely, we will give a formula for computing the character of $W(\lambda)$ in §8.5 and give a criterion for the vanishing of $W(\lambda)$ in §8.7. By Theorem 4.7.1, these results give a classification of indecomposable summands of mixed tensor space in $\text{Rep}(\mathfrak{gl}(m|n))$.

8.4. Composite supersymmetric Schur polynomials. In §8.5 we give a formula for computing the character of $W(\lambda)$ in $\text{Rep}(\mathfrak{gl}(m|n))$ (see Theorem 8.5.2). In the case $\lambda^\circ = \emptyset$ (resp. $\lambda^\bullet = \emptyset$), the character of $W(\lambda)$ was computed in [4] and [25] and is called a *covariant* (resp. *contravariant*) *supersymmetric Schur polynomial*. Our formula for $\text{ch } W(\lambda)$ for arbitrary λ is in terms of the numbers $D_{\lambda,\mu}$ (§6) and the so-called *composite supersymmetric Schur polynomials*⁸ (see for instance [22]). There are many equivalent definitions of composite supersymmetric Schur polynomials; we will use the determinantal formula found, for instance, in [22, (38)]. In order to state this formula we need a few preliminary definitions. As in §8.1, we work with the variables $x = \{x_i\}_{1 \leq i \leq m}$ and $y = \{y_i\}_{1 \leq i \leq n}$, and write $\bar{x} = \{x_i^{-1}\}_{1 \leq i \leq m}$ and $\bar{y} = \{y_i^{-1}\}_{1 \leq i \leq n}$. Now, we define the *complete supersymmetric polynomials* by

$$\mathbf{h}_k = \mathbf{h}_k(x|y) = \sum_{i=0}^k h_{k-i}(x)e_i(y)$$

where $h_k(x)$ and $e_k(y)$ are the complete and elementary symmetric polynomials respectively (see for instance [19, §I.2]). In particular, $\mathbf{h}_0 = 1$ and $\mathbf{h}_k = 0$ whenever $k < 0$. Next, we write $\bar{\mathbf{h}}_k = \mathbf{h}_k(\bar{x}|\bar{y})$. Now, given a bipartition λ , we define the *composite supersymmetric Schur polynomial* $\mathbf{s}_\lambda = \mathbf{s}_\lambda(x|y)$ as the following determinate (compare with [22, (38)]⁹):

$$\mathbf{s}_\lambda = \det \begin{pmatrix} \bar{\mathbf{h}}_{\lambda_q^\circ} & & & & \vdots \\ \bar{\mathbf{h}}_{\lambda_q^\circ-1} & \ddots & \bar{\mathbf{h}}_{\lambda_2^\circ+1} & & \vdots \\ \vdots & & \bar{\mathbf{h}}_{\lambda_2^\circ} & \bar{\mathbf{h}}_{\lambda_1^\circ+1} & \vdots \\ & & \bar{\mathbf{h}}_{\lambda_2^\circ-1} & \bar{\mathbf{h}}_{\lambda_1^\circ} & \mathbf{h}_{\lambda_1^\bullet-1} & \vdots \\ & & \vdots & \bar{\mathbf{h}}_{\lambda_1^\circ-1} & \mathbf{h}_{\lambda_1^\bullet} & \mathbf{h}_{\lambda_2^\bullet-1} \\ & & & \vdots & \mathbf{h}_{\lambda_1^\bullet+1} & \mathbf{h}_{\lambda_2^\bullet} & \vdots \\ & & & & \vdots & \mathbf{h}_{\lambda_2^\bullet+1} & \ddots & \mathbf{h}_{\lambda_p^\bullet-1} \\ & & & & & \vdots & & \mathbf{h}_{\lambda_p^\bullet} \end{pmatrix}$$

where p (resp. q) is any integer greater than or equal to $l(\lambda^\bullet)$ (resp. $l(\lambda^\circ)$).

⁸Also known as *composite supersymmetric S-polynomials*.

⁹In the literature $\mathbf{s}_\lambda(x|y)$ is sometimes denoted $s_{\overline{\lambda^\circ}; \lambda^\bullet}(x/y)$ or even $\{\overline{\lambda^\circ}; \lambda^\bullet\}$.

Example 8.4.1. Fix $m = 1$ and $n = 2$. Then

$$\begin{aligned} \mathbf{s}_{((1),(2))} &= \det \begin{pmatrix} \bar{\mathbf{h}}_2 & \mathbf{h}_0 \\ \bar{\mathbf{h}}_1 & \mathbf{h}_1 \end{pmatrix} \\ &= \det \begin{pmatrix} \frac{1}{x_1^2} + \frac{1}{x_1 y_1} + \frac{1}{x_1 y_2} + \frac{1}{y_1 y_2} & 1 \\ \frac{1}{x_1} + \frac{1}{y_1} + \frac{1}{y_2} & x_1 + y_1 + y_2 \end{pmatrix} \\ &= \frac{x_1}{y_1 y_2} + \frac{y_1}{x_1 y_2} + \frac{y_2}{x_1 y_1} + \frac{y_1}{x_1^2} + \frac{y_2}{x_1^2} + 2 \frac{1}{x_1} + \frac{1}{y_1} + \frac{1}{y_2}, \end{aligned}$$

whereas

$$\begin{aligned} \mathbf{s}_{((1^2),(3))} &= \det \begin{pmatrix} \bar{\mathbf{h}}_3 & \mathbf{h}_0 & \mathbf{h}_{-1} \\ \bar{\mathbf{h}}_2 & \mathbf{h}_1 & \mathbf{h}_0 \\ \bar{\mathbf{h}}_1 & \mathbf{h}_2 & \mathbf{h}_1 \end{pmatrix} \\ &= \det \begin{pmatrix} \frac{1}{x_1^3} + \frac{1}{x_1^2 y_1} + \frac{1}{x_1^2 y_2} + \frac{1}{x_1 y_1 y_2} & 1 & 0 \\ \frac{1}{x_1^2} + \frac{1}{x_1 y_1} + \frac{1}{x_1 y_2} + \frac{1}{y_1 y_2} & x_1 + y_1 + y_2 & 1 \\ \frac{1}{x_1} + \frac{1}{y_1} + \frac{1}{y_2} & x_1^2 + x_1 y_1 + x_1 y_2 + y_1 y_2 & x_1 + y_1 + y_2 \end{pmatrix} \\ &= \frac{y_1^2}{x_1^3} + \frac{y_1 y_2}{x_1^3} + \frac{y_2^2}{x_1^3} + \frac{y_1^2}{x_1^2 y_2} + \frac{y_2^2}{x_1^2 y_1} + 2 \frac{y_1}{x_1^2} + 2 \frac{y_2}{x_1^2} + \frac{1}{x_1} + \frac{y_1}{x_1 y_2} + \frac{y_2}{x_1 y_1} - \frac{x_1}{y_1 y_2}. \end{aligned}$$

8.5. The character of $W(\lambda)$. The following proposition lists some of the well-known properties of $\mathbf{s}_\lambda$ which will be useful for this paper.

Proposition 8.5.1. (1) $\text{ch } W(\lambda) = \mathbf{s}_\lambda$ whenever $\lambda^\bullet = \emptyset$ or $\lambda^\circ = \emptyset$.

$$(2) \mathbf{s}_{(\lambda^\bullet, \emptyset)} \mathbf{s}_{(\emptyset, \mu^\circ)} = \sum_\nu \sum_{\kappa \in \mathcal{P}} LR_{\kappa, \nu^\bullet}^{\lambda^\bullet} LR_{\kappa, \nu^\circ}^{\mu^\circ} \mathbf{s}_\nu.$$

Proof. (1) follows from the corresponding determinantal formula for the (non-composite) supersymmetric Schur polynomials (see [21, (6)] and references therein). For (2) see [13, (3.2)]. $\square$

We are now ready to prove our formula for computing the character of $W(\lambda)$.

Theorem 8.5.2. $\text{ch } W(\lambda) = \sum_\mu D_{\lambda, \mu}(m-n) \mathbf{s}_\mu$ for any bipartition λ .

Proof. We induct on the size of λ . The case $\lambda = (\emptyset, \emptyset)$ is easy to check. Now, by Proposition 4.6.4 we can write

$$(31) \quad \lambda = (\lambda^\bullet, \emptyset)(\emptyset, \lambda^\circ) - \sum_{\substack{\nu \\ |\nu| < |\lambda|}} a_{\lambda, \nu} \nu \in R_{m-n}$$

for some $a_{\lambda, \nu} \in \mathbb{Z}$. Write $D_{\lambda, \mu} = D_{\lambda, \mu}(m-n)$. Applying lift_{m-n} to (31) and using Example 6.2.1 we have

$$(32) \quad \sum_\mu D_{\lambda, \mu} \mu = (\lambda^\bullet, \emptyset)(\emptyset, \lambda^\circ) - \sum_{\substack{\nu \\ |\nu| < |\lambda|}} a_{\lambda, \nu} \sum_\mu D_{\nu, \mu} \mu \in R_t.$$

Using formula (26) we compute the coefficient of μ in (32) to be

$$(33) \quad D_{\lambda, \mu} = \sum_{\kappa \in \mathcal{P}} LR_{\kappa, \mu}^{\lambda^\bullet} LR_{\kappa, \mu^\circ}^{\lambda^\circ} - \sum_{\substack{\nu \\ |\nu| < |\lambda|}} a_{\lambda, \nu} D_{\nu, \mu}.$$

On the other hand, since $F_{m|n}$ is a tensor functor, it follows from (31) that

$$\text{ch } W(\lambda) = \text{ch } W((\lambda^\bullet, \emptyset)) \text{ch } W((\emptyset, \lambda^\circ)) - \sum_{\substack{\nu \\ |\nu| < |\lambda|}} a_{\lambda, \nu} \text{ch } W(\nu).$$

Hence, by Proposition 8.5.1(1) along with induction, we have

$$\text{ch } W(\lambda) = \mathbf{s}_{(\lambda^\bullet, \emptyset)} \mathbf{s}_{(\emptyset, \lambda^\circ)} - \sum_{\substack{\nu \\ |\nu| < |\lambda|}} a_{\lambda, \nu} \sum_{\mu} D_{\nu, \mu} \mathbf{s}_{\mu}.$$

Thus, by Proposition 8.5.1(2),

$$\text{ch } W(\lambda) = \sum_{\mu} \sum_{\kappa \in \mathcal{P}} LR_{\kappa, \mu}^{\lambda^\bullet} \cdot LR_{\kappa, \mu}^{\lambda^\circ} \mathbf{s}_{\mu} - \sum_{\substack{\nu \\ |\nu| < |\lambda|}} a_{\lambda, \nu} \sum_{\mu} D_{\nu, \mu} \mathbf{s}_{\mu},$$

and we are done by (33). $\square$

Example 8.5.3. Let $m = 1$, $n = 2$ and consider the bipartition $((1^2), (3))$. Since

$$c_{((1^2), (3))}(-1) = \cdots \quad \begin{array}{cccccccc} & & & \curvearrowright & & & & \\ \wedge & \wedge & & \wedge & & \circ & \vee & \vee \\ -3 & -2 & -1 & 0 & 1 & 2 & 3 & 4 \end{array} \cdots$$

and

$$x_{((1), (2))}(-1) = \cdots \quad \begin{array}{cccccccc} & & & \vee & & \circ & \vee & \vee \\ \wedge & \wedge & \wedge & \wedge & \wedge & \wedge & \wedge & \wedge \\ -3 & -2 & -1 & 0 & 1 & 2 & 3 & 4 \end{array} \cdots$$

by Corollary 6.4.2 we have $\text{lift}_{-1}(((1^2), (3)))) = ((1^2), (3)) + ((1), (2))$. Hence, by Theorem 8.5.2 and Example 8.4.1 we have

$$\begin{aligned} \text{ch } W(((1^2), (3)))) &= \mathbf{s}_{((1^2), (3))} + \mathbf{s}_{((1), (2))} \\ &= \frac{y_1^2}{x_1^3} + \frac{y_1 y_2}{x_1^3} + \frac{y_2^2}{x_1^3} + \frac{y_1^2}{x_1^2 y_2} + \frac{y_2^2}{x_1^2 y_1} + 3 \frac{y_1}{x_1^2} + 3 \frac{y_2}{x_1^2} + 2 \frac{y_1}{x_1 y_2} + 2 \frac{y_2}{x_1 y_1} \\ &\quad + 3 \frac{1}{x_1} + \frac{1}{y_1} + \frac{1}{y_2}. \end{aligned}$$

For any Laurent polynomial $f = f(z_1, \dots, z_k)$, write

$$\text{edeg } f = \max\{ \deg(f|_{z_i := z_i^{\varepsilon_i}, 1 \leq i \leq k}) \mid \varepsilon_i \in \{\pm 1\}, 1 \leq i \leq k \}$$

for the *extremal degree* of f , where $\deg$ defines the familiar total degree. Then one has $\text{edeg}(fg) \leq \text{edeg } f + \text{edeg } g$. For any bipartition λ , we consider $\mathbf{s}_{\lambda}$ and $\text{ch } W(\lambda)$ as Laurent polynomials in the variables $x = \{x_i\}$ and $y = \{y_j\}$. Then we have the following corollary to Theorem 8.5.2:

Corollary 8.5.4. *Let λ be a bipartition. Then $\text{edeg } \text{ch } W(\lambda) \leq |\lambda|$.*

Proof. That $\text{edeg } \mathbf{s}_{\mu} \leq |\mu|$ for any bipartition μ follows from e.g. [22, (32)]. The claim then follows from Theorems 6.2.3(2) and 8.5.2. $\square$

8.6. Dimensions. In this subsection we derive a formula for the $\mathbb{K}$ -dimension of $W(\lambda)$ in $\text{Rep}(\mathfrak{gl}(m|n))$. To do so, let $\mathbf{d}_k = \mathbf{d}_k(m|n)$ denote the result of setting $x_i = y_j = 1$ for all $1 \leq i \leq m$, $1 \leq j \leq n$ in $\mathbf{h}_k$. From the definition of $\mathbf{h}_k$ we have

$$(34) \quad \mathbf{d}_k = \sum_{0 \leq i \leq k} \binom{m+k-i-1}{m-1} \binom{n}{i}$$

whenever $m > 0$, and $\mathbf{d}_k = \binom{n}{k}$ when $m = 0$. Now, given a bipartition λ we let $\mathbf{d}_{\lambda} = \mathbf{d}_{\lambda}(m|n)$ denote the result of setting $x_i = y_j = 1$ for all $1 \leq i \leq m$, $1 \leq j \leq n$

in s_λ , so that

$$(35) \quad d_\lambda = \det \begin{pmatrix} d_{\lambda_q^\circ} & & \vdots & & \\ d_{\lambda_q^\circ-1} & \ddots & d_{\lambda_2^\circ+1} & \vdots & \\ \vdots & & d_{\lambda_2^\circ} & d_{\lambda_1^\circ+1} & \vdots \\ & & d_{\lambda_2^\circ-1} & d_{\lambda_1^\circ} & d_{\lambda_1^\bullet-1} & \vdots \\ & & \vdots & d_{\lambda_1^\circ-1} & d_{\lambda_1^\bullet} & d_{\lambda_2^\bullet-1} \\ & & & \vdots & d_{\lambda_1^\bullet+1} & d_{\lambda_2^\bullet} & \vdots \\ & & & & \vdots & d_{\lambda_2^\bullet+1} & \ddots & d_{\lambda_p^\bullet-1} \\ & & & & & \vdots & & d_{\lambda_p^\bullet} \end{pmatrix}$$

where p (resp. q) is any integer greater than or equal to $l(\lambda^\bullet)$ (resp. $l(\lambda^\circ)$). It follows immediately from Theorem 8.5.2 that

$$(36) \quad \dim_{\mathbb{K}} W(\lambda) = \sum_{\mu} D_{\lambda, \mu} d_{\mu}$$

in $\text{Rep}(\mathfrak{gl}(m|n))$. The following technical lemma concerning d_k will be useful later:

Lemma 8.6.1. *Assume $m > 0$ and let $g_k(u) = \prod_{1 \leq j < m} (k - u + j)$ (an $(m-1)$ -degree polynomial in the variable u). Then $d_k + (-1)^{m-1} d_{n-m-k} = \frac{1}{(m-1)!} \sum_{0 \leq i \leq n} g_k(i) \binom{n}{i}$ for all $k \in \mathbb{Z}$.*

Proof. Since $\binom{m+k-i-1}{m-1} = \frac{1}{(m-1)!} \prod_{1 \leq j < m} (k - i + j)$, by (34) we have

$$d_k = \frac{1}{(m-1)!} \sum_{0 \leq i \leq k} \prod_{1 \leq j < m} (k - i + j) \binom{n}{i} = \frac{1}{(m-1)!} \sum_{0 \leq i \leq k} g_k(i) \binom{n}{i}.$$

Moreover,

$$\begin{aligned} d_{n-m-k} &= \frac{1}{(m-1)!} \sum_{0 \leq i \leq n-m-k} \prod_{1 \leq j < m} (n-m-k-i+j) \binom{n}{i} \\ &= \frac{1}{(m-1)!} \sum_{k+m \leq i \leq n} \prod_{1 \leq j < m} (i-m-k+j) \binom{n}{i} \\ &= \frac{1}{(m-1)!} \sum_{k+m \leq i \leq n} \prod_{1 \leq j < m} (i-k-j) \binom{n}{i} \\ &= \frac{(-1)^{m-1}}{(m-1)!} \sum_{k+m \leq i \leq n} g_k(i) \binom{n}{i}. \end{aligned}$$

Since $g_k(i) = 0$ whenever $k < i < k+m$, the result follows. $\square$

8.7. A criterion for the vanishing of $W(\lambda)$. We now concern ourselves with the question of when $W(\lambda)$ is zero. In the case $\lambda^\circ = \emptyset$ or $\lambda^\bullet = \emptyset$, the answer is well known and due to [4] and [25]. To state their result, we say a partition α is $(m|n)$ -hook if $\alpha_{m+1} \leq n$. Pictorially, α is $(m|n)$ -hook if and only if its Young

diagram fits into an m -high, n -wide hook:

We now record the result on the vanishing of $W(\lambda)$ found in [4] and [25]:

Theorem 8.7.1. *Suppose λ is a bipartition with $\lambda^\circ = \emptyset$ (resp. $\lambda^\bullet = \emptyset$). Then $W(\lambda) \neq 0$ in $\text{Rep}(\mathfrak{gl}(m|n))$ if and only if $\lambda^\bullet$ (resp. λ°) is $(m|n)$ -hook.*

The goal of this subsection is to prove a theorem analogous to Theorem 8.7.1 which holds for arbitrary bipartitions λ (see Theorem 8.7.6). First, we generalize the notion of $(m|n)$ -hook as follows: We call a bipartition λ $(m|n)$ -cross if there exists k with $0 \leq k \leq m$ such that $\lambda_{k+1}^\bullet + \lambda_{m-k+1}^\circ \leq n$. Pictorially, λ is $(m|n)$ -cross if and only if its diagram (see §4.1) fits into an m -high, n -wide cross:

Indeed, if $\lambda_{k+1}^\bullet + \lambda_{m-k+1}^\circ \leq n$, then λ can be covered by an m -high, n -wide cross if the cross is placed so that the horizontal strip covers exactly the first k rows of $\lambda^\bullet$ and the vertical strip covers exactly the first $\lambda_{k+1}^\bullet$ columns of $\lambda^\bullet$:

For example, $((3,1),(2))$ is both $(2|1)$ -cross and $(1|3)$ -cross as shown below:

However, $((3, 1), (2))$ is not $(1|2)$ -cross since it is impossible to cover its diagram with a 1-high, 2-wide cross:

We will eventually show $W(\lambda) \neq 0$ in $\text{Rep}(\mathfrak{gl}(m|n))$ if and only if λ is $(m|n)$ -cross (see Theorem 8.7.6). Notice that for bipartitions λ with $\lambda^\circ = \emptyset$ (resp. $\lambda^\bullet = \emptyset$) we have that λ is $(m|n)$ -cross if and only if $\lambda^\bullet$ (resp. λ°) is $(m|n)$ -hook. Hence our criterion for the vanishing of $W(\lambda)$ will generalize Theorem 8.7.1. The following lemma gets us half way towards proving our criterion for the vanishing of $W(\lambda)$.

Lemma 8.7.2. *If λ is $(m|n)$ -cross, then $W(\lambda) \neq 0$ in $\text{Rep}(\mathfrak{gl}(m|n))$.*

Proof. Assume k is such that $0 \leq k \leq m$ and $\lambda_{k+1}^\bullet + \lambda_{m-k+1}^\circ \leq n$. For convenience, write $l = \lambda_{k+1}^\bullet$ and $l' = \lambda_{m-k+1}^\circ$. Now set $\eta_i^\bullet = (\lambda^\bullet)_i^t - k$ for all $1 \leq i \leq l$ and $\eta_i^\circ = (\lambda^\circ)_i^t - m + k$ for all $1 \leq i \leq l'$. Now, $\text{ch } W((\lambda^\bullet, \emptyset))$ (resp. $\text{ch } W((\emptyset, \lambda^\circ))$) is a covariant (resp. contravariant) supersymmetric Schur polynomial, which can be computed using so-called *supertableaux* (see [4]). From the supertableaux definition of supersymmetric Schur polynomials it is apparent that the coefficient of $x_1^{\lambda_1^\bullet} \cdots x_k^{\lambda_k^\bullet} y_1^{\eta_1^\bullet} \cdots y_l^{\eta_l^\bullet}$ (resp. $x_{k+1}^{-\lambda_{k+1}^\circ} \cdots x_m^{-\lambda_m^\circ} y_{n-l'+1}^{-\eta_1^\circ} \cdots y_n^{-\eta_{l'}^\circ}$) in $\text{ch } W((\lambda^\bullet, \emptyset))$ (resp. $\text{ch } W((\emptyset, \lambda^\circ))$) has non-zero coefficient. Hence the coefficient of

$$(37) \quad \frac{x_1^{\lambda_1^\bullet} \cdots x_k^{\lambda_k^\bullet} y_1^{\eta_1^\bullet} \cdots y_l^{\eta_l^\bullet}}{x_{k+1}^{\lambda_{k+1}^\circ} \cdots x_m^{\lambda_m^\circ} y_{n-l'+1}^{\eta_1^\circ} \cdots y_n^{\eta_{l'}^\circ}}$$

in $C := \text{ch}(W((\lambda^\bullet, \emptyset)) \otimes W((\emptyset, \lambda^\circ)))$ is non-zero. Thus

$$\text{edeg } C \geq \sum_{1 \leq i \leq k} \lambda_i^\bullet + \sum_{1 \leq i \leq l} \eta_i^\bullet + \sum_{1 \leq i \leq m-k} \lambda_i^\circ + \sum_{1 \leq i \leq l'} \eta_i^\circ = |\lambda|,$$

since $l + l' \leq n$. On the other hand, by Proposition 4.6.4 we have

$$(38) \quad C = \text{ch } W(\lambda) + \text{ch } W(\mu^{(1)}) + \cdots + \text{ch } W(\mu^{(s)})$$

for some bipartitions $\mu^{(1)}, \dots, \mu^{(s)}$ with $|\mu^{(i)}| < |\lambda|$ for all $i = 1, \dots, s$. Thus the coefficient of (37) is non-zero in $\text{ch } W(\lambda)$ by Corollary 8.5.4. $\square$

The rest of this subsection is devoted to proving the converse of Lemma 8.7.2. We begin by focusing our attention to a certain class of non- $(m|n)$ -cross bipartitions. More precisely, we call a bipartition λ *almost $(m|n)$ -cross* if $l(\lambda^\bullet), l(\lambda^\circ) \leq m+1$ and $\lambda_{k+1}^\bullet + \lambda_{m-k+1}^\circ = n+1$ whenever $0 \leq k \leq m$. Equivalently, λ is almost $(m|n)$ -cross if λ is not $(m|n)$ -cross, but any bipartition obtained from λ by removing a box is $(m|n)$ -cross. Pictorially, λ is almost $(m|n)$ -cross if and only if the Young diagrams of $\lambda^\bullet$ and λ° can be glued together to form a $(m+1) \times (n+1)$ rectangle:

The next proposition states that almost $(m|n)$ -cross bipartitions lift trivially from R_{m-n} , hence characters and dimensions of the corresponding representations of $\mathfrak{gl}(m|n)$ can be easily computed (see Theorem 8.5.2 and (36)).

Proposition 8.7.3. *If λ is almost $(m|n)$ -cross, then $\text{lift}_{m-n}(\lambda) = \lambda$.*

Proof. Assume λ is almost $(m|n)$ -cross. Then

$$\begin{aligned} I_{\wedge}(\lambda) &= \{\lambda_1^{\bullet}, \dots, \lambda_{m+1}^{\bullet} - m, -m-1, -m-2, \dots\}, \\ I_{\vee}(\lambda, m-n) &= \{1-m+n-\lambda_1^{\circ}, \dots, 1+n-\lambda_{m+1}^{\circ}, n+2, n+3, \dots\} \\ &= \{\lambda_{m+1}^{\bullet} - m, \dots, \lambda_1^{\bullet}, n+2, n+3, \dots\}. \end{aligned}$$

Hence,

$$x_{\lambda}(m-n) = \cdots \overbrace{\wedge \wedge \wedge}^{m} \underbrace{\quad \times \text{'s and } \bigcirc \text{'s only} \quad}_{\text{only}} \overbrace{\vee \vee \vee}^{n} \cdots$$

In particular, $c_{\lambda}(m-n)$ has no caps and we are done by Corollary 6.4.2. $\square$

Corollary 8.7.4. *If λ is almost $(m|n)$ -cross, then $\dim_{\mathbb{K}} W(\lambda) = d_{\lambda}$ in $\text{Rep}(\mathfrak{gl}(m|n))$.*

Proof. This follows immediately from Proposition 8.7.3 and (36). $\square$

We are now ready to prove the following lemma, which is the crucial step towards proving the converse of Lemma 8.7.2.

Lemma 8.7.5. *If λ is almost $(m|n)$ -cross, then $W(\lambda) = 0$ in $\text{Rep}(\mathfrak{gl}(m|n))$.*

Proof. By Corollary 8.7.4 it suffices to show $d_{\lambda} = 0$ whenever λ is almost $(m|n)$ -cross. If $m = 0$, then almost $(m|n)$ -cross bipartitions are exactly ones of the form $\lambda = ((l), (n+1-l))$ for some $0 \leq l \leq n+1$. In this case

$$d_{\lambda} = \det \begin{pmatrix} d_{n+1-l} & d_{l-1} \\ d_{n-l} & d_l \end{pmatrix} = \det \begin{pmatrix} \binom{n}{n+1-l} & \binom{n}{l-1} \\ \binom{n}{n-l} & \binom{n}{l} \end{pmatrix} = 0.$$

Hence we may assume $m > 0$. Assume λ is almost $(m|n)$ -cross so $l(\lambda^{\bullet}), l(\lambda^{\circ}) \leq m+1$ and $\lambda_i^{\bullet} = n+1-\lambda_{m-i+2}^{\circ}$ for all $1 \leq i \leq m+1$. Then, by (35), d_{λ} is the determinant of the $(2m+2) \times (2m+2)$ block matrix $\begin{pmatrix} A & B \end{pmatrix}$ where

$$A = \begin{pmatrix} d_{\lambda_{m+1}^{\circ}} & d_{\lambda_m^{\circ}-1} & \cdots & d_{\lambda_1^{\circ}+m} \\ d_{\lambda_{m+1}^{\circ}-1} & d_{\lambda_m^{\circ}} & \cdots & d_{\lambda_1^{\circ}+m-1} \\ \vdots & \vdots & \ddots & \vdots \\ d_{\lambda_{m+1}^{\circ}-2m-1} & d_{\lambda_m^{\circ}-2m} & \cdots & d_{\lambda_1^{\circ}-m-1} \end{pmatrix},$$

$$B = \begin{pmatrix} d_{n-m-\lambda_{m+1}^{\circ}} & d_{n-m+1-\lambda_m^{\circ}} & \cdots & d_{n-2m-\lambda_1^{\circ}} \\ d_{n-m+1-\lambda_{m+1}^{\circ}} & d_{n-m-\lambda_m^{\circ}} & \cdots & d_{n-2m+1-\lambda_1^{\circ}} \\ \vdots & \vdots & \ddots & \vdots \\ d_{n+m+1-\lambda_{m+1}^{\circ}} & d_{n+m-\lambda_m^{\circ}} & \cdots & d_{n+1-\lambda_1^{\circ}} \end{pmatrix}.$$

Hence, if we let C denote the $(2m+2) \times (m+1)$ matrix whose j th column is given by $\text{Col}_j(C) = \text{Col}_j(A) + (-1)^{m-1} \text{Col}_j(B)$, then by Lemma 8.6.1 we have

$$C = \frac{1}{(m-1)!} \sum_{0 \leq i \leq n} \binom{n}{i} \begin{pmatrix} g\lambda_{m+1}^\circ(i) & g\lambda_m^\circ(i) & \cdots & g\lambda_1^\circ(i) \\ g\lambda_{m+1}^\circ(i-1) & g\lambda_m^\circ(i-1) & \cdots & g\lambda_1^\circ(i-1) \\ \vdots & \vdots & & \vdots \\ g\lambda_{m+1}^\circ(i-2m-1) & g\lambda_m^\circ(i-2m) & \cdots & g\lambda_1^\circ(i-m-1) \end{pmatrix}$$

where $g_k(u)$ is the polynomial defined in Lemma 8.6.1. To show $d_\lambda = 0$ we will verify that the columns of C (and hence of $(A \ B)$) are linearly dependent. To do so, notice $g_{k-l}(u) = g_k(u+l)$ for any $k, l \in \mathbb{Z}$. Hence, the j th column of C is

$$(39) \quad \text{Col}_j(C) = \frac{1}{(m-1)!} \sum_{0 \leq i \leq n} \binom{n}{i} \begin{pmatrix} g\lambda_{m+2-j}^\circ(i+j-1) \\ g\lambda_{m+2-j}^\circ(i+j-1) \\ \vdots \\ g\lambda_{m+2-j}^\circ(i+2m+1) \end{pmatrix}.$$

The $m+1$ polynomials $g\lambda_{m+2-j}^\circ(i+j-1)(u)$ for $1 \leq j \leq m+1$ are each of degree $m-1$ in the variable u . Hence, there exist $a_1, \dots, a_{m+1} \in \mathbb{K}$ (not all zero) with $\sum_{j=1}^{m+1} a_j g\lambda_{m+2-j}^\circ(i+j-1)(u) = 0$. It follows from (39) that $\sum_{j=1}^{m+1} a_j \text{Col}_j(C) = 0$. $\square$

Finally, we are in position to prove our criterion for the vanishing of $W(\lambda)$.

Theorem 8.7.6. $W(\lambda) \neq 0$ in $\text{Rep}(\mathfrak{gl}(m|n))$ if and only if λ is $(m|n)$ -cross.

Proof. One direction is Lemma 8.7.2. To prove the other, assume λ is not $(m|n)$ -cross. We will proceed by inducting on $|\lambda| = (r, s)$. The non- $(m|n)$ -cross bipartitions of minimal size are exactly the almost $(m|n)$ -cross bipartitions, hence the base case of our induction is Lemma 8.7.5. If λ is not almost $(m|n)$ -cross, then there exists a non- $(m|n)$ -cross bipartition μ which is obtained from λ by removing a box. If $\mu \vdash (r-1, s)$ then by (27) we have $\mu(\square, \emptyset) = \lambda + \mu^{(1)} + \cdots + \mu^{(k)} \in R_t$ for some bipartitions $\mu^{(1)}, \dots, \mu^{(k)}$ with $|\mu^{(i)}| < |\lambda|$ for all $1 \leq i \leq k$. Hence, by Theorem 6.2.3(2), $\mu(\square, \emptyset) = \lambda + \nu^{(1)} + \cdots + \nu^{(l)} \in R_{m-n}$ for some bipartitions $\nu^{(1)}, \dots, \nu^{(l)}$, which implies $W(\mu) \otimes W((\square, \emptyset)) = W(\lambda) \oplus W(\nu^{(1)}) \oplus \cdots \oplus W(\nu^{(l)})$ in $\text{Rep}(\mathfrak{gl}(m|n))$. By induction, we know $W(\mu) = 0$, hence $W(\lambda) = 0$ too. For the case $\mu \vdash (r, s-1)$, one argues similarly using (28). $\square$

Remark 8.7.7. As mentioned above, if $\lambda^\circ = \emptyset$ (resp. $\lambda^\bullet = \emptyset$), then λ being $(m|n)$ -cross is equivalent to $\lambda^\bullet$ (resp. λ°) being $(m|n)$ -hook. Hence Theorem 8.7.6 is a generalization of Theorem 8.7.1.

On the other hand, suppose $n = 0$ and set $m = d$ so that $F_{m|n}$ is identified with F_d (see §5.2). In this case λ is $(m|n)$ -cross if and only if $l(\lambda) \leq d$. Hence, Theorem 8.7.6 also generalizes the vanishing criterion in Theorem 5.2.2.

8.8. Decomposing $W(\lambda) \otimes W(\mu)$. Since $F_{m|n}$ is a tensor functor, Theorem 8.7.6 along with our method of decomposing arbitrary tensor products in $\underline{\text{Rep}}(GL_\delta)$ from §7.2 immediately allow us to decompose tensor products of the form $W(\lambda) \otimes W(\mu)$.

Example 8.8.1. From Example 7.2.2 with $\delta = 0$ we have the following decomposition in $\text{Rep}(\mathfrak{gl}(m|m))$:

$$W((2), \emptyset) \otimes W((\square, \square)) \cong W(((2, 1), \square)) \oplus W(((3), \square)) \oplus W(((1^2), \emptyset)) \oplus W(((2), \emptyset))^{\oplus 2}.$$

If $m > 1$, all the bipartitions in the decomposition above are $(m|m)$ -cross. Hence by Theorem 8.7.6 the summands in the decomposition above are all non-zero. On the other hand, $((2, 1), \square)$ is the only bipartition above which is not $(1|1)$ -cross. Hence by Theorem 8.7.6, we have the following decomposition into non-zero indecomposables in $\text{Rep}(\mathfrak{gl}(1|1))$:

$$W((2), \emptyset) \otimes W((\square, \square)) \cong W(((3), \square)) \oplus W(((1^2), \emptyset)) \oplus W(((2), \emptyset))^{\oplus 2}.$$

ACKNOWLEDGEMENTS

The majority of the work for this paper was done while the authors shared an office at the Technische Universität München. The authors would like to thank the university for providing them with the time and freedom to explore this project. We are also grateful to Maud De Visscher for explaining the results of [11] in the case $\delta = 0$. The second author would like to thank Victor Ostrik and Jon Brundan for many valuable conversations concerning this paper.

REFERENCES

1. F. W. Anderson and K. R. Fuller, *Rings and categories of modules*, Graduate texts in mathematics, Springer-Verlag, 1992. MR1245487 (94i:16001)
2. Georgia Benkart, Manish Chakrabarti, Thomas Halverson, Robert Leduc, Chanyoung Lee, and Jeffrey Stroomer, *Tensor product representations of general linear groups and their connections with Brauer algebras*, J. Algebra **166** (1994), no. 3, 529–567. MR1280591 (95d:20071)
3. D. J. Benson, *Representations and cohomology. I*, Cambridge Studies in Advanced Mathematics, vol. 30, Cambridge University Press, Cambridge, 1991, Basic representation theory of finite groups and associative algebras. MR1110581 (92m:20005)
4. A. Berele and A. Regev, *Hook Young diagrams with applications to combinatorics and to representations of Lie superalgebras*, Adv. in Math. **64** (1987), no. 2, 118–175. MR884183 (88i:20006)
5. J. Brundan and C. Stroppel, *Gradings on walled Brauer algebras and Khovanov’s arc algebra*, Advances in Mathematics **231** (2012), 709–773. MR2955190
6. ———, *Highest weight categories arising from Khovanov’s diagram algebra II: Koszulity*, Transform. Groups **15** (2010), no. 1, 1–45. MR2600694 (2011b:17014)
7. ———, *Highest weight categories arising from Khovanov’s diagram algebra I: cellularity*, Mosc. Math. J. **11** (2011), no. 4, 685–722, 821–822. MR2918294
8. ———, *Highest weight categories arising from Khovanov’s diagram algebra III: category $\mathcal{O}$* , Represent. Theory **15** (2011), 170–243. MR2781018
9. ———, *Highest weight categories arising from Khovanov’s diagram algebra IV: the general linear supergroup*, J. Eur. Math. Soc. (JEMS) **14** (2012), no. 2, 373–419. MR2881300
10. Jonathan Comes and Victor Ostrik, *On blocks of Deligne’s category $\text{Rep}(S_t)$* , Advances in Mathematics **226** (2011), no. 2, 1331–1377. MR2737787 (2012b:20020)
11. Anton Cox and Maud De Visscher, *Diagrammatic Kazhdan-Lusztig theory for the (walled) Brauer algebra*, Journal of Algebra **340** (2011), no. 1, 151–181. MR2813567
12. Anton Cox, Maud De Visscher, Stephen Doty, and Paul Martin, *On the blocks of the walled Brauer algebra*, J. Algebra **320** (2008), no. 1, 169–212. MR2417984 (2009c:16094)
13. C. J. Cummins and R. C. King, *Composite young diagrams, supercharacters of $u(m/n)$ and modification rules*, Journal of Physics A: Mathematical and General **20** (1987), no. 11, 3121. MR914044 (88m:22050)
14. P. Deligne, *La catégorie des représentations du groupe symétrique S_t , lorsque t n’est pas un entier naturel*, Algebraic groups and homogeneous spaces, Tata Inst. Fund. Res. Stud. Math., Tata Inst. Fund. Res., Mumbai, 2007, pp. 209–273. MR2348906
15. Pierre Deligne, *La série exceptionnelle de groupes de Lie*, C. R. Acad. Sci. Paris Sér. I Math. **322** (1996), no. 4, 321–326. MR1378507 (96m:22012)
16. William Fulton and Joe Harris, *Representation theory*, Graduate Texts in Mathematics, vol. 129, Springer-Verlag, New York, 1991, A first course, Readings in Mathematics. MR1153249 (93a:20069)

17. J. J. Graham and G. I. Lehrer, *Cellular algebras*, Invent. Math. **123** (1996), no. 1, 1–34. MR1376244 (97h:20016)
18. Kazuhiko Koike, *On the decomposition of tensor products of the representations of the classical groups: by means of the universal characters*, Adv. Math. **74** (1989), no. 1, 57–86. MR991410 (90j:22014)
19. I. G. Macdonald, *Symmetric functions and Hall polynomials*, second ed., Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1995, With contributions by A. Zelevinsky, Oxford Science Publications. MR1354144 (96h:05207)
20. Saunders MacLane, *Categories for the Working Mathematician*, Graduate Texts in Mathematics, 2nd ed., Springer, September 1998. MR1712872 (2001j:18001)
21. E. M. Moens and J. Van der Jeugt, *A determinantal formula for supersymmetric Schur polynomials*, J. Algebraic Combin. **17** (2003), no. 3, 283–307. MR2001672 (2004f:05187)
22. ———, *Composite supersymmetric s-functions and characters of $\mathfrak{gl}(m|n)$ representations*, Lie theory and its applications in physics VI, World Sci. Publ., River Edge, NJ, 2006. MR2172173 (2006g:17013)
23. P. Selinger, *A survey of graphical languages for monoidal categories*, New structures for physics, Lecture Notes in Phys., vol. 813, Springer, Heidelberg, 2011, pp. 289–355. MR2767048
24. Vera Serganova, *Kazhdan-Lusztig polynomials and character formula for the Lie superalgebra $\mathfrak{gl}(n|m)$* , Selecta Math. (N.S.) **2** (1996), no. 4, 607–651. MR1443186 (98f:17007)
25. A. N. Sergeev, *Representations of the Lie superalgebras $\mathfrak{gl}(n, m)$ and $Q(n)$ in a space of tensors*, Funktsional. Anal. i Prilozhen. **18** (1984), no. 1, 80–81. MR739101 (86b:17005)
26. John R. Stembridge, *Rational tableaux and the tensor algebra of $\mathfrak{gl}_n$* , Journal of Combinatorial Theory, Series A **46** (1987), no. 1, 79–120. MR899903 (89a:05012)
27. V. G. Turaev, *Operator invariants of tangles, and R-matrices*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 5, 1073–1107, 1135. MR1024455 (91e:17011)
28. H. Weyl, *The classical groups: their invariants and representations*, Princeton Landmarks in Mathematics and Physics, Princeton University Press, 1997. MR1488158 (98k:01049)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OREGON 97403
E-mail address: jcomes@uoregon.edu

DIEFFENBACHSTRASSE 27, 10967 BERLIN, GERMANY
E-mail address: benjamin@asmusas.net