

THE 2-BLOCKS OF DEFECT 4

BURKHARD KÜLSHAMMER AND BENJAMIN SAMBALE

ABSTRACT. We show that the major counting conjectures of modular representation theory are satisfied for 2-blocks of defect at most 4 except one possible case. In particular, we determine the invariants of such blocks.

1. INTRODUCTION

Let B be a 2-block of a finite group G with defect group D . Then there are several open conjectures regarding the number $k(B)$ of irreducible ordinary characters of B and the number $l(B)$ of irreducible Brauer characters of B . The aim of this paper is to show that most of these conjectures are fulfilled if D is small. More precisely, we will assume that B has defect at most 4, i.e., D has order at most 16. We denote the number of irreducible ordinary characters of height i in B by $k_i(B)$ for $i \geq 0$.

An essential tool is the following recent theorem by Kessar and Malle [17].

Theorem 1.1 (Kessar, Malle, 2011). *For every p -block B of a finite group with abelian defect group we have $k(B) = k_0(B)$.*

For $|D| \leq 8$ the block invariants and conjectures for B are known by the work of Brauer [5], Olsson [23] and Kessar, Koshitani and Linckelmann [16]. So we assume that D has order 16.

2. THE ELEMENTARY ABELIAN CASE

Let $I(B)$ be the inertial quotient of B and set $e(B) := |I(B)|$.

Proposition 2.1. *Let B be a block of a finite group G with elementary abelian defect group D of order 16. Then one of the following holds:*

- (i) B is nilpotent. Then $e(B) = l(B) = 1$ and $k(B) = k_0(B) = 16$.
- (ii) $e(B) = l(B) = 3$, $C_D(I(B)) = 1$ and $k(B) = k_0(B) = 8$.
- (iii) $e(B) = l(B) = 3$, $|C_D(I(B))| = 4$ and $k(B) = k_0(B) = 16$.
- (iv) $e(B) = l(B) = 5$ and $k(B) = k_0(B) = 8$.
- (v) $e(B) = l(B) = 7$ and $k(B) = k_0(B) = 16$.
- (vi) $e(B) = l(B) = 9$ and $k(B) = k_0(B) = 16$.
- (vii) $e(B) = 9$, $l(B) = 1$ and $k(B) = k_0(B) = 8$.
- (viii) $e(B) = l(B) = 15$ and $k(B) = k_0(B) = 16$.
- (ix) $e(B) = 15$, $l(B) = 7$ and $k(B) = k_0(B) = 8$.
- (x) $e(B) = 21$, $l(B) = 5$ and $k(B) = k_0(B) = 16$.

Moreover, all cases except possibly case (ix) actually occur.

Received by the editors February 7, 2012.

2010 *Mathematics Subject Classification.* Primary 20C15, 20C20.

Key words and phrases. 2-blocks, Alperin's weight conjecture, OWC.

Proof. First of all, by Theorem 1.1 we have $k(B) = k_0(B)$. The inertial quotient $I(B)$ is a subgroup of $\text{Aut}(D) \cong \text{GL}(4, 2)$ of odd order. It follows that $e(B) \in \{1, 3, 5, 7, 9, 15, 21\}$ (this can be shown with GAP [13]). If $e(B) \neq 21$, the inertial quotient is necessarily abelian. Then by Corollary 1.2(ii) in [29] there is a nontrivial subsection (u, b) such that $l(b) = 1$. Hence, Corollary 2 in [6] implies that $|D| = 16$ is a sum of $k(B)$ odd squares. This shows $k(B) \in \{8, 16\}$ for these cases. In order to determine $l(B)$ we calculate the numbers $l(b)$ for all nontrivial subsections (u, b) . Here it suffices to consider a set of representatives of the orbits of D under $I(B)$, since B is a controlled block. If $e(B) = 1$, the block is nilpotent and the result is clear. We discuss the remaining cases separately:

Case 1 ($e(B) = 3$). Here by results of Usami and Puig (see [28, 40]) there is a perfect isometry between B and its Brauer correspondent in $N_G(D)$. According to two different actions of $I(B)$ on D , we get $k(B) = 8$ if $C_D(I(B)) = 1$ or $k(B) = 16$ if $|C_D(I(B))| = 4$. In both cases we have $l(B) = 3$.

Case 2 ($e(B) = 5$). Then there are four subsections $(1, B)$, (u_1, b_1) , (u_2, b_2) and (u_3, b_3) with $l(b_1) = l(b_2) = l(b_3) = 1$ up to conjugation. In [37] it was shown that $k(B) = 16$ is impossible. Hence, $k(B) = 8$ and $l(B) = 5$.

Case 3 ($e(B) = 7$). There are again four subsections $(1, B)$, (u_1, b_1) , (u_2, b_2) and (u_3, b_3) up to conjugation. But in this case $l(b_1) = l(b_2) = 1$ and $l(b_3) = 7$ by [16]. Thus, $k(B) = 16$ and $l(B) = 7$.

Case 4 ($e(B) = 9$). There are four subsections $(1, B)$, (u_1, b_1) , (u_2, b_2) and (u_3, b_3) such that $l(b_1) = 1$ and $l(b_2) = l(b_3) = 3$ up to conjugation. This gives the possibilities (vi) and (vii).

Case 5 ($e(B) = 15$). Here $I(B)$ acts regularly on $D \setminus \{1\}$. Thus, there are only two subsections $(1, B)$ and (u, b) such that $l(b) = 1$. This gives the possibilities (viii) and (ix).

Case 6 ($e(B) = 21$). Here $I(B)$ is nonabelian. Hence, we get four subsections $(1, B)$, (u_1, b_1) , (u_2, b_2) and (u_3, b_3) up to conjugation. We have $l(b_1) = l(b_2) = 3$ and $l(b_3) = 5$ by [16]. Since $I(B)$ has a fixed point on D , it follows that $l(B) = 5$ and $k(B) = 16$ by Theorem 1 in [45].

For all cases except (vii) and (ix) examples are given by the principal block of $D \rtimes I(B)$. In case (vii) we can take a nonprincipal block of the group $\text{SmallGroup}(432, 526) \cong D \rtimes E$ where E is the extraspecial group of order 27 and exponent 3 (see “small groups library”). \square

We will see later that case (ix) would contradict Alperin’s Weight Conjecture. Now we investigate the differences between the cases (vi) and (vii).

Lemma 2.2. *Let B be a block of a finite group G with elementary abelian defect group D of order 16. If $e(B) = l(B) = 9$, then the elementary divisors of the Cartan matrix of B are 1, 1, 1, 1, 4, 4, 4, 4, 16. Moreover, the two $I(B)$ -stable subgroups of D of order 4 are lower defect groups of B . Both occur with 1-multiplicity 2.*

Proof. Let C be the Cartan matrix of B . As in the proof of Proposition 2.1 there are four subsections $(1, B)$, (u_1, b_1) , (u_2, b_2) and (u_3, b_3) such that $l(b_1) = 1$ and $l(b_2) = l(b_3) = 3$ up to conjugation. In order to determine C up to basic sets, we need to investigate the generalized decomposition numbers $d_{rs}^{u_i}$ for $i = 1, 2, 3$. The

block b_2 dominates a block $\overline{b_2}$ of $C_G(u_2)/\langle u_2 \rangle$ with defect group $D/\langle u_2 \rangle$ and inertial index 3. Thus, as in the proof of Theorem 3 in [36] the Cartan matrix of b_2 has the form

$$\begin{pmatrix} 8 & 4 & 4 \\ 4 & 8 & 4 \\ 4 & 4 & 8 \end{pmatrix}$$

up to basic sets. Since $k(B) = 16$, we may assume that the numbers $d_{rs}^{u_2}$ take the form

$$\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & . & . & . & . & . & . & . & . \\ 1 & 1 & 1 & 1 & . & . & . & . & 1 & 1 & 1 & 1 & . & . & . & . \\ 1 & 1 & 1 & 1 & . & . & . & . & . & . & . & . & 1 & 1 & 1 & 1 \end{pmatrix}^T.$$

For the column of decomposition numbers $d_{rs}^{u_1}$ we have essentially the following possibilities:

$$\begin{aligned} (A) &: (1, 1, 1, 1, -1, -1, -1, -1, -1, -1, -1, -1, -1, -1, -1, -1)^T, \\ (B) &: (1, 1, 1, -1, 1, -1, -1, -1, 1, -1, -1, -1, 1, -1, -1, -1)^T, \\ (C) &: (1, 1, -1, -1, 1, 1, -1, -1, 1, 1, -1, -1, 1, 1, -1, -1)^T. \end{aligned}$$

Now we use a GAP program to enumerate the possible decomposition numbers $d_{rs}^{u_3}$. After that the ordinary decomposition matrix M can be calculated as the orthogonal space. Then $C = M^T M$ up to basic sets. It turns out that in some cases C has 2 as an elementary divisor. Using the notion of lower defect groups as described in [24] we show that these cases cannot occur. If 2 is an elementary divisor of C , then there exists a lower defect group $Q \leq D$ of order 2. With the notation of [24] we have $m_B^{(1)}(Q) > 0$. By Theorem 7.2 in [24] there is a block b_Q of $N_G(Q) = C_G(Q)$ such that $b_Q^G = B$ and $m_{b_Q}^{(1)}(Q) > 0$. In particular, the Cartan matrix of b_Q has 2 as elementary divisor. Hence, b_Q is conjugate to b_2 or b_3 . But we have seen above that all elementary divisors of the Cartan matrix of b_2 (and also b_3) must be divisible by 4. This contradiction shows that 2 does not occur as elementary divisor of C . After excluding these cases the GAP program reveals the following two possibilities for the elementary divisors of C : 1, 1, 1, 1, 4, 4, 4, 4, 16 or 1, 1, 4, 4, 4, 4, 4, 4, 16.

Now we have to look at the lower defect group multiplicities more carefully. The calculation above and (7G) in [4] imply

$$4 \leq \sum_{R \in \mathcal{R}} m_B^{(1)}(R)$$

where \mathcal{R} is a set of representatives for the G -conjugacy classes of subgroups of G of order 4. After combining this with the formula (2S) of [7] we get

$$4 \leq \sum_{(R, b_R) \in \mathcal{R}'} m_B^{(1)}(R, b_R)$$

where \mathcal{R}' is a set of representatives for the G -conjugacy classes of B -subpairs (R, b_R) such that R has order 4. Let b_D be a Brauer correspondent of B in $C_G(D)$. Then, after changing the representatives if necessary we may assume $(R, b_R) \leq (D, b_D)$ for $(R, b_R) \in \mathcal{R}'$. Then it is well known that $b_R = b_D^{C_G(R)}$ is uniquely determined

by R . Since the fusion of these subpairs is controlled by $N_G(D, b_D)$, we get

$$4 \leq \sum_{R \in \mathcal{R}''} m_B^{(1)}(R, b_R)$$

where \mathcal{R}'' is a set of representatives for the $I(B)$ -conjugacy classes of subgroups of D of order 4.

Now let $Q \leq D$ of order 4 such that $m_B^{(1)}(Q, b_Q) > 0$. Then by (2Q) in [7] we have $m_{B_Q}^{(1)}(Q) > 0$ where $B_Q := b_Q^{N_G(Q, b_Q)}$. If Q is not fixed under $I(B)$, then we would have the contradiction $e(B_Q) = l(B_Q) = 1$. Thus, we have shown that Q is stable under $I(B)$. Hence,

$$(1) \quad 4 \leq m_B^{(1)}(Q, b_Q) + m_B^{(1)}(P, b_P)$$

where $P \neq Q$ is the other $I(B)$ -stable subgroup of D of order 4. Since 16 is always an elementary divisor of C , we have $m_{B_Q}^{(1)}(D) = 1$. Observe that b_Q has defect group D and inertial index 3, so that $l(b_Q) = 3$ by Proposition 2.1. Now Theorem 5.11 in [24] and the remark following it imply

$$3 = l(b_Q) \geq m_{B_Q}^{(1)}(Q) + m_{B_Q}^{(1)}(D).$$

(Notice that in Theorem 5.11, loc. cit., it should read $B \in \text{Bl}(G)$ instead of $B \in \text{Bl}(Q)$.) Thus, $m_{B_Q}^{(1)}(Q) \leq 2$ and similarly $m_{B_P}^{(1)}(P) \leq 2$. Now equation (1) yields $m_B^{(1)}(Q, b_Q) = m_B^{(1)}(P, b_P) = 2$. In particular, 4 occurs as elementary divisor of C with multiplicity 4. It is easy to see that we also have $m_B^{(1)}(Q) = m_B^{(1)}(P) = 2$ which proves the last claim. \square

Proposition 2.3. *Let B be a block of a finite group G with elementary abelian defect group D of order 16. If $e(B) = 9$, then Alperin's Weight Conjecture holds for B .*

Proof. Let b_D be a Brauer correspondent of B in $C_G(D)$, and let B_D be the Brauer correspondent of B in $N_G(D, b_D)$. Then it suffices to show that $l(B) = l(B_D)$. By Proposition 2.1 we have to consider two cases $l(B) \in \{1, 9\}$. As in Lemma 2.2 we set $b_R := b_D^{C_G(R)}$ for $R \leq D$.

We start with the assumption $l(B) = 9$. Then by Lemma 2.2 there is an $I(B)$ -stable subgroup $Q \leq D$ of order 4 such that $m_{B_Q}^{(1)}(Q) = m_B^{(1)}(Q, b_Q) > 0$ where $B_Q := b_Q^{N_G(Q, b_Q)}$. In particular, $l(B_Q) = 9$. Let $P \leq D$ be the other $I(B)$ -stable subgroup of order 4. Moreover, let $b'_P := b_D^{N_G(Q, b_Q) \cap C_G(P)}$ such that (P, b'_P) is a B_Q -subpair. Then by the same argument we get

$$m_\beta^{(1)}(P) = m_{B_Q}^{(1)}(P, b'_P) > 0$$

where $\beta := (b'_P)^{N_G(Q, b_Q) \cap N_G(P, b'_P)}$ is a block with defect group D and $l(\beta) = 9$. Now $D = QP$ implies

$$N_G(D, b_D) \leq N_G(Q, b_Q) \cap N_G(P, b'_P) \leq N_G(D).$$

Since $B_D^{N_G(Q, b_Q) \cap N_G(P, b'_P)} = \beta$, it follows that $l(B_D) = 9$ as desired.

Now let us consider the case $l(B) = 1$. Here we can just follow the same lines except that we have $m_{B_Q}^{(1)}(Q) = 0$ and $m_\beta^{(1)}(P) = 0$. \square

We want to point out that Usami showed in [42] that in case $2 \neq p \neq 7$ there is a perfect isometry between a p -block with abelian defect group D and inertial index 9 and its Brauer correspondent in $N_G(D)$.

3. THE ORDINARY WEIGHT CONJECTURE

For most 2-blocks of defect 4, Robinson's Ordinary Weight Conjecture (OWC) [30] is known to hold. In this section we handle the remaining cases.

Proposition 3.1. *Let B be a block of a finite group G with minimal nonabelian defect group*

$$D := \langle x, y \mid x^{2^r} = y^2 = [x, y]^2 = [x, x, y] = [y, x, y] = 1 \rangle$$

of order $2^{r+2} \geq 16$. Then the Ordinary Weight Conjecture holds for B .

Proof. The block invariants of B were determined and several conjectures were verified in [35]. In order to prove the OWC we use the version in Conjecture 6.5 in [15]. Let \mathcal{F} be the fusion system of B . We may assume that \mathcal{F} is nonnilpotent. Let $z := [x, y]$. Then it was shown in [35] that $Q := \langle x^2, y, z \rangle \cong C_{2^{r-1}} \times C_2^2$ and D are the only \mathcal{F} -centric and \mathcal{F} -radical subgroups of D . Moreover, $\text{Out}_{\mathcal{F}}(Q) = \text{Aut}_{\mathcal{F}}(Q) \cong S_3$ and $\text{Out}_{\mathcal{F}}(D) = 1$. Hence, it follows easily that $\mathbf{w}(D, d) = k^d(D) = k^d(B)$ for all $d \in \mathbb{N}$ where $k^d(D)$ is the number of characters of defect d in D . Thus, it suffices to show $\mathbf{w}(Q, d) = 0$ for all $d \in \mathbb{N}$ by Theorem 3.6 in [35]. Since Q is abelian, we have $\mathbf{w}(Q, d) = 0$ unless $d = r + 1$. Thus, let $d = r + 1$. Up to conjugation \mathcal{N}_Q consists of the trivial chain $\sigma : 1$ and the chain $\tau : 1 < C$, where $C \leq \text{Out}_{\mathcal{F}}(Q)$ has order 2. We consider the chain σ first. Here $I(\sigma) = \text{Out}_{\mathcal{F}}(Q) \cong S_3$ acts faithfully on $\Omega(Q) \cong C_2^3$ and thus fixes a four-group. Hence, the characters in $\text{Irr}(Q)$ split in 2^{r-1} orbits of length 3 and 2^{r-1} orbits of length 1 under $I(\sigma)$. For a character $\chi \in \text{Irr}(D)$ lying in an orbit of length 3 we have $I(\sigma, \chi) \cong C_2$ and thus $w(Q, \sigma, \chi) = 0$. For the 2^{r-1} stable characters $\chi \in \text{Irr}(D)$ we get $w(Q, \sigma, \chi) = 1$, since $I(\sigma, \chi) = \text{Out}_{\mathcal{F}}(Q)$ has precisely one block of defect 0.

Now consider the chain τ . Here $I(\tau) = C$ and the characters in $\text{Irr}(Q)$ split in 2^{r-1} orbits of length 2 and 2^r orbits of length 1 under $I(\tau)$. For a character $\chi \in \text{Irr}(D)$ in an orbit of length 2 we have $I(\tau, \chi) = 1$ and thus $w(Q, \tau, \chi) = 1$. For the 2^r stable characters $\chi \in \text{Irr}(D)$ we get $I(\tau, \chi) = I(\tau) = C$ and $w(Q, \tau, \chi) = 0$.

Taking both chains together, we derive

$$\mathbf{w}(Q, d) = (-1)^{|\sigma|+1} 2^{r-1} + (-1)^{|\tau|+1} 2^{r-1} = 2^{r-1} - 2^{r-1} = 0.$$

This proves the OWC. \square

Now we consider the OWC for blocks with abelian defect groups D of order 2^d . Here of course D is the only \mathcal{F} -centric and \mathcal{F} -radical subgroup of D and $I(B) = \text{Out}_{\mathcal{F}}(D)$ has odd order. In particular, \mathcal{N}_D consists only of the trivial chain. Moreover, $\mathbf{w}(D, d') = 0$ unless $d' = d$. If we assume in addition that $I(B)$ is cyclic, then

$$(2) \quad \mathbf{w}(D, d) = \sum_{\chi \in \text{Irr}(D)/I(B)} |I(B) \cap I(\chi)|$$

where $I(B) \cap I(\chi) := \{\alpha \in I(B) : {}^\alpha \chi = \chi\}$. In connection with Theorem 1.1, the OWC predicts $k(B) = \mathbf{w}(D, d)$.

Now let us consider the case where D is elementary abelian of order 16. Then if $21 \neq e(B) \neq 9$, the OWC follows easily from Proposition 2.1 and equation (2)

except if case (ix) occurs (where OWC does not hold). Now assume $e(B) = 21$. Here the number of 2-blocks of defect 0 in $I(B)$ (which is denoted by $z(kI(B))$ in [15] where k is an algebraically closed field of characteristic 2) is 5. We have to insert this number for $|I(B) \cap I(\chi)|$ in equation (2) if χ is invariant under $I(B)$. Now the OWC also follows in this case. We will deal with the remaining case $e(B) = 9$ in the next section.

4. THE GENERAL CASE

Theorem 4.1. *Let B be a 2-block of a finite group G with defect group D of order at most 16. Then one of the following holds:*

(i) *The following conjectures are satisfied for B :*

- *Alperin's Weight Conjecture* [2],
- *Brauer's $k(B)$ -Conjecture* [3],
- *Brauer's Height-Zero Conjecture* [3],
- *Olsson's Conjecture* [25],
- *Alperin-McKay Conjecture* [1],
- *Robinson's Ordinary Weight Conjecture* [30],
- *Gluck's Conjecture* [14],
- *Eaton's Conjecture* [9],
- *Eaton-Moretó Conjecture* [11],
- *Malle-Navarro Conjecture* [22].

Moreover, the Gluing Problem [21] *for B has a unique solution.*

(ii) $D \cong C_2^4$, $e(B) = 15$, $k(B) = k_0(B) = 8$, $l(B) = 7$ and $D \notin \text{Syl}_2(G)$. The Cartan matrix of B is given by

$$\begin{pmatrix} 6 & 5 & 5 & 5 & 5 & 5 & 7 \\ 5 & 6 & 5 & 5 & 5 & 5 & 7 \\ 5 & 5 & 6 & 5 & 5 & 5 & 7 \\ 5 & 5 & 5 & 6 & 5 & 5 & 7 \\ 5 & 5 & 5 & 5 & 6 & 5 & 7 \\ 5 & 5 & 5 & 5 & 5 & 6 & 7 \\ 7 & 7 & 7 & 7 & 7 & 7 & 10 \end{pmatrix}$$

up to basic sets. Alperin's Weight Conjecture and the Alperin-McKay Conjecture are not satisfied for B .

Proof. As explained earlier we may assume that $|D| = 16$. Then the situation splits in the following possibilities:

- (a) D is metacyclic,
- (b) D is minimal nonabelian,
- (c) D is abelian, but nonmetacyclic,
- (d) $D \cong D_8 \times C_2$,
- (e) $D \cong Q_8 \times C_2$,
- (f) $D \cong D_8 * C_4$.

We start with a remark about Gluck's Conjecture which only applies to rational defect groups of nilpotency class at most 2. By Corollary 3.2 and Lemma 2.1 in [14] we may assume that D is nonabelian of exponent 4 in order to prove Gluck's Conjecture. Moreover, Gluck's Conjecture is satisfied for nilpotent blocks.

In case (a) the block invariants are known by [5], [23], [39]. From this most of the conjectures follow trivially. Observe here that the nonabelian metacyclic groups of

exponent 4 provide only nilpotent blocks. In particular, Gluck's Conjecture follows. For the OWC we refer to [32] and for the Gluing Problem to [26].

In case (b), D has the form $D \cong \langle x, y \mid x^4 = y^2 = [x, y]^2 = [x, x, y] = [y, x, y] = 1 \rangle$; in particular, D is not rational. Then the result follows from [35] (for the OWC see Proposition 3.1). Again we skip the elementary details for the three (less-known) conjectures in Theorem 4.1. The last three cases (d), (e) and (f) were handled in [31, 32, 38] (for Gluck's Conjecture see [37]). It remains to consider case (c). Here it is known that the Gluing Problem has a unique solution (see [21]). We have two possibilities: $D \cong C_4 \times C_2 \times C_2$ or D is elementary abelian. We may assume that B is nonnilpotent.

In case $D \cong C_4 \times C_2 \times C_2$, 3 is the only odd prime divisor of $|\text{Aut}(D)|$. Thus, by Usami and Puig (see [28], [40]) there is a perfect isometry between B and its Brauer correspondent in $N_G(D)$. Then it is easy to see that the conjectures are true.

Now we consider the elementary abelian case. By Proposition 2.1, Brauer's $k(B)$ -Conjecture, Brauer's Height-Zero Conjecture, Olsson's Conjecture, Eaton's Conjecture, the Eaton-Moretó Conjecture and the Malle-Navarro Conjecture are satisfied. For abelian defect groups, Alperin's Weight Conjecture is equivalent to $l(B) = l(b)$ where b is the Brauer correspondent of B in $N_G(D)$. For $e(B) = 9$ this was shown in Proposition 2.3. Thus, assume $e(B) \neq 9$. By the main result in [20], b is Morita equivalent to a twisted group algebra of $D \rtimes I(B)$. Since $e(B) \neq 9$, the corresponding 2-cocycle must be trivial so that b is Morita equivalent to the group algebra of $D \rtimes I(B)$. This gives $l(b) = k(I(B))$. Now it can be seen that Alperin's Weight Conjecture holds unless case (ix) in Proposition 2.1 occurs.

Since $k(B) - l(B) = k_0(B) - l(B)$ is determined locally, the Alperin-McKay Conjecture follows from Alperin's Weight Conjecture. Now consider the Ordinary Weight Conjecture. By the remarks in the last section it suffices to look at the case $e(B) = 9$. Here again b is Morita equivalent to a twisted group algebra of $D \rtimes I(B)$. If the corresponding 2-cocycle α is trivial we have $l(B) = 9$ and $l(B) = 1$ otherwise. Then with the notation in [15] we have $z(k_\alpha I(B)) = 9$ or $z(k_\alpha I(B)) = 1$, respectively. Now the OWC follows as in the last section.

Now we consider the situation $e(B) = 15$, $k(B) = k_0(B) = 8$ and $l(B) = 7$ more closely. The arguments above imply that Alperin's Weight Conjecture and thus also the Alperin-McKay Conjecture are not fulfilled. In particular, G is nonsolvable. The Cartan matrix C of B can be determined as in [37]. Here observe that $\det C = 16 = |D|$ a fact which is also predicted by Corollary 1 in [12].

Assume that $D \in \text{Syl}_2(G)$. We use the rest of the proof to derive a contradiction. By the first Fong reduction we may assume that B is quasiprimitive, i.e., that, for any normal subgroup N of G , B covers a unique block B_N of N . Note that $D \cap N$ is a Sylow 2-subgroup of N and a defect group of B_N .

Suppose now that $N = O(G)$. Then, by the second Fong reduction there exist a finite group G^* with a cyclic central subgroup N^* of odd order such that G^*/N^* is isomorphic to G/N , and a block B^* of G^* whose defect group D^* is isomorphic to D ; moreover, B^* is Morita equivalent to B ; in particular, we have $k(B^*) = k(B) = 8$ and $l(B^*) = l(B) = 7$.

Thus, Proposition 2.1 implies that $e(B^*) = 15$ as well, so that G^* , B^* is also a counterexample. So we may assume that $G = G^*$ and $B = B^*$. Then N is a central cyclic subgroup of odd order in G .

Let M/N be a minimal normal subgroup of G/N . Then $D \cap M$ is a Sylow 2-subgroup of M ; in particular, $D \cap M \neq 1$. Then $D \cap M$ is stable under the inertial subgroup $N_G(D, b)$ of B . Since $N_G(D, b)$ acts transitively on $D \setminus \{1\}$, we must have $D = D \cap M \subseteq M$. Thus M/N is the only minimal normal subgroup of G/N , and $|G : M|$ is odd.

If M/N is abelian, then $M = D \times N$; in particular, B has a normal defect group. But this is impossible since G, B is a counterexample.

Hence M/N is a direct product of isomorphic nonabelian finite simple groups which are transitively permuted under conjugation in G :

$$M/N = S_1/N \times \dots \times S_t/N.$$

Thus $D = (D \cap S_1) \times \dots \times (D \cap S_t)$ with isomorphic factors. Since $|D| = 2^4$, we must have $t = 1$, $t = 2$ or $t = 4$. Since $|G : M|$ is odd, this implies that $t = 1$. Hence M/N is a simple group with Sylow 2-subgroup D . By Walter's Theorem (see [44]), we must have $M/N = \text{PSL}(2, 16)$. Note also that $M = F^*(G)$. Since $\text{PSL}(2, 16)$ has a trivial Schur multiplier and an outer automorphism group of order 4, we conclude that $G = M = \text{PSL}(2, 16) \times N$. We may therefore clearly assume that $N = 1$. In this case B is the principal 2-block of $\text{PSL}(2, 16)$, and $l(B) = 15$, a final contradiction. \square

We remark that even more information about 2-blocks of defect 4 can be extracted from [37]. For example Cartan matrices and the number of 2-rational and 2-conjugate characters of these blocks are known in most cases.

5. INVARIANTS OF BLOCKS

In this section we give an overview in which cases the block invariants of p -blocks for arbitrary primes p are known. It should be pointed out that many p -groups provide only nilpotent fusion systems. For such defect groups all block invariants are known, and we will omit these cases. The extraspecial group of order p^3 and exponent p^2 for an odd prime p is denoted by p_-^{1+2} . More generally, let M_{p^n} be the (unique) nonabelian group of order p^n with exponent p^{n-1} .

p	D	$I(B)$	classification used?	references
arbitrary	cyclic	arbitrary	no	[8]
arbitrary	abelian	$e(B) \leq 4$	no	[27, 28, 40]
arbitrary	abelian	S_3	no	[41]
≥ 7	abelian	$C_4 \times C_2$	no	[43]
$\notin \{2, 7\}$	abelian	C_3^2	no	[42]
2	metacyclic	arbitrary	no	[5, 23, 28, 39]
2	maximal class * cyclic, incl. * = \times	arbitrary	only for $D \cong C_2^3$	[16, 31, 32, 38]
2	minimal nonabelian	arbitrary	only for one family where $ D = 2^{2r+1}$	[10, 35]
2	minimal nonmetacyclic	arbitrary	only for $D \cong C_2^3$	[37]
2	$ D \leq 16$	$\not\cong C_{15}$	yes	this paper
2	$C_4 \wr C_2$	arbitrary	no	[19]
2	$D_8 * Q_8$	C_5	yes	[34]
2	$C_{2^n} \times C_2^3, n \geq 2$	arbitrary	yes	[34]
3	C_3^2	$\notin \{C_8, Q_8\}$	no	[18, 46]
3, 5, 7, 11	p_-^{1+2}	$e(B) \leq 2$	no	[33]
3	M_{81}	arbitrary	no	[33]

ACKNOWLEDGMENTS

The second author is supported by the German Academic Exchange Service (DAAD). The second author is grateful to the University of California for their hospitality.

REFERENCES

- [1] J. L. Alperin, *The main problem of block theory*, Proceedings of the Conference on Finite Groups (Univ. Utah, Park City, Utah, 1975), Academic Press, New York, 1976, pp. 341–356. MR0404417 (53 #8219)
- [2] J. L. Alperin, *Weights for finite groups*, The Arcata Conference on Representations of Finite Groups (Arcata, Calif., 1986), Proc. Sympos. Pure Math., vol. 47, Amer. Math. Soc., Providence, RI, 1987, pp. 369–379. MR933373 (89h:20015)
- [3] Richard Brauer, *Representations of finite groups*, Lectures on Modern Mathematics, Vol. I, Wiley, New York, 1963, pp. 133–175. MR0178056 (31 #2314)
- [4] Richard Brauer, *Defect groups in the theory of representations of finite groups*, Illinois J. Math. **13** (1969), 53–73. MR0246979 (40 #248)
- [5] Richard Brauer, *On 2-blocks with dihedral defect groups*, Symposia Mathematica, Vol. XIII (Convegno di Gruppi e loro Rappresentazioni, INDAM, Rome, 1972), Academic Press, London, 1974, pp. 367–393. MR0354838 (50 #7315)
- [6] Michel Broué, *On characters of height zero*, The Santa Cruz Conference on Finite Groups (Univ. California, Santa Cruz, Calif., 1979), Proc. Sympos. Pure Math., vol. 37, Amer. Math. Soc., Providence, R.I., 1980, pp. 393–396. MR604610 (82g:20021)
- [7] Michel Broué and Jörn B. Olsson, *Subpair multiplicities in finite groups*, J. Reine Angew. Math. **371** (1986), 125–143. MR859323 (87j:20026)
- [8] E. C. Dade, *Blocks with cyclic defect groups*, Ann. of Math. (2) **84** (1966), 20–48. MR0200355 (34 #251)
- [9] Charles W. Eaton, *Generalisations of conjectures of Brauer and Olsson*, Arch. Math. (Basel) **81** (2003), no. 6, 621–626, DOI 10.1007/s00013-003-0832-y. MR2029237 (2004i:20012)
- [10] Charles W. Eaton, Burkhard Külshammer, and Benjamin Sambale, *2-blocks with minimal nonabelian defect groups II*, J. Group Theory **15** (2012), no. 3, 311–321, DOI 10.1515/jgt.2011.110. MR2920888
- [11] Charles W. Eaton and A. Moretó, *Extending Brauer’s Height Zero Conjecture to blocks with nonabelian defect groups*, preprint.
- [12] Mitsuo Fujii, *On determinants of Cartan matrices of p -blocks*, Proc. Japan Acad. Ser. A Math. Sci. **56** (1980), no. 8, 401–403. MR596014 (81m:20014)
- [13] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.4.12*; 2008, (<http://www.gap-system.org>).
- [14] David Gluck, *Rational defect groups and 2-rational characters*, J. Group Theory **14** (2011), no. 3, 401–412, DOI 10.1515/JGT.2010.063. MR2794375 (2012e:20012)
- [15] Radha Kessar, *Introduction to block theory*, Group representation theory, EPFL Press, Lausanne, 2007, pp. 47–77. MR2336637 (2008f:20020)
- [16] R. Kessar, S. Koshitani and M. Linckelmann, *Conjectures of Alperin and Broué for 2-blocks with elementary abelian defect groups of order 8*, J. Reine Angew. Math. **671** (2012), 85–130.
- [17] R. Kessar and G. Malle, *Quasi-isolated blocks and Brauer’s height zero conjecture*, arXiv:1112.2642v1.
- [18] Masao Kiyota, *On 3-blocks with an elementary abelian defect group of order 9*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **31** (1984), no. 1, 33–58. MR743518 (85k:20036)
- [19] Burkhard Külshammer, *On 2-blocks with wreathed defect groups*, J. Algebra **64** (1980), no. 2, 529–555, DOI 10.1016/0021-8693(80)90157-X. MR579072 (81i:20008)
- [20] Burkhard Külshammer, *Crossed products and blocks with normal defect groups*, Comm. Algebra **13** (1985), no. 1, 147–168, DOI 10.1080/00927878508823154. MR768092 (86c:20015)
- [21] Markus Linckelmann, *Fusion category algebras*, J. Algebra **277** (2004), no. 1, 222–235, DOI 10.1016/j.jalgebra.2003.12.010. MR2059628 (2005a:20017)
- [22] Gunter Malle and Gabriel Navarro, *Inequalities for some blocks of finite groups*, Arch. Math. (Basel) **87** (2006), no. 5, 390–399, DOI 10.1007/s00013-006-1769-8. MR2269921 (2007i:20073)

- [23] Jørn Børling Olsson, *On 2-blocks with quaternion and quasidihedral defect groups*, J. Algebra **36** (1975), no. 2, 212–241. MR0376841 (51 #13016)
- [24] Jørn B. Olsson, *Lower defect groups*, Comm. Algebra **8** (1980), no. 3, 261–288, DOI 10.1080/00927878008822458. MR558114 (81g:20024)
- [25] Jørn B. Olsson, *On the number of characters in blocks of finite general linear, unitary and symmetric groups*, Math. Z. **186** (1984), no. 1, 41–47, DOI 10.1007/BF01215489. MR735049 (85d:20008)
- [26] Sejong Park, *The gluing problem for some block fusion systems*, J. Algebra **323** (2010), no. 6, 1690–1697, DOI 10.1016/j.jalgebra.2010.01.003. MR2588132 (2011b:20027)
- [27] Lluís Puig and Yoko Usami, *Perfect isometries for blocks with abelian defect groups and Klein four inertial quotients*, J. Algebra **160** (1993), no. 1, 192–225, DOI 10.1006/jabr.1993.1184. MR1237084 (94g:20017)
- [28] Lluís Puig and Yoko Usami, *Perfect isometries for blocks with abelian defect groups and cyclic inertial quotients of order 4*, J. Algebra **172** (1995), no. 1, 205–213, DOI 10.1006/jabr.1995.1059. MR1320630 (96d:20012)
- [29] Geoffrey R. Robinson, *On Brauer’s $k(B)$ problem*, J. Algebra **147** (1992), no. 2, 450–455, DOI 10.1016/0021-8693(92)90215-8. MR1161303 (93b:20020)
- [30] Geoffrey R. Robinson, *Weight conjectures for ordinary characters*, J. Algebra **276** (2004), no. 2, 761–775, DOI 10.1016/j.jalgebra.2004.01.019. MR2058466 (2005c:20021)
- [31] B. Sambale, *Blocks with central product defect group $D_{2^n} * C_{2^m}$* , Proc. Amer. Math. Soc. (to appear), arXiv:1105.4977v1.
- [32] B. Sambale, *Blocks with defect group $Q_{2^n} \times C_{2^m}$ and $SD_{2^n} \times C_{2^m}$* , Algebr. Represent. Theory (to appear), DOI:10.1007/s10468-012-9379-6.
- [33] B. Sambale, *Brauer’s Height Zero Conjecture for metacyclic defect groups*, Pacific J. Math. (to appear), arXiv:1204.6651v1.
- [34] B. Sambale, *Further evidence for conjectures in block theory*, submitted.
- [35] Benjamin Sambale, *2-blocks with minimal nonabelian defect groups*, J. Algebra **337** (2011), 261–284, DOI 10.1016/j.jalgebra.2011.02.006. MR2796075 (2012d:20019)
- [36] Benjamin Sambale, *Cartan matrices and Brauer’s $k(B)$ -conjecture*, J. Algebra **331** (2011), 416–427, DOI 10.1016/j.jalgebra.2010.10.036. MR2774667 (2012c:20023)
- [37] Benjamin Sambale, *Cartan matrices and Brauer’s $k(B)$ -conjecture II*, J. Algebra **337** (2011), 345–362, DOI 10.1016/j.jalgebra.2011.03.023. MR2796080 (2012e:20018)
- [38] Benjamin Sambale, *Blocks with defect group $D_{2^n} \times C_{2^m}$* , J. Pure Appl. Algebra **216** (2012), no. 1, 119–125, DOI 10.1016/j.jpaa.2011.05.009. MR2826425 (2012f:20031)
- [39] Benjamin Sambale, *Fusion systems on metacyclic 2-groups*, Osaka J. Math. **49** (2012), no. 2, 325–329. MR2945751
- [40] Yoko Usami, *On p -blocks with abelian defect groups and inertial index 2 or 3. I*, J. Algebra **119** (1988), no. 1, 123–146, DOI 10.1016/0021-8693(88)90079-8. MR971349 (89i:20024)
- [41] Yoko Usami, *Perfect isometries for blocks with abelian defect groups and dihedral inertial quotients of order 6*, J. Algebra **172** (1995), no. 1, 113–125, DOI 10.1006/jabr.1995.1051. MR1320622 (96a:20018)
- [42] Yoko Usami, *Perfect isometries and isotypies for blocks with abelian defect groups and the inertial quotients isomorphic to $Z_3 \times Z_3$* , J. Algebra **182** (1996), no. 1, 140–164, DOI 10.1006/jabr.1996.0165. MR1388861 (97c:20016)
- [43] Yoko Usami, *Perfect isometries and isotypies for blocks with abelian defect groups and the inertial quotients isomorphic to $Z_4 \times Z_2$* , J. Algebra **181** (1996), no. 3, 727–759, DOI 10.1006/jabr.1996.0144. MR1386577 (97d:20010)
- [44] John H. Walter, *The characterization of finite groups with abelian Sylow 2-subgroups.*, Ann. of Math. (2) **89** (1969), 405–514. MR0249504 (40 #2749)
- [45] Atumi Watanabe, *Notes on p -blocks of characters of finite groups*, J. Algebra **136** (1991), no. 1, 109–116, DOI 10.1016/0021-8693(91)90069-K. MR1085125 (92i:20016)
- [46] A. Watanabe, *Appendix on blocks with elementary abelian defect group of order 9*, in Representation Theory of Finite Groups and Algebras, and Related Topics (Kyoto, 2008), 9–17, Kyoto University Research Institute for Mathematical Sciences, Kyoto, 2010.

MATHEMATISCHES INSTITUT, FRIEDRICH-SCHILLER-UNIVERSITÄT, GERMANY
E-mail address: **kuelshammer@uni-jena.de**

MATHEMATISCHES INSTITUT, FRIEDRICH-SCHILLER-UNIVERSITÄT, GERMANY
E-mail address: **benjamin.sambale@uni-jena.de**