A GEOMETRIC CONSTRUCTION OF TYPES FOR THE SMOOTH REPRESENTATIONS OF $\mathrm{PGL}(2)$ OVER A LOCAL FIELD

PAUL BROUSSOUS

Dedicated to Guy Henniart on his 60th birthday

ABSTRACT. We show that almost all (Bushnell and Kutzko) types of $\mathrm{PGL}(2, F)$, F a non-Archimedean locally compact field of odd residue characteristic, naturally appear in the cohomology of finite graphs.

1. INTRODUCTION

Let F be a non-Archimedean locally compact field and G the group $\mathrm{PGL}(2, F)$. We assume that *the residue characteristic of F is not 2*. In previous works ([2], [3]) we defined a tower of directed graphs $(\tilde{X}_n)_{n \geq 0}$ lying G -equivariantly over the Bruhat-Tits tree X of G . We proved the two following facts:

Theorem 1 ([3], Theorem (3.2.4), page 502). *Let $(\pi, \mathcal{V})$ be a non-spherical generic smooth irreducible representation. Then $(\pi, \mathcal{V})$ is a quotient of the cohomology space with compact support $H_c^1(\tilde{X}_{n(\pi)}, \mathbb{C})$, where $n(\pi)$ is the conductor of π .*

Theorem 2 ([3], Theorem (5.3.2), page 512). *If $(\pi, \mathcal{V})$ is a supercuspidal smooth irreducible representation of G , then we have*

$$\dim_{\mathbb{C}} \mathrm{Hom}_G [H_c^1(\tilde{X}_{n(\pi)}, \mathbb{C}), \mathcal{V}] = 1 .$$

In this paper we make the G -module structure of $H_c^1(\tilde{X}_n, \mathbb{C})$ more explicit for all $n \geq 0$, and draw some interesting consequences.

Let us fix an edge $[s_0, s_1]$ of X and denote by $\mathcal{K}_0$ and $\mathcal{K}_1$ the stabilizers in G of s_0 and $[s_0, s_1]$, respectively. Then $\mathcal{K}_0$ and $\mathcal{K}_1$ form a set of representatives of the two conjugacy classes of maximal compact subgroups in G . If n is even, we have a G -equivariant mapping $p_n : \tilde{X}_n \rightarrow X$ which respects the graph structures. We denote by Σ_n the subgraph $p_n^{-1}([s_0, s_1])$. If n is odd, then after passing to the first barycentric subdivisions, we have a G -equivariant mapping $p_n : \tilde{X}_n \rightarrow X$ which respects the graph structures. We denote by Σ_n the subgraph $p_n^{-1}(S(s_0, 1/2))$, where $S(s_0, 1/2)$ denotes the set of points x in X such that $d(x, s_0) \leq 1/2$ (here d is the natural distance on the standard geometric realization of X , normalized in such a way that $d(s_0, s_1) = 1$).

Then for all n , Σ_n is a finite graph, equipped with an action of $\mathcal{K}_1$ if n is even, and $\mathcal{K}_0$ if n is odd. So the cohomology spaces $H^1(\Sigma_n, \mathbb{C})$ provide finite dimensional smooth representations of $\mathcal{K}_1$ or $\mathcal{K}_0$, according to the parity of n .

Received by the editors March 26, 2012 and, in revised form, April 17, 2013.

2010 *Mathematics Subject Classification*. Primary 22E50; Secondary 20J05.

The author wants to thanks the anonymous referee whose remarks helped him to improve the presentation of this article.

For each $n \geq 0$, we define an finite set $\mathcal{P}_n$ of pairs $(\mathcal{K}, \lambda)$ formed of a maximal compact subgroup $\mathcal{K} \in \{\mathcal{K}_0, \mathcal{K}_1\}$ and of an irreducible smooth representation of $\mathcal{K}$. By definition we have $(\mathcal{K}, \lambda) \in \mathcal{P}_n$ if and only if there exists $k \in \{0, 1, \dots, n\}$ such that $(\mathcal{K}, \lambda)$ is an irreducible constituent of the representation $H^1(\Sigma_k, \mathbb{C})$. For $(\mathcal{K}, \lambda) \in \mathcal{P}_n$ and $k \leq n$, we denote by m_λ^k the multiplicity of λ in $H_c^1(\Sigma_k, \mathbb{C})$ and we set $m_{n,\lambda} = m_\lambda = m_\lambda^0 + \dots + m_\lambda^n$. Note that m_λ depends on $(\mathcal{K}, \lambda)$ and n .

The main results of this article are the following.

Theorem A. *For all $n \geq 0$, we have the direct sum decomposition*

$$H_c^1(\tilde{X}_n, \mathbb{C}) = \mathbf{St}_G \oplus \bigoplus_{(\mathcal{K}, \lambda) \in \mathcal{P}_n} (c\text{-ind}_{\mathcal{K}}^G \lambda)^{m_\lambda}.$$

(Here $\mathbf{St}_G$ denotes the Steinberg representation of G).

Theorem B. *For all $n \geq 0$, any element of $\mathcal{P}_n$ is:*

- a) *either a type in the sense of Bushnell and Kutzko's type theory [6], which is not a type for the unramified principal series,*
- b) *or a pair of the form $(\mathcal{K}_0, \chi \circ \det \otimes \mathbf{St}_{\mathcal{K}_0})$, where χ is a smooth character of $F^\times$ of order 2, trivial on the group of 1-units in $F^\times$, and $\mathbf{St}_{\mathcal{K}_0}$ is the representation inflated from the Steinberg representation of $\mathrm{PGL}(2)$ of the residue field of F ,*
- c) *or the pair $(\mathcal{K}_1, \mathbf{1}_{\mathcal{K}_1})$, where $\mathbf{1}$ denotes a trivial character.*

It is worth noting that if the pairs of cases a) and b) are not types, they are *typical* in the sense of [9].

Corollary C. *Let $n \geq 0$. If $(\mathcal{K}, \lambda) \in \mathcal{P}_n$ is a cuspidal type, then $m_{n,\lambda} = 1$.*

Indeed this follows from Theorems 2 and A using Frobenius reciprocity for compact induction.

By Theorem 1, any Bernstein component of G , different from the unramified principal series component, must have a type in $\mathcal{P}_n$ for n large enough. Hence the graphs $\tilde{X}_n$, $n \geq 0$, provide a geometric construction of types for almost all Bernstein components of G .

We conjecture that if $(\mathcal{K}, \lambda) \in \mathcal{P}_n$ is a type of G , then $m_\lambda = 1$.

Finally, let us observe that this construction gives a new proof that the irreducible supercuspidal representations of G are obtained by compact induction. Our proof differs from Kutzko's original proof ([10], also see [4]) only at the exhaustion steps. Indeed our "supercuspidal" types are the same as Kutzko's, but we prove that any irreducible supercuspidal representation contains such a type by using an argument based on [2] and [3], that is mainly on the existence of the new vector.

The article is organized as follows. The proof of Theorem A relies first on combinatorial properties of the graphs $\tilde{X}_n$ that are stated and proved in §2. Using these combinatorial properties and some homological arguments, we show in §3 how to relate the cohomology of $\tilde{X}_n$ to that of $\tilde{X}_{n-1}$. The irreducible components of $H^1(\Sigma_n)$ are determined in §4 when n is even, and in §5 and §6 when n is odd. A synthesis of the arguments of paragraphs 2 to 6, leading to a proof of Theorems A and B, is given in §7.

We shall assume that the reader is familiar with the language of Bushnell and Kutzko's type theory [6] and with the language of strata ([5], [4]).

2. NOTATION

We shall denote by

F a non-Archimedean non-discrete locally compact field,

$\mathfrak{o}$ its valuation ring,

$\mathfrak{p}$ the maximal ideal of $\mathfrak{o}$,

ϖ the choice of a uniformizer of $\mathfrak{o}$,

$\mathbf{k} = \mathfrak{o}/\mathfrak{p}$ the residue field of F ,

p the characteristic of $\mathbf{k}$,

$q = p^f$ the cardinal of $\mathbf{k}$,

G the group $\mathrm{PGL}(2, F)$.

t_ϖ the image of the matrix $\begin{pmatrix} 1 & 0 \\ 0 & \varpi \end{pmatrix}$ in G .

The results of this article are obtained by the following.

Hypothesis. *The characteristic of $\mathbf{k}$ is not 2*

We shall often define an element, a subset, or a subgroup of G by giving a (set of) representative(s) in $\mathrm{GL}(2, F)$.

We write T for the diagonal torus of G and $B \supset T$ for the upper standard Borel subgroup. We denote by T^0 the maximal compact subgroup of T , i.e., the set of matrices with coefficients in $\mathfrak{o}^\times$, and by T^n the subgroup of matrices with coefficients in $1 + \mathfrak{p}^n$, $n > 0$.

Let k, l be integers satisfying $k + l \geq 0$. Then

$$\mathfrak{A}(k, l) = \begin{pmatrix} \mathfrak{o} & \mathfrak{p}^l \\ \mathfrak{p}^k & \mathfrak{o} \end{pmatrix}$$

is an $\mathfrak{o}$ -order of $\mathrm{M}(2, F)$. We denote by $\Gamma_0(k, l)$ the image in G of its group of units. There are two conjugacy classes of maximal compact subgroups of G . The first one has representative $K = \Gamma_0(0, 0)$. A representative $\tilde{I}$ of the second one is the semidirect product of the cyclic group generated by $\Pi = \begin{pmatrix} 0 & 1 \\ \varpi & 0 \end{pmatrix}$ with the Iwahori subgroup $I = \Gamma_0(1, 0)$.

The group K is filtered by the normal compact open subgroups

$$K_n = \begin{pmatrix} 1 + \mathfrak{p}^n & \mathfrak{p}^n \\ \mathfrak{p}^n & 1 + \mathfrak{p}^n \end{pmatrix}, \quad n \geq 1.$$

The group I is filtered by the normal compact subgroups I_n , $n \geq 1$, defined by

$$I_{2k+2} = \begin{pmatrix} 1 + \mathfrak{p}^{k+1} & \mathfrak{p}^{k+1} \\ \mathfrak{p}^{k+2} & 1 + \mathfrak{p}^{k+1} \end{pmatrix}, \quad I_{2k+1} = \begin{pmatrix} 1 + \mathfrak{p}^{k+1} & \mathfrak{p}^k \\ \mathfrak{p}^{k+1} & 1 + \mathfrak{p}^{k+1} \end{pmatrix}, \quad k \geq 0.$$

The subgroups I_n , $n \geq 1$, are normalized by Π .

We denote by X the Bruhat-Tits building of G . This is a uniform tree with valency $q + 1$. As a G -set and as a simplicial complex X identifies with the following complex. Its vertices are the homothety classes $[L]$ of full $\mathfrak{o}$ -lattices L in the vector space $V = F^2$. Two vertices $[L]$ and $[M]$ define an edge if and only if there exists a basis (e_1, e_2) of V such that, up to homothety, we have $L = \mathfrak{o}e_1 \oplus \mathfrak{o}e_2$ and $M = \mathfrak{o}e_1 \oplus \mathfrak{p}e_2$.

The vertices of the standard apartment (i.e., the apartment stabilized by T) are the $s_k = [\mathfrak{o} \oplus \mathfrak{p}^k]$, $k \in \mathbb{Z}$. The element t_ϖ acts as $t_\varpi s_k = s_{k+1}$, $k \in \mathbb{Z}$. The maximal compact subgroup K is the stabilizer of s_0 and $\tilde{I}$ (resp. I) is the global stabilizer (resp. pointwise stabilizer) of the edge $[s_0, s_1]$. If $l \geq k$, the pointwise stabilizer of the segment $[s_k, s_l]$ is $\Gamma_0(l, -k)$.

3. COMBINATORICS OF $\tilde{X}_n$

We recall the construction of the directed graphs $\tilde{X}_n$, $n \geq 1$.

For any integer $k \geq 1$, an *oriented k -path* in X is an injective sequence of vertices $(s_i)_{i=0,\dots,k}$ in X such that, for $k = 0, \dots, k-1$, $\{s_i, s_{i+1}\}$ is an edge in X . We shall allow the index i to run over any interval of integers of length $k+1$. Let us fix an integer $n \geq 1$. The directed graph $\tilde{X}_n$ is constructed as follows. Its edge set (resp. vertex set) is the set of oriented $(n+1)$ -paths (resp. n -paths) in X . If $a = \{s_0, s_1, \dots, s_{n+1}\}$ is an edge of $\tilde{X}_n$, its head (resp. its tail) is $a^+ = \{s_1, s_2, \dots, s_{n+1}\}$ (resp. $a^- = \{s_0, s_1, \dots, s_n\}$). The graphs we obtain in this way are actually simplicial complex. The group G acts on $\tilde{X}_n$ in an obvious way; the action preserves the structure of directed graph.

When $n = 2m$ is even, we have a natural simplicial projection $p = p_n : \tilde{X}_n \rightarrow X$ given on vertices by $p(s_{-m}, \dots, s_0, \dots, s_m) = s_0$. It is G -equivariant. Let $e = \{s_0, t_0\}$ be an edge of X . We are going to describe the finite simplicial complex $p^{-1}(e)$. An edge in $\tilde{X}_n$ above the edge e corresponds to an oriented $(2m+1)$ -path of one of the following forms:

- i) $(s_{-m}, s_{-m+1}, \dots, s_0, t_0, \dots, t_{m-1}, t_m)$,
- ii) $(t_{-m}, t_{-m+1}, \dots, t_0, s_0, \dots, s_{m-1}, s_m)$.

Let $C_{2m-1}(e)$ be the set of $(2m-1)$ -paths $c = (s_{-m+1}, \dots, s_0, t_0, \dots, t_{m-1})$. We say that $c \in C_{2m-1}(e)$ *lies above* e . Fix $c \in C_{2m-1}(e)$ and consider the simplicial subcomplex $\tilde{X}_{2m}[e, c]$ of $\tilde{X}_{2m}$ whose edges correspond to the $(2m+1)$ -paths of the form

$$(a, s_{-m+1}, \dots, s_0, t_0, \dots, t_{m-1}, b).$$

So a (resp. b) can be any neighbour of s_{-m+1} (resp. t_{m-1}) different from s_{-m+2} (resp. t_{m-1}), with the convention that $s_1 = t_0$ and $t_{-1} = s_0$. The simplicial complex $\tilde{X}_{2m}[e, c]$ is connected. It is indeed isomorphic to the complete bipartite graph with sets of vertices:

$\{a ; a \text{ neighbour of } s_{-m+1}, a \neq s_{-m+2}\}$ and $\{b ; b \text{ neighbour of } t_{m-1}, b \neq t_{m-2}\}$.

Lemma 3.1. *Let e and e' be two edges of X and $c \in C_{2m-1}(e)$, $c' \in C_{2m-1}(e')$. Then $\tilde{X}_{2m}[e, c] \cap \tilde{X}_{2m}[e', c'] \neq \emptyset$ if and only if we are in one of the following cases:*

- i) $e = e'$ and $c = c'$ (so that $\tilde{X}_{2m}[e, c] = \tilde{X}_{2m}[e', c']$);
- ii) $e \cap e'$ is reduced to one vertex of X and $c \cup c'$ is an oriented $2m$ -path in X .

In that case $\tilde{X}_{2m}[e, c] \cap \tilde{X}_{2m}[e', c']$ is reduced to the vertex of $\tilde{X}_{2m}$ corresponding to the $2m$ -path $c \cup c'$.

Proof. If $\tilde{X}_{2m}[e, c] \cap \tilde{X}_{2m}[e', c'] \neq \emptyset$, then $e \cap e' = p(\tilde{X}_{2m}[e, c]) \cap p(\tilde{X}_{2m}[e', c']) \neq \emptyset$. Assume first that $e = e'$. Then $c = c'$, for if $c \neq c'$, then $\tilde{X}_{2m}[e, c] \cap \tilde{X}_{2m}[e', c'] = \emptyset$; indeed, if $\tilde{s}$ is a vertex of $\tilde{X}_{2m}[e, c]$, then it determines c uniquely. Now assume that $e \cap e'$ is a vertex. Let $\tilde{s} \in \tilde{X}_{2m}[e, c] \cap \tilde{X}_{2m}[e', c']$. Then $\tilde{s}$ contains c and c' as subsequences, with $c \neq c'$. So by a length argument $s = c \cup c'$. Conversely if $c \cup c'$ is an oriented $2n$ -path, then $c \cup c'$ is a vertex of $\tilde{X}$ lying in $\tilde{X}_{2m}[e, c] \cap \tilde{X}_{2m}[e', c']$. $\square$

Corollary 3.2. *For any edge e of X , the connected components of $p^{-1}(e)$ are the $\tilde{X}_{2m}[e, c]$, where c runs over $C_{2m-1}(e)$.*

Define a 1-dimensional simplicial complex Y_{2m-1} in the following way. Its vertices are the connected components $\tilde{X}_{2m}[e, c]$, where e runs over the edges of X and c

over $C_{2m-1}(e)$, and two vertices $\tilde{X}_{2m}[e, c]$ and $\tilde{X}_{2m}[e', c']$ are linked by an edge if they intersect. Note that Y_{2m-1} is naturally a G -simplicial complex.

Corollary 3.3. *As a G -simplicial complex, Y_{2m-1} is canonically isomorphic to the complex $\tilde{X}_{2m-1}$.*

Lemma 3.4. *For all $n \geq 0$, the simplicial complex $\tilde{X}_n$ is connected.*

Proof. When n is even this is [2, Lemma (4.1)]. For odd n , the proof is similar. $\square$

Assume that $m \geq 1$. We say that an edge of $\tilde{X}_{2m-1}$ lies above a vertex s_0 of X if as an oriented $2m$ -path it has the form $(s_{-m}, \dots, s_0, \dots, s_m)$. For any vertex s_0 of X we write $\tilde{X}_{2m-1}[s_0]$ for the subsimplicial complex of $\tilde{X}_{2m-1}$ formed of the edges lying above s_0 .

Lemma 3.5. *When $m = 1$ the simplicial complexes $\tilde{X}_{2m-1}[s_0] = \tilde{X}_1[s_0]$ are connected.*

Proof. We may identify the neighbour vertices of s_0 in X with the points of the projective line $\mathbb{P}^1(\bar{M}) \simeq \mathbb{P}^1(\mathbf{k})$, where $s_0 = [M]$ and $\bar{M} = M/\mathfrak{p}M$. The vertices of $\tilde{X}_1[s_0]$ are the oriented 1-paths (s_0, x) , (y, s_0) , $x, y \in \mathbb{P}^1(\bar{M})$. Two oriented 1-paths of the form (x, s_0) and (s_0, y) are linked by the edge (x, s_0, y) . Let (x, s_0) , (y, s_0) be two oriented 1-paths with $x \neq y$. Since $|\mathbb{P}^1(\mathbf{k})| \geq 3$, there exists $z \in \mathbb{P}^1(\bar{M})$ distinct from x and y . Then (x, s_0) is linked to (s_0, z) via the path (x, s_0, z) and (s_0, z) is linked to (y, s_0) via the path (y, s_0, z) . For vertices of the form (s_0, x) , (s_0, y) the proof is similar. $\square$

We now assume that $m > 1$. We write $C_{2m-2}(s_0)$ for the set $(2m-2)$ -paths of the form $(s_{-m+1}, \dots, s_0, \dots, s_{m-1})$. For any $c \in C_{2m-2}(s_0)$, we consider the subsimplicial complex $\tilde{X}_{2n-1}[s_0, c]$ of $\tilde{X}_{2m-1}$ whose edges corresponds to the $2m$ -paths of the form $(a, s_{-n+1}, \dots, s_0, s_{n-1}, b)$. We have results similar to Lemma 3.1 and Corollaries 3.2 and 3.3.

Lemma 3.6. i) *For any vertex s_0 of X and for $c \in C_{2m-2}(s_0)$, $\tilde{X}_{2m-1}[s_0, c]$ is connected. It is indeed isomorphic to a complete bipartite graph constructed on two sets of $q = |\mathbf{k}|$ elements.*

ii) *Let s and s' be vertices of X , $c \in C_{2m-2}(s)$ and $c' \in C_{2m-2}(s')$. Then $\tilde{X}_{2m-1}[s, c] \cap \tilde{X}_{2m-1}[s', c'] \neq \emptyset$ if and only if $s = s'$ and $c = c'$, or $\{s, s'\}$ is an edge in X and $c \cup c'$ is an oriented $2n-1$ -path. In this last case $\tilde{X}_{2m-1}[s, c] \cap \tilde{X}_{2m-1}[s', c'] = \{\tilde{s}\}$, where the vertex $\tilde{s}$ of $\tilde{X}_{2m-1}$ corresponds to the $(2n-1)$ -path $c \cup c'$.*

iii) *For any vertex s of X , the connected components of $\tilde{X}_{2m-1}[s]$ are $\tilde{X}_{2m-1}[s, c]$, c running over $C_{2m-2}(s)$.*

We can consider the 1-dimensional simplicial complex Z_{2m-2} whose vertices are the connected components $\tilde{X}_{2m-1}[s, c]$, s running over the vertices of X and c over $C_{2m-2}(s)$, and where two connected components define an edge if and only if they intersect. Note that Z_{2m-2} is naturally a G -simplicial complex.

Corollary 3.7. *As a G -simplicial complex Z_{2m-2} is isomorphic to X_{2n-2} .*

4. THE COHOMOLOGY OF $\tilde{X}_n$: FIRST REDUCTIONS

If Σ is a locally finite 1-dimensional simplicial complex, we write Σ^0 (resp. $\Sigma^{(1)}, \Sigma^{(1)}$) for its set of vertices (resp. non-oriented edges, oriented edges). We

let $C_0(\Sigma)$ (resp. $C_1(\Sigma)$) denote the $\mathbb{C}$ -vector space with basis Σ^0 (resp. Σ^1). We define the space $C_c^0(\Sigma, \mathbb{C}) = C_c^0(\Sigma)$ (resp. $C_c^1(\Sigma, \mathbb{C}) = C_c^1(\Sigma)$) of oriented simplicial 0-cochains (resp. 1-cochains) with compact support by:

$C_c^0(\Sigma)$ = space of all linear forms $f : C_0(\Sigma) \rightarrow \mathbb{C}$ such that $f(s) = 0$ except for a finite number of vertices s ;

$C_c^1(\Sigma)$ = space of all linear forms $\omega : C_1(\Sigma) \rightarrow \mathbb{C}$ such that $\omega([a, b]) = 0$ except for a finite number of oriented edges $[a, b]$ and $\omega([a, b]) = -\omega([b, a])$.

We set $C_c^k(\Sigma) = 0$ for $k \in \mathbb{Z} \setminus \{0, 1\}$ and define a coboundary map $d : C_c^0(\Sigma) \rightarrow C_c^1(\Sigma)$ by $df([a, b]) = f(b) - f(a)$. The cohomology of the cochain complex $(C_c^\bullet(\Sigma), d)$ computes the cohomology with compact support $H_c^\bullet(\Sigma, \mathbb{C}) = H_c^\bullet(\Sigma)$ of (the standard geometric realization of) Σ . If Σ is acted upon by a group H whose action is simplicial, then $(C_c^\bullet(\Sigma), d)$ is in a straightforward way a complex of H -modules, and its cohomology computes $H_c^1(\Sigma)$ as a H -module. When Σ is finite we drop the subscripts c .

Since the stabilizer of a finite number of vertices of X is open in G , we see that for $n \geq 1$, the G -modules $C_c^0(\tilde{X}_n)$, $C_c^1(\tilde{X}_n)$ and therefore $H_c^1(\tilde{X}_n)$ are smooth.

In the sequel we fix $m \geq 1$ and we abbreviate $\tilde{X}_{2m} = \tilde{X}$. The disjoint union $\tilde{X}^1 = \bigsqcup_{e \in X^{(1)}} \tilde{X}_e$, where $\tilde{X}_e = p^{-1}(e)$, induces an isomorphism:

$$(4.1) \quad \begin{array}{ccc} C_c^1(\tilde{X}) & \simeq & \bigoplus_{e \in X^{(1)}} C^1(\tilde{X}_e), \\ \omega & \mapsto & (\omega|_{C_1(\tilde{X}_e)})_{e \in X^{(1)}}. \end{array}$$

Similarly, the non-disjoint union $\tilde{X}^0 = \bigcup_{e \in X^{(1)}} \tilde{X}_e^0$ induces an injection:

$$(4.2) \quad \begin{array}{ccc} C_c^0(\tilde{X}) & \hookrightarrow & \bigoplus_{e \in X^{(1)}} C^0(\tilde{X}_e), \\ f & \mapsto & (f|_{C_0(\tilde{X}_e)})_{e \in X^{(1)}}. \end{array}$$

We have the following commutative diagram of G -modules:

$$\begin{array}{ccccccc} H_c^0(\tilde{X}) & \longrightarrow & \bigoplus_{e \in X^{(1)}} H^0(\tilde{X}_e) & \xrightarrow{\varphi} & \text{coker } j & & \\ \downarrow & & \downarrow & & \downarrow & & \\ 0 \longrightarrow & C_c^0(\tilde{X}) & \xrightarrow{j} & \bigoplus_{e \in X^{(1)}} C^0(\tilde{X}_e) & \longrightarrow & \text{coker } j & \longrightarrow 0 \\ \downarrow d & & \downarrow \oplus d_e & & \downarrow & & \\ 0 \longrightarrow & C_c^1(\tilde{X}) & \longrightarrow & \bigoplus_{e \in X^{(1)}} C^1(\tilde{X}_e) & \longrightarrow & 0 & \longrightarrow 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ H_c^1(\tilde{X}) & \longrightarrow & \bigoplus_{e \in X^{(1)}} H^1(\tilde{X}_e) & \longrightarrow & 0 & & \end{array}$$

Here, for $e \in X^{(1)}$, d_e denote the coboundary map $C^0(\tilde{X}_e) \rightarrow C^1(\tilde{X}_e)$. Since $\tilde{X}$ is connected ([2] Lemma 4.1) and non-compact, we have $H_c^0(\tilde{X}) = 0$. So the snake lemma gives the kernel-cokernel exact sequence:

$$0 \rightarrow \bigoplus_{e \in X^{(1)}} H^0(\tilde{X}_e) \rightarrow \text{coker } j \rightarrow H_c^1(\tilde{X}) \rightarrow \bigoplus_{e \in X^{(1)}} H^1(\tilde{X}_e) \rightarrow 0,$$

that is,

$$(4.3) \quad 0 \rightarrow \text{coker } j / \varphi \left(\bigoplus_{e \in X^{(1)}} H^0(\tilde{X}_e) \right) \rightarrow H_c^1(\tilde{X}) \rightarrow \bigoplus_{e \in X^{(1)}} H^1(\tilde{X}_e) \rightarrow 0.$$

Abbreviate $Y = Y_{2m-1}$.

Lemma 4.1. *We have a canonical isomorphism of G -modules*

$$\operatorname{coker} j / \varphi \left(\bigoplus_{e \in X^{(1)}} H^0(\tilde{X}_e) \right) \simeq H_c^1(Y).$$

Proof. From Corollary 3.2 we have

$$\bigoplus_{e \in X^{(1)}} C^0(\tilde{X}_e) = \bigoplus_{e \in X^{(1)}} \bigoplus_{c \in C_{2m-1}(e)} C^0(\tilde{X}_{2m}[e, c]).$$

So the map j is given by $f \mapsto \bigoplus_{e \in X^{(1)}} \bigoplus_{c \in C_{2m-1}(e)} f_{e,c}$, where $f_{e,c} = f|_{C_0(\tilde{X}_{2m}[e,c])}$. Consider the G -equivariant morphism of vector spaces

$$\psi : \bigoplus_{e \in X^{(1)}} \bigoplus_{c \in C_{2m-1}(e)} C^0(\tilde{X}_{2m}[e, c]) \rightarrow C_c^1(Y)$$

given as follows. If σ is an oriented edge of Y , then there exist uniquely determined edges e_o, e'_o of X , $c_o \in C_{2m-1}(e_o)$, $c'_o \in C_{2m-1}(e'_o)$, such that σ corresponds to the intersection $\tilde{X}_{2m}[e_o, c_o] \cap \tilde{X}_{2m}[e'_o, c'_o] = \{s_o\}$, $s_o \in \tilde{X}^0$. We then set

$$\psi[(f_{e,c})_{e,c}](\sigma) = f_{e'_o, c'_o}(s_o) - f_{e_o, c_o}(s_o).$$

Then ψ is surjective and its kernel is precisely $j(C_c^0(\tilde{X}))$. So we may identify $\operatorname{coker} j$ with $C_c^1(Y)$. From Corollary 3.2, we have

$$\bigoplus_{e \in X^{(1)}} H^0(\tilde{X}_e) = \bigoplus_{e \in X^{(1)}} \bigoplus_{c \in C_{2m-1}(e)} H^0(\tilde{X}_{2m}[e, c])$$

so that we may identify $\bigoplus_{e \in X^{(1)}} H^0(\tilde{X}_e)$ with $C_c^0(\tilde{Y})$. Under our identifications the map $\varphi : \bigoplus_{e \in X^{(1)}} H^0(\tilde{X}_e) \rightarrow \operatorname{coker} j$ corresponds to the coboundary map $d : C_c^0(Y) \rightarrow C_c^1(Y)$, and we are done since all our identifications are G -equivariant. $\square$

Proposition 4.2. *For $m \geq 1$, we have an isomorphism of G -modules:*

$$H_c^1(\tilde{X}_{2m}) \simeq H_c^1(\tilde{X}_{2m-1}) \oplus c\text{-ind}_{\mathcal{K}_{e_o}}^G H^1(\tilde{X}_{e_o})$$

for any edge e_o of X and where $\mathcal{K}_{e_o}$ denotes the stabilizer of e_o in G .

Proof. From the short exact sequence (4.3) and Lemma 4.1, we have the exact sequence of G -modules:

$$(4.4) \quad 0 \rightarrow H_c^1(Y) \rightarrow H_c^1(\tilde{X}) \rightarrow \bigoplus_{e \in X^{(1)}} H^1(\tilde{X}_e) \rightarrow 0.$$

Since G acts transitively on the edges of X , $\bigoplus_{e \in X^{(1)}} H^1(\tilde{X}_e)$ identifies with the compactly induced representation $c\text{-ind}_{\mathcal{K}_{e_o}}^G H^1(\tilde{X}_{e_o})$. Moreover, this induced representation is projective in the category of smooth complex representations of G . This is classical and follows from Frobenius reciprocity for compact induction together with the fact that the category of smooth $\mathcal{K}_{e_o}$ -modules is semisimple. So the sequence (4.4) splits. $\square$

We assume that $m \geq 1$ and we abbreviate $\tilde{X} = \tilde{X}_{2m-1}$. The disjoint union $\tilde{X}^1 = \bigsqcup_{s \in X^0} \tilde{X}_{2m-1}[s]^1$ induces an isomorphism:

$$(4.5) \quad \begin{array}{ccc} C_c^1(\tilde{X}) & \simeq & \bigoplus_{s \in X^0} C^1(\tilde{X}_{2m-1}[s]), \\ \omega & \mapsto & (\omega|_{C_1(\tilde{X}_{2m-1}[s])})_{s \in X^0}. \end{array}$$

Similarly, the non-disjoint union $\tilde{X}^0 = \bigcup_{s \in X^0} \tilde{X}_s^0$ induces an injection:

$$(4.6) \quad \begin{array}{ccc} j : C_c^0(\tilde{X}) & \hookrightarrow & \bigoplus_{s \in X^0} C^0(\tilde{X}_{2m-1}[s]), \\ f & \mapsto & (f|_{C_0(\tilde{X}_{2m-1}[s])})_{s \in X^0}. \end{array}$$

We have the following commutative diagram of G -modules:

$$\begin{array}{ccccccc} H_c^0(\tilde{X}) & \longrightarrow & \bigoplus_{s \in X^0} H^0(\tilde{X}_{2m-1}[s]) & \xrightarrow{\varphi} & \text{coker } j & & \\ \downarrow & & \downarrow & & \downarrow & & \\ 0 \longrightarrow & C_c^0(\tilde{X}) & \xrightarrow{j} & \bigoplus_{s \in X^0} C^0(\tilde{X}_{2m-1}[s]) & \longrightarrow & \text{coker } j & \longrightarrow 0 \\ \downarrow d & & \downarrow \oplus d_s & & \downarrow & & \\ 0 \longrightarrow & C_c^1(\tilde{X}) & \longrightarrow & \bigoplus_{s \in X^0} C^1(\tilde{X}_{2m-1}[s]) & \longrightarrow & 0 & \longrightarrow 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ H_c^1(\tilde{X}) & \longrightarrow & \bigoplus_{s \in X^0} H^1(\tilde{X}_{2m-1}[s]) & \longrightarrow & 0 & & \end{array}$$

Here, for $s \in X^0$, d_s denote the coboundary map $C^0(\tilde{X}_{2m-1}[s]) \rightarrow C^1(\tilde{X}_{2m-1}[s])$. By Lemma 3.4, $\tilde{X}$ is connected. So we have $H_c^0(\tilde{X}) = 0$ since $\tilde{X}$ is non-compact. The *snake lemma* gives the kernel-cokernel exact sequence:

$$(4.7) \quad 0 \rightarrow \text{coker } j / \varphi \left(\bigoplus_{s \in X^0} H^0(\tilde{X}_{2m-1}[s]) \right) \rightarrow H_c^1(\tilde{X}) \rightarrow \bigoplus_{s \in X^0} H^1(\tilde{X}_{2m-1}[s]) \rightarrow 0.$$

Lemma 4.3. *We have a canonical isomorphism of G -modules*

$$\text{coker } j / \varphi \left(\bigoplus_{s \in X^0} H^0(\tilde{X}_{2m-1}[s]) \right) \simeq H_c^1(\tilde{X}_{2m-2}).$$

Proof. It is similar to the proof of Lemma 4.1 and relies on Lemma 3.6 and Corollary 3.7. $\square$

Proposition 4.4. *For $m \geq 1$, we have an isomorphism of G -modules:*

$$H_c^1(\tilde{X}_{2m-1}) \simeq H_c^1(\tilde{X}_{2m-2}) \oplus c\text{-ind}_{\mathfrak{K}_{s_o}}^G H^1(\tilde{X}_{s_o})$$

for any vertex s_o and where $\mathfrak{K}_{s_o}$ denotes the stabilizer of s_o in G .

Proof. Similar to the proof of Proposition 4.2. $\square$

Recall [3] that $\tilde{X}_0$ is different from X . This is a directed graph whose set of vertices is isomorphic to X^0 as a G -set and whose set of edges is isomorphic to the G -set of oriented edges of X .

5. DETERMINATION OF THE INDUCING REPRESENTATIONS – I

Let $m \geq 0$ be a fixed integer and $e_0 = [s_0, s_1]$ the standard edge. The aim of this section is to determine the $\mathcal{K}_{e_0}$ -module $H^1(\tilde{X}_{2m}[e_0])$. Here we have $\mathcal{K}_{e_0} = \tilde{I}$, the normalizer in G of the standard Iwahori subgroup. We have the semidirect products

$$\tilde{I} = \left\langle \begin{pmatrix} 0 & 1 \\ \varpi & 0 \end{pmatrix} \right\rangle \ltimes I = E^\times I$$

for any totally ramified subfield extension $E/F \subset M(2, F)$ such that $E^\times$ normalizes I .

We first assume that $m \geq 1$. By Corollary 3.2, we have the disjoint union

$$\tilde{X}_{2m}[e_0] = \coprod_{c \in C_{2m-1}(e_0)} \tilde{X}_{2m}[e_0, c] .$$

The group $\tilde{I}$ acts transitively on $C_{2m-1}(e_0)$. This comes from the standard fact that I , the pointwise stabilizer of e_0 acts transitively on the apartments of X containing e_0 .

Let $c_0 \in C_{2m-1}(e_0)$ be the path

$$s_{-m+1}, \dots, s_0, s_1, \dots, s_m .$$

The global stabilizer of $\tilde{X}_{2m}[e_0, c_0]$ in $\tilde{I}$ is the pointwise stabilizer of c_0 in $\tilde{I}$, that is,

$$\Gamma_0(m, m-1) = \begin{pmatrix} \mathfrak{o}^\times & \mathfrak{p}^{m-1} \\ \mathfrak{p}^m & \mathfrak{o}^\times \end{pmatrix} = T^0 I_{2m-1} .$$

It follows that

$$(5.1) \quad H^1(\tilde{X}_{2m}[e_0]) = \text{ind}_{T^0 I_{2m-1}}^{\tilde{I}} H^1(\tilde{X}_{2m}[e_0, c_0]) .$$

On the other hand, an easy calculation shows that the pointwise stabilizer of $\tilde{X}_{2m}[e_0, c_0]$ is $T^1 I_{2m}$, where T^1 is the congruence subgroup of T given by

$$T^1 = \begin{pmatrix} 1 + \mathfrak{p} & 0 \\ 0 & 1 + \mathfrak{p} \end{pmatrix} .$$

So the $T^0 I_{2m-1}$ -module $H^1(\tilde{X}_{2m}[e_0, c_0])$ may be viewed as a representation of the finite group $T^0 I_{2m-1}/T^1 I_{2m}$, that is, a semidirect product of the cyclic group $\mathbf{k}^\times$ with the abelian group $I_{2m-1}/I_{2m} \simeq \mathbf{k} \oplus \mathbf{k}$.

Set $\Gamma = \tilde{X}_{2m}[e_0, c_0]$. This is a finite directed graph. Let Σ_{-m} (resp. Σ_{m+1}) denote the set of vertices of X that are neighbours of s_{-m+1} and different from s_{-m+2} (resp. neighbours of s_m and different from s_{m-1}). Then the vertex set of Γ is

$$\Gamma^0 = \{(a, s_{-m+1}, \dots, s_0, \dots, s_m); a \in \Sigma_{-m}\} \coprod \{(s_{-m+1}, \dots, s_0, \dots, s_m, b); b \in \Sigma_{m+1}\} \\ \simeq \Sigma_{-m} \coprod \Sigma_{m+1}$$

and its edge set is

$$\Gamma^1 = \{(a, s_{-m+1}, \dots, s_0, \dots, s_m, b) ; a \in \Sigma_{-m}, b \in \Sigma_{m+1}\} \simeq \Sigma_{-m} \times \Sigma_{m+1} .$$

In particular, Γ is a bipartite graph based on two sets of q elements. In particular, its Euler character is given by

$$\chi(\Gamma) = 1 - \dim_{\mathbb{C}} H^1(\Gamma) = 2q - q^2 ,$$

so that

$$(5.2) \quad \dim_{\mathbb{C}} H^1(\Gamma) = q^2 - 2q + 1 = (q-1)^2 .$$

Let $\mathbb{C}[\Gamma^1]$ be the space of complex function on Γ^1 and let $\mathcal{H}(\Gamma)$ be the space of harmonic 1-cochains on Γ :

$$\mathcal{H}(\Gamma) = \{f \in \mathbb{C}[\Gamma^1]; \sum_{a \in \Gamma^1, s \in a} [a : s] f(a) = 0, \text{ all } s \in \Gamma^0\} .$$

Here $[a : s]$ denote an incidence number. In our case:

$$(5.3) \quad f \in \mathcal{H}(\Gamma) \text{ iff } \begin{cases} \sum_{a \in \Sigma_{-m}} f(a, s_{-m+1}, \dots, s_m, b) = 0, & \text{for all } b, \\ \sum_{b \in \Sigma_{m+1}} f(a, s_{-m+1}, \dots, s_m, b) = 0, & \text{for all } a. \end{cases}$$

This is a standard result (see e.g. [3, Lemma (1.3.2)]) that, as a $T^0 I_{2m-1}/T^1 I_{2m-1}$ -module, $H^1(\Gamma)$ is isomorphic to the contragredient module of $\mathcal{H}(\Gamma)$.

An easy computation shows that we may identify Γ^1 with $\mathbf{k} \times \mathbf{k}$ in such a way that:

1) an element of $I_{2m-1} = \begin{pmatrix} 1 + \mathfrak{p}^m & \mathfrak{p}^{m-1} \\ \mathfrak{p}^m & 1 + \mathfrak{p}^m \end{pmatrix}$ acts as

$$\left(1 + \begin{pmatrix} \varpi^m a & \varpi^{m-1} b \\ \varpi^m c & \varpi^m d \end{pmatrix}\right) \cdot (x, y) = (x + \bar{b}, y + \bar{c})$$

for $a, b, c, d \in \mathfrak{o}$, $x, y \in \mathbf{k}$, and

2) an element of T^0 acts as

$$\begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \cdot (x, y) = (\bar{a} \bar{d}^{-1} x, \bar{d} \bar{a}^{-1} y)$$

and the condition (5.3) gives us:

$$f \in \mathcal{H}(\Gamma) \text{ iff } \begin{cases} \sum_{x \in \mathbf{k}} f(x, y) = 0, & \text{for all } y \in \mathbf{k}, \\ \sum_{y \in \mathbf{k}} f(x, y) = 0, & \text{for all } x \in \mathbf{k}. \end{cases}$$

A basis of $\mathbb{C}[\Gamma]$ is formed of the functions $\chi_1 \otimes \chi_2(x, y) = \chi_1(x)\chi_2(y)$, where, for $i = 1, 2$, χ_i runs over the characters of $(\mathbf{k}, +)$. It is clear that the $(q-1)^2$ -dimensional subspace of $\mathbb{C}[\Gamma]$ generated by the $\chi_1 \otimes \chi_2$, $\chi_1 \not\equiv 1$, $\chi_2 \not\equiv 1$, is contained in $\mathcal{H}(\Gamma)$. So using (5.2), we obtain:

$$(5.4) \quad \mathcal{H}(\Gamma) = \text{Span}\{\chi_1 \otimes \chi_2 ; \chi_i \in \widehat{\mathbf{k}}, \chi_i \not\equiv 1, i = 1, 2\}.$$

It follows from (5.4) that, as an I_{2m-1}/I_{2m-1} -module, the space $\mathcal{H}(\Gamma)$ is the direct sum of 1-dimensional representations corresponding to the characters $\alpha = \alpha(\chi_1, \chi_2)$, $\chi_i \not\equiv 1$, $i = 1, 2$, given by

$$\alpha\left(1 + \begin{pmatrix} \varpi^m a & \varpi^{m-1} b \\ \varpi^m c & \varpi^m d \end{pmatrix}\right) = \chi_1(b)\chi_2(c).$$

In particular, $\mathcal{H}(\Gamma)$ is isomorphic to its contragredient and therefore isomorphic to $H^1(\Gamma)$ as an I_{2m-1}/I_{2m-1} -module. In the language of strata (the reader may refer to [4, §4]), for $\chi_i \not\equiv 1$, $i = 1, 2$, the character $\alpha(\chi_1, \chi_2)$ corresponds to a stratum of the form $[\mathcal{I}, 2m, 2m-1, \beta]$, where $\mathcal{I}$ is the standard *Iwahori order* and $\beta \in M(2, F)$ is an element of the form $\Pi^{2m-1} \begin{pmatrix} u & 0 \\ 0 & v \end{pmatrix}$, $u, v \in \mathfrak{o}^\times$. In the terminology of [4, §4, page 98] this stratum is a *ramified simple stratum*.

We now have enough material to prove the following result.

Proposition 5.1. *Let λ be an irreducible constituent of*

$$H^1(\tilde{X}_{2m}[e_0]) = \text{ind}_{T^0 I_{2m-1}}^{\tilde{I}} H^1(\tilde{X}_{2m}[e_0, c_0]).$$

Then the compactly induced representation $c\text{-ind}_{\tilde{I}}^G \lambda$ is irreducible, whence supercuspidal.

Proof. It is a standard result that an irreducible compactly induced representation is supercuspidal (see e.g. [8, page 194]).

The proof of the irreducibility is also standard by an argument due to Kutzko. But we repeat it for convenience. By Frobenius reciprocity, the restriction of λ to I_{2m-1} contains a character $\alpha(\chi_1, \chi_2)$ corresponding to a (ramified) simple stratum. Since λ is irreducible and since $\tilde{I}$ normalizes I_{2m-1} , the restriction $\lambda|_{I_{2m-1}}$ is a direct sum $\alpha_1 \oplus \cdots \oplus \alpha_r$ of $\tilde{I}$ -conjugates of $\alpha(\chi_1, \chi_2)$. They all correspond to simple strata. Let $g \in G$ be an element intertwining λ with itself. Then by restriction it intertwines a character α_i with a character α_j for some $j = 1, \dots, r$. By [4, Lemma (16.1), page 111], such an element G must belong to $\tilde{I}$. It follows that the G -intertwining of λ is equal to $\tilde{I}$ and that the representation $c\text{-ind}_{\tilde{I}}^G \lambda$ is irreducible according to Mackey's irreducibility criterion ([8, Proposition (1.5), page 195]). $\square$

We finally consider the case $m = 0$. The directed graph $\tilde{X}_0$ has X^0 as vertex set. An edge $\{t, s\}$ in X gives rise to two edges $[s, t]$ and $[t, s]$ in $\tilde{X}_0$. Since the action of G on $\tilde{X}_0$ preserves the structure of the digraph, the G -module $H_c^1(\tilde{X}_0)$ may be computed using the complex

$$0 \longrightarrow C_c^0(\tilde{X}_0) \longrightarrow C_c^{(1)}(\tilde{X}_0),$$

where $C_c^{(1)}(\tilde{X}_0)$ is the space of (unoriented) 1-cochains, that is the space of maps from $\tilde{X}_0^{(1)}$ (unoriented edges) to $\mathbb{C}$ with finite support. The coboundary map is given here by $df(a) = f(a^+) - f(a^-)$. Consider the G -equivariant injection $j : C_c^1(X) \longrightarrow C_c^{(1)}(\tilde{X}_0)$ given by $j(\omega) : [s, t] \mapsto \omega([s, t])$. We have the commutative diagram of G -modules:

$$\begin{array}{ccccccc} 0 & \longrightarrow & C_c^0(X) & \xrightarrow{\text{id}} & C_c^0(\tilde{X}_0) & \longrightarrow & 0 \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & C_c^1(X) & \xrightarrow{j} & C_c^{(1)}(\tilde{X}_0) & \longrightarrow & C_c^{(1)}(\tilde{X}_0)/\text{Im } j \longrightarrow 0 \end{array}$$

The quotient $C_c^{(1)}(\tilde{X}_0)/\text{Im } j$ identifies with the subspace of $C_c^{(1)}(\tilde{X}_0)$ formed by those functions f satisfying $f([s, t]) = f([t, s])$ for all edges $\{s, t\}$ of X . This subspace is nothing more than the compactly induced representation $c - \text{Ind}_{\tilde{I}}^G \mathbf{1}_{\tilde{I}}$. The cokernel exact sequence gives us:

$$0 \longrightarrow H_c^1(X) \longrightarrow H_c^1(\tilde{X}_0) \longrightarrow c\text{-ind}_{\tilde{I}}^G \mathbf{1}_{\tilde{I}} \longrightarrow 0.$$

Now we use the following two facts to obtain Proposition 5.2:

- the representation $c\text{-ind}_{\tilde{I}}^G \mathbf{1}_{\tilde{I}}$ is a projective object of the category of smooth representations of G ,
- the G -module $H_c^1(X)$ is isomorphic to the Steinberg representation $\mathbf{St}_G$ of G ([7]).

Proposition 5.2. *The G -module $H_c^1(\tilde{X}_0)$ is isomorphic to $\mathbf{St}_G \oplus c\text{-ind}_{\tilde{I}}^G \mathbf{1}_{\tilde{I}}$.*

6. THE INDUCING REPRESENTATIONS – II

We now determine the $\mathcal{K}_{s_0}$ -module $H^1(\tilde{X}_{2m+1}[s_0])$. The arguments are very often similar to those of the previous section and we will not give all details. Since the case $m = 0$ requires slightly different techniques we postpone it to the end of the section and assume first that $m > 0$.

Recall that the stabilizer $\mathcal{K}_{s_0}$ of s_0 in G is the image K of $\mathrm{GL}(2, \mathfrak{o})$ in G .

Let $c_0 \in C_{2m}(s_0)$ be the path $(s_{-m}, \dots, s_0, \dots, s_m)$. Its pointwise stabilizer is $\Gamma_0(m, m) = T^0 K_m$. So as a K -module, $H^1(\tilde{X}_{2m+1}[s_0])$ is isomorphic to the induced representation $\mathrm{Ind}_{T^0 K_m}^K H^1(\tilde{X}_{2m+1}[s_0, c_0])$. Moreover, the pointwise stabilizer of $\tilde{X}_{2m+1}[s_0, c_0]$ is $T^1 K_{m+1}$ and $H^1(\tilde{X}_{2m+1}[s_0, c_0])$ may be viewed as a representation of $T^0 K_m / T^1 K_{m+1}$.

As in the previous section, one may consider the bipartite graph Ω whose both vertex sets identify with $\mathbf{k}$, equipped with an action of K_m on Ω^1 given by

$$[I_2 + \varpi^m \begin{pmatrix} a & b \\ c & d \end{pmatrix}] \cdot (x, y) = (x + \bar{b}, y + \bar{c}) ,$$

the action of T^0 being given by

$$\begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \cdot (x, y) = (\bar{a}\bar{d}^{-1}x, \bar{d}\bar{a}^{-1}y) .$$

Then the contragredient of the $T^0 K_m / T^1 K_{m+1}$ -module $H^1(\tilde{X}_{2m+1}[s_0, c_0])$ is isomorphic to the space $\mathcal{H}(\Omega)$ of harmonic cochains on Ω . As in the previous section the latter space is generated by the functions $\chi_1 \otimes \chi_2$, where χ_i , $i = 1, 2$, runs over the non-trivial characters of $(k, +)$. The line $\mathbb{C}\chi_1 \otimes \chi_2$ is acted upon by K_m via the character $\alpha(\chi_1, \chi_2)$ given by

$$\alpha(\chi_1, \chi_2)(I_2 + \varpi^m \begin{pmatrix} a & b \\ c & d \end{pmatrix}) = \chi_1(b)\chi_2(c) .$$

It follows that $\mathcal{H}(\Omega)$ is isomorphic to its contragredient and that $H^1(\tilde{X}_{2m+1}[s_0, c_0])$ is the direct sum of the characters $\alpha(\chi_1, \chi_2)$, $\chi_i \neq 1$, $i = 1, 2$.

For $\chi_i \neq 1$, $i = 1, 2$, the character $\alpha(\chi_1, \chi_2)$ corresponds to a stratum of the form $[\mathrm{M}(2, \mathfrak{o}), m, m-1, \beta]$, where $\beta \in \mathrm{M}(2, F)$ is given by $\varpi^{-m} \begin{pmatrix} 0 & u \\ v & 0 \end{pmatrix}$, $u, v \in \mathfrak{o}^\times$. This stratum is either simple and non-scalar or split fundamental according to whether $uv \bmod \mathfrak{p}$ is a square in $\mathbf{k}^\times$ or not (here we have used the fact that $\mathrm{Char}(\mathbf{k}) \neq 2$).

It is clear that T^0 leaves the set of characters corresponding to simple strata (resp. split fundamental strata) stable. So we may write

$$H^1(\tilde{X}_{2m+1}[s_0, c_0]) = H^1(\tilde{X}_{2m+1}[s_0, c_0])_{\mathrm{simple}} \oplus H^1(\tilde{X}_{2m+1}[s_0, c_0])_{\mathrm{split}} ,$$

where $H^1(\tilde{X}_{2m+1}[s_0, c_0])_{\mathrm{simple}}$ (resp. $H^1(\tilde{X}_{2m+1}[s_0, c_0])_{\mathrm{split}}$) is the $T^0 K_m$ -submodule which decomposes as a K_m / K_{m+1} -module as a direct sum of (characters corresponding to) simple non-scalar strata (resp. split fundamental strata).

We have a result similar to Proposition 5.1, whose proof uses the same arguments.

Proposition 6.1. *Let λ be an irreducible constituent of*

$$\mathrm{Ind}_{T^0 K_m}^K H^1(\tilde{X}_{2m+1}[s_0, c_0])_{\mathrm{simple}} \subset H^1(\tilde{X}_{2m+1}[s_0]) .$$

Then the compactly induced representation $c\text{-ind}_K^G \lambda$ is irreducible, whence supercuspidal.

The study of $\text{Ind}_{T^0 K_m}^K H^1(\tilde{X}_{2m+1}[s_0, c_0])_{\text{split}}$ is the aim of the next section.

We are now going to determine the K -module structure of $H^1(\tilde{X}_1[s_0])$. Set $\mathbf{G} = \text{PGL}(2, \mathbf{k}) \simeq K/K^1$ and write $\mathbf{B}$ and $\mathbf{T}$ for the upper Borel subgroup and diagonal torus of $\mathbf{G}$, respectively. Let $\mathbf{U}$ be the unipotent radical of $\mathbf{B}$. As a K -set the set of neighbour vertices of s_0 is isomorphic to $\mathbb{P}^1(\mathbf{k}) = \mathbf{G}/\mathbf{B}$.

The graph $\Omega = \tilde{X}_1[s_0]$ has for the vertex set the set of paths of the form (s, s_0) or (s_0, s) where s runs over the neighbour vertices of s_0 in X . So the space $C^0(\Omega)$ of 0-cochains identifies with the space $\mathcal{F}(\mathbb{P}^1(\mathbf{k}) \amalg \mathbb{P}^1(\mathbf{k}))$ of complex-valued functions on the disjoint union $\mathbb{P}^1(\mathbf{k}) \amalg \mathbb{P}^1(\mathbf{k})$, thus has a $\mathbf{G}$ -module $C^0(\Omega)$ that is isomorphic to $\mathbf{1}_{\mathbf{G}} \oplus \mathbf{St}_{\mathbf{G}} \oplus \mathbf{1}_{\mathbf{G}} \oplus \mathbf{St}_{\mathbf{G}}$, where $\mathbf{1}$ denotes a trivial representation and $\mathbf{St}$ a Steinberg representation.

The $\mathbf{G}$ -set Ω^1 is the set of paths of the form (s, s_0, t) , where s and t are two different neighbour vertices of s_0 . This $\mathbf{G}$ -set is isomorphic to the quotient $\mathbf{G}/\mathbf{T}$. The space $C^{(1)}(\Omega)$ of unoriented 1-cochains identifies as a G -module with the space $\mathcal{F}(\mathbf{G}/\mathbf{T})$.

Fix a non-trivial character ψ of $\mathbf{U}$. It is well known that the induced representation $\text{Ind}_{\mathbf{U}}^{\mathbf{G}} \psi$ is multiplicity free. Its irreducible constituent form is by definition the *generic* (irreducible) representations of $\mathbf{G}$. Moreover, an irreducible representation is generic if and only if it is not a character.

We have a natural G -equivariant map $\Phi : \mathcal{F}(\mathbf{G}/\mathbf{T}) \rightarrow \text{Ind}_{\mathbf{U}}^{\mathbf{G}} \psi$, given by

$$\Phi(f)(g) = \sum_{u \in \mathbf{U}} f(gu) \bar{\psi}(u), \quad f \in \mathcal{F}(\mathbf{G}/\mathbf{T}), \quad g \in \mathbf{G}.$$

If a function f lies in the kernel of Φ , then we have $\sum_{u \in \mathbf{U}} f(gu) \theta(u) = 0$, for all $g \in \mathbf{G}$ and all non-trivial character θ of $\mathbf{U}$. Indeed, it suffices to use the fact that the action of $\mathbf{T}$ on $\mathbf{U}$ by conjugation acts transitively on the non-trivial characters of $\mathbf{U}$ and the right invariance of f under the action of T . So the kernel of Φ consists of the function f such that $u \mapsto f(gu)$ is a constant function on U , for all $g \in G$. In other words, $\text{Ker } \Phi = \mathcal{F}(G/B) \simeq \mathbf{1}_{\mathbf{G}} \oplus \mathbf{St}_{\mathbf{G}}$. By a dimension argument, we see that Φ is surjective. It follows that

$$C^{(1)}(\Omega) \simeq \text{Ind}_{\mathbf{U}}^{\mathbf{G}} \psi \oplus \mathbf{1}_{\mathbf{G}} \oplus \mathbf{St}_{\mathbf{G}}.$$

We have the cochain complex of G -modules:

$$0 \longrightarrow C^0(\Omega) \longrightarrow C^{(1)}(\Omega) \longrightarrow 0.$$

Since Ω is connected the kernel of the coboundary operator is the trivial module $\mathbb{C}$. Hence in the Grothendieck groups of G -modules, we have $dC^0(\Omega) \simeq 2 \cdot \mathbf{1}_{\mathbf{G}} + 2 \cdot \mathbf{St}_{\mathbf{G}} - \mathbf{1}_{\mathbf{G}} = \mathbf{1}_{\mathbf{G}} + 2 \cdot \mathbf{St}_{\mathbf{G}}$. Therefore,

$$H^1(\Omega) = C^1(\Omega)/dC^0(\Omega) \simeq \text{Ind}_{\mathbf{U}}^{\mathbf{G}} \psi + \mathbf{1}_{\mathbf{G}} + \mathbf{St}_{\mathbf{G}} - \mathbf{1}_{\mathbf{G}} - 2 \cdot \mathbf{St}_{\mathbf{G}} = \text{Ind}_{\mathbf{U}}^{\mathbf{G}} \psi - \mathbf{St}_{\mathbf{G}}.$$

Since $q = |\mathbf{k}|$ is odd, there exists a unique non-trivial character of $\mathbf{k}^{\times}/(\mathbf{k}^{\times})^2$, that we denote by χ_0 . The irreducible constituents of the Gelfand-Graev representation $\text{Ind}_{\mathbf{U}}^{\mathbf{G}} \psi$ are the following:

- the irreducible cuspidal representations of $\mathbf{G}$,
- the principal series $\text{Ind}_{\mathbf{B}}^{\mathbf{G}} \chi \otimes \chi^{-1}$, where $\chi : \mathbf{k}^{\times} \rightarrow \mathbb{C}^{\times}$ is a character such that $\chi^2 \not\equiv 1$ (i.e. $\chi \notin \{1, \chi_0\}$).
- the steinberg representation $\mathbf{St}_{\mathbf{G}}$,
- (when q is odd) the twisted representation $\mathbf{St}_{\mathbf{G}} \otimes \chi_0$.

If σ is a cuspidal representation of $\mathbf{G} = K/K^1$, then the induced representation $c\text{-ind}_K^G \sigma$ is irreducible and supercuspidal ([4, (11.5), page 81]). Such a representation of G is called a *level 0 supercuspidal* representation.

A principal series of $\mathbf{G} = K/K^1$ may be written as $\text{Ind}_I^K \rho$, where ρ is a character of I/I^1 . The pair (I, ρ) is actually a type in the sense of Bushnell and Kutzko's type theory. For technical reasons we postpone definitions and references to the next section. Since the representation $\text{Ind}_I^K \rho$ is irreducible, it is a type of the same constituent as (I, ρ) .

To sum up, we have proved the following.

Proposition 6.2. *An irreducible constituent λ of $H^1(\tilde{X}_1[s_0])$ is of one of the following forms:*

- (i) *the inflation of a cuspidal representation of $\mathbf{G}$; in that case $c\text{-ind}_K^G \lambda$ is a level 0 irreducible supercuspidal representation of G ,*
- (ii) *the inflation to K of the representation $\mathbf{St}_{\mathbf{G}} \otimes \chi_0$,*
- (iii) *a type of the form $\text{Ind}_I^K \rho$, where the ρ is inflated from a character of $I/I^1 \simeq (\mathbf{k}^\times \times \mathbf{k}^\times)/\mathbf{k}^\times$ of the form $\chi \otimes \chi^{-1}$, $\chi^2 \neq 1$.*

Note that in (iii), the pair $(K, \text{Ind}_I^K \rho)$ is a principal series type.

7. THE INDUCING REPRESENTATIONS – III

We keep the notation as in the previous section. To determine the structure of $\text{Ind}_{T^0 K_m}^K H^1(\tilde{X}_{2m+1}[s_0, c_0])_{\text{split}}$, we first recall crucial facts on split strata and types for principal series representations. The basic reference for *type theory* is [6].

Let χ be a character of T , that we view as a character of T^0 by restriction. Assume that the conductor of χ is $n > 0$: $T^n \subset \text{Ker } \chi$ and n is minimal for this property. Set

$$J_\chi = \begin{pmatrix} \mathfrak{o}^\times & \mathfrak{o} \\ \mathfrak{p}^n & \mathfrak{o}^\times \end{pmatrix} = \Gamma_0(\mathfrak{p}^n).$$

If U and $\bar{U}$ denote the groups of upper and lower unipotent matrices respectively, then J_χ has an Iwahori decomposition,

$$J_\chi = (J_\chi \cap \bar{U}) \cdot (J_\chi \cap T) \cdot (J_\chi \cap U)$$

and one may define a character ρ_χ of J_χ by

$$\rho_\chi(\bar{u}t^0u) = \chi(t^0), \quad \bar{u} \in J_\chi \cap \bar{U}, \quad u \in J_\chi \cap U, \quad t^0 \in T^0.$$

Let $\mathcal{R}_{[T, \chi]}$ be the Bernstein component of the category of smooth representations of G whose objects are the representations $\mathcal{V}$ satisfying the following property: Any irreducible subquotient of $\mathcal{V}$ occurs in a parabolically induced representation $\text{Ind}_{\mathcal{B}}^G(\chi \otimes \chi_0)$, where $\mathcal{B}$ is a Borel subgroup with Levi component T and χ_0 an unramified character of T . We then have:

Theorem 7.1 (A. Roche). *The pair (J_χ, ρ_χ) is a type for $\mathcal{R}_{[T, \chi]}$.*

This is indeed Theorem (7.7) of [11]. Note that our J_χ is not exactly the same as Roche's, but a conjugate under an element of T (see [11, Example (3.5)]).

Proposition 7.2. *With the notation as before, assume that $\chi|_{T^0}$ is not of the form $\alpha \circ \text{Det}$, where α is a character of $\mathfrak{o}^\times$ (necessarily of order 2). Then the induced representation $\text{Ind}_{J_\chi}^K \rho_\chi$ is irreducible. In particular, it is a type for $\mathcal{R}_{[T, \chi]}$.*

Proof. Let W be the extended affine Weyl group of G w.r.t. T and set $W_\chi = \{w \in W ; w\chi = \chi\}$. Then by Theorem (4.14) of [11], the G -intertwining of ρ_χ is $J_\chi W_\chi J_\chi$. The hypothesis on χ forces $W_\chi = T/T^0$. So $(J_\chi W_\chi J_\chi) \cap K = J_\chi T^0 J_\chi = J_\chi$, and we may apply Mackey's criterion of irreducibility. $\square$

For $n > 0$ and $q \in \{0, \dots, n\}$, define compact open subgroups of G as follows:

$${}_q\mathfrak{h}_1 = \begin{pmatrix} 1 + \mathfrak{p}^n & \mathfrak{p}^q \\ \mathfrak{p}^{n+1} & 1 + \mathfrak{p}^n \end{pmatrix} \text{ and } {}_q\mathfrak{h}_2 = \begin{pmatrix} 1 + \mathfrak{p}^{n+1} & \mathfrak{p}^q \\ \mathfrak{p}^{n+1} & 1 + \mathfrak{p}^{n+1} \end{pmatrix}.$$

These groups are particular cases of groups considered in [1, §(2.3)]. The quotients ${}_q\mathfrak{h}_1/{}_q\mathfrak{h}_2$, $q = 0, \dots, n$, are abelian, and for $\alpha \in \mathbf{k}^\times$, one may define a character ψ_α of ${}_q\mathfrak{h}_1/{}_q\mathfrak{h}_2$ by the formula

$$\psi_\alpha(I_2 + \begin{pmatrix} \varpi^n a & \varpi^q b \\ \varpi^{n+1} c & \varpi^n d \end{pmatrix}) = \psi(\alpha(a - d)),$$

where ψ is a fixed non-trivial character of $(\mathbf{k}, +)$. We shall need the following result.

Lemma 7.3. *If a smooth representation of K contains $(\psi_\alpha)_{|_{{}_n\mathfrak{h}_1}}$ by restriction, then it contains the character $(\psi_\alpha)_{|_{{}_0\mathfrak{h}_1}}$.*

Proof. Since the characteristic of $\mathbf{k}$ is not 2, then $\alpha \neq -\alpha$ and $(\psi_\alpha)_{|_{{}_n\mathfrak{h}_1}}$ is the restriction to ${}_n\mathfrak{h}_1$ of a split fundamental stratum of K_n/K_{n+1} . Our lemma is then a particular case of [1, Lemma (2.4.5)]. $\square$

Proposition 7.4. *Let λ be an irreducible constituent of $\text{Ind}_{T^0 K_m}^K H^1(\tilde{X}_{2m+1}[s_0, c_0])_{\text{split}}$. Then with the notation as above, λ is of the form $\text{Ind}_{J_\chi}^K \rho_\chi$, for some principal series type (J_χ, ρ_χ) with χ of conductor $m + 1$.*

Proof. We know that such a λ contains a split fundamental stratum of the form $[M(2, \mathfrak{o}), m, m - 1, b]$, where $b = \varpi^{-m} \begin{pmatrix} 0 & u \\ v & 0 \end{pmatrix}$, $u, v \in \mathfrak{o}^\times$, and uv is a square modulo $\mathfrak{p}$. If $\alpha \in \mathfrak{o}$ is such that $\alpha^2 \equiv uv \pmod{\mathfrak{p}}$, then the stratum is equivalent to a K -conjugate of $[M(2, \mathfrak{o}), m, m - 1, b']$, where $b' = \varpi^{-m} \begin{pmatrix} \alpha & 0 \\ 0 & -\alpha \end{pmatrix}$. So we deduce that λ contains this latter stratum by restriction. Now consider the group ${}_n\mathfrak{h}_1$ for $n = m$. The representation λ contains the character $(\psi_\alpha)_{|_{{}_n\mathfrak{h}_1}}$ by restriction. By applying Lemma 7.3 we obtain that it contains the character $(\psi_\alpha)_{|_{{}_0\mathfrak{h}_1}}$. This character clearly extends to $T^0{}_0\mathfrak{h}_1 = \Gamma_0(m + 1, 0)$ and the quotient $T^0{}_0\mathfrak{h}_1/{}_0\mathfrak{h}_1$ is abelian. It follows that λ contains an extension of ψ_α to $\Gamma_0(m + 1, 0)$. Such an extension is of the form (J_χ, ρ_χ) , for some character χ of T of conductor $m + 1$. The fact that λ is induced from (J_χ, ρ_χ) follows from Proposition 7.2. $\square$

8. SYNTHESIS

We now prove Theorems A and B of the Introduction.

By Propositions 4.2 and 4.4, we have isomorphisms of G -modules:

$$(8.1) \quad H_c^1(\tilde{X}_{2m}) \simeq H_c^1(\tilde{X}_{2m-1}) \oplus c\text{-ind}_{\mathcal{K}_1}^G H^1(\Sigma_{2m}), \quad m \geq 1,$$

$$(8.2) \quad H_c^1(\tilde{X}_{2m+1}) \simeq H_c^1(\tilde{X}_{2m}) \oplus c\text{-ind}_{\mathcal{K}_0}^G H^1(\Sigma_{2m+1}), \quad m \geq 0.$$

Recall that with the notation of the introduction, we have:

- $\Sigma_{2m} = \tilde{X}_{2m}[e_0]$, $\Sigma_{2m+1} = \tilde{X}_{2m+1}[s_0]$,
- $\mathcal{K}_0 = \mathcal{K}_{s_0}$, $\mathcal{K}_1 = \mathcal{K}_{e_0}$.

Moreover, by Proposition 5.2, we have

$$(8.3) \quad H_c^1(\tilde{X}_0) \simeq \mathrm{St}_G \oplus c\text{-ind}_{\mathcal{K}_1}^G H^1(\Sigma_0)$$

so that (1) holds for $m = 0$. Hence, Theorem A follows from (8.1) and (8.2) by a straightforward inductive argument.

Theorem B follows from the description of the irreducible components of $H^1(\Sigma_n)$ given in Proposition 5.1 (n even and $n > 0$), Proposition 5.2 ($n = 0$), and Propositions 6.1 and 7.4 (n odd).

REFERENCES

- [1] Paul Broussous, *Minimal strata for $\mathrm{GL}(m, D)$* , J. Reine Angew. Math. **514** (1999), 199–236, DOI 10.1515/crll.1999.071. MR1711267 (2000m:22020)
- [2] Paul Broussous, *Simplicial complexes lying equivariantly over the affine building of $\mathrm{GL}(N)$* , Math. Ann. **329** (2004), no. 3, 495–511, DOI 10.1007/s00208-004-0516-3. MR2127987 (2005m:20117)
- [3] Paul Broussous, *Representations of $\mathrm{PGL}(2)$ of a local field and harmonic cochains on graphs*, Ann. Fac. Sci. Toulouse Math. (6) **18** (2009), no. 3, 495–513 (English, with English and French summaries). MR2582439 (2011e:22021)
- [4] Colin J. Bushnell and Guy Henniart, *The local Langlands conjecture for $\mathrm{GL}(2)$* , Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 335, Springer-Verlag, Berlin, 2006. MR2234120 (2007m:22013)
- [5] Colin J. Bushnell and Philip C. Kutzko, *The admissible dual of $\mathrm{GL}(N)$ via compact open subgroups*, Annals of Mathematics Studies, vol. 129, Princeton University Press, Princeton, NJ, 1993. MR1204652 (94h:22007)
- [6] Colin J. Bushnell and Philip C. Kutzko, *Smooth representations of reductive p -adic groups: structure theory via types*, Proc. London Math. Soc. (3) **77** (1998), no. 3, 582–634, DOI 10.1112/S0024611598000574. MR1643417 (2000c:22014)
- [7] Armand Borel and Jean-Pierre Serre, *Cohomologie à supports compacts des immeubles de Bruhat-Tits; applications à la cohomologie des groupes S -arithmétiques*, C. R. Acad. Sci. Paris Sér. A-B **272** (1971), A110–A113 (French). MR0274456 (43 #221)
- [8] H. Carayol, *Représentations cuspidales du groupe linéaire*, Ann. Sci. École Norm. Sup. (4) **17** (1984), no. 2, 191–225 (French). MR760676 (86f:22019)
- [9] G. Henniart, *Sur l’unicité des types pour GL_2* , appendix to *Multiplicité modulaires et représentations de $\mathrm{GL}(2, \mathbb{Z}_p)$ et $\mathrm{Gal}(\bar{\mathbb{Q}}_p/\mathbb{Q}_p)$ en $l = p$* , by C. Breuil and A. Mézard.
- [10] P. C. Kutzko, *On the supercuspidal representations of Gl_2* , Amer. J. Math. **100** (1978), no. 1, 43–60. MR0507253 (58 #22411a)
- [11] Alan Roche, *Types and Hecke algebras for principal series representations of split reductive p -adic groups*, Ann. Sci. École Norm. Sup. (4) **31** (1998), no. 3, 361–413, DOI 10.1016/S0012-9593(98)80139-0 (English, with English and French summaries). MR1621409 (99d:22028)

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE POITIERS, TÉLÉPORT 2 - BP 30179, BOULEVARD MARIE ET PIERRE CURIE, 86962 FUTUROSCEPE CHASSENEUIL CEDEX, FRANCE

E-mail address: paul.broussous@math.univ-poitiers.fr