CORRECTIONS TO: “A MURNAGHAN–NAKAYAMA RULE FOR VALUES OF UNIPOTENT CHARACTERS IN CLASSICAL GROUPS”

FRANK LÜBECK AND GUNTER MALLE

ABSTRACT. We settle a missing case in the proof of one of the main applications of our results in [Frank Lübeck and Gunter Malle, *A Murnaghan–Nakayama rule for values of unipotent characters in classical groups*, Represent. Theory **20** (2016), 139–161, MR3466537].

In [5] we derived a Murnaghan–Nakayama formula for the values of unipotent characters of finite classical groups. As one application, we showed that the first Cartan invariant $c_{11}^{(\ell)}$ of a finite simple classical group G in non-defining characteristic ℓ is never equal to 2 if Sylow ℓ -subgroups of G are non-cyclic. But contrary to what was claimed in the first line of the proof of [5, Thm. 7.1], the same assertion for the case of defining characteristic does not follow from the work of Koshitani, Külshammer, and Sambale [2]. We thank Shigeo Koshitani and Jürgen Müller for pointing this out to us; see [3].

Here we give a short argument for the missing case. The argument also covers the analogous statement in [4, Prop. 6.3] pertaining to simple groups of exceptional Lie type. The crucial observation is that Sylow p -subgroups in groups of Lie type of characteristic p are large. Except for a few small rank cases, we do not need any knowledge about the ordinary character table of the group G .

Note that if $c_{11}^{(p)} = 2$, then there exists $\chi \in \text{Irr}(G)$ such that $1_G + \chi$ is the character of a projective module; in particular, $\chi(g) = -1$ for all p -singular elements $g \in G$.

Lemma 1. *Let G be a finite group and p a prime. Assume that $H \leq G$ is a subgroup such that $|G : H|$ is divisible by p , but $|G : H| < |G|_p$. Then $c_{11}^{(p)} > 2$.*

Proof. The permutation character π of G on the cosets of H has exactly one trivial constituent 1_G . On the other hand, as $|G : H|$ is divisible by p , the modular reduction of the permutation representation has (at least) two trivial modular constituents: one in the socle, one in the head. Thus, the p -modular reduction of one of the non-trivial ordinary constituents χ of π must contain the trivial p -modular character of G . By Brauer reciprocity, χ then occurs in the projective cover Ψ of the trivial character. If we had $c_{11}^{(p)} = 2$, then necessarily $\Psi = 1_G + \chi$. Since the degree of any projective character is divisible by $|G|_p$, this is a contradiction to

$$|G|_p \leq \Psi(1) = 1_G(1) + \chi(1) \leq \pi(1) = |G : H|. \quad \square$$

Received by the editors October 25, 2016.

2010 *Mathematics Subject Classification.* Primary 20C20; Secondary 20C33.

Key words and phrases. Finite simple groups, Cartan invariants.

Proposition 2. *Let G be a finite simple group of Lie type in characteristic p . Then $c_{11}^{(p)} > 2$ unless $G = L_2(p)$.*

Proof. If the Lie-rank of G is not very small, then we can find a subgroup of G which fulfills the condition of Lemma 1. In Table 1 we give a subgroup $\hat{H}$ (see for example [6, §13] for the existence) of a suitable central extension $\hat{G}$ of G . From the order formulae for these groups (see for example [6, Tab. 24.1]) it is easy to check that $\hat{H} \leq \hat{G}$ fulfills the condition of Lemma 1. Since the center of $\hat{G}$ is of order prime to p , the same holds for $H \leq G$ where H is the image of $\hat{H}$ in G .

It remains to consider the groups of type A_1 , A_2 , 2A_2 , 3D_4 , 2B_2 , 2G_2 and 2F_4 . For all of these the (generic) character tables are known and available in CHEVIE [1]. Here we look for all irreducible characters χ which have value -1 on all p -singular (non-semisimple) elements, so that $\chi + 1_G$ is a candidate for the 1-PIM of G . It is easy to see that, apart from two exceptions, all characters either: vanish on some p -singular element, or that a value on a non-trivial unipotent class is positive, or that the values on two non-trivial unipotent classes are different.

The two exceptions occur in type A_1 , so $G = L_2(q)$, which has a family of characters of degree $q - 1$ with value -1 on all p -singular elements, and in type A_2 , so $G = L_3(q)$, which has a family of characters of degree $q^3 - 1$ which also take value -1 on all p -singular elements. By [7, Thm. 5.2] the 1-PIM of $G = L_2(q)$ has degree $(2^f - 1)q$, where $q = p^f$. Since the maximal character degree for G is $q + 1$, this shows that $c_{11}^{(p)} > 2$ unless $f = 1$; that is, unless $q = p$ is prime, in which case the Sylow p -subgroups of G are cyclic (and so indeed $c_{11}^{(p)} = 2$). Similarly, by [7, Thm. 5.3] the 1-PIM of $G = L_3(q)$, $q = p^f$, has degree $(12^f - 6^f + 1)q^3$, so none of the candidate $1_G + \chi$ mentioned above can be the 1-PIM. $\square$

TABLE 1. Large subgroups

$\hat{G}$		$\hat{H}$	$\hat{G}$	$\hat{H}$
$SL_4^\pm(q)$		$Sp_4(q)$	$G_2(q)$	${}^2A_2(q)$
$SL_n^\pm(q)$	$(n \geq 5)$	$GL_{n-1}^\pm(q)$	$F_4(q)$	$B_4(q)$
$Spin_{2n+1}(q)$	$(n \geq 2)$	$Spin_{2n}^-(q)$	$E_6^\pm(q)$	$F_4(q)$
$Sp_{2n}(q)$	$(n \geq 3)$	$Sp_{2n-2}(q)Sp_2(q)$	$E_7(q)$	$E_6(q)$
$Spin_{2n}^\pm(q)$	$(n \geq 4)$	$Spin_{2n-1}(q)$	$E_8(q)$	$E_7(q)$

Observe that the case $p = 2$ was already dealt with in [2, Thm. 4.5].

REFERENCES

- [1] Meinolf Geck, Gerhard Hiss, Frank Lübeck, Gunter Malle, and Götz Pfeiffer, *CHEVIE—a system for computing and processing generic character tables*, Appl. Algebra Engrg. Comm. Comput. **7** (1996), no. 3, 175–210, DOI 10.1007/BF01190329. Computational methods in Lie theory (Essen, 1994). MR1486215
- [2] Shigeo Koshitani, Burkhard Külshammer, and Benjamin Sambale, *On Loewy lengths of blocks*, Math. Proc. Cambridge Philos. Soc. **156** (2014), no. 3, 555–570, DOI 10.1017/S0305004114000103. MR3181640
- [3] S. Koshitani, J. Müller, *The projective cover of the trivial representation for a finite group of Lie type in defining characteristic*. *Algebra Colloq.*, to appear, ArXiv: 1609.08070.

- [4] Caroline Lassueur and Gunter Malle, *Simple endotrivial modules for linear, unitary and exceptional groups*, Math. Z. **280** (2015), no. 3-4, 1047–1074, DOI 10.1007/s00209-015-1465-0. MR3369366
- [5] Frank Lübeck and Gunter Malle, *A Murnaghan–Nakayama rule for values of unipotent characters in classical groups*, Represent. Theory **20** (2016), 139–161, DOI 10.1090/ert/480. MR3466537
- [6] Gunter Malle and Donna Testerman, *Linear algebraic groups and finite groups of Lie type*, Cambridge Studies in Advanced Mathematics, vol. 133, Cambridge University Press, Cambridge, 2011. MR2850737
- [7] Gunter Malle and Thomas Weigel, *Finite groups with minimal 1-PIM*, Manuscripta Math. **126** (2008), no. 3, 315–332, DOI 10.1007/s00229-008-0176-9. MR2411231

LEHRSTUHL D FÜR MATHEMATIK, RWTH AACHEN, PONTDRIESCH 14/16, 52062 AACHEN, GERMANY.

E-mail address: Frank.Luebeck@math.rwth-aachen.de

FB MATHEMATIK, TU KAISERSLAUTERN, POSTFACH 3049, 67653 KAISERSLAUTERN, GERMANY.

E-mail address: malle@mathematik.uni-kl.de