ALTERNATING SIGNS OF QUIVER COEFFICIENTS

ANDERS SKOVSTED BUCH

1. INTRODUCTION

Let X be a non-singular algebraic variety and $E_0 \rightarrow E_1 \rightarrow \cdots \rightarrow E_n$ a sequence of vector bundles and bundle maps over X . A set of *rank conditions* for this sequence is a collection $r = \{r_{ij}\}$ of non-negative integers, for $0 \leq i < j \leq n$. This data defines the quiver variety

$$\Omega_r(E_\bullet) = \{x \in X \mid \text{rank}(E_i(x) \rightarrow E_j(x)) \leq r_{ij} \ \forall i < j\},$$

which comes with a natural structure of subscheme of X , given as the intersection of the zero sections of the maps $\bigwedge^{r_{ij}+1} E_i \rightarrow \bigwedge^{r_{ij}+1} E_j$. We demand that the rank conditions can *occur* as the ranks over a point in X . If we set $r_{ii} = \text{rank}(E_i)$, then this is equivalent to the conditions that $r_{ij} \leq \min(r_{i,j-1}, r_{i+1,j})$ for all $0 \leq i < j \leq n$ and $r_{ij} + r_{i-1,j+1} \geq r_{i-1,j} + r_{i,j+1}$ for all $0 < i \leq j < n$. In this case the expected codimension of the quiver variety $\Omega_r(E_\bullet)$ is the integer $d(r) = \sum_{i < j} (r_{i,j-1} - r_{ij})(r_{i+1,j} - r_{ij})$.

In joint work with Fulton [10] we established a formula for the cohomology (or Chow) class of the quiver variety when this codimension is attained. This was generalized in [7] to the following formula for the structure sheaf of a quiver variety in the Grothendieck ring $K(X)$ of algebraic vector bundles on X :

$$(1) \quad [\mathcal{O}_{\Omega_r(E_\bullet)}] = \sum_{\mu} c_{\mu}(r) G_{\mu_1}(E_1 - E_0) G_{\mu_2}(E_2 - E_1) \cdots G_{\mu_n}(E_n - E_{n-1}).$$

Here the sum is over finitely many sequences $\mu = (\mu_1, \dots, \mu_n)$ of partitions μ_i such that the sum $\sum |\mu_i|$ of the weights is greater than or equal to $d(r)$. The stable Grothendieck polynomials $G_{\mu_i}(E_i - E_{i-1})$ are defined in Section 2. The *quiver coefficients* $c_{\mu}(r)$ appearing in this formula are integers which are uniquely determined by the condition that (1) is true for all varieties X and bundle sequences $E_\bullet$, together with the condition that $c_{\mu}(r) = c_{\mu}(r + m)$ holds for all $m \in \mathbb{N}$, where $r + m = \{r_{ij} + m\}$ are the rank conditions obtained by adding the integer m to the original rank conditions.

A formula for quiver coefficients was also given in [10, 7]; in the case of K -theory, this is based on the algebra of stable Grothendieck polynomials constructed in [8]. Although the original formulas for quiver coefficients do not keep track of their signs, it was conjectured that the cohomological quiver coefficients (given by sequences μ such that $\sum |\mu_i| = d(r)$) are non-negative, while the K -theoretic quiver coefficients

Received by the editors January 5, 2004.

2000 *Mathematics Subject Classification*. Primary 05E15; Secondary 14M15, 14M12, 19E08.

Key words and phrases. Quiver variety, quiver coefficient, degeneracy locus, Grothendieck polynomial, Zelevinsky permutation, factor sequence.

©2004 American Mathematical Society
 Reverts to public domain 28 years from publication

have signs that alternate with codimension, i.e. $(-1)^{\sum |\mu_i| - d(r)} c_\mu(r) \geq 0$. Special cases of these conjectures have been proved in [10, 5, 6, 7, 12, 13].

In their recent paper [24], Knutson, Miller, and Shimozono deliver a breakthrough within the theory of quiver formulas and prove at least two explicit combinatorial formulas for the cohomological quiver coefficients, which show that these coefficients are non-negative. One of the important ideas in their work is to reinterpret the *lace diagrams* of Abeasis and Del Fra [1] as sequences of partial permutations. This interpretation is explained by a Gröbner degeneration of a quiver variety in a matrix space into a union of products of matrix Schubert varieties. The *component formula* of [24] writes the cohomology class of a quiver variety as a sum, over all ‘*minimal*’ lace diagrams, of the products of the Schubert polynomials for the corresponding partial permutations. The proof that quiver coefficients are non-negative is obtained by proving a stable version of this component formula, where the Schubert polynomials are replaced with Stanley symmetric functions. This is sufficient because Stanley symmetric functions are known to be Schur positive [15, 28].

Knutson, Miller, and Shimozono also prove a *ratio formula*, which writes the class of a quiver variety as a quotient of two Schubert polynomials. This formula follows from a careful analysis of the Zelevinsky map [34, 25] and is in fact established for both cohomology and K -theory. The component formulas are proved using a combination of the Gröbner degeneration and the ratio formula, as well as a combinatorial study of a double version of the ratio formula. In particular, it is proved that a limit of the double ratio formula agrees with the *double quiver functions* introduced in [10, 7] and named in [24]. The authors of [24] have informed us that they can generalize their methods to work in K -theory, although, according to their own description, this approach is rather complicated.¹

In this paper we give simpler proofs of the above-mentioned formulas, using methods that work equally well in K -theory. In particular, we prove that the K -theoretic quiver coefficients have alternating signs, and we derive an explicit combinatorial formula for these coefficients. Starting from the ratio formula, we give combinatorial proofs of K -theoretic generalizations of the component formulas, where the Schubert polynomials and Stanley symmetric functions are replaced with ordinary and stable Grothendieck polynomials. These formulas are given in terms of sequences of partial permutations, which we call *KMS-factorizations* of the *Zelevinsky permutation* [34, 24]. To conclude that K -theoretic quiver coefficients have alternating signs and to derive the combinatorial formula for these coefficients, we use Lascoux’s result that stable Grothendieck polynomials are linear combinations with alternating signs of stable Grothendieck polynomials for partitions [27]. To make our paper self-contained, we also give a short proof of the ratio formula.

The *factor sequences conjecture* of [10] states that cohomological quiver coefficients count the number of sequences of semistandard Young tableaux, which can be generated by a sequence of plactic factorizations and multiplications of chosen tableaux arranged in a *tableaux diagram*. A special case of this conjecture, corresponding to a particular choice of tableaux diagram, was proved in [24]. However, so far there has been no progress in generalizing this conjecture to K -theory. In this paper we close this gap by showing that KMS-factorizations can be defined by

¹This work is now available in the paper [31], which proves a K -theoretic component formula and a double version of it, but only in the stable case and for large rank conditions.

the same algorithm as defines factor sequences, except that the tableau diagram is replaced with a diagram of permutations and the plactic product of tableaux is replaced with multiplication of permutations in the degenerate Hecke algebra. It is also possible to formulate a new tableau-based version of the factor sequences conjecture. We briefly explain how our description of KMS-factorizations imply that cohomological quiver coefficients are counted by a new type of factor sequences of tableaux, which are generated using the Coxeter-Knuth product rather than the plactic product. In a paper in preparation with Kresch, Shimozono, Tamvakis, and Yong [11], the new tableau-based factor sequences have been generalized to work in K -theory.

We remark that it was already known that the cohomological component formulas can be derived combinatorially from the ratio formula, by using a simplification of Yong [33]. However, Yong's method still requires the analysis of the double ratio formula and its limits from [24]. The approach presented here simplifies things further by working only with the single ratio formula, by applying Fomin and Kirillov's construction of Grothendieck polynomials based on solutions to Yang-Baxter equations [19, 18], and by observing that the stable component formula follows easily from the non-stable component formula.

Other simple proofs of the cohomological component formulas have also surfaced. For example, they can be deduced very easily from the Thom polynomial theory developed by Fehér and Rimányi [16] or deduced directly from the above-mentioned Gröbner degeneration with a symmetry argument. This is explained in [9]. While attempts to generalize these methods to K -theory have not been successful, they might hold more promise for quiver varieties of other types (see [16]).

Some of the results proved in [24] imply that the cohomological double ratio formula for large rank conditions satisfies nice properties, including *multisupersymmetry* and a double version of the component formula. In the last section of this paper, we establish these properties for the K -theoretic double ratio formula given by arbitrary rank conditions. In particular, we prove a conjecture from [24] stating that the double ratio formula satisfies a *rank stability* property. This conjecture is equivalent to the statement that the polynomials defined by the double ratio formula are specializations of the original quiver formulas [10, 7]. Even though the double ratio formula is not needed for the proof of alternating signs of quiver coefficients given in this paper, its multisupersymmetry property has some nice applications. For example, this property was used in [24] to prove the above-mentioned case of the original factor sequences conjecture. The multisupersymmetry property also implies that general quiver coefficients are special cases of the coefficients studied in [13]. In fact, quiver coefficients can be realized as Schubert structure constants on flag varieties [3, 30, 14].

This paper is organized as follows. In Section 2 we explain Fomin and Kirillov's construction of Grothendieck polynomials, which is a key ingredient in the combinatorial parts of this paper. Section 3 gives a new construction of the Zelevinsky permutation which is required for our proof that KMS-factorizations can be viewed as factor sequences. Section 4 contains the proof of the ratio formula. In Section 5 we prove a formula for double Grothendieck polynomials applied to certain rearrangements of the same set of variables, which in Section 6 is used to derive the non-stable component formula from the ratio formula. In Section 7 we establish the factor sequences definition of KMS-factorizations and discuss its consequences.

As a corollary we obtain a rank stability property for KMS-factorizations, which in Section 8 is used to derive the stable component formula and deduce that quiver coefficients have alternating signs. Section 9 finally proves the above-mentioned properties of the double ratio formula.

We are very grateful to Richárd Rimányi for discussions at the Banach Center in Warsaw, which led to our observation that the stable component formula can be deduced from the non-stable formula and which triggered our search for other simplifications to [24]. We also thank Fehér, Kresch, Sottile, Tamvakis, and Yong for inspiring collaboration on related papers, and Fulton, Miller, and Sottile for helpful comments about our paper. Finally, we thank Martin Guest, Anatol Kirillov, and the Research Institute for Mathematical Sciences in Kyoto for their hospitality while this paper was written.

2. GROTHENDIECK POLYNOMIALS

The *degenerate Hecke algebra* $\mathcal{H}$ over a commutative ring R is the free R -algebra generated by symbols $s_1, s_2, \dots$, modulo the relations

$$\begin{aligned} s_i s_j &= s_j s_i & \text{if } |i - j| \geq 2, \\ s_i s_{i+1} s_i &= s_{i+1} s_i s_{i+1}, \\ s_i^2 &= -s_i. \end{aligned}$$

In this paper, R will be a ring of Laurent polynomials. The algebra $\mathcal{H}$ is a free R -module with a basis of permutations corresponding to reduced expressions in the generators.

Given permutations $u_1, u_2, \dots, u_n$, the product $u_1 \cdot u_2 \cdots u_n$ in $\mathcal{H}$ of these permutations is equal to plus or minus a single permutation w . We will call this permutation w for the *absolute Hecke product* of the u_i . Notice that the descent positions of w , i.e. the indices i for which $w(i) > w(i+1)$, include the descent positions of u_n , while the descent positions of u_1^{-1} are also descent positions of w^{-1} . Notice also that if $u_1 u_2 \cdots u_n$ is a *reduced product* of permutations in the sense that $\ell(u_1 u_2 \cdots u_n) = \sum \ell(u_i)$, then the absolute Hecke product w agrees with the usual product of permutations.

We will need Fomin and Kirillov's construction [19, 18] of the Grothendieck (Laurent) polynomials $\mathfrak{G}_w(a; b)$ of Lascoux and Schützenberger [28, 26]. Consider a diagram $\mathfrak{D}$ of strings going from a top horizontal border to a left vertical border. Each string must be composed of line segments, each of which is labeled with a variable and has a direction between due south and due west. Furthermore, strings may only cross each other transversally, at inner points of the line segments. Of particular importance is the following diagram $\mathfrak{D}_N$, which contains only horizontal and vertical line segments.

For the diagram $\mathfrak{D}_N$, we identify the crossing point of the horizontal line labeled b_p and the vertical line labeled a_q , with the point $(p, q) \in \mathbb{N} \times \mathbb{N}$. We then have $\nu(p, q) = p + q - 1$, and the FK-product of $\mathfrak{D}_N$ is given by

$$\mathfrak{G}(\mathfrak{D}_N) = \prod_{q=1}^{N-1} \prod_{p=N-q}^1 \left(1 + \left(1 - \frac{b_p}{a_q} \right) s_{p+q-1} \right) \in \mathcal{H}.$$

Theorem 2.1 (Fomin and Kirillov). *In $\mathcal{H}$ we have the identity*

$$\mathfrak{G}(\mathfrak{D}_N) = \sum_{w \in S_N} \mathfrak{G}_w(a; b) \cdot w$$

Suppose D is a subset of the crossing positions $C(\mathfrak{D})$ of a diagram $\mathfrak{D}$. We let $w(D)$ be the absolute Hecke product of the simple reflections $s_{\nu(P)}$ for $P \in D$, in south-west to north-east order as above. We say that D is an *FK-graph* for this permutation $w(D)$ and that D is *reduced* if $|D| = \ell(w(D))$. We can picture an FK-graph D by replacing the crossing positions of $\mathfrak{D}$ which belong to this graph with the symbol “ $\frac{\downarrow}{\uparrow}$ ”, while the remaining crossing positions are replaced with the symbol “ $\swarrow\searrow$ ”. Notice that if D is reduced, then the string entering the resulting diagram at column i at the top will exit at row $w(D)(i)$ at the left-hand side. Notice also that any FK-graph D contains a reduced FK-graph $D' \subset D$ such that $w(D') = w(D)$. In fact, D' can be found by simply skipping the points $P \in D$ for which $s_{\nu(P)}$ does not increase the length when the product $w(D)$ is formed.

$$D = \begin{array}{c} \text{---} \diagup \text{---} \diagup \text{---} \diagup \\ | \quad | \quad | \\ \text{---} \text{---} \diagup \\ | \quad | \\ \text{---} \text{---} \diagup \\ | \\ \text{---} \text{---} \diagup \\ | \\ \text{---} \end{array}$$

It follows from the definitions that the coefficient of a permutation w in the FK-product of a diagram $\mathfrak{D}$ is equal to

$$(2) \quad \sum_{D \subset C(\mathfrak{D}), w(D)=w} (-1)^{|D|-\ell(w)} \prod_{P \in D} h(P).$$

Theorem 2.1 therefore has the following corollary. See also [23] for an alternative proof and [2] for the case of Schubert polynomials.

Corollary 2.2. *For any permutation w we have*

$$\mathfrak{G}_w(a; b) = \sum_{w(D)=w} (-1)^{|D|-\ell(w)} \prod_{(p,q) \in D} \left(1 - \frac{b_p}{a_q}\right)$$

where the sum is over all FK-graphs $D \subset \mathbb{N} \times \mathbb{N}$ for w .

Let us remark that Theorem 2.1 is more flexible than its corollary, as amply demonstrated in [19]. The point is that many operations can be performed on a diagram $\mathfrak{D}$ without changing the corresponding FK-product. We will write $\mathfrak{D} \approx \mathfrak{D}'$ if $\mathfrak{G}(\mathfrak{D}) = \mathfrak{G}(\mathfrak{D}')$. The two key examples of this are

$$(3) \quad \begin{array}{c} x \\ | \\ y \text{---} \text{---} z \end{array} \approx \begin{array}{c} x \\ | \\ y \text{---} \text{---} \end{array} \quad \text{and} \quad \begin{array}{c} x \\ | \\ y \text{---} \text{---} z \end{array} \approx \begin{array}{c} x \\ | \\ y \text{---} \text{---} z \end{array}$$

Notice that when $x = y$, the last diagram is also equivalent to a north to west hook labeled x together with a disjoint north-east to south-west line labeled z . Although most diagrams in this paper contain only horizontal and vertical line segments, the availability of slanted lines often makes it more natural to manipulate these diagrams using the rules of (3).

Lascoux and Schützenberger's original definition of Grothendieck polynomials says that $\mathfrak{G}_{w_0}(a; b) = \prod_{p+q \leq N} (1 - b_p/a_q)$ where $w_0 \in S_N$ is the longest permutation, and that $(a_i - a_{i+1})\mathfrak{G}_w(a; b) = a_i\mathfrak{G}_{ws_i}(a; b) - a_{i+1}\mathfrak{G}_{ws_i}(a_{s_i}; b)$ when $w(i) < w(i+1)$. Here we let a_v denote the sequence of variables $a_{v(1)}, \dots, a_{v(N)}$ for any permutation $v \in S_N$. It follows that if i is not a descent position for w , then $\mathfrak{G}_w(a; b)$ is symmetric in the variables a_i and a_{i+1} . In particular, if k is the last descent position for w , then the variables a_i for $i > k$ do not occur in $\mathfrak{G}_w(a; b)$. A similar relationship holds between the descent positions of w^{-1} and the variables b_i . We need the identity

$$(4) \quad \mathfrak{G}_w(a_{v(1)}, \dots, a_{v(N)}; a_1, \dots, a_N) = \mathfrak{G}_{w_0 w^{-1} w_0}(a_N, \dots, a_1; a_{v(N)}, \dots, a_{v(1)}).$$

In fact, using that $\mathfrak{G}_w(a; b) = \mathfrak{G}_{w^{-1}}(b_1^{-1}, \dots, b_N^{-1}; a_1^{-1}, \dots, a_N^{-1})$ and the above definition, equation (4) follows by descending induction on $\ell(w)$ from the calculation

$$\begin{aligned} (a_{v(i)} - a_{v(i+1)})\mathfrak{G}_w(a_v; a) &= a_{v(i)}\mathfrak{G}_{ws_i}(a_v; a) - a_{v(i+1)}\mathfrak{G}_{ws_i}(a_{vs_i}; a) \\ &= a_{v(i)}\mathfrak{G}_{w_0 s_i w^{-1} w_0}(a_{w_0}; a_{vw_0}) - a_{v(i+1)}\mathfrak{G}_{w_0 s_i w^{-1} w_0}(a_{w_0}; a_{vs_i w_0}) \\ &= (a_{v(i)} - a_{v(i+1)})\mathfrak{G}_{w_0 w^{-1} w_0}(a_{w_0}; a_{vw_0}). \end{aligned}$$

The identity (4) also follows from the results in [29].

We also need the *stable double Grothendieck polynomials* $G_w(a; b)$ of Fomin and Kirillov [18]. These polynomials are characterized by the property that

$$G_w(a_1, \dots, a_q; b_1, \dots, b_p) = \mathfrak{G}_{1^m \times w}(a_1, \dots, a_q, 1, \dots, 1; b_1, \dots, b_p, 1, \dots, 1)$$

for all $m \geq \max(p, q)$. Here the permutation $1^m \times w$ is the identity on $\{1, \dots, m\}$ while it maps j to $w(j - m) + m$ for $j > m$. The polynomial $G_w(a; b)$ is separately symmetric in its two sets of variables.

If $\lambda = (\lambda_1 \geq \dots \geq \lambda_k)$ is a partition, we set $G_\lambda(a; b) = G_{w_\lambda}(a; b)$ where w_λ is the Grassmannian permutation for λ , defined by $w_\lambda(i) = i + \lambda_{k+1-i}$ for $1 \leq i \leq k$

and $w_\lambda(i) < w_\lambda(i+1)$ for $i \neq k$. It is proved in [8] that any stable Grothendieck polynomial $G_w(a; b)$ can be written as an integral linear combination

$$(5) \quad G_w(a; b) = \sum_{\lambda} c_{w, \lambda} G_{\lambda}(a; b)$$

of stable Grothendieck polynomials for partitions. We will need a combinatorial formula of Lascoux for the coefficients $c_{w, \lambda}$ in this expansion, which shows that they have alternating signs.

Let r be the last descent position of w , i.e. r is maximal such that $w(r) > w(r+1)$. Set $w' = w\tau_{rk}$ where $k > r$ is maximal such that $w(r) > w(k)$, and τ_{rk} is the transposition interchanging r and k . We also set $I(w) = \{i < r \mid \ell(w'\tau_{ir}) = \ell(w)\}$. Then define a relation $\triangleright$ on the set of all permutations as follows. If $I(w) = \emptyset$, then we write $w \triangleright v$ if and only if $v = 1 \times w$. Otherwise we write $w \triangleright v$ if and only if there exist elements $i_1 < \dots < i_p$ of $I(w)$, $p \geq 1$, such that $v = w'\tau_{i_1 r} \dots \tau_{i_p r}$. The following result was proved in [27, Thm. 4].

Theorem 2.3 (Lascoux). *The coefficient $c_{w, \lambda}$ of (5) is equal to $(-1)^{|\lambda| - \ell(w)}$ times the number of paths $w = w_1 \triangleright w_2 \triangleright \dots \triangleright w_m$, such that $w_m = w_\lambda$ is a Grassmannian permutation for λ and w_i is not Grassmannian for $i < m$.*

Given vector bundles $F = L_1 \oplus \dots \oplus L_p$ and $H = M_1 \oplus \dots \oplus M_q$ over X which are direct sums of line bundles, we write $G_\lambda(H - F) = G_\lambda(M_1, \dots, M_q; L_1, \dots, L_p) \in K(X)$. By the symmetry of G_λ , this is a polynomial in the exterior powers of F and $H^\vee$, so the notation $G_\lambda(H - F)$ also makes sense for bundles which are not sums of line bundles. This explains the notation used in the quiver formula (1).

3. THE ZELEVINSKY PERMUTATION

In this section we give a new construction of the Zelevinsky permutation [34, 24], which is needed for our proof of the K -theoretic analogue of the factor sequences conjecture. To be precise, we construct the conjugate of the Zelevinsky permutation defined in [24], which turns out to have a simpler relationship with the geometry of quiver varieties and KMS-factorizations. However, the Zelevinsky permutation itself is necessary to obtain nice combinatorial properties of the ratio formula.

Extend the set of rank conditions r by setting $r_{ij} = e_j + e_{j+1} + \dots + e_i$ for $j \leq i$, where $(e_0, \dots, e_n) = (r_{00}, \dots, r_{nn})$ is the dimension vector corresponding to r . For $i < 0$ or $j > n$ we set $r_{ij} = 0$, and we set $N = r_{n0} = e_0 + \dots + e_n$. For each $0 \leq i < n$ and $0 < j \leq n$ we define a permutation $W_{ij} \in S_{r_{i+1, j-1}}$ by the expression

$$W_{ij}(p) = \begin{cases} p + r_{i, j-1} - r_{ij} & \text{if } r_{ij} < p \leq r_{i+1, j}, \\ p - r_{i+1, j} + r_{ij} & \text{if } r_{i+1, j} < p \leq r_{i+1, j} + r_{i, j-1} - r_{ij}, \\ p & \text{otherwise.} \end{cases}$$

When $i < j$, this is the Grassmannian permutation for the rectangular partition R_{ij} from [10], with descent at position $r_{i+1, j}$. Now define the *conjugate Zelevinsky permutation* $z(r) \in S_N$ for the rank conditions r to be the south-west to north-east product of the matrix of permutations

$$(6) \quad \begin{array}{cccc} W_{n-1,1} & W_{n-1,2} & \cdots & W_{n-1,n} \\ W_{n-2,1} & W_{n-2,2} & \cdots & W_{n-2,n} \\ \vdots & \vdots & \ddots & \vdots \\ W_{0,1} & W_{0,2} & \cdots & W_{0,n} \end{array},$$

that is, $z(r) = \prod_{j=1}^n \prod_{i=0}^{n-1} W_{ij}$. Notice that a ‘south-west to north-east product’ makes sense because W_{ij} commutes with $W_{i'j'}$ whenever $i < i'$ and $j' < j$. Notice also that W_{ij} is the conjugate Zelevinsky permutation for the set of rank conditions consisting of the integers $r_{i,j-1}$, $r_{i+1,j}$, and r_{ij} . Let $w_0^{(N)}$ denote the longest permutation in S_N . We will call the permutation $\hat{z}(r) = w_0^{(N)} z(r)^{-1} w_0^{(N)}$ the *Zelevinsky permutation* for r , although the definition in [24] assigns this name to the inverse of $\hat{z}(r)$. The action of $z(r)$ can be described explicitly as follows.

Lemma 3.1. *Given any integer $1 \leq p \leq N$, there are unique integers i and j with $0 \leq i \leq j+1 \leq n+1$, such that $r_{n,j+1} + r_{i-1,j} - r_{i-1,j+1} < p \leq r_{n,j+1} + r_{i,j} - r_{i,j+1}$. We then have $z(r)(p) = p - r_{n,j+1} + r_{i,j+1} + r_{i-1,0} - r_{i-1,j}$.*

Proof. To find i and j , one first chooses j such that $r_{n,j+1} < p \leq r_{nj}$, after which i is uniquely determined. Now, when $z(r)$ is applied to p , the permutation $W_{k,j+1}$ subtracts $r_{k+1,j+1} - r_{k,j+1}$ from its argument for $k = n-1, n-2, \dots, i$, after which $W_{i-1,k}$ adds $r_{i-1,k-1} - r_{i-1,k}$ to its argument for $k = j, j-1, \dots, 1$. All other factors of $z(r)$ preserve their argument. $\square$

The relation of the Zelevinsky permutation with quiver varieties is based on the following lemma, which is equivalent to [24, Prop. 1.6].

Lemma 3.2. *The conjugate Zelevinsky permutation $z(r)$ is the unique permutation in S_N such that*

- (i) *all descent positions of $z(r)$ are contained in the set $\{r_{nj} \mid 0 < j \leq n\}$,*
- (ii) *all descent positions of $z(r)^{-1}$ are contained in $\{r_{i0} \mid 0 \leq i < n\}$, and*
- (iii) *for all $0 \leq i, j \leq n$, there are r_{ij} integers $p \leq r_{nj}$ for which $z(r)(p) \leq r_{i0}$.*

Proof. Part (i) follows directly from Lemma 3.1, and (ii) follows by observing that $z(r)^{-1}$ is the conjugate Zelevinsky permutation for the mirrored rank conditions r' given by $r'_{ij} = r_{n-j,n-i}$. Notice that if p, i, j are chosen as in Lemma 3.1, then $r_{i-1,0} < z(r)(p) \leq r_{i,0}$. Therefore there are exactly $r_{i,j} - r_{i,j+1} - r_{i-1,j} + r_{i-1,j+1}$ integers p such that $r_{n,j+1} < p \leq r_{n,j}$ and $r_{i-1,0} < z(r)(p) \leq r_{i,0}$. This proves (iii) since $r_{ij} = \sum_{k=0}^i \sum_{l=j}^n (r_{kl} - r_{k,l+1} - r_{k-1,l} + r_{k-1,l+1})$. The uniqueness statement is not needed in this paper, and its easy proof is left to the reader. $\square$

Lemma 3.2 implies that the length of $z(r)$ is given by

$$(7) \quad \ell(z(r)) = \sum_{0 \leq i < n, 0 < j \leq n} (r_{i,j-1} - r_{ij})(r_{i+1,j} - r_{ij}).$$

In fact, there are $(r_{i,j-1} - r_{ij})(r_{i+1,j} - r_{ij})$ pairs (p, q) for which $p \leq r_{nj} < q \leq r_{n,j-1}$ and $z(r)(q) \leq r_{i0} < z(r)(p) \leq r_{i+1,0}$. In particular, $z(r)$ is a reduced product of the permutations W_{ij} of the matrix (6).

We let $z(e) = z(r^e)$ be the conjugate Zelevinsky permutation for the maximal rank conditions r^e given by $r_{ij}^e = \min\{e_i, e_{i+1}, \dots, e_j\}$ for $i \leq j$. Similarly we write $\hat{z}(e) = w_0^{(N)} z(e)^{-1} w_0^{(N)}$. The inverse of this permutation is called $v(Hom)$ in [24].

It follows from Lemma 3.1 that for all i and $p > r_{ni}$ we have $z(r)(p) \leq r_{i0}$. This implies that for $q \leq r_{n,i+1}$ we have $\hat{z}(r)(q) = N + 1 - z(r')(N + 1 - q) > r_{i-1,0}$, so any FK-graph for $\hat{z}(r)$ must contain the set $D_e = \bigcup_{i=1}^{n-1} [1, r_{i-1,0}] \times [1, r_{n,i+1}]$. Since $|D_e| = \sum_{j-i \geq 2} e_i e_j = \ell(\hat{z}(e))$, it follows that D_e is the unique FK-graph for $\hat{z}(e)$. Corollary 2.2 therefore implies that the polynomial $\mathfrak{G}_{\hat{z}(e)}(a; b) = \prod_{(p,q) \in D_e} (1 - \frac{b_p}{a_q})$ divides $\mathfrak{G}_{\hat{z}(r)}(a; b)$ (cf. [24, §5.1]).

4. THE RATIO FORMULA

In this section we give a coordinate-free proof of the ratio formula [24, Thm. 2.7]. The underlying geometry is similar to the original proof, but it becomes slightly simpler by working with the conjugate Zelevinsky permutation.

Suppose $F_1 \subset F_2 \subset \cdots \subset F_N \rightarrow H_N \twoheadrightarrow \cdots \twoheadrightarrow H_2 \twoheadrightarrow H_1$ is a flag of vector bundles over X with a morphism to a dual flag, such that $\text{rank}(F_i) = \text{rank}(H_i) = i$. For a permutation $w \in S_N$, Fulton [20] defines the degeneracy locus

$$\Omega_w = \{x \in X \mid \text{rank}(F_q(x) \rightarrow H_p(x)) \leq r_w(p, q) \ \forall p, q\}$$

where $r_w(p, q) = \#\{k \leq p \mid w(k) \leq q\}$. This locus does not depend on the bundles H_p for which $w(p) < w(p+1)$, or on the bundles F_q such that $w^{-1}(q) < w^{-1}(q+1)$. The expected codimension of the locus is the length $\ell(w)$. We need the following formula for the Grothendieck class of Ω_w , which generalizes Fulton's formula for its cohomology class [20]. The K -theory formula was proved in [7, Thm. 2.1] as an application of [21, Thm. 3]. An equivalent statement was given in [22, Thm. A].

Theorem 4.1. *If the codimension of Ω_w in X equals $\ell(w)$, then*

$$[\mathcal{O}_{\Omega_w}] = \mathfrak{G}_w(L_1, \dots, L_N; M_1, \dots, M_N) \in K(X)$$

where $L_i = \ker(H_i \rightarrow H_{i-1})$ and $M_i = F_i/F_{i-1}$.

It follows from Lemma 3.2 that the Grothendieck polynomial $\mathfrak{G}_{\hat{z}(r)}(a; b)$ is symmetric in each interval of variables $\{a_p : r_{n,i+1} < p \leq r_{n,i}\}$ and $\{b_p : r_{i-1,0} < p \leq r_{i,0}\}$. We therefore allow the vector bundle E_i to be substituted for these intervals, and we interpret the result as if imaginative line bundle summands of E_i had been inserted. We use the same notation in the quotient of polynomials $\mathfrak{G}_{\hat{z}(r)}(a; b)/\mathfrak{G}_{\hat{z}(e)}(a; b)$.

Theorem 4.2 (Knutson, Miller, Shimozono). *If the codimension of $\Omega_r(E_\bullet)$ in X is equal to $d(r)$, then the class of its structure sheaf in $K(X)$ is given by*

$$[\mathcal{O}_{\Omega_r(E_\bullet)}] = \frac{\mathfrak{G}_{\hat{z}(r)}(E_n, \dots, E_0; E_0, \dots, E_n)}{\mathfrak{G}_{\hat{z}(e)}(E_n, \dots, E_0; E_0, \dots, E_n)}.$$

Proof. By replacing X with $\bigoplus_{j-i \geq 2} \text{Hom}(E_{i-1}, E_i)$, we may assume that there are general maps $\phi_{ij} : E_i \rightarrow E_j$ for all $i < j$, such that each map $\phi_{i-1,i}$ comes from the given bundle sequence $E_\bullet$. We may then construct the bundle sequence

$$E_0 \subset E_0 \oplus E_1 \subset \cdots \subset E_0 \oplus \cdots \oplus E_n \xrightarrow{\phi} E_0 \oplus \cdots \oplus E_n \twoheadrightarrow \cdots \twoheadrightarrow E_{n-1} \oplus E_n \twoheadrightarrow E_n$$

where the middle map ϕ is composed of maps $E_i \rightarrow E_j$ which is ϕ_{ij} for $i < j$, the identity for $i = j$, and zero for $i > j$. All other maps are the obvious embeddings or projections. It follows from Lemma 3.2 that Fulton's locus $\Omega_{z(r)}$ for this sequence consists of the points in X where each composed map $E_0 \oplus \cdots \oplus E_i \rightarrow E_j \oplus \cdots \oplus E_n$ has rank at most r_{ij} .

We claim that $\Omega_{z(r)}$ is the intersection of the quiver variety $\Omega_r(E_\bullet)$ with the subset of X where each map ϕ_{ij} is equal to the composition $\phi_{j-1,j} \phi_{j-2,j-1} \cdots \phi_{i,i+1}$. To see this, notice that the condition that the rank of the map $E_0 \oplus \cdots \oplus E_k \rightarrow E_k \oplus \cdots \oplus E_n$ is at most e_k is equivalent to demanding that for all $i < k < j$ we have $\phi_{ij} = \phi_{kj} \phi_{ik}$. Given that this holds for all k , we obtain that for each $i < j$ the map $E_0 \oplus \cdots \oplus E_i \rightarrow E_j \oplus \cdots \oplus E_n$ can be factored as a surjection followed by ϕ_{ij} followed by an injection, so the rank condition on this map is equivalent to

$\text{rank}(\phi_{ij}) \leq r_{ij}$. In particular, $\Omega_{z(e)}$ is the locus where $\phi_{ij} = \phi_{j-1,j} \cdots \phi_{i,i+1}$ for all $i < j$, and we have $\Omega_{z(r)} = \Omega_r(E_\bullet) \cap \Omega_{z(e)}$ as a scheme-theoretic intersection.

The above description also shows that the codimension of $\Omega_{z(r)}$ in X is $\ell(z(r)) = d(r) + \ell(z(e))$, so by Theorem 4.1 and (4) we have $[\mathcal{O}_{\Omega_{z(r)}}] = \mathfrak{G}_{z(r)}(\check{E}_\bullet; E_\bullet) = \mathfrak{G}_{\hat{z}(r)}(\check{E}_\bullet; E_\bullet)$, where we write $\check{E}_\bullet$ for the reversed sequence $E_n, E_{n-1}, \dots, E_0$. Similarly we have $[\mathcal{O}_{\Omega_{z(e)}}] = \mathfrak{G}_{\hat{z}(e)}(\check{E}_\bullet; E_\bullet)$. Since all of the degeneracy loci are Cohen-Macaulay [25] we obtain the identity

$$(8) \quad \mathfrak{G}_{\hat{z}(r)}(\check{E}_\bullet; E_\bullet) = [\mathcal{O}_{\Omega_r(E_\bullet)}] \cdot \mathfrak{G}_{\hat{z}(e)}(\check{E}_\bullet; E_\bullet).$$

Comparing with (1), it follows that $\mathfrak{G}_{\hat{z}(r)}(\check{E}_\bullet; E_\bullet) = (\sum_\mu c_\mu(r) \prod_i \mathfrak{G}_{\mu_i}(E_i - E_{i-1})) \cdot \mathfrak{G}_{\hat{z}(e)}(\check{E}_\bullet; E_\bullet)$ on all varieties X , so this must hold as an identity of polynomials in the exterior powers of the bundles E_i . The theorem follows from this. $\square$

Remark. The above proof cites the main theorem of [7] for the existence of a universal polynomial that expresses the class of a quiver variety in the Grothendieck ring of an arbitrary variety X . When X has an ample line bundle L , one can also deduce the theorem directly from (8) as follows. By twisting the sequence $E_\bullet$ with a power of L , one may assume that all bundles E_i are globally generated. In this case, one can construct a bundle $Y = \bigoplus_{i=1}^n \text{Hom}(F_{i-1}, F_i)$ over a product of Grassmannians $\prod_{i=0}^n \text{Gr}^{e_i}(\mathbb{C}^k)$ with tautological quotient bundles $F_0, \dots, F_n$, such that the sequence $E_\bullet$ is the pullback of the universal sequence $F_0 \rightarrow \dots \rightarrow F_n$ on Y along a map of varieties $f : X \rightarrow Y$. On Y we know that $\mathfrak{G}_{\hat{z}(r)}(\check{F}_\bullet; F_\bullet) = [\mathcal{O}_{\Omega_r(F_\bullet)}] \cdot \mathfrak{G}_{\hat{z}(e)}(\check{F}_\bullet; F_\bullet)$. By taking k sufficiently large, we may therefore assume that the Grothendieck class of $\Omega_r(F_\bullet)$ agrees with $\mathfrak{G}_{\hat{z}(r)}(\check{F}_\bullet; F_\bullet) / \mathfrak{G}_{\hat{z}(e)}(\check{F}_\bullet; F_\bullet)$ in $K(Y)/I$, where $I \subset K(Y)$ is the ideal generated by the classes of subvarieties of codimension $\dim(X)$ or higher. The theorem follows from this because $f^*[\mathcal{O}_{\Omega_r(F_\bullet)}] = [\mathcal{O}_{\Omega_r(E_\bullet)}]$ and $f^*(I) = 0$.

5. RESTRICTED FK-GRAPHS

We will say that an FK-graph $D \subset \mathbb{N} \times \mathbb{N}$ is *restricted* w.r.t. the dimension vector $e = (e_0, \dots, e_n)$ if for every point $(p, q) \in D$ and $0 \leq i \leq n$ we have $p \leq r_{i-1,0}$ or $q \leq r_{n,i+1}$.

For each $0 \leq i \leq n$ we let $a^i = (a_1^i, \dots, a_{e_i}^i)$ be a set of e_i variables. Set $a = (a^0, a^1, \dots, a^n) = (a_1^0, \dots, a_{e_0}^0, \dots, a_1^n, \dots, a_{e_n}^n)$ and $\check{a} = (a^n, a^{n-1}, \dots, a^0) = (a_1^n, \dots, a_{e_n}^n, \dots, a_1^0, \dots, a_{e_0}^0)$. We need the following variation of Corollary 2.2.

Corollary 5.1. *For $w \in S_N$ we have*

$$\mathfrak{G}_w(\check{a}; a) = \sum_D (-1)^{|D| - \ell(w)} \prod_{(p,q) \in D} \left(1 - \frac{a_p}{\check{a}_q}\right)$$

where the sum is over all FK-graphs D for w , which are restricted w.r.t. e .

Proof. By Theorem 2.1, the Grothendieck polynomial $\mathfrak{G}_w(\check{a}; a)$ is the coefficient of w in the FK-product of the diagram $\mathfrak{D}'_N$ obtained from $\mathfrak{D}_N$ by replacing the top variables with $\check{a}$ and the left side variables with a . For each crossing position (p, q) in this diagram such that $a_p = \check{a}_q$, the corresponding factor $(1 + (1 - \frac{a_p}{\check{a}_q})s_{p+q-1})$ of $\mathfrak{G}(\mathfrak{D}'_N)$ is equal to one. Therefore $\mathfrak{G}(\mathfrak{D}'_N)$ is equal to the FK-product of the

diagram obtained from $\mathfrak{D}'_N$ by replacing these crossings with “ $\swarrow$ ” symbols. Now this diagram is the first of the following two equivalent diagrams:

The equivalence follows by using the “ $x = y$ ” case of the second transformation of (3) to move the thickened line segments in the south-east direction. The corollary follows from (2) because $\mathfrak{G}_w(\check{a}; a)$ is the coefficient of w in the FK-product of the second diagram. $\square$

6. A K -THEORETIC COMPONENT FORMULA

For each $1 \leq j \leq n-1$ we set $\delta_j = W_{jj}W_{j+1,j} \cdots W_{n-1,j} \in S_{r_{n,j-1}}$. This is the Grassmannian permutation given by $\delta_j(p) = p + e_{j-1}$ for $e_j < p \leq r_{nj}$ and $\delta_j(p) = p - r_{n,j+1}$ for $r_{nj} < p \leq r_{n,j-1}$. We define a *KMS-factorization* for the rank conditions r to be a sequence $(w_1, \dots, w_n)$ of permutations with $w_i \in S_{e_{i-1}+e_i}$, such that the conjugate Zelevinsky permutation $z(r)$ is equal to the absolute Hecke product

$$(9) \quad w_1 \cdot \delta_1 \cdot w_2 \cdot \delta_2 \cdots \delta_{n-1} \cdot w_n.$$

In the reduced case, these factorizations are equivalent to the minimal lace diagrams of Knutson, Miller, and Shimozono [24].²

A *partial permutation* contained in the rectangle $k \times l$ with k rows and l columns is a permutation $u \in S_{k+l}$, such that all descent positions of u are less than or equal to l , while all descent positions of u^{-1} are less than or equal to k . If this is true, then all FK-graphs for u will be contained in $[1, k] \times [1, l]$.

Lemma 6.1. *If $(w_1, \dots, w_n)$ is a KMS-factorization for the rank conditions r , then each permutation w_j is a partial permutation contained in $e_{j-1} \times e_j$.*

Proof. Since the absolute Hecke product $\alpha = w_{j+1} \cdot \delta_{j+1} \cdot w_{j+2} \cdots \delta_{n-1} \cdot w_n$ is a permutation in $S_{r_{nj}}$, we have $\delta_j \cdot \alpha(p) = \delta_j(p) = p - r_{n,j+1}$ for $r_{nj} < p \leq r_{n,j-1}$. If w_j had a descent in the interval $[e_j + 1, e_j + e_{j-1} - 1]$, then the product (9) would obtain a descent in the interval $[r_{nj} + 1, r_{n,j-1} - 1]$, a contradiction. By using that $(w_n^{-1}, \dots, w_1^{-1})$ is a KMS-factorization for the mirrored rank conditions $r'_{ij} = r_{n-j,n-i}$, we similarly deduce that w_j^{-1} has no descent positions in the interval $[e_{j-1} + 1, e_{j-1} + e_j - 1]$. $\square$

²All permutations w_i are inverted in the notation of [24].

If u is a partial permutation, and if the rectangle $k \times l$ is understood, we set $\hat{u} = w_0^{(k+l)} u^{-1} w_0^{(k+l)}$, where $w_0^{(k+l)}$ is the longest element in S_{k+l} . Notice that the 180° rotation of an FK-graph D for u will be an FK-graph for $\hat{u}$. We will denote this rotated FK-graph by $\hat{D}$, that is, $\hat{D} = \{(k+1-p, l+1-q) \mid (p, q) \in D\}$.

Given a sequence $(P_1, \dots, P_n)$ of FK-graphs such that each P_i is contained in $[1, e_{i-1}] \times [1, e_i]$, we let $\hat{\Phi}(P_1, \dots, P_n)$ denote the FK-graph which is the union of the sets $\{(p + r_{i-2,0}, q + r_{n,i+1}) \mid (p, q) \in \hat{P}_i\}$, for all $1 \leq i \leq n$, together with the unique FK-graph D_e for the minimal Zelevinsky permutation $\hat{z}(e)$.

This construction was used in [24] (in the reverse direction) and in [33] (for certain special RC-graphs). Notice that the crossing positions to the left of $\hat{P}_i$ in $\hat{\Phi}(P_1, \dots, P_n)$ form a reduced FK-graph for $w_0^{(N)} \delta_i^{-1} w_0^{(N)}$, and D_e is the union of these positions.

Lemma 6.2. *Let r be a set of rank conditions. The map $\hat{\Phi}$ gives a bijection of the set of sequences $(P_1, \dots, P_n)$ of FK-graphs for which $(w(P_1), \dots, w(P_n))$ is a KMS-factorization for r , with the set of all restricted FK-graphs for $\hat{z}(r)$.*

Proof. The south-west to north-east absolute Hecke product of the simple reflections s_{p+q-1} for (p, q) in the block of $\hat{P}_i$ is equal to $1^{N-e_i-e_{i-1}} \times w(\hat{P}_i) = w_0^{(N)} w(P_i)^{-1} w_0^{(N)}$. It follows from this that

$$w_0^{(N)} w(\hat{\Phi}(P_1, \dots, P_n))^{-1} w_0^{(N)} = w(P_1) \cdot \delta_1 \cdot w(P_2) \cdot \delta_2 \cdots \delta_{n-1} \cdot w(P_n).$$

The lemma therefore follows from Lemma 6.1 together with the definition (9) of KMS-factorizations. $\square$

We can now prove the following K -theoretic generalization of [24, Cor. 6.15].

Theorem 6.3. *For any set of rank conditions r we have*

$$\frac{\mathfrak{G}_{\hat{z}(r)}(\check{a}; a)}{\mathfrak{G}_{\hat{z}(e)}(\check{a}; a)} = \sum_{(w_1, \dots, w_n)} (-1)^{\ell(w_1 \cdots w_n) - d(r)} \mathfrak{G}_{w_1}(a^1; a^0) \mathfrak{G}_{w_2}(a^2; a^1) \cdots \mathfrak{G}_{w_n}(a^n; a^{n-1})$$

where the sum is over all KMS-factorizations $(w_1, \dots, w_n)$ for r .

Proof. Corollary 5.1 and Lemma 6.2 imply that $\mathfrak{G}_{\hat{z}(r)}(\check{a}; a) / \mathfrak{G}_{\hat{z}(e)}(\check{a}; a)$ is equal to

$$\sum (-1)^{\ell(w_1 \cdots w_n) - d(r)} \mathfrak{G}_{w_1}(a_{e_1}^1, \dots, a_1^1; a_{e_0}^0, \dots, a_1^0) \cdot \mathfrak{G}_{w_2}(a_{e_2}^2, \dots, a_1^2; a_{e_1}^1, \dots, a_1^1) \cdots \mathfrak{G}_{w_n}(a_{e_n}^n, \dots, a_1^n; a_{e_{n-1}}^{n-1}, \dots, a_1^{n-1}).$$

The theorem follows because $\mathfrak{G}_{\hat{z}(r)}(\check{a}; a)$ is symmetric in each set of variables a^i . $\square$

Remark. Let $(w_1, \dots, w_n)$ be a sequence of partial permutations with w_i contained in $e_{i-1} \times e_i$. Suppose $1 \leq j < n$ and $1 \leq k < e_j$ are given such that $w_j(k) < w_j(k+1)$ and $w_{j+1}^{-1}(k) < w_{j+1}^{-1}(k+1)$. If any of the sequences $(w_1, \dots, w_j s_k, w_{j+1}, \dots, w_n)$, $(w_1, \dots, w_j, s_k w_{j+1}, \dots, w_n)$, or $(w_1, \dots, w_j s_k, s_k w_{j+1}, \dots, w_n)$ is a KMS-factorization, then the definition shows that all three are KMS-factorizations. These transformations were first observed during an attempt to generalize the symmetry arguments of [9] to K -theory. In fact, in [9] it is proved that all KMS-factorizations of a given Zelevinsky permutation are connected by these transformations, which gives an easy way to find all of them.

7. KMS-FACTORIZATIONS ARE FACTOR SEQUENCES

We define the *permutation diagram* for the rank conditions r to be the part of the matrix (6) which is on or below the antidiagonal.

$$(10) \quad \begin{array}{cccc} & & & W_{n-1,n} \\ & & \ddots & \vdots \\ & W_{12} & \dots & W_{1n} \\ W_{01} & W_{02} & \dots & W_{0n} \end{array}$$

This diagram can also be obtained from the rectangle diagram of [10, §2.1] by replacing each rectangle R_{ij} with the corresponding Grassmannian permutation W_{ij} (and rotating the result 45 degrees counterclockwise.)

For $0 \leq k \leq n$ we let $r^{(k)}$ denote the rank conditions obtained from r by dropping all integers r_{ij} with $j - i < k$, that is, $r^{(k)} = \{r_{ij}^{(k)} \mid 0 \leq i \leq j \leq n - k\}$ where $r_{ij}^{(k)} = r_{i,j+k}$. The permutation diagram for these rank conditions is obtained from (10) by dropping the top k diagonals of matrices W_{ij} with $j - i \leq k$.

In this section we prove that all KMS-factorizations for r can be obtained from the permutation diagram in the following way. First, if $n = 1$, then the diagram has only one permutation W_{0n} , and the only KMS-factorization is (W_{0n}) . Otherwise any KMS-factorization can be obtained by first constructing a KMS-factorization $(\alpha_1, \dots, \alpha_{n-1})$ for $r^{(1)}$ and choosing arbitrary factorizations $\alpha_i = u_i \cdot v_i$ w.r.t. the absolute Hecke product. Then the sequence $(W_{01} \cdot u_1, v_1 \cdot W_{12} \cdot u_2, \dots, v_{n-1} \cdot W_{n-1,n})$ is a KMS-factorization for the rank conditions r . The exact statement that we prove is the following theorem, which also includes a criterion for KMS-factorizations similar to the criterion for factor sequences proved in [5].

Theorem 7.1. (a) *If $(w_1, \dots, w_n)$ is a KMS-factorization for r , then each permutation w_i has a reduced factorization $w_i = v_{i-1} \cdot W_{i-1,i} \cdot u_i$ with $v_{i-1} \in S_{e_{i-1}}$ and $u_i \in S_{e_i}$, such that $v_0 = u_n = 1$.*

(b) *Let $u_1, v_1, \dots, u_{n-1}, v_{n-1}$ be permutations with $u_i, v_i \in S_{e_i}$. Then the sequence $(W_{01} \cdot u_1, v_1 \cdot W_{12} \cdot u_2, \dots, v_{n-1} \cdot W_{n-1,n})$ is a KMS-factorization for r if and only if $(u_1 \cdot v_1, u_2 \cdot v_2, \dots, u_{n-1} \cdot v_{n-1})$ is a KMS-factorization for $r^{(1)}$.*

Based on this result, one can formulate a combinatorial formula for quiver coefficients in terms of a new type of factor sequences of tableaux. For cohomological quiver coefficients, this relies on Fomin and Greene's formula for Stanley coefficients [17] and uses a diagram of tableaux obtained from (10) by replacing W_{ij} with the unique row and column *decreasing* tableau T_{ij} of shape R_{ij} (i.e. with $r_{i+1,j} - r_{ij}$ rows and $r_{i,j-1} - r_{ij}$ columns), such that the reading word of T_{ij} is a reduced word

for W_{ij} . Factor sequences are generated using the Coxeter-Knuth product, and it follows from Theorem 7.1, Corollary 8.2, and [17, Thm. 1.2] that a cohomological quiver coefficient $c_\mu(r)$ counts the number of factor sequences of shape μ . A generalization of this formula to K -theoretic quiver coefficients has recently been proved with Kresch, Shimozone, Tamvakis, and Yong [11]. This is based on a new formula for the coefficients $c_{w,\lambda}$ in the expansion of stable Grothendieck polynomials (5), which generalizes Fomin and Greene's result. We refer to [11] for details.

Let $0 \leq k < n$ and consider permutations $w_1, \dots, w_{n-k}$ such that $w_i \in S_{r_{i,k+i-1}}$ for all i . We let $\Phi_k(w_1, \dots, w_{n-k}) \in S_N$ denote the south-west to north-east product of the matrix obtained from (6) by replacing $W_{i-1,k+i}$ with w_i for $1 \leq i \leq n-k$, and by replacing W_{ij} with the identity for $j-i \geq k+2$. When $n=4$ and $k=1$ this matrix looks as follows:

$$\begin{array}{cccc} W_{31} & W_{32} & W_{33} & W_{34} \\ W_{21} & W_{22} & W_{23} & w_3 \\ W_{11} & W_{12} & w_2 & 1 \\ W_{01} & w_1 & 1 & 1 \end{array}$$

Notice that $(w_1, \dots, w_n)$ is a KMS-factorization for r if and only if $\Phi_0(w_1, \dots, w_n) = z(r)$. Furthermore, if $k > 0$ and $u_1, v_1, \dots, u_{n-k}, v_{n-k}$ are permutations such that $u_i, v_i \in S_{r_{i,k+i-1}}$, then

$$\Phi_k(u_1 \cdot v_1, \dots, u_{n-k} \cdot v_{n-k}) = \Phi_{k-1}(W_{0,k} \cdot u_1, v_1 \cdot W_{1,k+1} \cdot u_2, \dots, v_{n-k} \cdot W_{n-k,n}).$$

Lemma 7.2. *If $\Phi_k(w_1, \dots, w_{n-k}) = z(r)$, then each permutation w_i has a reduced factorization $w_i = v \cdot W_{i-1,k+i} \cdot u$, where $v \in S_{r_{i-1,k+i-1}}$ and $u \in S_{r_{i,k+i}}$. Furthermore, v is trivial when $i=1$ while u is trivial when $i=n-k$.*

Proof. Fix i and set $j = k+i-1$, $a = r_{i-1,j}$, $b = r_{i,j+1}$, and $c = r_{i-1,j+1}$. For the first assertion it is enough to show that $w_i \in S_{a+b-c}$, that $w_i(p) \leq a$ for $b < p \leq a+b-c$, and that $w_i^{-1}(p) \leq b$ for $a < p \leq a+b-c$. In fact, vw_iu will then satisfy the same properties for every $v \in S_a$ and $u \in S_b$. If we choose v and u such that $v^{-1}w_iu^{-1}$ has no descents in the interval $[1, b-1]$, $uw_i^{-1}v$ has no descents in the interval $[1, a-1]$, and $\ell(v^{-1}w_iu^{-1}) = \ell(w_i) - \ell(v) - \ell(u)$, then we must have $v^{-1}w_iu^{-1} = W_{i-1,j+1}$, so we can use the factorization $w_i = v \cdot W_{i-1,j+1} \cdot u$.

Let $\sigma = W_{i-1,1}W_{i-1,2} \cdots W_{i-1,j}$ be the product of the permutations west of w_i in the matrix (6), and let $\tau = \prod_{s=i}^{n-1} \prod_{t=1}^{j+1} W_{st}$ be the product of the permutations weakly west and strictly north of w_i in this matrix. It follows by induction on j that all descent positions of σ are greater than or equal to r_{ij} . A similar argument shows that the descent positions of τ^{-1} are greater than or equal to r_{ij} . Furthermore, the product $\sigma\tau$ is reduced, since it is part of the defining reduced factorization of the conjugate Zelevinsky permutation for rank conditions obtained by replacing $r_{i-1,j+1}$ with $\min(r_{i-1,j}, r_{i,j+1})$. Since $w_i \in S_{r_{ij}}$, it follows that $\sigma w_i \tau$ is also a reduced product. We can therefore write $z(r) = \alpha \sigma w_i \tau \beta$ as a product of permutations, where $\alpha \in S_{r_{i-1,0}}$ and $\beta \in S_{r_{n,j+1}}$.

Notice that for $b < p \leq r_{ij}$ we have $\tau\beta(r_{n,j+1} - b + p) = \tau(r_{n,j+1} - b + p) = p$, and for $a < p \leq r_{ij}$ we have $\alpha\sigma(p) = \alpha(r_{i-1,0} - a + p) = r_{i-1,0} - a + p$. For $a+b-c < p \leq r_{ij}$ it follows from Lemma 3.1 that $z(r)(r_{n,j+1} - b + p) = r_{i-1,0} - a + p$, so we must have $w_i(p) = p$, that is, $w_i \in S_{a+b-c}$. The lemma also shows that $z(r)(r_{n,j+1} - b + p) \leq r_{i-1,0}$ for all $b < p \leq a+b-c$. This implies that $w_i(p) \leq a$, since otherwise we would have $\alpha\sigma w_i \tau\beta(r_{n,j+1} - b + p) = \alpha\sigma w_i(p) = r_{i-1,0} - a + w_i(p) >$

$r_{i-1,0}$. A symmetric argument shows that $w_i^{-1}(p) \leq b$ for $a < p \leq a + b - c$, which completes the proof of the first assertion.

When $i = n - k$, β is trivial, so we have $\tau\beta(p) = p$ for all $p \leq b = r_{in}$. If $u \in S_b$ was not the identity, then $z(r)$ would get a descent in the interval $[1, e_n - 1]$, a contradiction. A similar argument shows that v must be trivial when $i = 1$. $\square$

Proof of Theorem 7.1. Part (a) is a special case of Lemma 7.2 so we prove part (b). By induction on n we may assume that the theorem is true for all rank conditions $r^{(k)}$ with $k \geq 1$. We claim that $(w_1, \dots, w_{n-k})$ is a KMS-factorization for $r^{(k)}$ if and only if $\Phi_k(w_1, \dots, w_{n-k}) = z(r)$. If either is true, then we can write $w_i = v_{i-1} \cdot W_{i-1,k+i} \cdot u_i$ for each i , where $u_i, v_i \in S_{r_{i,k+i}}$, and $v_0 = u_{n-k} = 1$. This proves the claim when $k = n - 1$. For $1 \leq k \leq n - 2$ we know by induction on k that $\Phi_k(w_1, \dots, w_{n-k}) = \Phi_{k+1}(u_1 \cdot v_1, \dots, u_{n-k-1} \cdot v_{n-k-1})$ equals $z(r)$ if and only if $(u_1 \cdot v_1, \dots, u_{n-k-1} \cdot v_{n-k-1})$ is a KMS-factorization for $r^{(k+1)}$, which by the theorem for $r^{(k)}$ is equivalent to $(w_1, \dots, w_{n-k})$ being a KMS-factorization for $r^{(k)}$. This proves the claim, and the theorem follows from the claim with $k = 1$. $\square$

In the next section we need the following generalization of [24, Cor. 4.12].

Corollary 7.3. *The KMS-factorizations for $r + m$ are precisely the sequences $(1^m \times w_1, \dots, 1^m \times w_n)$ for which $(w_1, \dots, w_n)$ is a KMS-factorization for r .*

Proof. This is immediate from Theorem 7.1 because the permutation diagram for $r + m$ is obtained from (10) by replacing each permutation W_{ij} with $1^m \times W_{ij}$. $\square$

8. ALTERNATING SIGNS OF QUIVER COEFFICIENTS

We can now prove the stable version of the component formula. We note that Theorem 9.3 of the next section can be substituted for the reference to geometry in its proof.

Theorem 8.1. *For any set of rank conditions r we have*

$$\frac{\mathfrak{G}_{\hat{z}(r)}(\check{a}; a)}{\mathfrak{G}_{\hat{z}(e)}(\check{a}; a)} = \sum_{(w_1, \dots, w_n)} (-1)^{\ell(w_1 \dots w_n) - d(r)} G_{w_1}(a^1; a^0) G_{w_2}(a^2; a^1) \cdots G_{w_n}(a^n; a^{n-1})$$

where the sum is over all KMS-factorizations for r .

Proof. It follows from (1) and Theorem 4.2 that $\mathfrak{G}_{\hat{z}(r)}(\check{a}; a) / \mathfrak{G}_{\hat{z}(e)}(\check{a}; a)$ is equal to $\sum_{\mu} c_{\mu}(r) G_{\mu_1}(a^1; a^0) \cdots G_{\mu_n}(a^n; a^{n-1})$. Since $c_{\mu}(r) = c_{\mu}(r + m)$, we deduce that

$$\frac{\mathfrak{G}_{\hat{z}(r)}(\check{a}; a)}{\mathfrak{G}_{\hat{z}(e)}(\check{a}; a)} = \frac{\mathfrak{G}_{\hat{z}(r+m)}(a^n, 1^m, \dots, a^0, 1^m; a^0, 1^m, \dots, a^n, 1^m)}{\mathfrak{G}_{\hat{z}(e+m)}(a^n, 1^m, \dots, a^0, 1^m; a^0, 1^m, \dots, a^n, 1^m)}$$

for all integers $m \geq 0$. For $m \geq \max(e_0, \dots, e_n)$ it follows from Theorem 6.3 and Corollary 7.3 that the right-hand side of this identity equals

$$\begin{aligned} & \sum (-1)^{\ell(w_1 \dots w_n) - d(r)} \mathfrak{G}_{1^m \times w_1}(a^1, 1^m; a^0, 1^m) \cdots \mathfrak{G}_{1^m \times w_n}(a^n, 1^m; a^{n-1}, 1^m) \\ &= \sum (-1)^{\ell(w_1 \dots w_n) - d(r)} G_{w_1}(a^1; a^0) \cdots G_{w_n}(a^n; a^{n-1}) \end{aligned}$$

as required. $\square$

It was conjectured in [7] that quiver coefficients have signs which alternate with codimension. This conjecture is a consequence of the following explicit formula for quiver coefficients, which follows from Theorem 8.1 and Lascoux's Theorem 2.3.

Corollary 8.2. *The quiver coefficient $c_\mu(r)$ is given by*

$$c_\mu(r) = (-1)^{\sum |\mu_i| - d(r)} \sum_{(w_1, \dots, w_n)} |c_{w_1, \mu_1} c_{w_2, \mu_2} \cdots c_{w_n, \mu_n}|$$

where the sum is over all KMS-factorizations for the rank conditions r , and c_{w_i, μ_i} is defined by (5). Equivalently, $c_\mu(r)$ is equal to $(-1)^{\sum |\mu_i| - d(r)}$ times the number of sequences $(\gamma_1, \dots, \gamma_n)$ of paths $\gamma_i = \{w_i = w_{i,1} \triangleright \cdots \triangleright w_{i,m_i}\}$, such that $(w_1, \dots, w_n)$ is a KMS-factorization for r , $w_{i,m_i} = w_{\mu_i}$ is a Grassmannian permutation for μ_i , and $w_{i,j}$ is not Grassmannian for $1 \leq j < m_i$.

9. DOUBLE QUIVER POLYNOMIALS

Let r be a set of rank conditions. For each $0 \leq i \leq n$ we let $b^i = (b_1^i, \dots, b_{e_i}^i)$ be a set of $e_i = r_{ii}$ variables, and we set $b = (b^0, b^1, \dots, b^n)$. The K -theoretic analogue of the double ratio formula from [24] defines the polynomial

$$K_r(a; b) = \frac{\mathfrak{G}_{\widehat{z}(r)}(\check{a}; b)}{\mathfrak{G}_{\widehat{z}(e)}(\check{a}; b)},$$

which is named a *double quiver polynomial* in [24]. In this section we prove some facts about such polynomials. In particular, we show that $K_r(a; b)$ is a specialization of the quiver formula constructed in [7]. When the rank conditions have the form $r + m$ for m large, the cohomology versions of Theorems 9.1, 9.3, and 9.5 below follow from the results in [24] (see Cor. 6.13 and Thm. 6.20 in loc. cit.).

It follows from (i) and (ii) of Lemma 3.2 that $K_r(a; b)$ is separately symmetric in each set of variables a^i and b^i (and that the variables a^0 and b^n do not occur). In addition we have

Theorem 9.1. *The polynomial $K_r(a; b)$ is multisupersymmetric, that is, if one sets $a_1^{i+1} = b_1^i$ in $K_r(a; b)$, then the result is independent of these variables.*

Since $\mathfrak{G}_{\widehat{z}(e)}(\check{a}; b) = \prod_{(p,q) \in D_e} (1 - \frac{b_p}{\check{a}_q})$, this theorem is an immediate consequence of the following proposition, applied to $\widehat{z}(r)$ with $k = r_{i-1,0} + 1$ and $l = r_{n,i+2} + 1$.

Proposition 9.2. *Let $w \in S_N$ be a permutation and let $k, l \in \mathbb{N}$. Assume that $w(i) > k$ for $1 \leq i < l$ and that $w^{-1}(i) > l$ for $1 \leq i < k$. Then the polynomial*

$$\frac{\mathfrak{G}_w(a_1, \dots, a_{l-1}, b_k, a_{l+1}, \dots, a_N; b_1, \dots, b_N)}{\left(\prod_{i=1}^{k-1} (1 - \frac{b_i}{b_k})\right) \left(\prod_{i=1}^{l-1} (1 - \frac{b_k}{a_i})\right)}$$

is independent of b_k .

Proof. By Theorem 2.1, $\mathfrak{G}_w(a_1, \dots, a_{l-1}, b_k, a_{l+1}, \dots, a_N; b_1, \dots, b_N)$ is the coefficient of w in the FK-product of the first of the following three equivalent diagrams:

The first diagram is obtained from $\mathfrak{D}_N$ by replacing a_l with b_k and the crossing at position (k, l) with “ $\searrow$ ”, and the others are the result of moving the thick line segments labeled b_k in the south-east direction, using the rules of (3).

Notice that if D is a subset of the crossing positions of the third diagram such that $w(D) = w$, then D must contain all crossings involving the line segments labeled b_k . The proposition therefore follows from (2). $\square$

Our next theorem is a K -theoretic generalization of Conjecture 6.14 from [24]. It says that the double ratio formula satisfies rank stability.

Theorem 9.3. *For any rank conditions r and non-negative integer m we have*

$$K_{r+m}(a^0, 1^m, \dots, a^n, 1^m; b^0, 1^m, \dots, b^n, 1^m) = K_r(a^0, \dots, a^n; b^0, \dots, b^n).$$

Proof. Using symmetry, it is enough to prove that

$$(11) \quad \begin{aligned} &K_{r+1}(a^0, 1, c_1, a^1, \dots, c_n, a^n; c_1, b^0, \dots, c_n, b^{n-1}, b^n, 1) \\ &= K_r(a^0, \dots, a^n; b^0, \dots, b^n). \end{aligned}$$

The polynomial $\mathfrak{G}_{\hat{z}(r+1)} = \mathfrak{G}_{\hat{z}(r+1)}(c_n, a^n, \dots, c_1, a^1, a^0, 1; c_1, b^0, \dots, c_n, b^{n-1}, b^n, 1)$ is the coefficient of $\hat{z}(r+1)$ in the FK-product of the following two equivalent diagrams (of size $N + n + 1$):

As indicated in the first diagram, the line segments extending furthest north and west are labeled with the variable sets a^i and b^i , respectively. The equivalence is obtained by moving the thickened line segments toward the borders of the diagram. Since $\hat{z}(r+1)$ fixes $N + n + 1$, it follows that $\mathfrak{G}_{\hat{z}(r+1)}$ is also the coefficient of $\hat{z}(r+1)$

in the FK-product of the diagram:

We claim that the FK-graphs for $\widehat{z}(r+1)$ w.r.t. this diagram are exactly those obtained by placing an FK-graph for $\widehat{z}(r)$ w.r.t. $\mathfrak{D}_N$ in the triangular region of $\mathfrak{D}$ and including all crossing positions P outside this region. The theorem follows from this claim because it shows that $\mathfrak{G}_{\widehat{z}(r+1)} = \mathcal{Q} \cdot \mathfrak{G}_{\widehat{z}(r)}(a^n, a^{n-1}, \dots, a^0; b^0, b^1, \dots, b^n)$, where $\mathcal{Q}$ is the product of the factors $h(P)$ for the crossing positions P outside the triangular region, and (11) is immediate from this identity.

We will say that a permutation $w \in S_N$ satisfies the condition (*) for the set of rank conditions r , if for all i and $p \leq r_{n,i+1}$ we have $w(p) > r_{i-1,0}$. As noted in Section 3, $\widehat{z}(r)$ has this property. Notice that if $D \subset C(\mathfrak{D})$ is any FK-graph such that $w(D)$ satisfies (*) for $r+1$, then D must contain all the crossing positions outside the triangular region. In particular, all FK-graphs for $\widehat{z}(r+1)$ must contain these crossing positions.

The crossing positions in the south-west region of $\mathfrak{D}$ represent the Grassmannian permutation α defined by $\alpha(i) = r_{i-2,0} + i$ for $1 \leq i \leq n$ and $\alpha(i) < \alpha(i+1)$ for $i \neq n$, while the permutation β represented by the north-east crossings is given by $\beta^{-1}(i) = r_{n,n+2-i} + i$ for $1 \leq i \leq n$ and $\beta^{-1}(i) < \beta^{-1}(i+1)$ for $i \neq n$. If we let $w \in S_N$ be the permutation of an FK-graph placed in the triangular region of $\mathfrak{D}$, then the claim states that the absolute Hecke product $\alpha \cdot (w_0^{(n)} \times w) \cdot \beta$ equals $\widehat{z}(r+1)$ if and only if $w = \widehat{z}(r)$.

If $\alpha \cdot (w_0^{(n)} \times w) \cdot \beta$ satisfies (*) for $r+1$, then this product must be reduced. Otherwise one could skip a simple reflection factor of α or β , which could be exploited to construct an FK-graph for the product that missed the corresponding crossing outside the triangular region in $\mathfrak{D}$. Since a similar argument applies if the (usual) product of permutations $\alpha(w_0^{(n)} \times w)\beta$ satisfies (*) for $r+1$, it is enough to check that $\alpha(w_0^{(n)} \times w)\beta$ equals $\widehat{z}(r+1)$ if and only if $w = \widehat{z}(r)$.

Equivalently, we must check that the *inverse* of $\alpha(w_0^{(n)} \times \widehat{z}(r))\beta$ maps $N+n+2-p$ to $N+n+2-z(r+1)(p)$ for every $2 \leq p \leq N+n+1$. We use Lemma 3.1 to do this. Set $r^+ = r+1$. This means that $r_{ij}^+ = r_{ij} + 1$ for $i \leq j$ while $r_{ij}^+ = r_{ij} + 1 + i - j$ when $i > j$. Now choose $0 \leq i \leq j+1 \leq n+1$ such that $r_{n,j+1}^+ + r_{i-1,j}^+ - r_{i-1,j+1}^+ < p \leq r_{n,j+1}^+ + r_{i,j}^+ - r_{i,j+1}^+$.

If $0 \leq j \leq n-1$ and $p = r_{n,j}^+$, then $i = j+1$ and $z(r^+)(p) = r_{j+1,0}^+$. Furthermore, $N+n+2-p = r_{j-1,0} + j+1$ is mapped to $j+1$ by α^{-1} , which in turn is mapped to $n-j$ by $w_0^{(n)}$, and $\beta^{-1}(n-j) = r_{n,j+2} + n-j = N+n+2-r_{j+1,0}^+$, as required.

Otherwise we have $p \neq r_{n,j}^+$ or $j = n$. Set $i' = \max(i, 1)$ and $j' = \min(j, n-1)$. Then $z(r^+)(p) = p + \delta + i' + j' - n$, where $\delta = -r_{n,j+1} + r_{i,j+1} + r_{i-1,0} - r_{i-1,j}$. Since $j' \leq n-1$ is maximal such that $\alpha(j'+1) < N+n+2-p$, we get $\alpha^{-1}(N+n+2-p) = (N+n+2-p) + n - j' - 1$. One now checks that $r_{n,j+1} + r_{i-1,j} - r_{i-1,j+1} < p - n + j' \leq r_{n,j+1} + r_{ij} - r_{i,j+1}$. Using this, it follows that the inverse of $w_0^{(n)} \times \widehat{z}(r)$ maps $\alpha^{-1}(N+n+2-p) = n + N + 1 - (p - n + j')$ to $n + N + 1 - z(r)(p - n + j') = n + N + 1 - (p - n + j' + \delta)$, and this number furthermore lies in the interval $[n + r_{n,i+1} + 1, n + r_{ni}]$. Thus β^{-1} subtracts $i' - 1$, so the inverse of $\alpha(w_0^{(n)} \times \widehat{z}(r))\beta$ maps $N+n+2-p$ to $N+2n+2-p-\delta-i'-j'$, as required. $\square$

We will finish this paper by proving that the double quiver polynomial $K_r(a; b)$ is a specialization of the quiver formula of [7]. We need the following statement (about power series in the variables $x_i = 1 - a_i$ and $y_i = 1 - b_i$).

Lemma 9.4. *Let $f(a; b) \in \mathbb{Z}[a_1^{-1}, \dots, a_q^{-1}, b_1, \dots, b_p]$ be a Laurent polynomial that is separately symmetric in the variables $\{a_i\}$ and $\{b_i\}$. Assume furthermore that $f(c, a_2, \dots, a_q; c, b_2, \dots, b_p)$ is independent of the variable c . Then f is a (possibly infinite) linear combination of double stable Grothendieck polynomials $G_\lambda(a; b)$.*

Proof. If we set $b_i = 1 + y_i$ and $a_i = 1 + x_i$, that is, $(a_i)^{-1} = \sum_{k \geq 0} (-x_i)^k$, then the resulting power series $f(1 + x_i; 1 + y_i)$ is *supersymmetric*, i.e. if one sets $x_1 = y_1$, then $f(1 + x_i; 1 + y_i)$ becomes independent of these variables. Since the lowest term of $G_\lambda(1 + x_i; 1 + y_i)$ is the double Schur polynomial $s_\lambda(x; y)$, it follows from [32, Thm. 1] that there are coefficients $d_\lambda \in \mathbb{Z}$ such that the lowest term of

$$f(1 + x_i; 1 + y_i) - \sum d_\lambda G_\lambda(1 + x_i; 1 + y_i)$$

has higher degree than the lowest term of $f(1 + x_i; 1 + y_i)$. The lemma follows from this because each $G_\lambda(1 + x_i; 1 + y_i)$ is also supersymmetric [19]. $\square$

We will prove in [4] that the linear combination in this lemma is always finite, but this fact is not needed for the proof of our last theorem.

Theorem 9.5. *For any rank conditions r we have*

$$K_r(a; b) = \sum_{\mu} c_{\mu}(r) G_{\mu_1}(a^1; b^0) G_{\mu_2}(a^2; b^1) \cdots G_{\mu_n}(a^n; b^{n-1}),$$

where the sum is over all sequences $\mu = (\mu_1, \dots, \mu_n)$ of partitions.

Proof. Theorem 9.1 and Lemma 9.4 imply that we can write

$$K_{r+m}(a; b) = \sum c_{\mu} G_{\mu_1}(a^1; b^0) G_{\mu_2}(a^2; b^1) \cdots G_{\mu_n}(a^n; b^{n-1})$$

where the sum is over (possibly infinitely many) sequences of partitions μ . Theorem 9.3 implies that the coefficients c_{μ} are independent of m . By setting $b^i = a^i$ for all i , it therefore follows from Theorem 4.2 and the definition (1) of quiver coefficients that $c_{\mu} = c_{\mu}(r)$ for all μ . The result is obtained by setting $m = 0$. $\square$

By using Corollary 8.2, the above theorem can also be interpreted as a double component formula, that is,

$$K_r(a; b) = \sum_{(w_1, \dots, w_n)} (-1)^{\ell(w_1 \cdots w_n) - d(r)} G_{w_1}(a^1; b^0) G_{w_2}(a^2; b^1) \cdots G_{w_n}(a^n; b^{n-1}),$$

where the sum is over all KMS-factorizations for r .

Theorem 9.3 makes it possible to extend the quiver polynomial $K_r(a; b)$ to infinite sets of variables a^i and b^i , by taking the limit of $K_{r+m}(a; b)$ as m tends to infinity. Such limits are called *double quiver functions* in [24]. In the K -theory case it is preferable to change to the variables $x_j^i = 1 - (a_j^i)^{-1}$ and $y_j^i = 1 - b_j^i$ in order to obtain a nice formal power series. This recovers the quiver formula P_r constructed in [7, §4]. In fact, Theorem 9.5 shows that the function $\lim_{m \rightarrow \infty} K_{r+m}(a; b)$ is obtained from P_r by setting $1^{\otimes i-1} \otimes x_j \otimes 1^{\otimes n-i} = 1 - (a_j^i)^{-1}$ and $1^{\otimes i-1} \otimes y_j \otimes 1^{\otimes n-i} = 1 - b_j^{i-1}$. In particular, the cohomological double quiver function used in [24] is equal to the lowest term of P_r . Equivalently, this cohomological quiver function is a specialization of the original quiver formula from [10] (see also the construction of this formula given in [5, §2]).

By setting the variables b_j^i equal to 1 in Theorem 9.5, one can deduce that general quiver coefficients are special cases of the coefficients studied in [13]. More details about this will be given in [14], together with a proof that quiver coefficients are special cases of Schubert structure constants (see also [3, 30]).

REFERENCES

- [1] S. Abeasis and A. Del Fra, *Degenerations for the representations of an equioriented quiver of type A_m* , Boll. Un. Mat. Ital. Suppl. (1980), no. 2, 157–171. MR0675498 (84e:16019)
- [2] N. Bergeron and S. Billey, *RC-graphs and Schubert polynomials*, Experiment. Math. **2** (1993), no. 4, 257–269. MR1281474 (95g:05107)
- [3] N. Bergeron and F. Sottile, *Schubert polynomials, the Bruhat order, and the geometry of flag manifolds*, Duke Math. J. **95** (1998), no. 2, 373–423. MR1652021 (2000d:05127)
- [4] A. S. Buch, *Supersymmetry of Grothendieck polynomial*, in preparation.
- [5] ———, *On a conjectured formula for quiver varieties*, J. Algebraic Combin. **13** (2001), no. 2, 151–172. MR1826950 (2002g:14074)
- [6] ———, *Stanley symmetric functions and quiver varieties*, J. Algebra **235** (2001), no. 1, 243–260. MR1807664 (2001m:05257)
- [7] ———, *Grothendieck classes of quiver varieties*, Duke Math. J. **115** (2002), no. 1, 75–103. MR1932326 (2003m:14018)
- [8] ———, *A Littlewood-Richardson rule for the K -theory of Grassmannians*, Acta Math. **189** (2002), no. 1, 37–78. MR1946917 (2003j:14062)
- [9] A. S. Buch, L. M. Fehér, and R. Rimányi, *Positivity of quiver coefficients through Thom polynomials*, to appear in Adv. Math., 2003.
- [10] A. S. Buch and W. Fulton, *Chern class formulas for quiver varieties*, Invent. Math. **135** (1999), no. 3, 665–687. MR1669280 (2000f:14087)
- [11] A. S. Buch, A. Kresch, M. Shimozono, H. Tamvakis, and A. Yong, *Stable Grothendieck polynomials and K -theoretic factor sequences*, in preparation.
- [12] A. S. Buch, A. Kresch, H. Tamvakis, and A. Yong, *Schubert polynomials and quiver formulas*, Duke Math. J. **122** (2004), no. 1, 125–143. MR2046809
- [13] ———, *Grothendieck polynomials and quiver formulas*, To appear in Amer. J. Math., 2003.
- [14] A. S. Buch, F. Sottile, and A. Yong, *Quiver coefficients are Schubert structure constants*, preprint, 2003.
- [15] P. Edelman and C. Greene, *Balanced tableaux*, Adv. in Math. **63** (1987), no. 1, 42–99. MR0871081 (88b:05012)
- [16] L. Fehér and R. Rimányi, *Classes of degeneracy loci for quivers: the Thom polynomial point of view*, Duke Math. J. **114** (2002), 193–213. MR1920187 (2003j:14005)

- [17] S. Fomin and C. Greene, *Noncommutative Schur functions and their applications*, Discrete Math. **193** (1998), no. 1-3, 179–200, Selected papers in honor of Adriano Garsia (Taormina, 1994). MR1661368 (2000c:05149)
- [18] S. Fomin and A. N. Kirillov, *Grothendieck polynomials and the Yang-Baxter equation*, Proc. Formal Power Series and Alg. Comb. (1994), 183–190.
- [19] ———, *The Yang-Baxter equation, symmetric functions, and Schubert polynomials*, Discrete Math. **153** (1996), 123–143. MR1394950 (98b:05101)
- [20] W. Fulton, *Flags, Schubert polynomials, degeneracy loci, and determinantal formulas*, Duke Math. J. **65** (1992), no. 3, 381–420. MR1154177 (93e:14007)
- [21] W. Fulton and A. Lascoux, *A Pieri formula in the Grothendieck ring of a flag bundle*, Duke Math. J. **76** (1994), no. 3, 711–729. MR1309327 (96j:14036)
- [22] A. Knutson and E. Miller, *Gröbner geometry of Schubert polynomials*, to appear in Ann. of Math. (2), 2003.
- [23] ———, *Subword complexes in Coxeter groups*, Adv. Math. **184** (2004), 161–176. MR2047852
- [24] A. Knutson, E. Miller, and M. Shimozono, *Four positive formulas for type A quiver polynomials*, preprint, 2003.
- [25] V. Lakshmibai and P. Magyar, *Degeneracy schemes, quiver schemes, and Schubert varieties*, Internat. Math. Res. Notices **1998**, no. 12, 627–640. MR1635873 (99g:14065)
- [26] A. Lascoux, *Anneau de Grothendieck de la variété de drapeaux*, The Grothendieck Festschrift, Vol. III, Birkhäuser Boston, Boston, MA, 1990, pp. 1–34. MR1106909 (92j:14064)
- [27] ———, *Transition on Grothendieck polynomials*, Physics and combinatorics, 2000 (Nagoya), World Sci. Publishing, River Edge, NJ, 2001, pp. 164–179. MR1872255 (2002k:14082)
- [28] A. Lascoux and M.-P. Schützenberger, *Structure de Hopf de l’anneau de cohomologie et de l’anneau de Grothendieck d’une variété de drapeaux*, C. R. Acad. Sci. Paris Sér. I Math. **295** (1982), no. 11, 629–633. MR0686357 (84b:14030)
- [29] ———, *Décompositions dans l’algèbre des différences divisées*, Discrete Math. **99** (1992), no. 1-3, 165–179. MR1158787 (93c:20030)
- [30] C. Lenart, S. Robinson, and F. Sottile, *Grothendieck polynomials via permutation patterns and chains in the Bruhat order*, preprint, 2003.
- [31] E. Miller, *Alternating formulae for K-theoretic quiver polynomials*, to appear in Duke Math. J., 2003.
- [32] J. R. Stembridge, *A characterization of supersymmetric polynomials*, J. Algebra **95** (1985), no. 2, 439–444. MR0801279 (87a:11022)
- [33] A. Yong, *On combinatorics of quiver component formulas*, to appear in J. Algebraic Combin., 2003.
- [34] A. V. Zelevinskii, *Two remarks on graded nilpotent classes*, Uspekhi Mat. Nauk **40** (1985), no. 1(241), 199–200. MR0783619 (86e:14027)

MATEMATISK INSTITUT, AARHUS UNIVERSITET, NY MUNKEGADE, 8000 ÅRHUS C, DENMARK
E-mail address: abuch@imf.au.dk