p -ADIC PERIODS AND DERIVED DE RHAM COHOMOLOGY

A. BEILINSON

To Irene

INTRODUCTION

For a smooth variety X over a base field of characteristic 0 we have its algebraic de Rham cohomology $H_{\mathrm{dR}}^i(X) := H^i(X_{\mathrm{Zar}}, \Omega_X^i)$; for nonsmooth X , one defines $H_{\mathrm{dR}}^i(X)$ using cohomological descent as in Deligne [D]. If the base field is $\mathbb{C}$, then one has the Betti cohomology $H_{\mathrm{B}}^i(X) := H^i(X_{\mathrm{cl}}, \mathbb{Q})$ and a canonical period isomorphism (“integration of algebraic differential forms over topological cycles”)

$$(0.1) \quad \rho : H_{\mathrm{dR}}^i(X) \xrightarrow{\sim} H_{\mathrm{B}}^i(X) \otimes \mathbb{C}$$

compatible with the $\mathrm{Gal}(\mathbb{C}/\mathbb{R})$ -conjugation. To define ρ , consider the analytic de Rham cohomology $H_{\mathrm{dR}}^i(X_{\mathrm{an}})$. There are evident maps

$$(0.2) \quad H_{\mathrm{dR}}^i(X) \xrightarrow{\alpha} H_{\mathrm{dR}}^i(X_{\mathrm{an}}) \xleftarrow{\beta} H_{\mathrm{B}}^i(X) \otimes \mathbb{C}.$$

Then β is an isomorphism due to the Poincaré lemma, and $\rho := \beta^{-1}\alpha$ (the fact that ρ is an isomorphism was established by Grothendieck [Gr]).

Suppose our base field is an algebraic closure $\bar{K}$ of a p -adic field K (say, $K = \mathbb{Q}_p$). The role of $H_{\mathrm{B}}^i(X)$ is now played by the p -adic étale cohomology $H_{\mathrm{ét}}^i(X, \mathbb{Q}_p)$, and Fontaine conjectured in [F1]¹ the existence of a natural p -adic period isomorphism

$$(0.3) \quad \rho : H_{\mathrm{dR}}^i(X) \otimes_{\bar{K}} \mathrm{B}_{\mathrm{dR}} \xrightarrow{\sim} H_{\mathrm{ét}}^i(X, \mathbb{Q}_p) \otimes \mathrm{B}_{\mathrm{dR}}.$$

Here B_{dR} is Fontaine’s p -adic periods field ([F1], [F3]). Recall that it is a complete discretely-valued field whose ring of integers $\mathrm{B}_{\mathrm{dR}}^+$ contains $\bar{K}$, the residue field $\mathrm{B}_{\mathrm{dR}}^+/\mathfrak{m}_{\mathrm{dR}}$ is Tate’s field $\mathbb{C}_p$, the cotangent line $\mathfrak{m}_{\mathrm{dR}}/\mathfrak{m}_{\mathrm{dR}}^2$ is the Tate twist $\mathbb{C}_p(1)$. Both sides of (0.3) carry natural filtrations (coming from the filtration of B_{dR} by powers of $\mathfrak{m}_{\mathrm{dR}}$ and the Hodge-Deligne filtration on $H_{\mathrm{dR}}^i(X)$), and ρ is compatible with them and with the $\mathrm{Gal}(\bar{K}/K)$ -conjugation. Moreover, as was envisioned by Fontaine and Jannsen ([F4], [J]), the matrix coefficients of ρ lie in the subring $\bar{K}\mathrm{B}_{\mathrm{st}}$ of B_{dR} ,² and ρ is compatible with the extra symmetries of log crystalline cohomology.

Received by the editors February 22, 2011 and, in revised form, November 16, 2011, and January 5, 2012.

2010 *Mathematics Subject Classification*. Primary 14F30, 14F40; Secondary 14F20.

Key words and phrases. p -Adic periods, derived de Rham cohomology, h-topology, alterations.

The author was supported in part by NSF grant DMS-1001660.

¹The assumption of loc. cit. that X is proper and smooth is redundant.

²This assertion is true for a general reason, since, by Berger’s theorem [Ber], de Rham Galois modules are potentially log crystalline.

The p -adic period map was defined in three different ways in works of, respectively, Faltings, Niziol, and Tsuji (with prior crucial input of Bloch, Fontaine, Hyodo, Kato, Kurihara, and Messing; the nonproper setting was treated by Yamashita [Y]), see [Fa1], [Fa2], [N1], [N2], [Ts1], [Ts2]; the three ρ 's coincide by [N3].

In the article we give another construction of ρ which is fairly direct and has the same flavor as the classical picture (0.2). The tools are derived de Rham cohomology of Illusie [Ill2, Ch. VIII] and de Jong's alterations. The companion paper [B] treats the Fontaine-Jannsen side of the story; another approach was developed by Bhatt [Bh2]. It would be very interesting to see if these methods can help to understand the Riemann-Hilbert correspondence in the p -adic setting.

An outline of the construction: First we realize B_{dR}^+ as the ring of de Rham p -adic constants in the sense of *derived* algebraic geometry. Namely, let A_{dR} be the derived de Rham algebra $L\Omega_{\hat{O}_{\bar{K}}/O_K}$ completed with respect to the Hodge filtration $F^\cdot$; see [Ill2, Ch. VIII (2.1.3.3)]. Here $O_K, O_{\bar{K}}$ are the rings of integers in $K, \bar{K}$. Now B_{dR}^+ identifies canonically with $A_{\mathrm{dR}} \hat{\otimes} \mathbb{Q}_p$, where $\hat{\otimes}$ is the derived completed tensor product, so that $\mathfrak{m}_{\mathrm{dR}}^i \xrightarrow{\sim} F^i A_{\mathrm{dR}} \hat{\otimes} \mathbb{Q}_p$. This fact was observed independently by Fargues [Far].

Let $\mathcal{V}ar_F$ be the category of varieties over a field F , and $\mathcal{V}ar_F^{\mathrm{nc}}$ be the category of regular F -varieties U equipped with a regular compactification $\bar{U}$ with normal crossings divisor at infinity. As follows from de Jong's theorem [dJ1], the forgetful functor $\mathcal{V}ar_F^{\mathrm{nc}} \rightarrow \mathcal{V}ar_F, (U, \bar{U}) \mapsto U$, makes $\mathcal{V}ar_F^{\mathrm{nc}}$ a base for the h-topology on $\mathcal{V}ar_F$, so h-sheaves on $\mathcal{V}ar_F$ are the same as sheaves on $\mathcal{V}ar_F^{\mathrm{nc}}$ for the induced topology. For $F = \bar{K}$ as above, there is a finer category $\mathcal{V}ar_{\bar{K}}^{\mathrm{ss}}$ of *ss-pairs* $(V, \bar{V})$, i.e., smooth $\bar{K}$ -varieties V equipped with a semi-stable compactification $\bar{V}$ (that includes compactification in the arithmetic direction). Again by de Jong [dJ1], $\mathcal{V}ar_{\bar{K}}^{\mathrm{ss}}$ is a base for the h-topology on $\mathcal{V}ar_{\bar{K}}$.

Consider the presheaf on $\mathcal{V}ar_{\bar{K}}^{\mathrm{ss}}$ which assigns to $(V, \bar{V})$ the derived de Rham algebra with log singularities $R\Gamma(\bar{V}, L\Omega_{(\hat{V}, \bar{V})/O_K})$ (see [Ol]). Its h-sheafification $\mathcal{A}_{\mathrm{dR}}^{\natural}$ is an h-sheaf of filtered dg algebras on $\mathcal{V}ar_{\bar{K}}$ that contains the constant subsheaf A_{dR} . The key *p -adic Poincaré lemma* says that the map $A_{\mathrm{dR}} \otimes^L \mathbb{Z}/p^n \rightarrow \mathcal{A}_{\mathrm{dR}}^{\natural} \otimes^L \mathbb{Z}/p^n$ is a filtered quasi-isomorphism. It comes from the next assertion: The h-sheafification of the presheaf $(V, \bar{V}) \mapsto H^b(\bar{V}, \Omega_{(V, \bar{V})/O_{\bar{K}}}^a)$, where $\Omega_{(V, \bar{V})/O_{\bar{K}}}^a$ is the usual locally free $\mathcal{O}_{\bar{V}}$ -module of forms with log singularities, is an h-sheaf of $\mathbb{Q}$ -vector spaces for $(a, b) \neq (0, 0)$. The case $a = 0$ is essentially theorem 8.0.1 from Bhatt's thesis [Bh1]; the general result is obtained by a similar method (which uses coverings of families of stable curves that come from the multiplication by p isogeny of the generalized Jacobians).

Set $R\Gamma_{\mathrm{dR}}^{\natural}(X) := R\Gamma(X_{\mathrm{h}}, \mathcal{A}_{\mathrm{dR}}^{\natural})$; this is the *arithmetic de Rham complex* of X . By the above, $H^*(R\Gamma_{\mathrm{dR}}^{\natural}(X) \hat{\otimes} \mathbb{Q}_p)$ is a B_{dR}^+ -algebra. One has a diagram

$$(0.4) \quad H_{\mathrm{dR}}^*(X) \xrightarrow{\alpha} H^*(R\Gamma_{\mathrm{dR}}^{\natural}(X) \hat{\otimes} \mathbb{Q}_p) \xleftarrow{\beta} H_{\mathrm{\acute{e}t}}^*(X, \mathbb{Q}_p) \otimes B_{\mathrm{dR}}^+,$$

where α is the composition $H_{\mathrm{dR}}^*(X) \xrightarrow{\sim} H^*(R\Gamma_{\mathrm{dR}}^{\natural}(X) \otimes \mathbb{Q}) \rightarrow H^*(R\Gamma_{\mathrm{dR}}^{\natural}(X) \hat{\otimes} \mathbb{Q}_p)$ and β is the B_{dR}^+ -linear extension of the evident map (which comes from the embeddings $\mathbb{Z}/p^n \rightarrow \mathcal{A}_{\mathrm{dR}}^{\natural} \otimes^L \mathbb{Z}/p^n$ and the fact that the h-topology is stronger than the étale one). Since the étale and h-cohomology with torsion coefficients coincide, the Poincaré lemma implies that β is an isomorphism. Now the p -adic period map ρ is

the B_{dR} -linear extension of $\beta^{-1}\alpha$. An explicit computation for $X = \mathbb{G}_m$ followed by usual tricks of the trade shows that ρ is a filtered isomorphism.

1. A DERIVED DE RHAM CONSTRUCTION OF B_{dR}

1.1. The derived p -adic completion. Throughout the article we use (not too heavily) E_∞ algebras, for which we refer to, say, [HS].³ Recall that E_∞ algebras are dg algebras whose product is commutative and associative up to coherent higher homotopies (more formally, E_∞ algebras are dg algebras for a resolution of the commutative algebra operad). A key fact: for any commutative (more generally, E_∞) cosimplicial dg algebra the corresponding total complex is naturally an E_∞ algebra. Thus the homotopy limit of a diagram of E_∞ algebras is an E_∞ algebra.

For a projective system of complexes of abelian groups $\dots \xrightarrow{\phi_2} C_2 \xrightarrow{\phi_1} C_1$, one has $\text{holim } C_n = \text{Cone}(\text{id} - \phi : \prod C_n \rightarrow \prod C_n)[-1]$, where $\phi((c_n)) = (\phi_n(c_{n+1}))$. There is an embedding $\varprojlim C_n = \text{Ker}(\text{id} - \phi) \hookrightarrow \text{holim } C_n$. If all ϕ_n 's are surjective, then $\text{id} - \phi$ is surjective; hence $\hookrightarrow$ is a quasi-isomorphism. So holim , being an exact functor, is the right derived functor of $\varprojlim$.

If C is a projective system of dg algebras, then $\text{holim } C_n$ is naturally a dg algebra (and the above embedding is an embedding of algebras); if the C_n are commutative (or, more generally, E_∞) algebras, then $\text{holim } C_n$ is an E_∞ algebra.

Let p be a prime. Consider the projective system of commutative dg algebras $C_n := \text{Cone}(\mathbb{Z} \xrightarrow{p^n} \mathbb{Z})$. It is a resolution of the projective system $\dots \rightarrow \mathbb{Z}/p^2 \rightarrow \mathbb{Z}/p$, so $\mathbb{Z}_p^b := \text{holim } C_n$ is an E_∞ algebra with $H^0 \mathbb{Z}_p^b = \mathbb{Z}_p$ and acyclic in nonzero degrees. Set $\mathbb{Q}_p^b := \mathbb{Z}_p^b \otimes \mathbb{Q}$. For any complex F of abelian groups set

$$(1.1.1) \quad F \hat{\otimes} \mathbb{Z}_p := \text{holim } (F \otimes C_n), \quad F \hat{\otimes} \mathbb{Q}_p := (F \hat{\otimes} \mathbb{Z}_p) \otimes \mathbb{Q}.$$

These are dg $\mathbb{Z}_p^b$ and $\mathbb{Q}_p^b$ -modules, so their cohomology groups are $\mathbb{Z}_p$ - and $\mathbb{Q}_p$ -modules, and $F \mapsto F \hat{\otimes} \mathbb{Z}_p, F \hat{\otimes} \mathbb{Q}_p$ are exact functors. If F is an (E_∞) dg algebra, then so are $F \hat{\otimes} \mathbb{Z}_p$ and $F \hat{\otimes} \mathbb{Q}_p$.

Remark. One has an evident projective system $F_{p^n}[1] \rightarrow F \otimes C_n \rightarrow F/p^n F$ of exact triangles; applying holim , we get a canonical exact triangle $\text{holim } F_{p^n}[1] \rightarrow F \hat{\otimes} \mathbb{Z}_p \rightarrow \text{holim}(F/p^n F)$. Let $\hat{F} := \varprojlim F/p^n F$ be the p -adic completion of F and $T_p F := \varprojlim F_{p^n}$ be the Tate module of F . By above, we have a quasi-isomorphism $\hat{F} \xrightarrow{\sim} \text{holim}(F/p^n F)$. Thus if F has no p -torsion, then $F \hat{\otimes} \mathbb{Z}_p \xrightarrow{\sim} \hat{F}$. Similarly, if all components of F are p -divisible, then one has quasi-isomorphisms $T_p F \xrightarrow{\sim} \text{holim } F_{p^n}$ and $T_p F[1] \xrightarrow{\sim} F \hat{\otimes} \mathbb{Z}_p$. We see that $\cdot \hat{\otimes} \mathbb{Z}_p$ is the left derived functor of the p -adic completion functor and the right derived functor of $T_p[1]$.

Example. For a scheme X , its étale $\mathbb{Z}_p$ - and $\mathbb{Q}_p$ -cohomology are $R\Gamma_{\text{ét}}(X, \mathbb{Z}_p) := \text{holim } R\Gamma(X_{\text{ét}}, \mathbb{Z}/p^n) = R\Gamma(X_{\text{ét}}, \mathbb{Z}) \hat{\otimes} \mathbb{Z}_p$, $R\Gamma_{\text{ét}}(X, \mathbb{Q}_p) := R\Gamma(X_{\text{ét}}, \mathbb{Z}) \hat{\otimes} \mathbb{Q}_p$.⁴

1.2. The derived de Rham algebra. For a morphism of commutative rings $A \rightarrow B$ we denote by $\Omega_{B/A}$ the relative de Rham complex of B over A . This is a commutative dg A -algebra with components $\Omega_{B/A}^i = \Lambda_B^i \Omega_{B/A}$, where $\Omega_{B/A}$ is the B -module of relative Kähler differentials; it carries a ring filtration $F^n = \Omega_{B/A}^{\geq n}$.

³In loc. cit. E_∞ algebras are called “May algebras”.

⁴Since $R\Gamma(X_{\text{ét}}, \mathbb{Z}/p^n) = R\Gamma(X_{\text{ét}}, \mathbb{Z}) \otimes^L \mathbb{Z}/p^n$.

We will use the F -completed version $L\Omega_{\hat{B}/A}$ of Illusie's derived de Rham algebra defined in [Ill2, Ch. VIII, (2.1.3.3)]. To construct it, consider the canonical simplicial resolution $P = P_A(B)$ of B from [Ill1, Ch. I, (1.5.5.6)]. This is a simplicial commutative A -algebra such that each P_i is a polynomial A -algebra. The de Rham complexes $\Omega_{P./A}$ form a simplicial filtered commutative dg A -algebra, so the corresponding total complex $L\Omega_{B/A}$ is a filtered commutative dg A -algebra (see [Ill1, Ch. I, 3.1.3]). Now $L\Omega_{\hat{B}/A}$ is its completion with respect to the filtration $F^\cdot$. Here “completion” is understood as a mere projective system of quotients modulo F^i . One has a natural identification $\mathrm{gr}_F^i L\Omega_{\hat{B}/A} \xrightarrow{\sim} (L\Lambda_B^i(L_{B/A}))[-i]$ compatible with the product; here $L_{B/A} := \Omega_{P./A} \otimes_P B$ is the relative cotangent complex and $L\Lambda_B^i$ is the nonabelian left derived functor of the exterior power functor (see Ch. II and Ch. I of [Ill1]). For A -flat B 's, the construction is compatible with base change. It is compatible with direct limits. If in the above definition we replace P by any simplicial A -algebra resolution of B whose terms are polynomial A -algebras, then the output is naturally quasi-isomorphic to $L\Omega_{\hat{B}/A}$.

The next lemma is a particular case of [Ill1, Ch. I, 4.3.2.1(ii)]. For a flat B -module T we denote by $B\langle T \rangle$ its divided powers symmetric algebra.

Lemma. *The complex $L\Lambda_B^i(T[1])$ is acyclic outside degree $-i$. There is a canonical isomorphism of graded B -algebras compatible with base change*

$$(1.2.1) \quad H^\cdot L\Lambda_B^\cdot(T[1]) \xrightarrow{\sim} B\langle T \rangle.$$

1.3. Let K be a p -adic field, i.e., a complete discretely-valued field of characteristic zero with perfect residue field k of characteristic $p > 0$, $\bar{K}$ be an algebraic closure of K , and $O_K, O_{\bar{K}}$ be rings of integers in $K, \bar{K}$. Let $K_0 \subset K$ be the field of fractions of the Witt vectors $W(k) = O_{K_0}$, and let $\mathfrak{a}$ be the fractional ideal in $\bar{K}$ generated by $p^{-\frac{1}{p-1}} \mathfrak{D}_{K/K_0}^{-1}$, where $\mathfrak{D}_{K/K_0}$ is the different. For an O_K -algebra B we often write $\Omega_B := \Omega_{B/O_K}$, $L\Omega_{\hat{B}} := L\Omega_{\hat{B}/O_K}$, $L_B = L_{B/O_K}$, etc.

The next key result is due to Fontaine [F2, Th 1]; we include a proof for completeness sake. Consider the map $\mu_{p^\infty} \subset O_K^\times \xrightarrow{d\log} \Omega_{O_{\bar{K}}}$ and its $O_{\bar{K}}$ -linear extension

$$(1.3.1) \quad (\bar{K}/O_{\bar{K}})(1) = O_{\bar{K}} \otimes \mu_{p^\infty} \rightarrow \Omega_{O_{\bar{K}}}.$$

Theorem. *One has $L_{O_{\bar{K}}} \xrightarrow{\sim} \Omega_{O_{\bar{K}}}$, and (1.3.1) is surjective with kernel $(\mathfrak{a}/O_{\bar{K}})(1)$.*

Proof. If K'/K is a finite extension, then $O_{K'}/O_K$ is a complete intersection. So, if π is a generator of $O_{K'}/O_K$, $f(t)$ its minimal polynomial, then $L_{O_{K'}}$ is the cone of multiplication by $f'(\pi)$ endomorphism of $O_{K'}$; hence $L_{O_{K'}} \xrightarrow{\sim} \Omega_{O_{K'}}$. Passing to the limit, we get the first assertion. Let us prove the second one.

(i) By the above, $\Omega_{O_{K'}} \simeq O_{K'}/\mathfrak{D}_{K'/K}$. If K''/K' is another finite extension, then the standard exact triangle of the cotangent complexes reduces to a short exact sequence $0 \rightarrow O_{K''} \otimes_{O_{K'}} \Omega_{O_{K'}/O_K} \rightarrow \Omega_{O_{K''}/O_K} \rightarrow \Omega_{O_{K''}/O_{K'}} \rightarrow 0$.

(ii) Replacing K', K by $\bar{K}, K_0$ and passing to the limit, we get a short exact sequence $0 \rightarrow O_{\bar{K}} \otimes_{O_K} \Omega_{O_K/O_{K_0}} \rightarrow \Omega_{O_{\bar{K}}/O_{K_0}} \rightarrow \Omega_{O_{\bar{K}}/O_K} \rightarrow 0$. Thus it suffices to prove the theorem for $K = K_0$, which we now assume.

(iii) Set $T := \mathrm{Ker}((\bar{K}/O_{\bar{K}})(1) \rightarrow \Omega)$, $F := K(\mu_p)$. The set of $O_{\bar{K}}$ -submodules of $(\bar{K}/O_{\bar{K}})(1)$ is totally ordered by inclusion. Thus, since $O_{\bar{K}} \otimes_{O_F} \Omega_{O_F} \subset \Omega_{O_{\bar{K}}}$ is a nonzero $O_{\bar{K}}$ -module generated by $d\log(\mu_p)$, one has $T \subset (p^{-1}O_{\bar{K}}/O_{\bar{K}})(1) = O_{\bar{K}} \otimes \mu_p$. Since Ω_{O_F} is isomorphic to $O_F/p^{1-\frac{1}{p-1}}O_F$, one has $T = (p^{-\frac{1}{p-1}}O_{\bar{K}}/O_{\bar{K}})(1)$.

(iv) It remains to prove surjectivity of $(\bar{K}/O_{\bar{K}})(1) \rightarrow \Omega_{O_{\bar{K}}}$. Let $K' \subset \bar{K}$ be any finite extension of K ; we want to check that $\Omega_{O_{K'}} \subset \Omega_{O_{\bar{K}}}$ lies in $O_{\bar{K}} d \log(\mu_{p^\infty})$. Suppose p^n kills $\Omega_{O_{K'}}$. Let us show that $\Omega_{O_{K'}} \subset O_{\bar{K}} d \log(\mu_{p^{n+1}})$. Set $K'' := K'(\mu_{p^{n+1}})$. The set of $O_{K''}$ -submodules of $\Omega_{O_{K''}}$ is totally ordered. Thus, since $p^n d \log(\mu_{p^{n+1}}) \neq 0$ by (iii), $\Omega_{O_{K'}}$ lies in $O_{K''} d \log(\mu_{p^{n+1}})$, q.e.d. $\square$

1.4. For a complex P acyclic in degrees $\neq 0$, we often write P instead of $H^0 P$.

Consider the filtered commutative dg O_K -algebra $A_{dR} = A_{dR} \bar{K}/K := L\Omega_{O_{\bar{K}}/O_K}$ and the corresponding filtered E_∞ O_K -algebra $A_{dR} \hat{\otimes} \mathbb{Z}_p$ (see §1.1). Let us describe the graded $O_{\bar{K}}$ -algebras $\mathrm{gr}_F A_{dR}$, $\mathrm{gr}_F A_{dR} \hat{\otimes} \mathbb{Z}_p$.

Proposition. (i) The complexes $\mathrm{gr}_F^i A_{dR} \hat{\otimes} \mathbb{Z}_p$ are acyclic in nonzero degrees, and there is a canonical isomorphism of graded algebras

$$(1.4.1) \quad \mathrm{gr}_F A_{dR} \hat{\otimes} \mathbb{Z}_p \xrightarrow{\sim} \hat{O}_{\bar{K}} \langle \hat{\mathbf{a}}(1) \rangle.$$

(ii) One has $\mathrm{gr}_F^0 A_{dR} = A_{dR}/F^1 = O_{\bar{K}}$, and the complexes $\mathrm{gr}_F^i A_{dR}$ for $i > 0$ are acyclic in degrees $\neq 1$. There are natural isomorphisms of $O_{\bar{K}}$ -modules

$$(1.4.2) \quad \Omega^{(i)} := H^1 \mathrm{gr}_F^i A_{dR} \xrightarrow{\sim} (\bar{K}/i!^{-1} \mathbf{a}^i)(i) = (\mathbb{Q}_p/\mathbb{Z}_p) \otimes i!^{-1} \hat{\mathbf{a}}^i(i).$$

Proof. (i) By the theorem in §1.3, one has $L_{O_{\bar{K}}/O_K} \xrightarrow{\sim} \Omega_{O_{\bar{K}}} \xrightarrow{\sim} (\bar{K}/\mathbf{a})(1) = (\mathbb{Q}_p/\mathbb{Z}_p) \otimes \mathbf{a}(1)$; hence $\mathrm{gr}_F^i A_{dR} \xrightarrow{\sim} L\Lambda_{\mathbb{Z}}^i(\mathbb{Q}_p/\mathbb{Z}_p)[-i] \otimes \mathbf{a}^i(i)$. One has $L\Lambda_{\mathbb{Z}}^i(\mathbb{Q}_p/\mathbb{Z}_p) \otimes^L (\mathbb{Z}/p^n) = L\Lambda_{\mathbb{Z}/p^n}^i((\mathbb{Q}_p/\mathbb{Z}_p) \otimes^L (\mathbb{Z}/p^n)) = L\Lambda_{\mathbb{Z}/p^n}^i((\mathbb{Z}/p^n)[1])$, which identifies with $i!^{-1}(\mathbb{Z}/p^n)[i]$ in a way compatible with the product by (1.2.1). Therefore $\mathrm{gr}_F^i A_{dR} \otimes^L (\mathbb{Z}/p^n) \xrightarrow{\sim} \mathbb{Z}/p^n \langle (\mathbf{a}/p^n \mathbf{a})(1) \rangle$, which yields (1.4.1).

(ii) follows from (i) by the next observation (applied to $C = \mathrm{gr}_F^i A_{dR}$, with (1.4.2) defined by the condition that $T_p(1.4.2) = (1.4.1)$): If a complex C of abelian groups has p -torsion cohomology and $H^{\neq 0}(C \otimes^L \mathbb{Z}/p) = 0$, then $H^1 C$ is p -divisible and $H^{\neq 1} C = 0$.⁵ $\square$

1.5. By 1.4(i), the algebras $(A_{dR}/F^i) \hat{\otimes} \mathbb{Z}_p$, hence $(A_{dR}/F^i) \hat{\otimes} \mathbb{Q}_p$, are acyclic in non-zero degree. By loc. cit., $(A_{dR}/F^{i+1}) \hat{\otimes} \mathbb{Q}_p$ is an i -truncated dvr with residue field $\mathbb{C}_p := \hat{O}_{\bar{K}} \otimes \mathbb{Q}$, so $A_{dR} \hat{\otimes} \mathbb{Q}_p := \varprojlim (A_{dR}/F^i) \hat{\otimes} \mathbb{Q}_p$ is a dvr. Let $\mathfrak{m}_{dR}$ be its maximal ideal; (1.4.1) yields a canonical identification $\mathfrak{m}_{dR}/\mathfrak{m}_{dR}^2 = \mathrm{gr}_F^1 A_{dR} \hat{\otimes} \mathbb{Q}_p \xrightarrow{\sim} \mathbb{C}_p(1)$.

Proposition. There is a canonical ring isomorphism of filtered rings

$$(1.5.1) \quad u_{\mathbb{Q}} : B_{dR}^+ \xrightarrow{\sim} A_{dR} \hat{\otimes} \mathbb{Q}_p.$$

Proof. The ring $(A_{dR}/F^{i+1}) \hat{\otimes} \mathbb{Z}_p$ is an infinitesimal p -adic O_K -thickening of $\hat{O}_K = (A_{dR}/F^1) \hat{\otimes} \mathbb{Z}_p$ of order $\leq i$ (see [F3, 1.1]). Let A_{inf}/F^{i+1} be the universal thickening ([F3, 1.3]); we have a canonical map $u_i : A_{\mathrm{inf}}/F^{i+1} \rightarrow (A_{dR}/F^{i+1}) \hat{\otimes} \mathbb{Z}_p$. Since $B_{dR}^+/F^{i+1} := (A_{\mathrm{inf}}/F^{i+1}) \otimes \mathbb{Q}$ is an i -truncated dvr and u_1 is an isomorphism by [F3, 1.4.3], $u_{i\mathbb{Q}} : B_{dR}^+/F^{i+1} \xrightarrow{\sim} (A_{dR}/F^{i+1}) \hat{\otimes} \mathbb{Q}_p$. Set $u_{\mathbb{Q}} := \varprojlim u_{i\mathbb{Q}}$. $\square$

Remarks. (i) The map $A_{dR} \rightarrow A_{dR}/F^1 = O_{\bar{K}}$ yields an isomorphism $A_{dR} \otimes \mathbb{Q} \xrightarrow{\sim} \bar{K}$. Thus the morphism $A_{dR} \otimes \mathbb{Q} \rightarrow A_{dR} \hat{\otimes} \mathbb{Q}_p$ equals the usual embedding $\bar{K} \hookrightarrow B_{dR}^+$.

⁵Use the fact that every complex of abelian groups splits, i.e., is quasi-isomorphic to a complex with zero differential.

(ii) For a finite extension K'/K , $K' \subset \bar{K}$, the evident map $A_{dR} \bar{K}/K \rightarrow A_{dR} \bar{K}/K'$ yields an isomorphism $A_{dR} \bar{K}/K \hat{\otimes} \mathbb{Q}_p \xrightarrow{\sim} A_{dR} \bar{K}/K' \hat{\otimes} \mathbb{Q}_p$ compatible with (1.5.1).

1.6. The next result, which will not be used in the rest of the article, is a reinterpretation of Colmez's theorem [Col]. It would be nice to find a simpler direct proof.

Proposition. *The complexes A_{dR}/F^i are acyclic in nonzero degrees; the maps $H^0(A_{dR}/F^{i+1}) \rightarrow H^0(A_{dR}/F^i)$ are injective. Set $O^{(i)} := H^0(A_{dR}/F^{i+1})$; thus $O_{\bar{K}} = O^{(0)} \supset O^{(1)} \supset \dots$ and $(A_{dR}/F^{i+1}) \hat{\otimes} \mathbb{Z}_p$ is equal to the p -adic completion $\hat{O}^{(i)}$ of $O^{(i)}$.*

Proof. By 1.4(ii), the exact cohomology sequence for $0 \rightarrow \mathrm{gr}_F^i A_{dR} \rightarrow A_{dR}/F^{i+1} \rightarrow A_{dR}/F^i \rightarrow 0$ reduces to $0 \rightarrow O^{(i)} \rightarrow O^{(i-1)} \xrightarrow{d^{(i)}} \Omega^{(i)} \rightarrow H^1(A_{dR}/F^{i+1}) \rightarrow H^1(A_{dR}/F^i) \rightarrow 0$. So $O_{\bar{K}} = O^{(0)} \supset O^{(1)} \supset \dots$, and the vanishing of $H^1(A_{dR}/F^{i+1})$ amounts to that of $H^1(A_{dR}/F^i)$ combined with surjectivity of $d^{(i)} : O^{(i-1)} \rightarrow \Omega^{(i)}$. It remains to prove that all $d^{(i)}$ are surjective.

Recall that Colmez [Col] considers a sequence of subalgebras $O_{\bar{K}} = O^{(0)} \supset O^{(1)} \supset \dots$ and derivations $d^{(i)} : O^{(i-1)} \rightarrow \Omega^{(i)}$ defined by induction: $d^{(i)}$ is a universal O_K -linear derivation with values in an $O_{\bar{K}}$ -module, and $O^{(i)} := \mathrm{Ker} d^{(i)}$. An induction by i shows that $O^{(i)} \supset O^{(i)}$: Indeed, $\Omega^{(i)}$ are $O_{\bar{K}}$ -modules and $d^{(i)} : O^{(i-1)} \rightarrow \Omega^{(i)}$ is a derivation; so, if $O^{(i-1)} \supset O^{(i-1)}$, then $d^{(i)}|_{O^{(i-1)}} = a^{(i)} d^{(i)}$ for some $O_{\bar{K}}$ -linear map $a^{(i)} : \Omega^{(i)} \rightarrow \Omega^{(i)}$; thus $O^{(i)} \supset O^{(i)}$.

Let i be the smallest number such that $d^{(i)}$ is not surjective. Since $E := \Omega^{(i)}/d^{(i)}(O^{(i-1)})$ is p -torsion p -divisible, one has $E \hat{\otimes} \mathbb{Q}_p = T_p E \otimes \mathbb{Q} \neq 0$. Applying $\hat{\otimes} \mathbb{Z}_p$ to the exact triangle $O^{(i)} \rightarrow A_{dR}/F^{i+1} \rightarrow E[-1]$, we get a short exact sequence $0 \rightarrow \hat{O}^{(i)} \rightarrow (A_{dR}/F^{i+1}) \hat{\otimes} \mathbb{Z}_p \rightarrow T_p E \rightarrow 0$. By [Col], $A_{\mathrm{inf}}/F^{i+1} = \hat{O}^{(i)}$. By universality, the map $u_i : A_{\mathrm{inf}}/F^{i+1} \rightarrow (A_{dR}/F^{i+1}) \hat{\otimes} \mathbb{Z}_p$ equals the composition $\hat{O}^{(i)} \rightarrow \hat{O}^{(i)} \hookrightarrow (A_{dR}/F^{i+1}) \hat{\otimes} \mathbb{Z}_p$, so its composition with the projection onto $T_p E$ vanishes. This cannot happen since $u_{i\mathbb{Q}}$ is an isomorphism (see §1.5), q.e.d. $\square$

2. h-TOPOLOGY AND SEMI-STABLE COMPACTIFICATIONS

2.1. *A topological digression.* The next proposition is a generalization of [V2, 4.1].

Let $\mathcal{V}$ be an essentially small site. As in [V1], we denote by $\mathcal{V}^\sim$ the corresponding topos (the category of sheaves of sets on $\mathcal{V}$).

For us, a *base for $\mathcal{V}$* is a pair $(\mathcal{B}, \phi)$, where $\mathcal{B}$ is an essentially small category and $\phi : \mathcal{B} \rightarrow \mathcal{V}$ is a *faithful* functor, that satisfies the next property:

(*) *For any $V \in \mathcal{V}$ and a finite family of pairs (B_α, f_α) , $B_\alpha \in \mathcal{B}$, $f_\alpha : V \rightarrow \phi(B_\alpha)$, there exists a set of objects $B'_\beta \in \mathcal{B}$ and a covering family $\{\phi(B'_\beta) \rightarrow V\}$ such that every composition $\phi(B'_\beta) \rightarrow V \rightarrow \phi(B_\alpha)$ lies in $\mathrm{Hom}(B'_\beta, B_\alpha) \subset \mathrm{Hom}(\phi(B'_\beta), \phi(B_\alpha))$.*

Remarks. (i) Property (*) for an empty set of (B_α, f_α) 's means that every $V \in \mathcal{V}$ has a covering by objects $\phi(B)$, $B \in \mathcal{B}$. If ϕ is fully faithful, then (*) amounts to this assertion.⁶

(ii) If $\mathcal{B}$ admits finite products and ϕ commutes with finite products, then it suffices to check (*) for families (B_α, f_α) having ≤ 1 element.

⁶The proposition below in this situation amounts to [V2, 4.1].

(iii) In the general case, it suffices to check $(*)$ for families (B_α, f_α) having ≤ 2 elements.

Suppose $(\mathcal{B}, \phi)$ is a base for $\mathcal{V}$. Define a covering sieve in $\mathcal{B}$ as a sieve whose ϕ -image is a covering family in $\mathcal{V}$.

Proposition. (i) *Covering sieves in $\mathcal{B}$ form a Grothendieck topology on $\mathcal{B}$.*

(ii) *The functor $\phi : \mathcal{B} \rightarrow \mathcal{V}$ is continuous (see [V2, 1.1]).*

(iii) *ϕ yields an equivalence of the toposes: one has $\mathcal{B}^\sim \xrightarrow{\sim} \mathcal{V}^\sim$.*

We call the above topology on $\mathcal{B}$ the ϕ -induced topology.⁷

Proof. (i) Let us check that covering sieves in $\mathcal{B}$ are stable with respect to pullback; the rest of the axioms from [V1, 1.1] are evident. For a morphism $g : B' \rightarrow B$ in $\mathcal{B}$ and a covering sieve $\mathfrak{s}$ on B , let us find a covering family on B' that belongs to the g -pullback of $\mathfrak{s}$. The $\phi(g)$ -pullback of $\phi(\mathfrak{s})$ is a covering sieve in $\mathcal{V}$, so there is a covering family $\{\pi_\gamma : V_\gamma \rightarrow \phi(B')\}$ such that every composition $V_\gamma \rightarrow \phi(B') \rightarrow \phi(B)$ can be factored as $V_\gamma \xrightarrow{g_\gamma} \phi(B_\gamma) \xrightarrow{\phi(p_\gamma)} \phi(B)$, where $p_\gamma : B_\gamma \rightarrow B$ belong to $\mathfrak{s}$. Applying $(*)$ to V_γ and (B', π_γ) , (B_γ, g_γ) , we find a covering family $\{\phi(B'_{\beta_\gamma}) \rightarrow V_\gamma\}$ as in $(*)$. The composite covering $\{\phi(B'_{\beta_\gamma}) \rightarrow \phi(B')\}$ comes then from a covering family $\{B'_{\beta_\gamma} \rightarrow B'\}$ in $\mathcal{B}$ which lies in the g -pullback of $\mathfrak{s}$.

(ii) We know that ϕ sends covering families to covering families, so it suffices to show that for any given $p_\alpha : B_\alpha \rightarrow B$ in $\mathcal{B}$ and $f_\alpha : V \rightarrow \phi(B_\alpha)$, $\alpha = 1, 2$, such that $\phi(p_1)f_1 = \phi(p_2)f_2$ there is a covering $\{\pi_\beta : V_\beta \rightarrow V\}$ and morphisms $\xi_{\alpha\beta} : B'_\beta \rightarrow B_\alpha$, $g_\beta : V_\beta \rightarrow \phi(B'_\beta)$ such that $p_1\xi_{1\beta} = p_2\xi_{2\beta}$ and $\phi(\xi_{\alpha\beta})g_\beta = f_\alpha\pi_\beta$. Such a datum (with g_β the identity map) comes from $(*)$ applied to V and $(B_1, f_1), (B_2, f_2)$.

(iii) By (ii), one has the usual adjoint functors between the categories of sheaves $(\phi^s, \phi_s) : \mathcal{B}^\sim \rightleftarrows \mathcal{V}^\sim$. To prove that they are mutually inverse equivalences, we will check that for $\mathcal{F} \in \mathcal{B}^\sim$ and $\mathcal{G} \in \mathcal{V}^\sim$ the adjunction maps $a_{\mathcal{F}} : \mathcal{F} \rightarrow \phi_s\phi^s\mathcal{F}$, $b_{\mathcal{G}} : \phi^s\phi_s\mathcal{G} \rightarrow \mathcal{G}$ are isomorphisms.

Recall that $\phi^s\mathcal{F} = (\phi^*\mathcal{F})^\sim$, where ϕ^* is the pullback of presheaves and $^\sim$ is the sheafification functor. For $V \in \mathcal{V}$ one has $(\phi^*\mathcal{F})(V) = \text{colim}_{\mathcal{C}(V)} \mathcal{F}$, where $\mathcal{C}(V)$ is the category of pairs (B, f) , $B \in \mathcal{B}$, $f : V \rightarrow \phi(B)$, with $\text{Hom}_{\mathcal{C}(V)}((B, f), (B', f')) := \{g \in \text{Hom}(B', B) : \phi(g)f' = f\}$, and we set $\mathcal{F}(B, f) := \mathcal{F}(B)$.

(a) To show that $a_{\mathcal{F}}$ is an isomorphism, we check that it is injective and surjective:

$a_{\mathcal{F}}$ is injective: Suppose we have $B \in \mathcal{B}$ and $\xi_1, \xi_2 \in \mathcal{F}(B)$ such that $a_{\mathcal{F}}(\xi_1) = a_{\mathcal{F}}(\xi_2)$; let us show that the ξ_i coincide. One has

$$a_{\mathcal{F}}(\xi_i) \in (\phi_s\phi^s\mathcal{F})(B) = (\phi^s\mathcal{F})(\phi(B)),$$

and the equality means that there is a covering $\{\pi_\gamma : V_\gamma \rightarrow \phi(B)\}$ such that the images of ξ_i in $(\phi^*\mathcal{F})(V_\gamma) = \text{colim}_{\mathcal{C}(V_\gamma)} \mathcal{F}$ coincide. Thus for some *finite* subdiagram $\mathcal{C}(V_\gamma)' \subset \mathcal{C}(V_\gamma)$ that contains (B, π_γ) the images of ξ_i in $\text{colim}_{\mathcal{C}(V_\gamma)'} \mathcal{F}$ coincide. Applying $(*)$ to V_γ and pairs from $\mathcal{C}(V_\gamma)'$, we get a covering $\{\phi(B'_{\beta_\gamma}) \rightarrow V_\gamma\}$ such that the image of $\mathcal{C}(V_\gamma)'$ in each $\mathcal{C}(\phi(B'_{\beta_\gamma}))$ comes from a diagram in $\mathcal{B}^\circ/B'_{\beta_\gamma}$. The composite covering $\{\phi(B'_{\beta_\gamma}) \rightarrow \phi(B)\}$ comes then from a covering $\{B'_{\beta_\gamma} \rightarrow B\}$ in $\mathcal{B}$, and the images of ξ_i in $\mathcal{F}(B'_{\beta_\gamma})$ coincide. Then $\xi_1 = \xi_2$ since $\mathcal{F}$ is a sheaf, q.e.d.

⁷The terminology is compatible with that of [V2, 3.1].

$a_{\mathcal{F}}$ is surjective: For $B \in \mathcal{B}$, $\chi \in (\phi_s \phi^s \mathcal{F})(B)$ we look for a covering $\{B'_\beta \rightarrow B\}$ in $\mathcal{B}$ such that $\chi|_{B'_\beta}$ lies in the image of $\mathcal{F}(B'_\beta) \rightarrow (\phi_s \phi^s \mathcal{F})(B'_\beta)$. To find it, consider χ as an element of $(\phi^s \mathcal{F})(\phi(B))$. There is a covering $\{\pi_\gamma : V_\gamma \rightarrow \phi(B)\}$ such that $\chi|_{V_\gamma}$ lies in the image of $(\phi^s \mathcal{F})(V_\gamma) \rightarrow (\phi^s \mathcal{F})(\phi(B))$; i.e., one has $f_\gamma : V_\gamma \rightarrow \phi(B_\gamma)$ such that $\chi|_{V_\gamma}$ lies in the image of the composition $\mathcal{F}(B_\gamma) \rightarrow (\phi^s \mathcal{F})(\phi(B_\gamma)) \rightarrow (\phi^s \mathcal{F})(V_\gamma)$, where the second arrow comes from f_γ . Applying $(*)$ to V_γ and (B, π_γ) , (B_γ, f_γ) , we find a covering $\{\phi(B'_{\beta_\gamma}) \rightarrow V_\gamma\}$ as in $(*)$; the composite covering $\{\phi(B'_{\beta_\gamma}) \rightarrow \phi(B)\}$ comes then from a covering $\{B'_{\beta_\gamma} \rightarrow B\}$ that satisfies the promised property.

(b) $b_{\mathcal{G}}$ is an isomorphism: Since $\phi_s(b_{\mathcal{G}})a_{\phi_s \mathcal{G}} = \text{id}_{\phi_s \mathcal{G}}$ and we already know that $a_{\phi_s \mathcal{G}}$ is an isomorphism, we see that $\phi_s(b_{\mathcal{G}}) : \phi_s \phi^s \phi_s(\mathcal{G}) \rightarrow \phi_s \mathcal{G}$ is an isomorphism. Thus $b_{\mathcal{G}}(B) : \phi^s \phi_s \mathcal{G}(\phi(B)) \rightarrow \mathcal{G}(\phi(B))$ is an isomorphism for every $B \in \mathcal{B}$. Since every $V \in \mathcal{V}$ admits a covering by objects $\phi(B)$, $B \in \mathcal{B}$, this implies that $b_{\mathcal{G}}$ is both injective and surjective, hence an isomorphism, q.e.d. $\square$

Exercises. (i) For any presheaf $\mathcal{J}$ on $\mathcal{V}$ one has $\phi_s(\mathcal{J}^\sim) = (\phi_* \mathcal{J})^\sim$.
(ii) Suppose $(\mathcal{B}, \phi)$ is a base for $\mathcal{V}$ and $(\mathcal{B}', \phi')$ is a base for the ϕ -induced topology on $\mathcal{B}$. Then $(\mathcal{B}', \phi\phi')$ is a base for $\mathcal{V}$.

2.2. For a field K , let Var_K be the category of K -varieties, i.e., reduced separated K -schemes of finite type. We will consider categories $\mathcal{B}$ formed by varieties equipped with appropriate compactifications, referred to as *pairs*:

(a) *Geometric setting*: Let $j : U \hookrightarrow \bar{U}$ be an open embedding such that $\bar{U}$ is proper and U is dense in $\bar{U}$. We call such a datum a *geometric pair over K* , or *geometric K -pair*, and denote it by $(U, \bar{U})$. We say that $(U, \bar{U})$ is a regular normal crossings pair, *nc-pair* for short, if $\bar{U}$ is a regular scheme and $\bar{U} \setminus U$ is a divisor with normal crossings in $\bar{U}$; it is a *strict nc-pair* if the irreducible components of $\bar{U} \setminus U$ are regular. A morphism $f : (U', \bar{U}') \rightarrow (U, \bar{U})$ of pairs is a map $\bar{U}' \rightarrow \bar{U}$ which sends U' to U . We denote the category of geometric K -pairs by Var_K^c ; let Var_K^{nc} be the full subcategory of nc-pairs.

(b) *Arithmetic K -setting*: Suppose K is a p -adic field as in §1.3. An *arithmetic pair over K* , a.k.a. *arithmetic K -pair*, is an open embedding $j : U \hookrightarrow \bar{U}$ with dense image of a K -variety U into a reduced proper flat O_K -scheme $\bar{U}$.

For such a $(U, \bar{U})$ we set $O_{K_U} := \Gamma(\bar{U}, \mathcal{O}_{\bar{U}})$, $K_U := \Gamma(\bar{U}_K, \mathcal{O}_{\bar{U}})$. Then K_U is the product of several finite extensions of K (labeled by the connected components of $\bar{U}_K$; if $\bar{U}$ is normal, then O_{K_U} is the product of the corresponding rings of integers. The closed fiber $\bar{U}_s$ of $\bar{U}$ is the union of fibers over the closed points of O_{K_U} .

We say that $(U, \bar{U})$ is a *semi-stable pair*, or simply *ss-pair*, if (i) $\bar{U}$ is a regular scheme, (ii) $\bar{U} \setminus U$ is a divisor with normal crossings on $\bar{U}$, and (iii) the closed fiber $\bar{U}_s$ is reduced. Our ss-pair is *strict* if the irreducible components of $\bar{U} \setminus U$ are regular. Arithmetic K -pairs form a category Var_K^{cc} ; let Var_K^{ss} be the full subcategory of ss-pairs.

(c) *Arithmetic $\bar{K}$ -setting*: For K as in (b), let $\bar{K}$ be its algebraic closure. An *arithmetic pair over $\bar{K}$* , a.k.a. *arithmetic $\bar{K}$ -pair*, is an open embedding $j : V \hookrightarrow \bar{V}$ with dense image of a $\bar{K}$ -variety V into a reduced proper flat $O_{\bar{K}}$ -scheme $\bar{V}$. A connected $(V, \bar{V})$ is said to be *semi-stable*, a.k.a. *ss-pair*, if there exists an ss-pair $(U, \bar{U})$ over K and a $\bar{K}$ -point $\alpha : K_U \rightarrow \bar{K}$ (see (b)) such that $(V, \bar{V})$ is isomorphic to $(U, \bar{U})_\alpha = (U_{\bar{K}}, \bar{U}_{O_{\bar{K}}}) := (U \otimes_{K_U} \bar{K}, \bar{U} \otimes_{O_{K_U}} O_{\bar{K}})$. Then $\bar{V}$ is normal (say, by Serre's criterion). An arbitrary $(V, \bar{V})$ is semi-stable if such are all its connected

components. Denote by $\mathcal{V}ar_K^{\text{cc}}$ the category of all arithmetic pairs over $\bar{K}$, and by $\mathcal{V}ar_K^{\text{ss}} = \mathcal{V}ar_{\bar{K}/K}^{\text{ss}}$ its full subcategory of ss-pairs.

Remark. If K' is a finite extension of K contained in $\bar{K}$, then $\mathcal{V}ar_{\bar{K}/K'}^{\text{ss}} \subset \mathcal{V}ar_{\bar{K}/K}^{\text{ss}}$. For all the constructions below the difference between them is irrelevant.

These categories are connected by commutative diagrams of functors

$$(2.2.1) \quad \begin{array}{ccccccc} \mathcal{V}ar_K^{\text{cc}} & \rightarrow & \mathcal{V}ar_K^{\text{c}} & \rightarrow & \mathcal{V}ar_K, & \mathcal{V}ar_K^{\text{cc}} & \rightarrow & \mathcal{V}ar_K^{\text{c}} & \rightarrow & \mathcal{V}ar_{\bar{K}}, \\ \uparrow & & \uparrow & & & \uparrow & & \uparrow & & \\ \mathcal{V}ar_K^{\text{ss}} & \rightarrow & \mathcal{V}ar_K^{\text{nc}} & & & \mathcal{V}ar_K^{\text{ss}} & \rightarrow & \mathcal{V}ar_K^{\text{nc}} & & \end{array}$$

where the vertical arrows are the fully faithful embeddings, and the upper horizontal lines are faithful forgetful functors of passing to the generic fiber and $(U, \bar{U}) \mapsto U$. The K - and $\bar{K}$ -settings are connected by base change functors

$$(2.2.2) \quad \begin{array}{ccccc} \mathcal{V}ar_{\bar{K}}^{\text{ss}} & \rightarrow & \mathcal{V}ar_{\bar{K}}^{\text{nc}} & \rightarrow & \mathcal{V}ar_{\bar{K}} \\ \uparrow & & \uparrow & & \uparrow \\ \mathcal{V}ar_K^{\text{ss}} & \rightarrow & \mathcal{V}ar_K^{\text{nc}} & \rightarrow & \mathcal{V}ar_K. \end{array}$$

Here the two right vertical arrows are the evident base change $\cdot \otimes_K \bar{K}$, and the left one assigns to a semi-stable K -pair $(U, \bar{U})$ the disjoint sum of pairs $(U, \bar{U})_\alpha$ for all $\bar{K}$ -points $\alpha : K_U \rightarrow \bar{K}$.

2.3. A morphism $f : (V, \bar{V}) \rightarrow (U, \bar{U})$ of pairs in either of the settings of §2.2 is called an *alteration* (of $(U, \bar{U})$) if $f^{-1}(U) = V$, the generic fibers of f are zero-dimensional, and their union is dense in V . In setting (a), f is a *(strict) nc-alteration* if $(V, \bar{V})$ is a (strict) nc-pair; in settings (b), (c), f is a *(strict) ss-alteration* if $(V, \bar{V})$ is a (strict) ss-pair.

If f is an alteration, then $f|_V : V \rightarrow U$ is proper and surjective; the composition of alterations is an alteration.

Here is a key result of de Jong [dJ1, 4.1, 6.5]:

Theorem. *Every geometric pair admits a strict nc-alteration. Every arithmetic pair, either over K or over $\bar{K}$, admits a strict ss-alteration. The alterations can be chosen so that $\bar{V}$ is projective.*

Remark. Our conventions slightly differ from de Jong's: he understands varieties to be irreducible and semi-stable K -pairs $(U, \bar{U})$ to have property $K_U = K$; his notation for $(U, \bar{U})$ is $(\bar{U}, Z)$, $Z := \bar{U} \setminus U$.

2.4. For a field K , the *h-topology* (see [SV]) on $\mathcal{V}ar_K$ is generated by the pretopology whose coverings are finite families of maps $\{Y_\alpha \rightarrow X\}$ such that $Y := \coprod Y_\alpha \rightarrow X$ is a universal topological epimorphism.⁸ It is stronger than the étale and proper topologies.⁹ We denote the h-site by $\mathcal{V}ar_{K\text{h}}$; the h-site of X is denoted by X_{h} .

Exercise. Let $f : Y \rightarrow X$ be a morphism in $\mathcal{V}ar_K$.

(i) f is an h-covering if (and only if) for every irreducible curve $C \subset X$ the base change $Y_{\tilde{C}} \rightarrow \tilde{C}$, where $\tilde{C}$ is the normalization of C , is an h-covering.¹⁰

⁸This means that a subset of X is Zariski open if (and only if) its preimage in Y is open, and the same is true after any base change.

⁹The latter is generated by a pretopology whose coverings are proper surjective maps.

¹⁰Hint: For an open $V \subset Y$ its image in X is constructible (EGA IV 1.8.4), so to show that $f(V)$ is open it suffices to check that for any curve $C \subset X$ the intersection $C \cap f(V)$ is open in C .

(ii) If X is a regular curve, then f is an h -covering if (and only if) the closure of the generic fiber of f maps onto X , or, equivalently, f is a covering for the flat topology.

Remark. By [SV, 10.4], every h -covering is a Zariski covering locally in proper topology. Therefore (see [D], [SD], or [Con]) h -coverings are morphisms of universal cohomological descent for torsion étale sheaves; if $K = \mathbb{C}$, then h -coverings are morphisms of universal cohomological descent for arbitrary sheaves on the classical topology. In particular, for any h -hypercovering Y of X and an abelian group A the canonical map $R\Gamma(X_{\text{ét}}, A) \rightarrow R\Gamma(Y_{\text{ét}}, A)$ ($:=$ the total complex of the cosimplicial system of complexes $R\Gamma(Y_{i\text{ét}}, A)$) is a quasi-isomorphism if A is a torsion group. Passing to the limit, we see that $R\Gamma_{\text{ét}}(X, \mathbb{Z}_p) \xrightarrow{\sim} R\Gamma_{\text{ét}}(Y, \mathbb{Z}_p)$. If $K = \mathbb{C}$, then $R\Gamma(X_{\text{cl}}, A) \xrightarrow{\sim} R\Gamma(Y_{\text{cl}}, A)$ for any A . Since h -topology is stronger than the étale one, we see that $R\Gamma(X_{\text{ét}}, A) \xrightarrow{\sim} R\Gamma(X_h, A)$ if A is a torsion group (see [SV, 10.7] for a direct proof).

2.5. Let ϕ be the forgetful functor $(U, \bar{U}) \mapsto U$ on any of the categories $\mathcal{V}ar^?$ in §2.2.

Proposition. *If $\mathcal{V}ar^?$ is either of the categories $\mathcal{V}ar_K^c$, $\mathcal{V}ar_K^{nc}$, $\mathcal{V}ar_K^{cc}$, $\mathcal{V}ar_K^{ss}$, then $(\mathcal{V}ar^?, \phi)$ is a base for $\mathcal{V}ar_{Kh}$. If $\mathcal{V}ar^?$ is $\mathcal{V}ar_K^{cc}$ or $\mathcal{V}ar_K^{ss}$, then $(\mathcal{V}ar^?, \phi)$ is a base for $\mathcal{V}ar_{Kh}$.*

Proof. We consider the arithmetic K -setting, leaving the other two settings for the reader. Let us show that $(\mathcal{V}ar_K^{cc}, \phi)$ satisfies condition $(*)$ from §2.1. Our datum is a K -variety V and a finite collection of arithmetic K -pairs $(U_\alpha, \bar{U}_\alpha)$ and maps $f_\alpha : V \rightarrow U_\alpha$. We need to find an h -covering $\pi : V' \rightarrow V$ and an arithmetic pair $(V', \bar{V}')$ such that $f_\alpha \pi$ extend to maps $(V', \bar{V}') \rightarrow (U_\alpha, \bar{U}_\alpha)$. First we find an h -covering $V' \rightarrow V$ such that V' sits in some arithmetic K -pair $(V', \tilde{V}')$: let V'/V be a quasi-projective modification of V provided by the Chow lemma, and take for $\tilde{V}'$ the closure of V' in a projective space.¹¹ Then take $\bar{V}'$ to be the closure of the image of V' by the embedding $V' \hookrightarrow \tilde{V}' \times \Pi \bar{U}_\alpha$, and we are done.

To show that $(\mathcal{V}ar_K^{ss}, \phi)$ is a base for $\mathcal{V}ar_{Kh}$, it suffices to check that $(\mathcal{V}ar_K^{ss}, \iota)$, where ι is the embedding $\mathcal{V}ar_K^{ss} \hookrightarrow \mathcal{V}ar_K^{cc}$, is a base for the ϕ -induced topology on $\mathcal{V}ar_K^{cc}$ (see Exercise (ii) in §2.1). Since ι is fully faithful, it suffices to check that for every $(U, \bar{U}) \in \mathcal{V}ar_K^{cc}$ there exists a map $(U', \bar{U}') \rightarrow (U, \bar{U})$ such that $U' \rightarrow U$ is an h -covering and $(U', \bar{U}')$ is semi-stable. Such a datum is provided by the de Jong theorem in §2.3, and we are done. $\square$

We call the ϕ -induced topology on either of the categories $\mathcal{V}ar^?$ the h -topology.

Remarks. (i) Any h -covering of $(U, \bar{U}) \in \mathcal{V}ar_K^{ss}$ has a refinement with terms of the same dimension as U (indeed, the same assertion in $\mathcal{V}ar_K$ is true by [SV, 10.4]; to pass to $\mathcal{V}ar_K^{ss}$, we apply the constructions from the proof above, and they preserve the dimension).

(ii) The proposition remains true if we replace the category of ss- or nc-pairs by its subcategory of strict pairs $(U, \bar{U})$ with projective $\bar{U}$.

(iii) For any functor in (2.2.1) its source is a base for the h -topology of the target, and the induced topology on the source is the h -topology.

(iv) The functors in (2.2.2) are continuous for the h -topologies.

¹¹In fact, every V sits in a K -pair due to Nagata's theorem.

2.6. By §2.1 and §2.5, ϕ identifies h-sheaves on $\mathcal{V}ar_K$, resp. $\mathcal{V}ar_{\bar{K}}$, with h-sheaves on $\mathcal{V}ar_K^c$, $\mathcal{V}ar_K^{nc}$, $\mathcal{V}ar_K^{cc}$, $\mathcal{V}ar_K^{ss}$, resp. $\mathcal{V}ar_{\bar{K}}^c$, $\mathcal{V}ar_{\bar{K}}^{nc}$, $\mathcal{V}ar_{\bar{K}}^{cc}$, $\mathcal{V}ar_{\bar{K}}^{ss}$. Thus we have the *h-localization* functors

$$(2.6.1) \quad \mathcal{P}Sh(\mathcal{V}ar_K^?) \rightarrow \mathcal{V}ar_{\bar{K}h}, \quad \mathcal{P}Sh(\mathcal{V}ar_{\bar{K}}^?) \rightarrow \mathcal{V}ar_{\bar{K}h}$$

which assign to any presheaf $\mathcal{F}$ on pairs the corresponding h-sheaf $\mathcal{F}^\sim$ viewed as an h-sheaf on varieties.

Remark. For any presheaf on $\mathcal{V}ar_K^c$, $\mathcal{V}ar_K^{cc}$ or $\mathcal{V}ar_{\bar{K}}^{cc}$, its h-sheafification coincides with h-sheafification of its restriction to resp. $\mathcal{V}ar_K^{nc}$, $\mathcal{V}ar_K^{ss}$ or $\mathcal{V}ar_{\bar{K}}^{ss}$. For a presheaf on $\mathcal{V}ar_{\bar{K}/K}^{ss}$, its h-sheafification is the same as h-sheafification of its restriction to $\mathcal{V}ar_{\bar{K}/K'}^{ss}$, where $K' \subset \bar{K}$ is any finite extension of K (see Remark in §2.2).

3. THE p -ADIC PERIOD MAP

3.1. *The derived de Rham algebra in logarithmic setting.* We refer to [K1] for log scheme basics. There are two (in general, nonequivalent) ways to define the cotangent complex for log schemes due, respectively, to Gabber and Olsson; see [Ol].¹² Gabber's approach ([Ol, §8]) is more direct and precise;¹³ we recall it briefly.

For a commutative ring A , a prelog structure on A is a homomorphism of monoids $\alpha : L \rightarrow A$, where L is a commutative integral monoid (written multiplicatively) and A is viewed as a monoid with respect to the product. Rings equipped with prelog structures form a category in an evident way; denote its objects simply by (A, L) . For a fixed (A, L) , let $\mathcal{C}_{(A,L)}$ be the category of morphisms $(A, L) \rightarrow (B, M)$; we denote such an object by $(B, M)/(A, L)$. Let $\Omega_{(B,M)/(A,L)}$ be the B -module of relative Kähler log differentials: it is generated by $\Omega_{B/A}$ and elements $d \log m$, $m \in M$, subject to relations $d \log(m_1 m_2) = d \log m_1 + d \log m_2$, $\alpha(m) d \log m = d\alpha(m)$, and $d \log m = 0$ if m is in the image of L . The de Rham dg algebra of relative log forms $\Omega_{(B,M)/(A,L)}^i$ has components $\Omega_{(B,M)/(A,L)}^i := \Lambda_B^i \Omega_{(B,M)/(A,L)}$; elements $d \log m$ are degree 1 cycles. It carries the Hodge filtration $F^n = \Omega_{(B,M)/(A,L)}^{\geq n}$.

A pair of sets I, J yields a free object $P_{(A,L)}[I, J]$ in $\mathcal{C}_{(A,L)}$: the corresponding ring is a polynomial algebra $A[t_i, t_j]_{i \in I, j \in J}$, the monoid is $L \oplus \mathbb{N}[I]$, where $\mathbb{N}[I]$ is the free monoid generated by I , and the structure map sends the generator m_i of $\mathbb{N}[I]$, $i \in I$, to t_i . The de Rham algebra $\Omega_{P_{(A,L)}[I,J]/(A,L)}$ is freely generated, as a graded commutative A -algebra, by elements t_i, t_j of degree 0 and $d \log t_i := d \log m_i, dt_j$ of degree 1, where $i \in I, j \in J$.

Every $(B, M)/(A, L) \in \mathcal{C}_{(A,L)}$ admits a canonical simplicial resolution $P. = P_{(A,L)}(B, M)$. This is a simplicial object of $\mathcal{C}_{(A,L)}$ augmented over the object $(B, M)/(A, L)$ and such that every P_i is a free object as above. Thus we have the simplicial dg algebra $\Omega_{P./ (A,L)}$ filtered by the Hodge filtration F . Denote by $L\Omega_{(B,M)/(A,L)}$ the corresponding total complex, $L\Omega_{(B,M)/(A,L)}^a = \bigoplus_{j-i=a} \Omega_{P_i/(A,L)}^j$; this is a filtered commutative dg algebra. Let $L\Omega_{(\hat{B},M)/(A,L)}$ be its F -completion; as in §1.2, we understand it as a mere projective system of quotients $L\Omega_{(\hat{B},M)/(A,L)}/F^n$. One has a natural quasi-isomorphism of graded dg algebras $\mathrm{gr}_F^* L\Omega_{(\hat{B},M)/(A,L)} \xrightarrow{\sim} (L\Lambda_B^*(L_{(B,M)/(A,L)}))[-*]$. Here $L_{(B,M)/(A,L)} := \Omega_{P./ (A,L)} \otimes_P B$ is the relative

¹²In all situations that we will consider, the two versions coincide by [Ol, 8.34].

¹³It produces a true complex, while Olsson's construction yields a mere compatible datum of the canonical filtration truncations.

log cotangent complex; it is acyclic in positive degrees, and $H^0 L_{(B,M)/(A,L)} = \Omega_{(B,M)/(A,L)}$. The constructions are compatible with direct limits. If in the above definition we replace P by any free simplicial resolution of $(B, M)/(A, L)$, then the output is naturally quasi-isomorphic to $L\Omega_{(\hat{B}, M)/(A, L)}$. The plain cotangent complex and derived de Rham algebra for B/A map naturally to logarithmic ones.

For any map $(X, \mathcal{M}) \rightarrow (S, \mathcal{L})$ of integral log schemes, the above construction, being étale sheafified, yields the log cotangent complex $L_{(X, \mathcal{M})/(S, \mathcal{L})}$, the derived log de Rham algebra $L\Omega_{(X, \mathcal{M})/(S, \mathcal{L})}$, and its F -completion $L\Omega_{(\hat{X}, \mathcal{M})/(S, \mathcal{L})}$, which are complexes of sheaves on $X_{\text{ét}}$. We use only the completed complex $L\Omega^\wedge$.

3.2. Let $(U, \bar{U})$ be a pair as in §2.2. We view $\bar{U}$ as a log scheme with the usual integral log structure $\mathcal{O}_{\bar{U}} \cap j_* \mathcal{O}_U^\times \rightarrow \mathcal{O}_{\bar{U}}$; by abuse of notation, let us denote this log scheme again by $(U, \bar{U})$. Any morphism of pairs $(U, \bar{U}) \rightarrow (V, \bar{V})$ is a morphism of log schemes, so we have the relative log cotangent complex $L_{(U, \bar{U})/(V, \bar{V})}$, the derived log de Rham algebra $L\Omega_{(\hat{U}, \bar{U})/(V, \bar{V})}$, etc., as above. There is a canonical morphism $L\Omega_{\mathcal{U}/\bar{V}} \rightarrow L\Omega_{(\hat{U}, \bar{U})/(V, \bar{V})}$. We also have “absolute” complexes: in the arithmetic K - or $\bar{K}$ -setting, these are $L_{(U, \bar{U})} := L_{(U, \bar{U})/O_K}$, $L\Omega_{(\hat{U}, \bar{U})} := L\Omega_{(\hat{U}, \bar{U})/O_K}$, where O_K is considered with the trivial log structure $O_K^\times$; for the geometric K - or $\bar{K}$ -setting, replace O_K by K , resp. $\bar{K}$.

Remark. For $(V, \bar{V}) \in \text{Var}_{\bar{K}}^{\text{nc}}$ one has $L\Omega_{(\hat{V}, \bar{V})} \xrightarrow{\sim} \Omega_{(V, \bar{V})}$. Hence for $(U, \bar{U}) \in \text{Var}_{\bar{K}}^{\text{ss}}$ one has $R\Gamma(\bar{U}, L\Omega_{(\hat{U}, \bar{U})}) \otimes \mathbb{Q} \xrightarrow{\sim} R\Gamma(\bar{U}_{\bar{K}}, \Omega_{(U, \bar{U}_{\bar{K}})})$. Ditto for pairs over K .

Consider now the arithmetic $\bar{K}$ -pair $\text{Spec}(\bar{K}, O_{\bar{K}}) := (\text{Spec } \bar{K}, \text{Spec } O_{\bar{K}})$:

Lemma. *The cotangent complex $L_{(\bar{K}, O_{\bar{K}})}$ is acyclic in nonzero degrees, and the canonical map $\Omega_{O_{\bar{K}}} \rightarrow \Omega_{(\bar{K}, O_{\bar{K}})} := H^0 L_{(\bar{K}, O_{\bar{K}})}$ is an isomorphism. Therefore the canonical map $A_{dR} := L\Omega_{\hat{O}_{\bar{K}}} \rightarrow L\Omega_{(\hat{\bar{K}}, O_{\bar{K}})}$ is a filtered quasi-isomorphism.*

Proof. For a finite extension K' of K consider the log scheme $\text{Spec}(K', O_{K'}) := (\text{Spec } K', \text{Spec } O_{K'})$. It is a log complete intersection over O_K (see [Ol], 6.8). If π is a generator of $O_{K'}/O_K$, $f(t)$ its minimal polynomial, then, by [Ol, 6.9], $L_{(K', O_{K'})}$ is quasi-isomorphic to the cone of the multiplication by $f'(\pi)$ map $O_{K'} \rightarrow O_{K'} \subset \mathfrak{m}_{K'}^{-1}$. Thus $L_{(K', O_{K'})}$ is acyclic in nonzero degrees, $\Omega_{(K', O_{K'})} := H^0 L_{(K', O_{K'})}$ is a cyclic $O_{K'}$ -module, and the canonical map $\Omega_{O_{K'}} \rightarrow \Omega_{(K', O_{K'})}$ is an embedding with cokernel isomorphic to the residue field $O_{K'}/\mathfrak{m}_{K'}$. Now pass to the inductive limit, and use the fact that $\Omega_{O_{\bar{K}}}$ is p -divisible (see §1.3). $\square$

3.3. Consider the presheaf $(U, \bar{U}) \mapsto R\Gamma_{dR}^\natural(U, \bar{U}) := R\Gamma(\bar{U}, L\Omega_{(\hat{U}, \bar{U})})$ of filtered E_∞ dg O_K -algebras on $\text{Var}_{\bar{K}}^{\text{ss}}$. Denote by $\mathcal{A}_{dR}^\natural$ its h -sheafification (2.6.1); this is an h -sheaf of filtered E_∞ O_K -algebras on $\text{Var}_{\bar{K}}$ (as above, we see it as the projective system of quotients modulo F^i). Since $A_{dR} = \mathcal{A}_{dR}^\natural(\text{Spec } \bar{K})$ by Lemma in §3.2, A_{dR} , viewed as a constant filtered h -sheaf, maps into $\mathcal{A}_{dR}^\natural$.

Theorem (p -adic Poincaré lemma). *The maps $A_{dR} \otimes^L \mathbb{Z}/p^n \rightarrow \mathcal{A}_{dR}^\natural \otimes^L \mathbb{Z}/p^n$ are filtered quasi-isomorphisms of h -sheaves on $\text{Var}_{\bar{K}}$.*

For a proof, see §4. Assuming it, let us define the p -adic period map ρ .

3.4. The Hodge-Deligne filtration. For this subsection, K is any field of characteristic 0. Consider the presheaf $(V, \bar{V}) \mapsto R\Gamma_{\mathrm{dR}}(V, \bar{V}) := R\Gamma(\bar{V}, \Omega_{(\bar{V}, \bar{V})})$ of filtered E_∞ dg K -algebras on $\mathcal{V}ar_K^{\mathrm{nc}}$. Let $\mathcal{A}_{\mathrm{dR}}$ be its h-sheafification (2.6.1), which is an h-sheaf of filtered E_∞ K -algebras on $\mathcal{V}ar_K$ (viewed as the projective system of quotients modulo F^i). For any $X \in \mathcal{V}ar_K$ set

$$(3.4.1) \quad R\Gamma_{\mathrm{dR}}(X) := R\Gamma(X_{\mathrm{h}}, \mathcal{A}_{\mathrm{dR}}).$$

This is Deligne's de Rham complex of X equipped with Deligne's Hodge filtration.

Proposition. (i) For $(V, \bar{V}) \in \mathcal{V}ar_K^{\mathrm{nc}}$ the canonical map $R\Gamma_{\mathrm{dR}}(V, \bar{V}) \rightarrow R\Gamma_{\mathrm{dR}}(V)$ is a filtered quasi-isomorphism.

(ii) The differential of $R\Gamma_{\mathrm{dR}}(X)$ is strictly compatible with the filtration. $H_{\mathrm{dR}}^i(X) := H^i R\Gamma_{\mathrm{dR}}(X)$ are K -vector spaces of dimension equal to $\dim H_{\mathrm{et}}^i(X_{\bar{K}}, \mathbb{Q}_p)$.

(iii) For any smooth variety X there is a canonical (nonfiltered) quasi-isomorphism $R\Gamma(X, \Omega_X) \xrightarrow{\sim} R\Gamma_{\mathrm{dR}}(X)$.

Proof. By Lefschetz's principle, we can assume that $K = \mathbb{C}$. For $(V, \bar{V}) \in \mathcal{V}ar_{\mathbb{C}}^{\mathrm{nc}}$ the maps $R\Gamma_{\mathrm{dR}}(V, \bar{V}) \rightarrow R\Gamma_{\mathrm{dR}}(V, \Omega_{\bar{V}}) \rightarrow R\Gamma(V_{\mathrm{cl}}, \mathbb{C})$ are quasi-isomorphisms by [Gr]. Thus for any h-hypercovering¹⁴ $(Y, \bar{Y})/X$ of X in $\mathcal{V}ar_K^{\mathrm{nc}}$ the cohomological descent (see Remark in §2.4) yields a canonical quasi-isomorphism $R\Gamma(\bar{Y}, \Omega_{(\bar{Y}, \bar{Y})}) \xrightarrow{\sim} R\Gamma(X_{\mathrm{cl}}, \mathbb{C})$. If we equip $R\Gamma(X_{\mathrm{cl}}, \mathbb{C})$ with the Hodge-Deligne filtration of mixed Hodge theory [D], then this is a filtered quasi-isomorphism. Therefore we have a canonical filtered quasi-isomorphism $R\Gamma_{\mathrm{dR}}(X) \xrightarrow{\sim} R\Gamma(X_{\mathrm{cl}}, \mathbb{C})$. Now (i) and the second assertion of (ii) are clear; the first assertion of (ii) follows from mixed Hodge theory. The quasi-isomorphism in (iii) is $R\Gamma(X, \Omega_X) \xrightarrow{\sim} R\Gamma(Y, \Omega_Y) \xleftarrow{\sim} R\Gamma(\bar{Y}, \Omega_{(\bar{Y}, \bar{Y})})$, where the arrows are quasi-isomorphisms by the cohomological descent (since $R\Gamma(X, \Omega_X) \xrightarrow{\sim} R\Gamma(X_{\mathrm{cl}}, \mathbb{C})$). $\square$

3.5. We return to the setting of §3.3, so K is our p -adic field. Let X be any variety over $\bar{K}$. It yields a filtered E_∞ O_K -algebra

$$(3.5.1) \quad R\Gamma_{\mathrm{dR}}^{\natural}(X) := R\Gamma(X_{\mathrm{h}}, \mathcal{A}_{\mathrm{dR}}^{\natural}).$$

Since $\mathcal{A}_{\mathrm{dR}} \otimes \mathbb{Q} = \bar{K}$ (see Remark (i) in §1.5), $R\Gamma_{\mathrm{dR}}^{\natural}(X) \otimes \mathbb{Q}$ is a $\bar{K}$ -algebra. By Remark in §3.2, we have a filtered quasi-isomorphism of E_∞ $\bar{K}$ -algebras

$$(3.5.2) \quad R\Gamma_{\mathrm{dR}}^{\natural}(X) \otimes \mathbb{Q} \xrightarrow{\sim} R\Gamma_{\mathrm{dR}}(X).$$

Let us compute $R\Gamma_{\mathrm{dR}}^{\natural}(X) \hat{\otimes} \mathbb{Z}_p$. Consider the morphisms of filtered complexes $R\Gamma(X_{\mathrm{et}}, \mathbb{Z}) \otimes^L \mathcal{A}_{\mathrm{dR}} \xrightarrow{\sim} R\Gamma(X_{\mathrm{et}}, \mathcal{A}_{\mathrm{dR}}) \rightarrow R\Gamma(X_{\mathrm{h}}, \mathcal{A}_{\mathrm{dR}}) \rightarrow R\Gamma(X_{\mathrm{h}}, \mathcal{A}_{\mathrm{dR}}^{\natural}) = R\Gamma_{\mathrm{dR}}^{\natural}(X)$. After applying $\cdot \otimes^L \mathbb{Z}/p^n$, the arrows become filtered quasi-isomorphisms (the first one by Remark in §2.4, the second one by the Poincaré lemma in §3.3), so we get a filtered quasi-isomorphism $R\Gamma(X_{\mathrm{et}}, \mathbb{Z}/p^n) \otimes^L \mathcal{A}_{\mathrm{dR}} \xrightarrow{\sim} R\Gamma_{\mathrm{dR}}^{\natural}(X) \otimes^L \mathbb{Z}/p^n$. Since $R\Gamma_{\mathrm{et}}(X, \mathbb{Z}_p) = \mathrm{holim} R\Gamma(X_{\mathrm{et}}, \mathbb{Z}/p^n) = R\Gamma(X_{\mathrm{et}}, \mathbb{Z}) \hat{\otimes} \mathbb{Z}_p$ (see §1.1) is a perfect $\mathbb{Z}_p$ -complex and $R\Gamma(X_{\mathrm{et}}, \mathbb{Z}/p^n) = R\Gamma_{\mathrm{et}}(X, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p}^L \mathbb{Z}/p^n$, one has, passing to the homotopy limit as in §1.1, $R\Gamma_{\mathrm{et}}(X, \mathbb{Z}_p) \otimes_{\mathbb{Z}_p}^L (\mathcal{A}_{\mathrm{dR}} \hat{\otimes} \mathbb{Z}_p) \xrightarrow{\sim} R\Gamma_{\mathrm{dR}}^{\natural}(X) \hat{\otimes} \mathbb{Z}_p$. Tensoring by $\mathbb{Q}$, we get a filtered quasi-isomorphism of filtered E_∞ B_{dR}^+ -algebras (see (1.5.1))

$$(3.5.3) \quad \beta : R\Gamma_{\mathrm{et}}(X, \mathbb{Q}_p) \otimes B_{\mathrm{dR}}^+ \xrightarrow{\sim} R\Gamma_{\mathrm{dR}}^{\natural}(X) \hat{\otimes} \mathbb{Q}_p.$$

¹⁴Here we view X as an h-sheaf on $\mathcal{V}ar_K^{\mathrm{nc}}$, so $(Y, \bar{Y})$ is a simplicial object of $\mathcal{V}ar_K^{\mathrm{nc}}$ equipped with an augmentation map $Y \rightarrow X$ that makes Y an h-hypercovering of X .

Let $\alpha : R\Gamma_{\mathrm{dR}}(X) \otimes_{\bar{K}} B_{\mathrm{dR}}^+ \rightarrow R\Gamma_{\mathrm{dR}}^{\natural}(X) \widehat{\otimes} \mathbb{Q}_p$ be the B_{dR}^+ -linear extension of the composition $R\Gamma_{\mathrm{dR}}(X) \xrightarrow{\sim} R\Gamma_{\mathrm{dR}}^{\natural}(X) \otimes \mathbb{Q} \rightarrow R\Gamma_{\mathrm{dR}}^{\natural}(X) \widehat{\otimes} \mathbb{Q}_p$, where the first arrow is inverse to (3.5.2) and the second one comes from the canonical map $? \rightarrow ? \widehat{\otimes} \mathbb{Z}_p$. We get a morphism of filtered E_{∞} B_{dR}^+ -algebras

$$(3.5.4) \quad \rho = \rho_{\mathrm{dR}} := \beta^{-1} \alpha : R\Gamma_{\mathrm{dR}}(X) \otimes_{\bar{K}} B_{\mathrm{dR}}^+ \rightarrow R\Gamma_{\mathrm{ét}}(X, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} B_{\mathrm{dR}}^+.$$

Remarks. (i) The Galois group $\mathrm{Gal}(\bar{K}/K)$ acts on $\mathrm{Var}_{\bar{K}\mathfrak{h}}$ and on both sides of (3.5.4) by transport of structure, and ρ^+ is evidently compatible with this action. In particular, if X is defined over K , i.e., $X = X_K \otimes_K \bar{K}$, then $R\Gamma_{\mathrm{dR}}(X) = R\Gamma_{\mathrm{dR}}(X_K) \otimes_K \bar{K}$, and we can rewrite (3.5.4) as a $\mathrm{Gal}(\bar{K}/K)$ -equivariant morphism

$$(3.5.5) \quad \rho : R\Gamma_{\mathrm{dR}}(X_K) \otimes_K B_{\mathrm{dR}}^+ \rightarrow R\Gamma_{\mathrm{ét}}(X, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} B_{\mathrm{dR}}^+.$$

(ii) The map ρ does not change if we replace K by any of its finite extensions that are contained in $\bar{K}$ (see Remark in §2.6).

3.6. Theorem. *The B_{dR} -linear extension of ρ is a filtered quasi-isomorphism: for any $X \in \mathrm{Var}_{\bar{K}}$ one has*

$$(3.6.1) \quad \rho : R\Gamma_{\mathrm{dR}}(X) \otimes_{\bar{K}} B_{\mathrm{dR}} \xrightarrow{\sim} R\Gamma_{\mathrm{ét}}(X, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} B_{\mathrm{dR}}.$$

Proof. (a) *The case of $X = \mathbb{G}_m = \mathbb{G}_{m\bar{K}}$:* The $\bar{K}$ -line $H_{\mathrm{dR}}^1(\mathbb{G}_m) = \mathrm{gr}_F^1 H_{\mathrm{dR}}^1(\mathbb{G}_m)$ is generated by $d \log t$. The $\mathbb{Z}_p$ -line $H_{\mathrm{ét}}^1(\mathbb{G}_m, \mathbb{Z}_p)(1) = H_{\mathrm{ét}}^1(\mathbb{G}_m, \mathbb{Z}_p(1))$ is generated by the class $cl(\mathfrak{k})$ of the Kummer $\mathbb{Z}_p(1)$ -torsor $\mathfrak{k} = \varprojlim \mathfrak{k}_n$, $\mathfrak{k}_n := (t^{1/p^n})$. Due to the canonical identification $\mathbb{C}_p(1) \xrightarrow{\sim} \mathfrak{m}_{\mathrm{dR}}/\mathfrak{m}_{\mathrm{dR}}^2 = \mathrm{gr}_F^1 B_{\mathrm{dR}}$, see §1.4, §1.5, we can view $cl(\mathfrak{k})$ as a generator of the $\mathbb{C}_p$ -line $H_{\mathrm{ét}}^1(\mathbb{G}_m, \mathbb{Q}_p) \otimes \mathrm{gr}_F^1 B_{\mathrm{dR}}$.

Lemma. *One has $\mathrm{gr}_F^1(\rho)(d \log t) = cl(\mathfrak{k}) \in H_{\mathrm{ét}}^1(\mathbb{G}_m, \mathbb{Q}_p) \otimes \mathrm{gr}_F^1 B_{\mathrm{dR}}$.*

Proof of Lemma. We make a mod p^n computation. Consider the ss-pair $(\mathbb{G}_m, \bar{\mathbb{G}}_m)$, $\bar{\mathbb{G}}_m := \mathbb{P}_{O_{\bar{K}}}^1$. One has $\mathrm{gr}_F^1 L\Omega_{(\mathbb{G}_m, \bar{\mathbb{G}}_m)}^1 = \Omega_{(\mathbb{G}_m, \bar{\mathbb{G}}_m)}^1[-1]$, so $d \log t \in \Gamma(\mathbb{G}_m, \Omega_{(\mathbb{G}_m, \bar{\mathbb{G}}_m)}^1)$ is a 1-cocycle in $\mathrm{gr}_F^1 R\Gamma_{\mathrm{dR}}^{\natural}(\mathbb{G}_m)$. As in §1.1, set $C_n := \mathrm{Cone}(p^n : \mathbb{Z} \rightarrow \mathbb{Z})$. Let $d \log cl(\mathfrak{k}_n)$ be the image of the class $cl(\mathfrak{k}_n)$ of $\mathfrak{k}_n$ by the composition $H^1(\mathbb{G}_m, \mu_{p^n}) \rightarrow H^1(\mathbb{G}_m, \mathrm{gr}_F^1 A_{\mathrm{dR}} \otimes C_n) \rightarrow \mathrm{gr}_F^1 R\Gamma_{\mathrm{dR}}^{\natural}(\mathbb{G}_m) \otimes C_n$, where the first arrow comes from the coefficient maps $\mu_{p^n} \xrightarrow{d \log} \Omega_{O_{\bar{K}} p^n} \hookrightarrow \Omega_{O_{\bar{K}}}[-1] \otimes C_n = \mathrm{gr}_F^1 A_{\mathrm{dR}} \otimes C_n$. To prove the lemma, we will show that the image of $d \log t$ by the embedding $\mathrm{gr}_F^1 R\Gamma_{\mathrm{dR}}^{\natural}(\mathbb{G}_m) \hookrightarrow \mathrm{gr}_F^1 R\Gamma_{\mathrm{dR}}^{\natural}(\mathbb{G}_m) \otimes C_n$ is homologous to $d \log cl(\mathfrak{k}_n)$.

Let $\mathbb{G}_{\tilde{m}}$ be a copy of $\mathbb{G}_m$ with parameter $\tilde{t}$, and $\pi : \mathbb{G}_{\tilde{m}} \rightarrow \mathbb{G}_m$ be the projection $t = \tilde{t}^{p^n}$. Thus $\mathbb{G}_{\tilde{m}}/\mathbb{G}_m$ is our μ_{p^n} -torsor $\mathfrak{k}_n$, so $cl(\mathfrak{k}_n)$ is represented by a Čech μ_{p^n} -cocycle $c(\mathfrak{k}_n)$ for the étale covering $\mathbb{G}_{\tilde{m}}/\mathbb{G}_m$. The corresponding Čech hypercovering is the twist of $\mathbb{G}_{\tilde{m}}$ by the universal μ_{p^n} -torsor $\mathfrak{k}_n$ over the classifying simplicial space $B_{\mu_{p^n}}$, so for any sheaf $\mathcal{F}$ the Čech complex of $\mathbb{G}_{\tilde{m}}/\mathbb{G}_m$ with coefficients in $\mathcal{F}$ is the cochain complex $C^*(\mu_{p^n}, \Gamma(\mathbb{G}_{\tilde{m}}, \mathcal{F}))$ for μ_{p^n} acting on sections by the translations. The 1-cocycle $c(\mathfrak{k}_n)$ is the identity map $\mu_{p^n} \rightarrow \mu_{p^n} = \Gamma(\mathbb{G}_{\tilde{m}}, \mu_{p^n})$.

Our π extends to the h-covering of semi-stable pairs $(\mathbb{G}_{\tilde{m}}, \bar{\mathbb{G}}_{\tilde{m}}) \rightarrow (\mathbb{G}_m, \bar{\mathbb{G}}_m)$, and the Čech hypercovering extends to a hypercovering in $\mathrm{Var}_{\bar{K}}^{\mathrm{ss}}$ which is the $\mathfrak{k}_n$ -twist of $(\mathbb{G}_{\tilde{m}}, \bar{\mathbb{G}}_{\tilde{m}})$. So one has a canonical map $C^*(\mu_{p^n}, \Gamma(\mathbb{G}_{\tilde{m}}, \Omega_{(\mathbb{G}_{\tilde{m}}, \bar{\mathbb{G}}_{\tilde{m}})}^1))[-1] \rightarrow \mathrm{gr}_F^1 R\Gamma_{\mathrm{dR}}^{\natural}(\mathbb{G}_m)$; hence $C^*(\mu_{p^n}, \Gamma(\mathbb{G}_{\tilde{m}}, \Omega_{(\mathbb{G}_{\tilde{m}}, \bar{\mathbb{G}}_{\tilde{m}})}^1))[-1] \otimes C_n \rightarrow \mathrm{gr}_F^1 R\Gamma_{\mathrm{dR}}^{\natural}(\mathbb{G}_m) \otimes C_n$. Both $d \log t$ and $d \log c(\mathfrak{k}_n)$ are 1-cocycles in $C^*(\mu_{p^n}, \Gamma(\mathbb{G}_{\tilde{m}}, \Omega_{(\mathbb{G}_{\tilde{m}}, \bar{\mathbb{G}}_{\tilde{m}})}^1))[-1] \otimes C_n$: namely, $d \log t \in C^0(\mu_{p^n}, \Gamma(\mathbb{G}_{\tilde{m}}, \Omega_{(\mathbb{G}_{\tilde{m}}, \bar{\mathbb{G}}_{\tilde{m}})}^1))[-1]$ and $d \log c(\mathfrak{k}_n) \in C^1(\mu_{p^n}, \Omega_{O_{\bar{K}} p^n})$.

$\subset C^1(\mu_{p^n}, \Gamma(\tilde{\mathbb{G}}_m, \Omega^1_{(\tilde{\mathbb{G}}_m, \tilde{\mathbb{G}}_m)})[-1] \otimes C_n)$. Their difference is the differential of the 0-cochain $d \log t \in C^0(\mu_{p^n}, \Gamma(\tilde{\mathbb{G}}_m, \Omega^1_{(\tilde{\mathbb{G}}_m, \tilde{\mathbb{G}}_m)})) \subset C^0(\mu_{p^n}, \Gamma(\tilde{\mathbb{G}}_m, \Omega^1_{(\tilde{\mathbb{G}}_m, \tilde{\mathbb{G}}_m)})[-1] \otimes C_n)$, q.e.d. $\square$

We see that for $X = \mathbb{G}_m$ the map ρ of (3.6.1) is a filtered quasi-isomorphism. It provides a canonical generator $\rho(d \log t)/cl(\mathfrak{k})$ of $\mathfrak{m}_{\mathrm{dR}}(-1)$. Thus we have a canonical identification between $H_{\mathrm{\acute{e}t}}^i(X, \mathbb{Q}_p(n)) \otimes B_{\mathrm{dR}}$ and $H_{\mathrm{\acute{e}t}}^i(X, \mathbb{Q}_p) \otimes B_{\mathrm{dR}}$ with the filtration shifted by n , etc.

Remark. The above generator is equal to Fontaine's generator from [F3, 1.5.4]. Indeed, they coincide modulo $\mathfrak{m}_{\mathrm{dR}}^2(-1)$ by the lemma, and both are $\mathrm{Gal}(\bar{K}/K)$ -invariant (see Remark in §3.5). Since $H^0(\mathrm{Gal}(\bar{K}/K), \mathfrak{m}_{\mathrm{dR}}^2(-1)) = 0$, we are done.

(b) *Compatibility with the Gysin maps:* Let $i : Y \hookrightarrow X$ be a closed codimension 1 embedding of smooth varieties. It yields the Gysin isomorphisms $i_{*\mathrm{dR}} : R\Gamma_{\mathrm{dR}}(Y) \xrightarrow{\sim} R\Gamma_{\mathrm{dR} Y}(X)(1)[2] := \mathrm{Cone}(R\Gamma_{\mathrm{dR}}(X) \rightarrow \Gamma_{\mathrm{dR}}(X \setminus Y))(1)[1]$, $i_{*\mathbb{Q}_p} : R\Gamma_{\mathrm{\acute{e}t}}(Y, \mathbb{Q}_p) \xrightarrow{\sim} R\Gamma_{\mathrm{\acute{e}t} Y}(X, \mathbb{Q}_p)(1)[2]$. Let us show that ρ commutes with the Gysin maps.

Consider the deformation to the normal cone diagram

$$(3.6.2) \quad \begin{array}{ccccc} \mathcal{L} & \hookrightarrow & X_{\mathbb{A}^1} & \hookleftarrow & X \\ \uparrow & & \uparrow & & \uparrow \\ Y & \hookrightarrow & Y_{\mathbb{A}^1} & \hookleftarrow & Y. \end{array}$$

Here $Y_{\mathbb{A}^1} = Y \times \mathbb{A}^1$, $X_{\mathbb{A}^1}$ is $X \times \mathbb{A}^1$ with $Y \times \{0\}$ blown up, the left arrow is the zero section of the normal bundle $\mathcal{L}$ to Y in X , and the bottom embeddings are $y \mapsto (y, 0), (y, 1)$. It yields a commutative diagram of the de Rham cohomology

$$(3.6.3) \quad \begin{array}{ccccc} R\Gamma_{\mathrm{dR} Y}(\mathcal{L})(1)[2] & \leftarrow & R\Gamma_{\mathrm{dR} Y_{\mathbb{A}^1}}(X_{\mathbb{A}^1})(1)[2] & \rightarrow & R\Gamma_{\mathrm{dR} Y}(X)(1)[2] \\ \uparrow & & \uparrow & & \uparrow \\ R\Gamma_{\mathrm{dR}}(Y) & \leftarrow & R\Gamma_{\mathrm{dR}}(Y_{\mathbb{A}^1}) & \rightarrow & R\Gamma_{\mathrm{dR}}(Y), \end{array}$$

where the vertical arrows are the Gysin isomorphisms and the horizontal ones are pullbacks. There is a similar diagram for the $\mathbb{Q}_p$ -cohomology. The horizontal maps are filtered quasi-isomorphisms, so, since ρ is compatible with pullbacks, we see that the Gysin compatibility for $Y \hookrightarrow X$ amounts to one for $Y \hookrightarrow \mathcal{L}$.

So we can assume that X is a line bundle $\mathcal{L}$ over Y and i its zero section. Now the source of both i_* 's are dg algebras, the targets are modules over them (due to the projection $\mathcal{L} \rightarrow Y$), and i_* 's are morphisms of modules. Thus it suffices to check that ρ identifies the images of 1. The assertion is local with respect to Y ; hence we can assume that $\mathcal{L}$ is trivial. By base change, we reduced to the case when Y is a point, where we are done by (a).

(c) *The case of a smooth projective X :* Let us check that the morphism of bigraded rings $\mathrm{gr}_F^* \rho^* : \mathrm{gr}_F^* H_{\mathrm{dR}}^*(X) \otimes_{\bar{K}} \mathrm{gr}_F^* B_{\mathrm{dR}} \rightarrow H_{\mathrm{\acute{e}t}}^*(X, \mathbb{Q}_p) \otimes_{\mathbb{Q}_p} \mathrm{gr}_F^* B_{\mathrm{dR}}$ is an isomorphism. It is an isomorphism for $*$ = 0. By (b), $\mathrm{gr}_F^1 \rho^2$ identifies the classes c of a hyperplane section. Since the product with $c^{\dim X}$ identifies H^0 and $H^{2 \dim X}$, $\mathrm{gr}_F^2 \rho^2$ is an isomorphism. Therefore, since $\mathrm{gr}_F^* \rho^*$ is compatible with the Poincaré pairing for classes of opposite degrees and the latter is nondegenerate, $\mathrm{gr}_F^* \rho^*$ is injective. Since $\dim_{\bar{K}} H_{\mathrm{dR}}^*(X) = \dim_{\mathbb{Q}_p} H_{\mathrm{\acute{e}t}}^*(X, \mathbb{Q}_p)$, we are done.

(d) *The case of $X = \bar{X} \setminus D$, where $\bar{X}$ is smooth projective, D is a strict normal crossings divisor:* Let Y be an irreducible component of D , D' be the union of the other components; set $X' := \bar{X} \setminus D'$, $Y' := Y \setminus D'$. By induction by the number

of components D (starting with (c)), we can assume that the theorem holds for X' and Y' . By (b), ρ provides a morphism between the exact Gysin triangles for (Y', X') . It is a filtered quasi-isomorphism on the X' and Y' terms; hence it is a filtered quasi-isomorphism on the X term, q.e.d.

(e) *The case of arbitrary X :* If Y/X is any h-hypercovering of X , then the canonical map $R\Gamma_{\text{dR}}(X) \rightarrow R\Gamma_{\text{dR}}(Y)$ (which is the total complex of the cosimplicial system of filtered complexes $R\Gamma_{\text{dR}}(Y_i)$) is a filtered quasi-isomorphism by the construction of $R\Gamma_{\text{dR}}$, and $R\Gamma_{\text{ét}}(X, \mathbb{Q}_p) \xrightarrow{\sim} R\Gamma_{\text{ét}}(Y, \mathbb{Q}_p)$ by cohomological descent (see Remark in 2.4). Thus if ρ is a filtered quasi-isomorphism for every Y_i , then it is a filtered quasi-isomorphism for X . We are done, since, by de Jong (or Hironaka), one can find Y/X with Y_i as in (d). $\square$

Remark. ρ is compatible with Chern classes of vector bundles: Indeed, $c_i(E)$ are determined in the usual way by $c_1(\mathcal{O}(1)_{\mathbb{P}(E)})$, so it suffices to show that ρ identifies c_1 's of line bundles. Notice that the construction of ρ extends tautologically to simplicial schemes. By (a) above, ρ identifies the de Rham and étale Chern classes of the universal line bundle over the classifying simplicial scheme $B_{\mathbb{G}_m}$. For a line bundle $\mathcal{L}$ on X , choose a finite open covering $\{U_i\}$ of X such that $\mathcal{L}$ is trivial on U_i ; let $\pi : X^\sim \rightarrow X$ be the Čech hypercovering. Since π yields an isomorphism between the cohomology, it suffices to check that ρ identifies the Chern classes of $\pi^*\mathcal{L}$. This is true since $\pi^*\mathcal{L}$ is the pullback of the universal line bundle by a map $X^\sim \rightarrow B_{\mathbb{G}_m}$.

4. PROOF OF THE POINCARÉ LEMMA

4.1. Pick any $(V, \bar{V}) \in \text{Var}_{\bar{K}}^{\text{ss}}$.

Proposition. *One has $L_{(V, \bar{V})} \xrightarrow{\sim} \Omega_{(V, \bar{V})}$, the $\mathcal{O}_{\bar{V}}$ -module $\Omega_{(V, \bar{V})} := \Omega_{(V, \bar{V})}/(\bar{K}, \mathcal{O}_{\bar{K}})$ is locally free of finite rank, and there is a canonical short exact sequence*

$$(4.1.1) \quad 0 \rightarrow \mathcal{O}_{\bar{V}} \otimes_{\mathcal{O}_{\bar{K}}} \Omega_{\mathcal{O}_{\bar{K}}} \rightarrow \Omega_{(V, \bar{V})} \rightarrow \Omega_{(V, \bar{V})} \rightarrow 0.$$

Proof. We can assume that V is connected, so $(V, \bar{V})$ is the base change of a semi-stable K -pair $(U, \bar{U})$ as in 2.2(c), i.e., $(V, \bar{V}) = (U_{\bar{K}}, \bar{U}_{\mathcal{O}_{\bar{K}}})$. For any finite extension K' of K_U , consider an arithmetic K -pair $(U_{K'}, \bar{U}_{\mathcal{O}_{K'}}) := (U \otimes_{K_U} K', \bar{U} \otimes_{\mathcal{O}_{K_U}} \mathcal{O}_{K'})$. Set $\Omega_{(U, \bar{U})} := \Omega_{(U, \bar{U})}/(K_U, \mathcal{O}_{K_U})$, $\Omega_{(U_{K'}, \bar{U}_{\mathcal{O}_{K'}})} := \Omega_{(U_{K'}, \bar{U}_{\mathcal{O}_{K'}})}/(K', \mathcal{O}_{K'})$.

Lemma. *The log scheme $(U_{K'}, \bar{U}_{\mathcal{O}_{K'}})$ coincides with the pullback of $(U, \bar{U})$ by the map $\text{Spec}(K', \mathcal{O}_{K'}) \rightarrow \text{Spec}(K_U, \mathcal{O}_{K_U})$ in the category of log schemes.*

Assume the lemma for a moment. The map $(U, \bar{U}) \rightarrow \text{Spec}(\mathcal{O}_{K_U}, K_U)$ is log smooth and integral; by the lemma, $(U_{K'}, \bar{U}_{\mathcal{O}_{K'}}) \rightarrow \text{Spec}(\mathcal{O}_{K'}, K')$ enjoys the same properties. So, by [Ol, 8.34], $L_{(U_{K'}, \bar{U}_{\mathcal{O}_{K'}})}/(K', \mathcal{O}_{K'}) \xrightarrow{\sim} \Omega_{(U_{K'}, \bar{U}_{\mathcal{O}_{K'}})} = \mathcal{O}_{K'} \otimes_{\mathcal{O}_{K_U}} \Omega_{(U, \bar{U})}$, which is a locally free $\mathcal{O}_{\bar{U}_{\mathcal{O}_{K'}}}$ -module of finite rank. Since $L_{(K', \mathcal{O}_{K'})} \xrightarrow{\sim} \Omega_{(K', \mathcal{O}_{K'})}$ (see the proof of Lemma in §3.2), the canonical exact triangle ([Ol, 8.18]) $\mathcal{O}_{\bar{U}_{\mathcal{O}_{K'}}} \otimes_{\mathcal{O}_{K'}} L_{(K', \mathcal{O}_{K'})} \rightarrow L_{(U_{K'}, \bar{U}_{\mathcal{O}_{K'}})} \rightarrow L_{(U_{K'}, \bar{U}_{\mathcal{O}_{K'}})}/(K', \mathcal{O}_{K'})$ reduces to the short exact sequence $0 \rightarrow \mathcal{O}_{\bar{U}_{\mathcal{O}_{K'}}} \otimes_{\mathcal{O}_{K'}} \Omega_{(K', \mathcal{O}_{K'})} \rightarrow \Omega_{(U_{K'}, \bar{U}_{\mathcal{O}_{K'}})} \rightarrow \Omega_{(U_{K'}, \bar{U}_{\mathcal{O}_{K'}})} \rightarrow 0$. Pass to the limit by all $K' \subset \bar{K}$ and use the lemma in §3.2; we are done. $\square$

Proof of Lemma. The underlying scheme of the pullback log scheme is $\bar{U}_{\mathcal{O}_{K'}}$. Let us show that its log structure map $\mathcal{M} \rightarrow \mathcal{O}_{\bar{U}_{\mathcal{O}_{K'}}} \cap j_* \mathcal{O}_{U_{K'}}^\times$ is an isomorphism. The assertion is étale local, so we can assume that $\bar{U}$ is étale over $\text{Spec } \mathcal{O}_{K_U}[t_a, t_b, t_c]/$

$(\Pi t_a - \pi_{K_U})$, where a, b, c are in finite sets A, B, C , π_{K_U} is a uniformizing parameter in $\mathcal{O}_{K_U}$, and U is the subscheme where all t_a, t_b are invertible. The log structure of $(U, \bar{U})$ is fine with a chart $\mathbb{N}[A \sqcup B] \rightarrow \mathcal{O}_{\bar{U}}$, which sends generators m_a, m_b of $\mathbb{N}[A \sqcup B]$ to t_a, t_b . Therefore¹⁵ $\bar{U}_{\mathcal{O}_{K'}}$ is étale over $\text{Spec } \mathcal{O}_{K'}[t_a, t_b, t_c]/(\Pi t_a - \pi_{K'}^e)$, where e is the ramification index of K'/K_U , $\pi_{K'}$ is a uniformizing parameter in $\mathcal{O}_{K'}$, and the log structure $\mathcal{M}$ has a chart $M_{A,B} \rightarrow \mathcal{O}_{\bar{U}_{\mathcal{O}_{K'}}}$, where $M_{A,B}$ is the quotient of $\mathbb{N}[A \sqcup B] \oplus \mathbb{N}$ modulo the relation $\Pi m_a = m_\pi^e$ (m_π is the generator of the last summand $\mathbb{N}$), the chart is $m_a, m_b, m_\pi \mapsto t_a, t_b, \pi_{K'}$. Consider an embedding $M_{A,B} \hookrightarrow M_{A,B}^w := e^{-1}\mathbb{N}[A] \oplus \mathbb{N}[B]$, $m_a, m_b, m_\pi \mapsto m_a, m_b, \Pi m_a^{1/e}$. Its image is formed by those $\Pi m_a^{n_a/e} \Pi m_b^{n_b}$, $n_a, n_b \in \mathbb{N}$, such that $n_a - n_{a'} \in e\mathbb{Z}$ for any $a, a' \in A$; thus $M_{A,B}$ is saturated. Now the log scheme $(\bar{U}_{\mathcal{O}_{K'}}, \mathcal{M})$ is evidently log regular in the sense of [K2, 2.1]; hence $\mathcal{M} \xrightarrow{\sim} \mathcal{O}_{\bar{U}_{\mathcal{O}_{K'}}} \cap j_* \mathcal{O}_{U_{K'}}^\times$ by [K2, 11.6], q.e.d.¹⁶

The reference to [K2] can be replaced by the next explicit argument: It suffices to show that the map of sheaves $\mathcal{M}/\mathcal{O}_{\bar{U}_{\mathcal{O}_{K'}}}^\times \rightarrow (\mathcal{O}_{\bar{U}_{\mathcal{O}_{K'}}} \cap j_* \mathcal{O}_{U_{K'}}^\times)/\mathcal{O}_{\bar{U}_{\mathcal{O}_{K'}}}^\times$ is an isomorphism. The r.h.s. is the sheaf $\mathcal{D}$ of effective Cartier divisors supported on $\bar{U}_{\mathcal{O}_{K'}} \setminus U_{K'}$. Let $\mathcal{D}^w \supset \mathcal{D}$ be the sheaf of the corresponding effective Weil divisors. For $x \in \bar{U}_{\mathcal{O}_{K'}}$, the fiber $(\mathcal{M}/\mathcal{O}_{\bar{U}_{\mathcal{O}_{K'}}}^\times)_x$ is the quotient M_{A_x, B_x} of $M_{A,B}$, where $A_x \subset A$, $B_x \subset B$ consist of those a, b such that t_a, t_b vanish at x . The map $M_{A_x, B_x} \rightarrow \mathcal{D}_x$ extends to an isomorphism $M_{A_x, B_x}^w \xrightarrow{\sim} \mathcal{D}_x^w$, which identifies a generator $m_a^{1/e}$ with the *reduced* divisor D_a of t_a , m_b with $D_b := \text{div}(t_b)$. Thus $M_{A_x, B_x} \hookrightarrow \mathcal{D}_x$. To show that $\hookrightarrow$ is an isomorphism, we need to check that if $D = \sum n_a D_a + \sum n_b D_b$ is a Cartier divisor at x , then $n_a - n_{a'} \in e\mathbb{Z}$ for any $a, a' \in A_x$. We can assume that $A = \{a, a'\}$, $B = C = \emptyset$, so $\bar{U}_{\mathcal{O}_{K'}}$ is a semi-stable curve over $\mathcal{O}_{K'}$. The exceptional divisor of its minimal desingularization $\tilde{U}_{\mathcal{O}_{K'}}$ is a chain of $e-1$ projective lines $P_1, \dots, P_{e-1}$ with self-intersection indices $(P_i, P_i) = -2$. Let $\tilde{D} = n_a D_a^b + n_1 P_1 + \dots + n_{e-1} P_{e-1} + n_{a'} D_{a'}^b$ be the pullback of D to $\tilde{U}_{\mathcal{O}_{K'}}$; here $D_a^b, D_{a'}^b$ are strict transforms of $D_a, D_{a'}$. One has $(\tilde{D}, P_i) = 0$, i.e., $n_{i-1} - 2n_i + n_{i+1} = 0$ or $n_i - n_{i-1} = n_{i+1} - n_i$, where $n_0 := n_a, n_e := n_{a'}$. Thus $n_{a'} - n_a = e(n_1 - n_a) \in e\mathbb{Z}$, and we are done. $\square$

4.2. Set $\Omega_{\langle V, \bar{V} \rangle}^a := \Lambda_{\mathcal{O}_{\bar{V}}}^a \Omega_{\langle V, \bar{V} \rangle}$. Consider (4.1.1) as a 2-step filtration on $\Omega_{\langle V, \bar{V} \rangle}$; it splits locally since $\Omega_{\langle V, \bar{V} \rangle}$ is locally free. Passing to derived exterior powers, we get for any m a finite increasing filtration I on $\text{gr}_F^m L\Omega_{\langle \hat{V}, \bar{V} \rangle} = (L\Lambda_{\mathcal{O}_{\bar{V}}}^m \Omega_{\langle V, \bar{V} \rangle})[-m]$ with $\text{gr}_a^I \text{gr}_F^m L\Omega_{\langle \hat{V}, \bar{V} \rangle} = \Omega_{\langle V, \bar{V} \rangle}^a \otimes_{\mathcal{O}_{\bar{K}}} \text{gr}_F^{m-a} \mathbf{A}_{\text{dR}}[-a]$, hence on $\text{gr}_F^m R\Gamma_{\text{dR}}^{\mathfrak{h}}(V, \bar{V})$ with

$$(4.2.1) \quad \text{gr}_a^I \text{gr}_F^m R\Gamma_{\text{dR}}^{\mathfrak{h}}(V, \bar{V}) = R\Gamma(\bar{V}, \Omega_{\langle V, \bar{V} \rangle}^a) \otimes_{\mathcal{O}_{\bar{K}}}^L \text{gr}_F^{m-a} \mathbf{A}_{\text{dR}}[-a].$$

Let $\mathcal{G}^a$ be the h-sheafification (see (2.6.1)) of the complex of presheaves $(V, \bar{V}) \mapsto R\Gamma(\bar{V}, \Omega_{\langle V, \bar{V} \rangle}^a)$ on $\mathcal{V}ar_{\bar{K}}^{\text{ss}}$. This is a complex of h-sheaves of $\mathcal{O}_{\bar{K}}$ -modules on $\mathcal{V}ar_{\bar{K}}$. Its cohomology $H^b \mathcal{G}^a$ is h-sheafification of the presheaf $(V, \bar{V}) \mapsto H^b(\bar{V}, \Omega_{\langle V, \bar{V} \rangle}^a)$ on $\mathcal{V}ar_{\bar{K}}^{\text{ss}}$. Our I is a filtration on the presheaf $(V, \bar{V}) \mapsto \text{gr}_F^m R\Gamma_{\text{dR}}^{\mathfrak{h}}(V, \bar{V})$; passing to h-sheafification, we get a finite filtration I on $\text{gr}_F^m \mathcal{A}_{\text{dR}}^{\mathfrak{h}}$ with span $[0, m]$ and

$$(4.2.2) \quad \text{gr}_a^I \text{gr}_F^m \mathcal{A}_{\text{dR}}^{\mathfrak{h}} = \mathcal{G}^a \otimes_{\mathcal{O}_{\bar{K}}}^L \text{gr}_F^{m-a} \mathbf{A}_{\text{dR}}[-a].$$

¹⁵We replace one $t_a \in \mathcal{O}(\bar{U}_{\mathcal{O}_{K'}})$ by $t_a \pi_{K'}^e / \pi_{K_U}$.

¹⁶I am grateful to Luc Illusie for the proof.

Notice that the bottom cohomology H^0 of the bottom term $I_0 = \mathrm{gr}_0^I$ is the constant sheaf $\mathcal{O}_{\bar{K}}$ and $\mathrm{Cone}(\mathrm{gr}_F^m \mathcal{A}_{\mathrm{dR}} \rightarrow \mathrm{gr}_F^m \mathcal{A}_{\mathrm{dR}}^\natural) = \mathrm{gr}_F^m \mathcal{A}_{\mathrm{dR}}^\natural / H^0 I_0$. Therefore, by (4.2.2), the Poincaré lemma follows from the next assertion:

Theorem. *The cohomology $H^b \mathcal{G}^a$ are h -sheaves of $\mathbb{Q}$ - (hence $\bar{K}$ -) vector spaces for $(a, b) \neq (0, 0)$.*

Remark. The p -divisibility of $H^b \mathcal{G}^0$, $b \neq 0$, was first proved by Bhatt [Bh1, 8.0.1].

Exercise. Consider a presheaf $(V, \bar{V}) \mapsto R\Gamma(\bar{V}, L\Omega_{(\hat{V}, \bar{V})/(\bar{K}, \mathcal{O}_{\bar{K}})})$; let $\mathcal{A}_{\mathrm{dR}}^{\mathrm{naive}}$ be its h -sheafification. One has an evident map $\mathrm{Cone}(F^1 \mathcal{A}_{\mathrm{dR}} \rightarrow \mathcal{A}_{\mathrm{dR}}^\natural) \rightarrow \mathcal{A}_{\mathrm{dR}}^{\mathrm{naive}}$. Show that the theorem implies that it is a filtered quasi-isomorphism; i.e., the triangle $F^1 \mathcal{A}_{\mathrm{dR}} \rightarrow \mathcal{A}_{\mathrm{dR}}^\natural \rightarrow \mathcal{A}_{\mathrm{dR}}^{\mathrm{naive}}$ is exact in the filtered derived category of h -sheaves.

4.3. We deduce the above theorem from a more concrete assertion. As in §4.1, for an ss-pair $(U, \bar{U})$ over K we have the locally free $\mathcal{O}_{\bar{U}}$ -module of log differentials $\Omega_{\langle U, \bar{U} \rangle} := \Omega_{(U, \bar{U})/(K_U, \mathcal{O}_{K_U})}$ and its exterior powers $\Omega_{\langle U, \bar{U} \rangle}^a := \Lambda^a \Omega_{\langle U, \bar{U} \rangle}$.

Let $f : (U', \bar{U}') \rightarrow (U, \bar{U})$ be a map in $\mathcal{V}ar_K^{\mathrm{ss}}$ or $\mathcal{V}ar_{\bar{K}}^{\mathrm{ss}}$. We say that f is (*Hodge*) p -negligible if the morphisms $(\tau_{>0} R\Gamma(\bar{U}, \mathcal{O}_{\bar{U}})) \otimes^L \mathbb{Z}/p \rightarrow (\tau_{>0} R\Gamma(\bar{U}', \mathcal{O}_{\bar{U}'})) \otimes^L \mathbb{Z}/p$ and $R\Gamma(\bar{U}, \Omega_{\langle U, \bar{U} \rangle}^a) \otimes^L \mathbb{Z}/p \rightarrow R\Gamma(\bar{U}', \Omega_{\langle U', \bar{U}' \rangle}^a) \otimes^L \mathbb{Z}/p$, $a > 0$, in $D^b(\mathcal{O}_{K_U}/p)$, resp. $D^b(\mathcal{O}_{\bar{K}}/p)$, vanish.

Remark. For $(U, \bar{U}) \in \mathcal{V}ar_K^{\mathrm{ss}}$ and a point $K_U \rightarrow \bar{K}$, one has $R\Gamma(\bar{U}_{\mathcal{O}_{\bar{K}}}, \Omega_{\langle U_{\bar{K}}, \bar{U}_{\mathcal{O}_{\bar{K}}} \rangle}^a) = R\Gamma(\bar{U}, \Omega_{\langle U, \bar{U} \rangle}^a) \otimes_{\mathcal{O}_{\bar{K}}}^L \mathcal{O}_{\bar{K}}$. Therefore the base change functor $\mathcal{V}ar_K^{\mathrm{ss}} \rightarrow \mathcal{V}ar_{\bar{K}}^{\mathrm{ss}}$ (see (2.2.2)) preserves p -negligible maps.

Theorem. *Every $U \in \mathcal{V}ar_K^{\mathrm{ss}}$ admits a p -negligible h -covering. Ditto for $\bar{K}$ -pairs.*

The theorem implies the one in §4.2: Indeed, the $\bar{K}$ -assertion shows that one has $(\tau_{>0} \mathcal{G}^0) \otimes^L \mathbb{Z}/p = 0$ and $\mathcal{G}^a \otimes^L \mathbb{Z}/p = 0$ for $a > 0$; since for a complex $\mathcal{G}$ the multiplication by p on $H^* \mathcal{G}$ is invertible if and only if $\mathcal{G} \otimes^L \mathbb{Z}/p = 0$, we are done. Thus it yields the Poincaré lemma.

The above remark shows that the K -version of the theorem implies the $\bar{K}$ -one. The proof of the K -version takes the rest of the section.

4.4. For the rest of §4, “pair” means “arithmetic K -pair” (see §2.2). We need further input from de Jong. A morphism $f : (C, \bar{C}) \rightarrow (S, \bar{S})$ of pairs is said to be a *family of pointed curves* (over $(S, \bar{S})$) if the map $\bar{C}_S := f^{-1}(S) \rightarrow S$ is smooth of relative dimension 1 with irreducible geometric fibers, and $D_{fS} := \bar{C}_S \setminus C$, viewed as a reduced scheme, is étale over S . Such an f is *semi-stable* if, in addition, $\bar{C}/\bar{S}$ is a semi-stable family of curves, and the closure D_f of D_{fS} in $\bar{C}$ (the *horizontal divisor*), viewed as a reduced scheme, is étale over $\bar{S}$ and intersects each fiber of f at smooth points. A section $e : (S, \bar{S}) \rightarrow (C, \bar{C})$ of f is said to be *nice* if $e(\bar{S})$ intersects fibers of f at smooth points and $D_f \cap e(\bar{S}) = \emptyset$. Families of pointed curves over $(S, \bar{S})$ form a category $\mathcal{C}_{(S, \bar{S})}$ in the obvious manner, and a morphism of bases $\psi : (S', \bar{S}') \rightarrow (S, \bar{S})$ yields an evident pullback functor $\mathcal{C}_{(S, \bar{S})} \rightarrow \mathcal{C}_{(S', \bar{S}')}$ which preserves semi-stable families. A morphism $f' \rightarrow f$ in $\mathcal{C}_{(S, \bar{S})}$ is called an *alteration* if $(C', \bar{C}')$ is an alteration of $(C, \bar{C})$; it is a *semi-stable alteration* (of f) if, in addition, f' is semi-stable.

Theorem. (a) *Any family $f : (C, \bar{C}) \rightarrow (S, \bar{S})$ of pointed curves with $f : \bar{C} \rightarrow \bar{S}$ projective admits a semi-stable alteration f' h -locally over $(S, \bar{S})$.*

(b) One can find f' as above which has a nice section e . Moreover, for a given closed subscheme $P \subset \bar{C}$ such that $f(P) = \bar{S}$ and $P \cap \bar{C}_S \subset C$, one can find e such that the map $\bar{C}' \rightarrow \bar{C}$ sends $e(\bar{S})$ to P .

(c) For any semi-stable family of pointed curves $f : (C, \bar{C}) \rightarrow (S, \bar{S})$ with $(S, \bar{S})$ a strict ss-pair, there exists a semi-stable alteration $m : (C, \tilde{C}) \rightarrow (C, \bar{C})$ of f with $m|_C = \text{id}_C$ such that $m : \tilde{C} \rightarrow \bar{C}$ is an isomorphism over smooth points of f and $(C, \tilde{C})$ is an ss-pair.

Proof. (c) is [dJ1, 3.6]. (a) follows from [dJ2, 2.4 (i),(ii)] except that de Jong does not care to control the domain of smoothness of the semi-stable alteration of f . A miniscule modification of his argument permits us to do this; see Appendix 1. Alternatively, (a) follows directly from a far more precise result of Temkin [T, 1.5].¹⁷

Let us check (b). Every pair has a canonical alteration by the union of normalizations of its irreducible components, so we assume all the way that $\bar{S}$ is normal and irreducible. Since P as in (b) exists h -locally on $(S, \bar{S})$,¹⁸ we can assume it is given. Replacing $(S, \bar{S})$ by its alteration (P_S, P) , we get a section e of f with image in P . Set $C^b := C \setminus e(S)$. Then $(C^b, \bar{C}) \rightarrow (S, \bar{S})$ is a family of pointed curves; let $f^b : (C^b, \bar{C}) \rightarrow (S, \bar{S})$ be its semi-stable alteration as in (a). Let D_e be the closure in $\bar{C}'$ of the preimage D_{eS} of $e(S)$. Then D_e is an étale covering of $\bar{S}$.¹⁹ Let C' be the preimage of $C \subset \bar{C}$ in $\bar{C}'$; then $(C', \bar{C}') \rightarrow (S, \bar{S})$ is a semi-stable alteration of $(C, \bar{C}) \rightarrow (S, \bar{S})$. Replacing $(S, \bar{S})$ by its alteration (D_{eS}, D_e) , we get a nice section of $(C', \bar{C}')$ which sits over e , hence over P . $\square$

Remark. In (c), every nice section of $(C, \bar{C})$ lifts to a nice section of $(C, \tilde{C})$.

Corollary. Any pair $(U, \bar{U})$ has an h -covering by ss-pairs $(C, \bar{C})$, $\dim C = \dim U$, for which there is a semi-stable family of pointed curves $f : (C, \bar{C}) \rightarrow (S, \bar{S})$ with a nice section such that $(S, \bar{S})$ is an ss-pair and C is affine over S (i.e., $f(D_f) = \bar{S}$).

Proof. It suffices to find an h -covering of $(U, \bar{U})$ by pairs $(C, \bar{C})$ with $\dim C = \dim U$ for which there exists a family of pointed curves $f : (C, \bar{C}) \rightarrow (S, \bar{S})$ with C affine over S and projective $\bar{S}, \bar{C}$. The theorem transforms it then, with an input from Remark (i) in §2.5 to preserve the dimension and de Jong's theorem in §2.3 to alter $(S, \bar{S})$ from (b) into a strict ss-pair, into a datum with all promised properties.

By de Jong's theorem in §2.3, we can assume that $(U, \bar{U})$ is an ss-pair and $\bar{U}$ is projective and irreducible;²⁰ set $d = \dim U$. Pick any closed point $u \in U$. It suffices to find an open neighborhood $U' \subset U$ of U , an alteration $(C, \bar{C})$ of $(U', \bar{U})$, and a family of curves $f : (C, \bar{C}) \rightarrow (S, \bar{S})$ such that $f(D_f) = \bar{S}$.

Embed $\bar{U}$ into a projective space $\mathbb{P}_{O_K}^N$. By Bertini, there is a plane $H \subset \mathbb{P}^N$ defined over K of codimension d such that $u \notin H$, H intersects $\bar{U}_K$ transversally, $H \cap \bar{U}_K \subset U$, and the codimension $d-1$ plane which contains H and u is transversal to $\bar{U}_K$ and $\bar{U}_K \setminus U$. Let $m : \bar{C} \rightarrow \bar{U}$ be the blowup at $\bar{U} \cap H_{O_K}$, $p : \bar{C} \rightarrow \mathbb{P}_{O_K}^{d-1}$ be the projection defined by H , and $\bar{C} \xrightarrow{f} \bar{S} \rightarrow \mathbb{P}_{O_K}^{d-1}$ be the Stein factorization of p (so $\bar{S} = \text{Spec } p_* \mathcal{O}_{\bar{C}}$). Let $D \subset \bar{C}$ be the union of $m^{-1}(\bar{U} \setminus U)$ and the exceptional divisor (viewed as a reduced scheme), and $S \subset \bar{S}_K$ be the maximal open subset

¹⁷To bring our datum to Temkin's setting, one flattens $\bar{C}/\bar{S}$ and $D_f/\bar{S}$ using [RG, 5.2.2] and replaces $\bar{S}$ by its normalization.

¹⁸Indeed, one can cover S by open subsets $\{S_\alpha\}$ such that P exists for $(C_{S_\alpha}, \bar{C}) \rightarrow (S_\alpha, \bar{S})$.

¹⁹Since D_{f^b} is étale over $\bar{S}$, D_{eS} is open and closed in $D_{f^b S}$, $f^b(D_{eS}) = S$, and $\bar{S}$ is normal.

²⁰We only need that $\bar{U}$ is projective and normal, and that U is smooth.

over which f is smooth and $f|_D$ is étale. Set $C := f^{-1}(S) \setminus D$ and $U' := m(C)$; notice that $m|_C : C \xrightarrow{\sim} U'$. Then $U', (C, \bar{C}), f$ satisfy the promised properties (one has $f(D_f) = \bar{S}$ since D_f contains the exceptional divisor), q.e.d. $\square$

4.5. Let us return to the proof of the theorem in §4.3. We use induction by $\dim U$. By the corollary in §4.4, we can replace $(U, \bar{U})$ by $(C, \bar{C})$ as in loc. cit., so we have $f : (C, \bar{C}) \rightarrow (S, \bar{S})$ with a nice section e and C affine over S . Notice that $(C, \bar{C})$ is log smooth over $(S, \bar{S})$ and the line bundle $\omega_f := \Omega_{(C, \bar{C})/(S, \bar{S})}$ equals $f^!(\mathcal{O}_{\bar{S}})[-1] \otimes \mathcal{O}_{\bar{C}}(D_f)$.

Key lemma. *h-locally over $(S, \bar{S})$, one can find a semi-stable alteration $\phi : f' \rightarrow f$ together with a nice section e' that lifts e such that $(C', \bar{C}')$ is an ss-pair and the pullback maps $\phi^* : R^1 f_* \mathcal{O}_{\bar{C}} \rightarrow R^1 f'_* \mathcal{O}_{\bar{C}'}, f_* \omega_f \rightarrow f'_* \omega_{f'}$ are divisible by p .²¹*

For a proof, see §4.6. Assuming it for the moment, let us finish the proof of the theorem in §4.3. By Remark (i) in §2.5, we can assume that the h-localization of $(S, \bar{S})$ in the Key Lemma does not change $\dim S$. We will show that for some h-covering $(S', \bar{S}')$ of $(S, \bar{S})$ the composition $(C', \bar{C}')_{(S', \bar{S}')} \rightarrow (C', \bar{C}') \xrightarrow{\phi} (C, \bar{C})$ is p -negligible.

For any a consider the exact sequence

$$(4.5.1) \quad 0 \rightarrow f^* \Omega_{(S, \bar{S})}^a \rightarrow \Omega_{(C, \bar{C})}^a \rightarrow (f^* \Omega_{(S, \bar{S})}^{a-1}) \otimes \omega_f \rightarrow 0.$$

The section e splits off $\Omega_{(S, \bar{S})}^a \hookrightarrow Rf_* \Omega_{(C, \bar{C})}^a$ as a direct summand whose complement is $\text{Cone}(\partial_C)$, where $\partial_C : \Omega_{(S, \bar{S})}^{a-1} \otimes f_* \omega_f \rightarrow \Omega_{(S, \bar{S})}^a \otimes R^1 f_* \mathcal{O}_{\bar{C}}$ is the boundary map for (4.5.1) (one has $R^1 f_* \omega_f = 0$ since $f(D_f) = \bar{S}$). There is a similar splitting in case of f' provided by e' , and the map $\phi^* : Rf_* \Omega_{(C, \bar{C})}^a \rightarrow Rf'_* \Omega_{(C', \bar{C}')}^a$ is compatible with the direct sum decompositions. Now ϕ^* is divisible by p on the second summand: Indeed, the Key Lemma asserts that the morphism of two-term complexes $\phi^* : \text{Cone}(\partial_C) \rightarrow \text{Cone}(\partial_{C'})$ is divisible by p on each term; since these are morphisms of vector bundles on $\mathcal{O}_K$ -flat $\bar{S}$, our $p^{-1}\phi^*$ is uniquely defined and commutes with the differentials. Thus the map $\phi^* \otimes \text{id}_{C_1} : \text{Cone}(\partial_C) \otimes C_1 \rightarrow \text{Cone}(\partial_{C'}) \otimes C_1$, where $C_1 := \text{Cone}(p : \mathbb{Z} \rightarrow \mathbb{Z})$, is homotopic to zero. Apply $R\Gamma(\bar{S}, \cdot)$ and use the induction assumption to treat the first summand $R\Gamma(\bar{S}, \Omega_{(S, \bar{S})}^a)$; we are done.

4.6. *Proof of Key Lemma.* Consider the relative Picard $\bar{S}$ -schemes $J := \text{Pic}^0(\bar{C}/\bar{S})$ and $J^b := \text{Pic}^0((\bar{C}, D_f)/\bar{S})$: the first scheme parametrizes line bundles $\mathcal{L}$ on $\bar{C}$ such that the restriction of $\mathcal{L}$ to the normalization of each irreducible component of any geometric fiber of f has degree 0; the second one parametrizes pairs $(\mathcal{L}, \gamma)$, where $\mathcal{L}$ is as above and γ is a trivialization of $\mathcal{L}|_{D_f}$. Since $(\bar{C}, D_f)$ is a semi-stable $\bar{S}$ -family of d -pointed curves, $d := \deg(D_f)$, our J and J^b are semi-abelian schemes (see [R]), and J^b is an extension of J by a torus $\mathbb{G}_m^{D_f}/\mathbb{G}_m$.

Over S our J^b is a generalized Jacobian; let $i : C \rightarrow J_S^b$ be the Abel-Jacobi map $i : C \rightarrow J_S^b, x \mapsto \mathcal{O}_{\bar{C}}(x - e)$. Let $C^\sim \rightarrow C$ be the i -pullback of the multiplication by p isogeny $p_{J^b} : J^b \rightarrow J^b$, and $\bar{C}^\sim \rightarrow \bar{C}$ be the normalization of $\bar{C}$ in $C^\sim$. Then $f^\sim : (C^\sim, \bar{C}^\sim) \rightarrow (S, \bar{S})$ is a family of pointed curves, which is an alteration of f . By the theorem in §4.4, h-locally over $(S, \bar{S})$ there is a semi-stable alteration f' of $f^\sim$

²¹As elements of the groups $\text{Hom}_{\mathcal{O}_{\bar{S}}} (R^1 f_* \mathcal{O}_{\bar{C}}, R^1 f'_* \mathcal{O}_{\bar{C}'}), \text{Hom}_{\mathcal{O}_{\bar{S}}} (f_* \omega_f, f'_* \omega_{f'})$.

with $(C', \bar{C}')$ semi-stable and equipped with a nice section e' which lies over e . Let us check that the alteration $\phi : f' \rightarrow f$ satisfies the conditions of the Key Lemma.

Set $J' := \text{Pic}^0(\bar{C}'/\bar{S})$ and $J^b := \text{Pic}^0((\bar{C}', D_f)/\bar{S})$. We have the pullback morphisms $\phi^* : J \rightarrow J'$, $J^b \rightarrow J'^b$ of our semi-abelian schemes; over S we have the norm maps $\phi_{*S} : J'_S \rightarrow J_S$, $J'^b_S \rightarrow J^b_S$. Both are compatible with the projections $J^b \rightarrow J$, $J'^b \rightarrow J'$.

Since $\bar{S}$ is normal, for any semi-abelian $\bar{S}$ -schemes A, B one has (see [FC, I 2.7])

$$(4.6.1) \quad \text{Hom}(A, B) \xrightarrow{\sim} \text{Hom}(A_S, B_S).$$

Thus ϕ_{*S} extends to morphisms $\phi_* : J' \rightarrow J$, $J'^b \rightarrow J^b$.

Notice that $R^1 f_* \mathcal{O}_{\bar{C}}$ is the Lie algebra of J , and, by Serre duality, $f_* \omega_f$ is dual to the Lie algebra of J^b ; the same is true for f' . Our $\phi^* : R^1 f_* \mathcal{O}_{\bar{C}} \rightarrow R^1 f'_* \mathcal{O}_{\bar{C}'}$ is the Lie algebra map for $\phi^* : J \rightarrow J'$, and $\phi^* : f_* \omega_f \rightarrow f'_* \omega_{f'}$ is the map between the duals to the Lie algebras for $\phi_* : J^b \rightarrow J'^b$ (this is true over S , hence everywhere since $\bar{S}$ is $\mathcal{O}_K$ -flat).

By construction, $\phi_{*S} : J'^b_S \rightarrow J^b_S$ factors through p_{J^b} over S ; i.e., it is divisible by p in $\text{Hom}(J'^b_S, J^b_S)$. By (4.6.1), ϕ_* is divisible by p in $\text{Hom}(J^b, J^b)$. Passing to Lie algebras, we see that $\phi^* : f_* \omega_f \rightarrow f'_* \omega_{f'}$ is divisible by p . Similarly, $\phi_{*S} : J'_S \rightarrow J_S$ is divisible by p . Notice that J_S, J'_S , being Jacobians of smooth projective curves, are self-dual abelian schemes, and $\phi^*_S : J_S \rightarrow J'_S$ is dual to ϕ_{*S} . Hence ϕ^*_S is divisible by p . So, by (4.6.1), $\phi^* : J \rightarrow J'$ is divisible by p . Passing to Lie algebras, we see that $\phi^* : R^1 f_* \mathcal{O}_{\bar{C}} \rightarrow R^1 f'_* \mathcal{O}_{\bar{C}'}$ is divisible by p , q.e.d. $\square$

APPENDIX

Below is a proof of part (a) in the theorem from §4.4. It follows closely de Jong's argument from §§2–3 of [dJ2] with a minor change of the lemma below; we refer the reader to sections of [dJ2] for details.

(i) ([dJ2, 2.10]) One can assume that $\bar{S}$ is irreducible. By [RG, 5.2.2], there is a canonical modification of $\bar{S}$, which is projective and is an isomorphism over S , such that the strict transforms of $\bar{C}$ and D_f are flat over $\bar{S}$. Passing to them, we can assume that *all fibers of f have dimension 1, of $f|_{D_f}$ have dimension 0*.

(ii) ([dJ2, 3.4–3.5]) We say that a family of pointed curves is *good* if irreducible components of all its geometric fibers are curves whose normalization has genus ≥ 2 . A good alteration is an alteration with a good source.

Lemma. *f admits a good alteration h -locally over $(S, \bar{S})$.*

Proof of Lemma. It suffices to find for any closed point s in S its open neighborhood $S_{(s)} \subset S$ and an alteration $(S'_{(s)}, \bar{S}')$ of $(S_{(s)}, \bar{S})$ such that the pullback of f to $(S'_{(s)}, \bar{S}')$ admits a good alteration. To do this, we define by induction a strictly increasing sequence of open subsets $\emptyset = V_0 \subset V_1 \subset \dots$ of $\bar{S}$ and a sequence of finite extensions $F = F_0 \subset F_1 \subset \dots$ of the field F of rational functions on $\bar{C}$ such that the normalization $\bar{C}_i$ of $\bar{C}$ in F_i has the following properties: (a) the map $\bar{C}_i \rightarrow \bar{S}$ is smooth at s , (b) the map $\pi_i : \bar{C}_i \rightarrow \bar{C}$ is étale at D_{fs} , (c) the normalizations of irreducible components of geometric fibers of $\bar{C}_i$ over points of V_i have genus ≥ 2 . There is an open neighborhood $U_i \subset S$ of s over which $\bar{C}_i$ is smooth and π_i is étale at D_f . The induction stops when $V_n = \bar{S}$; set $S_{(s)} = U_n$. The promised good alteration is $(\pi_n^{-1}(C), \bar{C}_n)$ fibered over the normalization $(S'_{(s)}, \bar{S}')$ of $(S_{(s)}, \bar{S})$ in F_n .

Let x be the closed point of the closure of s in $\bar{S}$. The induction produces simultaneously an auxiliary sequence of finite subsets $T_0 \subset T_1 \subset \dots$ of closed points of $\bar{C}_x$; it starts with $T_0 :=$ the union of D_{fx} and the set of nonregular points of $\bar{C}_x$. The induction step: suppose we have $V_{i-1}, F_{i-1}, T_{i-1}$; let us construct V_i, F_i, T_i assuming that $V_{i-1} \neq \bar{S}$. Let y be any closed point in $\bar{S} \setminus V_{i-1}$. Since $\bar{S}$ is projective, there is an affine open V which contains x and y . Let $\mathcal{L}$ be a very ample line bundle on $\bar{C}$. Replacing it by a sufficiently high power, we can assume that $\Gamma(\bar{C}_V, \mathcal{L}) \twoheadrightarrow \Gamma(\bar{C}_x, \mathcal{L}) \times \Gamma(\bar{C}_y, \mathcal{L})$. One can find a finite unramified extension²² K' of K with residue field k' and two sections $\gamma_1, \gamma_2 \in \Gamma(\bar{C}_V, \mathcal{L}) \otimes_{O_K} O_{K'}$ which do not vanish at the generic points of irreducible components of $\bar{C}_x, \bar{C}_y$, such that $t = \gamma_1/\gamma_2$ yields generically étale finite maps $t_x : \bar{C}_x \otimes k' \rightarrow \mathbb{P}_x^1 \otimes k', t_y : \bar{C}_y \otimes k' \rightarrow \mathbb{P}_y^1 \otimes k'$ étale over $\{0, 1, \infty\}$ and such that $t_x(T_{i-1}) \cap \{0, 1, \infty\} = \emptyset$. Pick $\ell \geq 5$ prime to p , and let F_i be an extension of F_{i-1} generated by $K', \mu_\ell, t^{1/\ell}$, and $(1-t)^{1/\ell}$. Let T_i be the union of T_{i-1} and the set of ramification points of t_x . The normalization $\bar{C}_i$ of $\bar{C}$ in F_i satisfies (a), (b), and satisfies (c) over some open set V_i which contains V_{i-1} and y . We are done. $\square$

(iii) It remains to show that *every good f admits a semi-stable alteration after a possible alteration of the base*. The genus of the generic fiber of f is ≥ 2 , so $(\bar{C}_S, D_{fS})$ is a stable n -pointed curve over S (where n is the degree of D_{fS} over S). The Deligne-Mumford stack of stable n -pointed curves is proper, so, after replacing $(S, \bar{S})$ by an alteration, we can assume that $(\bar{C}_S, D_{fS})$ extends to a stable n -pointed curve $(\bar{C}', D_{f'})$ over $\bar{S}$ (see [dJ2, 3.8]). We have a semi-stable family of pointed curves $f' : (C', \bar{C}') \rightarrow (S, \bar{S})$, $C' := \bar{C}' \setminus D_{f'}$. By [dJ2, 3.10], the goodness of f implies that, after a possible alteration of $\bar{S}$, the evident morphism $\bar{C}'_S \rightarrow \bar{C}_S$ extends to a morphism $f' \rightarrow f$, and we are done. $\square$

ACKNOWLEDGMENTS

This work was prompted by a discussion with Kobi Kremnizer (and watching the moon over a meadow in Oxford). The author is very grateful to him and to Bhargav Bhatt, Volodya Drinfeld, John Francis, Luc Illusie, David Kazhdan, Jacob Lurie, Madhav Nori, and Misha Temkin for explanations, questions, and corrections of mistakes.

REFERENCES

- [B] A. Beilinson, *On the crystalline period map*, math. AG 1111.3316 (2011).
- [Ber] L. Berger, *Représentations p -adiques et équations différentielles*, Inv. Math. **148** (2) (2002), 219–284. MR1906150 (2004a:14022)
- [Bh1] B. Bhatt, *Derived direct summands*, Thesis, Princeton University (2010). MR2753219
- [Bh2] B. Bhatt, *p -adic derived de Rham cohomology*, 2012.
- [Col] P. Colmez, *Les nombres algébriques sont denses dans B_{dR}^+* , Périodes p -adiques, Astérisque 223, Soc. Math. France, 1994, pp. 103–111.
- [Con] B. Conrad, *Cohomological descent*, <http://math.stanford.edu/~conrad/>.
- [D] P. Deligne, *Théorie de Hodge III*, Publ. Math. IHES **44** (1974), 5–77. MR0498552 (58:16653b)
- [dJ1] A. J. de Jong, *Smoothness, semi-stability and alterations*, Publ. Math. IHES **83** (1996), 51–93. MR1423020 (98e:14011)
- [dJ2] A. J. de Jong, *Families of curves and alterations*, Ann. Inst. Fourier **47** (2) (1997), 599–621. MR1450427 (98f:14019)

²²If the residue field of K is infinite, one can take $K' = K$.

- [Fa1] G. Faltings, *p-adic Hodge theory*, J. Amer. Math. Soc. **1** (1988), 255–288. MR924705 (89g:14008)
- [Fa2] G. Faltings, *Almost étale extensions*, Cohomologies p -adiques et applications arithmétiques (II), Astérisque 279, Soc. Math. France, 2002, pp. 185–270. MR1922831 (2003m:14031)
- [FC] G. Faltings, C.-L. Chai, *Degeneration of abelian varieties*, Springer-Verlag, 1990. MR1083353 (92d:14036)
- [Far] L. Fargues, *Letter to L. Illusie*, 2010.
- [F1] J.-M. Fontaine, *Sur certains types de représentations p -adiques du groupe de Galois d'un corps local; construction d'un anneau de Barsotti-Tate*, Ann. of Math. (2) **115** (1982), 529–577. MR657238 (84d:14010)
- [F2] J.-M. Fontaine, *Formes différentielles et modules de Tate des variétés abéliennes sur les corps locaux*, Inv. Math. **65** (1982), 379–409. MR643559 (83k:14023)
- [F3] J.-M. Fontaine, *Le corps de périodes p -adiques*, Périodes p -adiques, Astérisque 223, Soc. Math. France, 1994, pp. 59–101. MR1293971 (95k:11086)
- [F4] J.-M. Fontaine, *Représentations p -adiques semi-stables*, Périodes p -adiques, Astérisque 223, Soc. Math. France, 1994, pp. 113–184. MR1293972 (95g:14024)
- [Gr] A. Grothendieck, *On the de Rham cohomology of algebraic varieties*, Publ. Math. IHES **29** (1966), 95–103. MR0199194 (33:7343)
- [GV] A. Grothendieck, J.-L. Verdier, *Conditions de finitude. Topos et sites fibrés. Applications aux questions de passage à la limite*, Théorie des topos et cohomologie étale de schémas (SGA 4), Tome 2, Lect. Notes in Math. 270, Springer-Verlag, 1972, pp. 163–340. MR0354653 (50:7131)
- [HS] V. Hinich, V. Schechtman, *On homotopy limit of homotopy algebras*, K -theory, Arithmetic and Geometry, Yu. I. Manin (Ed.), Lect. Notes in Math. 1289, Springer-Verlag, 1987, pp. 240–264. MR923138 (89d:55052)
- [Ill1] L. Illusie, *Complexe Cotangent et Déformations I*, Lect. Notes in Math. 239, Springer-Verlag, 1971. MR0491680 (58:10886a)
- [Ill2] L. Illusie, *Complexe Cotangent et Déformations II*, Lect. Notes in Math. 283, Springer-Verlag, 1972. MR0491681 (58:10886b)
- [J] U. Jannsen, *On the ℓ -adic cohomology of varieties over number fields and its Galois cohomology*, Galois groups over $\mathbb{Q}$, Y. Ihara, K. Ribet, J.-P. Serre (Eds.), MSRI Publ. 16, Springer-Verlag, 1989, pp. 315–360. MR1012170 (90i:11064)
- [K1] K. Kato, *Logarithmic structures of Fontaine-Illusie*, Algebraic Analysis, Geometry, and Number Theory, The Johns Hopkins University Press, 1988, pp. 191–224. MR1463703 (99b:14020)
- [K2] K. Kato, *Toric singularities*, American Jour. of Math. **116** (5) (1994), 1073–1099. MR1296725 (95g:14056)
- [N1] W. Niziol, *Semistable conjecture via K -theory*, Duke Math. J. **141** (1) (2008), 151–178. MR2372150 (2008m:14035)
- [N2] W. Niziol, *p -adic motivic cohomology in arithmetic*, International Congress of Mathematicians, vol. II (2006), 459–472. MR2275605 (2007k:14029)
- [N3] W. Niziol, *On uniqueness of p -adic period morphisms*, Pure Appl. Math. Q. **5** (1) (2009), 163–212. MR2520458 (2011c:14058)
- [Ol] M. Olsson, *The logarithmic cotangent complex*, Math. Ann. **333** (2005), 859–931. MR2195148 (2006j:14017)
- [R] M. Raynaud, *Spécialisation du foncteur de Picard*, Publ. Math. IHES **38** (1970), 27–76. MR0282993 (44:227)
- [RG] M. Raynaud, L. Gruson, *Critères de platitude et de projectivité. Techniques de “platification” d’un module*, Inv. Math. **13** (1971), 1–89. MR0308104 (46:7219)
- [SD] B. Saint-Donat, *Téchniques de descente cohomologique*, Théorie des topos et cohomologie étale de schémas (SGA 4), Tome 2, Lect. Notes in Math. 270, Springer-Verlag, 1972, pp. 83–162. MR0354653 (50:7131)
- [SV] A. Suslin, V. Voevodsky, *Singular homology of abstract algebraic varieties*, Inv. Math. **123** (1996), 61–94. MR1376246 (97e:14030)
- [T] M. Temkin, *Stable modification of relative curves*, J. Algebraic Geom. **19** (4) (2010), 603–677. MR2669727 (2011j:14064)
- [Ts1] T. Tsuji, *p -adic étale cohomology and crystalline cohomology in the semi-stable reduction case*, Inv. Math. **137** (1999), 233–411. MR1705837 (2000m:14024)

- [Ts2] T. Tsuji, *Semi-stable conjecture of Fontaine-Jannsen: a survey*, Cohomologies p -adiques et applications arithmétiques (II), Astérisque 279, Soc. Math. France, 2002, pp. 323–370. MR1922833 (2003g:14028)
- [V1] J.-L. Verdier, *Topologies et faisceaux*, Théorie des topos et cohomologie étale de schémas (SGA 4), Tome 1, Lect. Notes in Math. 269, Springer-Verlag, 1972, pp. 219–264.
- [V2] J.-L. Verdier, *Fonctorialité de catégories de faisceaux*, Théorie des topos et cohomologie étale de schémas (SGA 4), Tome 1, Lect. Notes in Math. 269, Springer-Verlag, 1972, pp. 265–298.
- [V3] J.-L. Verdier, *Cohomologie dans les topos*, Théorie des topos et cohomologie étale de schémas (SGA 4), Tome 2, Lect. Notes in Math. 270, Springer-Verlag, 1972, pp. 1–82.
- [Y] G. Yamashita, *Théorie de Hodge p -adique pour les variétés ouvertes*, C. R. A. S. **349** (21–22) (2011), 1127–1130.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, ILLINOIS 60637
E-mail address: `sasha@math.uchicago.edu`