

## ON THE INTRODUCTION OF THE NOTION OF HYPERBOLIC FUNCTIONS.\*

BY PROFESSOR M. W. HASKELL.

THE difficulties in the way of a satisfactory geometrical deduction of the fundamental formulæ of the hyperbolic functions seem to be due to the lack of a definition of these functions which shall be independent of the particular position of the argument area. A general definition of this kind can, however, readily be found in terms of the ratios of certain *areas*, instead of *lines*. From this definition the addition-theorem and other characteristics can be easily deduced by the methods of analytic geometry; and the definitions hold, furthermore, not merely for the rectangular, but for *any* hyperbola.

I. *The circular functions.* In order to bring out clearly the analogy with the circular functions, I will first indicate briefly how the latter would be defined according to this method.

In a circle of radius  $a$  (Fig. 1) let  $\phi$  be the angle between the radii  $OP$  and  $OQ$ , and let  $OP'$  be drawn perpendicular to




FIG. 1.

\* Read before the AMERICAN MATHEMATICAL SOCIETY, December 28, 1894. For various geometrical definitions of these functions, see Professor A. Macfarlane's paper: "On the definition of the trigonometric functions," 1894.—EDITORS.

$OP$ . The following areas are either well known or easily found (the sign  $\Delta$  denoting triangle):

$$\begin{aligned} \text{sector } OPQ &= \frac{1}{2}a^2\phi, & \Delta OPQ &= \frac{1}{2}a^2 \sin \phi, \\ \Delta OQP' &= \frac{1}{2}a^2 \cos \phi, & \Delta OPP' &= \frac{1}{2}a^2, \end{aligned}$$

from which follow immediately:

$$\phi = \frac{\text{sector } OPQ}{\Delta OPP'}, \quad \sin \phi = \frac{\Delta OPQ}{\Delta OPP'}, \quad \cos \phi = \frac{\Delta OQP'}{\Delta OPP'}.$$

Similarly, if the tangent at  $Q$  meet  $OP$  in  $T$  and  $OP'$  in  $T'$ , it is not difficult to show that

$$\begin{aligned} \tan \phi &= \frac{\Delta OTQ}{\Delta OPP'}, & \text{ctn } \phi &= \frac{\Delta OQT'}{\Delta OPP'}, \\ \sec \phi &= \frac{\Delta OTP'}{\Delta OPP'}, & \text{csc } \phi &= \frac{\Delta OPT'}{\Delta OPP'}, \end{aligned}$$

Now, these formulæ might be taken as *definitions* of the argument  $\phi$  and of its various functions. They can then be immediately extended to any ellipse, the only modification necessary being that  $OP$  and  $OP'$  shall be conjugate semi-diameters. The area of the triangle  $OPP'$  is then  $= \frac{1}{2}ab$ , and  $\phi$  is equal to the difference between the eccentric angles  $P$  and  $Q$ .

II. *Definition of the hyperbolic functions.* Since of two conjugate diameters only one meets the hyperbola in real points, the conjugate hyperbola must be employed also, and  $P'$  is the point where the diameter conjugate to  $OP$  meets the conjugate hyperbola. We shall then define the argument  $u$  and its functions in strict analogy with the preceding results, as follows (see Fig. 2):

$$u = \frac{\text{sector } OPQ}{\Delta OPP'}, \quad \sinh u = \frac{\Delta OPQ}{\Delta OPP'}, \quad \cosh u = \frac{\Delta OQP'}{\Delta OPP'},$$

etc.

If now the hyperbola be referred to its principal axes as axes of coördinates, its equation may be written

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1. \quad (1)$$

Let  $x_1, y_1$  be the coördinates of  $P$ , and  $x_2, y_2$  those of  $Q$ . Then the coördinates of  $P'$  will be  $\frac{ay_1}{b}, \frac{bx_1}{a}$ , and the area of the triangle  $OPP'$  is equal to  $\frac{1}{2}ab$ . The definitions of  $\sinh u$  and  $\cosh u$  become

$$\sinh u = \frac{x_1y_2 - x_2y_1}{ab}, \quad \cosh u = \frac{x_1x_2}{a^2} - \frac{y_1y_2}{b^2}. \quad (2)$$

Interchanging the coördinates of  $P$  and  $Q$ , we have

$$\sinh(-u) = -\sinh u, \quad \cosh(-u) = \cosh u. \quad (3)$$

Also, if  $P$  and  $Q$  coincide,  $u = 0$  and

$$\sinh 0 = 0, \quad \cosh 0 = 1. \quad (4)$$




FIG. 2.

Let  $u_1 = \frac{\text{sector } OAP}{\Delta OPP'}$  and  $u_2 = \frac{\text{sector } OAQ}{\Delta OPP'}$ , so that  $u = u_2 - u_1$ . The coördinates of  $A$  being  $a, 0$ , we have

$$\sinh u_1 = \frac{y_1}{b}, \quad \cosh u_1 = \frac{x_1}{a}; \quad \sinh u_2 = \frac{y_2}{b}, \quad \cosh u_2 = \frac{x_2}{a}. \quad (5)$$

Comparing with (1), we see at a glance that

$$\cosh^2 u_1 - \sinh^2 u_1 = 1, \quad (6)$$

and that this result is general may be instantly verified by (2), for

$$\begin{aligned} \cosh^2 u - \sinh^2 u &= \frac{(b^2 x_1 x_2 - a^2 y_1 y_2)^2 - a^2 b^2 (x_1 y_2 - x_2 y_1)^2}{a^4 b^4} \\ &= \frac{b^2 x_1^2 - a^2 y_1^2}{a^2 b^2} \cdot \frac{b^2 x_2^2 - a^2 y_2^2}{a^2 b^2} = 1. \end{aligned}$$

III. *The Addition-Theorem.* Substituting (5) in the definitions of (2), we have

$$\left. \begin{aligned} \sinh (u_2 - u_1) &= \sinh u_2 \cosh u_1 - \cosh u_2 \sinh u_1 \\ \cosh (u_2 - u_1) &= \cosh u_2 \cosh u_1 - \sinh u_2 \sinh u_1 \end{aligned} \right\} \quad (7)$$

Writing  $+ u_1$  instead of  $- u_1$ , we have by (3)

$$\left. \begin{aligned} \sinh (u_2 + u_1) &= \sinh u_2 \cosh u_1 + \cosh u_2 \sinh u_1 \\ \cosh (u_2 + u_1) &= \cosh u_2 \cosh u_1 + \sinh u_2 \sinh u_1 \end{aligned} \right\} \quad (8)$$

The generality of these formulæ is easily verified in the manner just exemplified in formula (6).

IV. It is clear from (5) that the definitions we have given reduce to the ordinary form for the special position there considered. It remains to be shown that in the form given  $\sinh u$  and  $\cosh u$  are really functions of  $u$  alone. To this end let us choose the asymptotes as axes of coördinates. Let  $\alpha$  and  $\beta$  be the coördinates of any point, and let  $\omega$  be the angle between the asymptotes; the equation of the hyperbola is then

$$\alpha\beta = \frac{1}{2}ab \csc \omega. \quad (9)$$

The coördinates of  $P$  being  $\alpha_1 = OL$  and  $\beta_1 = LP$ , those of  $Q$  being  $\alpha_2$  and  $\beta_2$ , those of  $P'$  will be  $\alpha_1$  and  $-\beta_1$ , and we have

$$\sinh u = \frac{\alpha_2 \beta_1 - \alpha_1 \beta_2}{ab \csc \omega}, \quad \cosh u = \frac{\alpha_2 \beta_1 + \alpha_1 \beta_2}{ab \csc \omega}, \quad (10)$$

or, if we write  $\frac{\alpha_2}{\alpha_1} = \lambda$  and therefore  $\frac{\beta_2}{\beta_1} = \lambda^{-1}$ ,

$$\sinh u = \frac{1}{2}(\lambda - \lambda^{-1}), \quad \cosh u = \frac{1}{2}(\lambda + \lambda^{-1}). \quad (11)$$

If we now apply to the plane the linear transformation

$$\alpha' = k\alpha, \quad \beta' = k^{-1}\beta, \tag{12}$$

the hyperbola is transformed into itself. The points  $P$  and  $Q$  are moved to any arbitrary position on the curve, but the ratio  $\alpha_2 : \alpha_1$  is unaltered. It is easily shown that in this transformation the area of any triangle, and hence the area of any figure in the plane, is unchanged; so that  $u$ ,  $\sinh u$  and  $\cosh u$  are unchanged. They are therefore all functions of the ratio  $\alpha_2 : \alpha_1$  alone. Hence  $\sinh u$  and  $\cosh u$  are functions of  $u$  alone, and the definitions here given are a proper generalization of the usual definitions.

V. *The exponential formulæ.* The sector  $OPQ$  may be regarded as the limit of a circumscribed polygon and hence  $u$  may be regarded as the limit of the sum of a series of hyperbolic sines. To make each of the terms in this series equal, we have evidently only to put  $\lambda = \rho^n$ , where  $n$  may be any whole number. Then, writing  $lt$  briefly for *limit*,

$$u = \frac{1}{2} lt \ n(\rho - \rho^{-1}) = lt \ \frac{n}{2} (\lambda^{\frac{1}{n}} - \lambda^{-\frac{1}{n}}) \tag{13}$$

$$= lt \ \frac{\lambda^{\frac{2}{n}} - 1}{2} \cdot \lambda^{-\frac{1}{n}}, \text{ as } n \text{ increases indefinitely.}$$

The limit of  $\lambda^{-\frac{1}{n}}$  is equal to 1, and the limit of  $\frac{\lambda^{\frac{2}{n}} - 1}{2}$  is

the natural logarithm of  $\lambda$ . Hence  $\lambda = e^u$ . Introducing this value of  $\lambda$  in (12), we have

$$\sinh u = \frac{1}{2}(e^u - e^{-u}), \quad \cosh u = \frac{1}{2}(e^u + e^{-u}). \tag{14}$$

BERKELEY, CAL., November 26, 1894.