

ON THE SIMULTANEOUS APPROXIMATION OF TWO REAL NUMBERS¹

RAPHAEL M. ROBINSON

If $\xi_1, \xi_2, \dots, \xi_n$ are any real numbers and t is a positive integer, then it is well known that integers a_1, a_2, \dots, a_n, b can be found, such that $0 < b \leq t^n$ and

$$|b\xi_k - a_k| < 1/t, \quad k = 1, 2, \dots, n.$$

The proof is briefly the following.² Consider the $t^n + 1$ points $(r\xi_1, r\xi_2, \dots, r\xi_n)$, where $r = 0, 1, \dots, t^n$. Reduce mod 1 to congruent points in the unit cube ($0 \leq x_1 < 1, \dots, 0 \leq x_n < 1$). If this cube is divided into t^n cubes of edge $1/t$ (including the lower boundaries), then at least one of these small cubes must contain two of the reduced points, say those with $r = r'$ and $r = r''$. With $b = |r' - r''|$ and suitable a 's, we evidently satisfy the required inequalities.

For $n = 1$, the inequality can be sharpened to

$$|b\xi - a| \leq 1/(t + 1),$$

b satisfying the condition $0 < b \leq t$. For if we consider the points $r\xi$ ($r = 0, 1, \dots, t$), and mark the points in the interval $0 \leq x \leq 1$ which are congruent to them mod 1, we have at least $t + 2$ points marked, since corresponding to $r = 0$ we mark both 0 and 1. Some two of the marked points must lie within a distance $1/(t + 1)$ from each other, so that the desired conclusion follows. This is the best result, as the example $\xi = 1/(t + 1)$ shows.

The present note solves the corresponding problem for $n = 2$. For larger values of n the problem appears more difficult.

THEOREM. *If ξ_1 and ξ_2 are any real numbers, and s is a positive integer, then integers a_1, a_2, b can be found, such that $0 < b \leq s$, and*

$$|b\xi_k - a_k| \leq \max\left(\frac{[s^{1/2}]}{s + 1}, \frac{1}{[s^{1/2}] + 1}\right), \quad k = 1, 2.$$

For every s , values of ξ_1 and ξ_2 can be found for which the inequalities could not both be satisfied if the equality sign were omitted.

¹ Presented to the Society, November 23, 1940.

² The method used in this proof (*Schubfachprinzip* or "pigeonhole principle") was first used by Dirichlet in connection with a similar problem. We sketch the proof here in order to compare it with the proof of the theorem below, which also uses that method.

The inequalities may also be written

$$|b\xi_k - a_k| \leq \begin{cases} t/(s+1) & \text{for } t^2 - 1 \leq s \leq t(t+1) - 1, \\ 1/(t+1) & \text{for } t(t+1) - 1 \leq s \leq (t+1)^2 - 1. \end{cases}$$

It will be noted that in some intervals the bound does not decrease as s increases.

We show first that the theorem is the best possible. We shall think of the inequalities in the form just given. If $s < (t+1)^2$, then it is evident that $\xi_1 = 1/(t+1)$, $\xi_2 = 1/(t+1)^2$ are a pair of real numbers which cannot be approximated simultaneously with an error less than $1/(t+1)$; this settles the second case. For the first case, consider the pair of real numbers $\xi_1 = 1/(s+1)$, $\xi_2 = t/(s+1)$. We are to show that not both errors can be made less than $t/(s+1)$. We note first that $b\xi_1$ and $b\xi_2$ differ from integers by the same amount as $(s+1-b)\xi_1$ and $(s+1-b)\xi_2$; hence we may suppose that $b \leq (s+1)/2$, and therefore $0 < b\xi_1 \leq 1/2$. In order to make $|b\xi_1 - a_1| < t/(s+1)$, we must have $0 < b < t$. Then $0 < b\xi_2 < 1$. Since $b\xi_2 \geq \xi_2 = t/(s+1)$ and $1 - b\xi_2 \geq 1 - (t-1)\xi_2 = 1 - (t-1)t/(s+1) \geq t/(s+1)$, we see that the inequality $|b\xi_2 - a_2| < t/(s+1)$ cannot be satisfied.

The theorem evidently follows from the lemma below, by putting $t = [s^{1/2}]$.

LEMMA. *Let s and t be positive integers with $s \geq t$. If ξ_1 and ξ_2 are any real numbers, then integers a_1, a_2, b can be found, such that $0 < b \leq s$, and*

$$|b\xi_1 - a_1| \leq t/(s+1), \quad |b\xi_2 - a_2| \leq 1/(t+1).$$

PROOF. Consider the points $(r\xi_1, r\xi_2)$ with $r = 0, 1, \dots, s$. Mark all the points congruent to these mod 1 which fall in the rectangle $0 \leq x_1 \leq t, 0 \leq x_2 < 1$. There are $(s+1)t$ points to be marked with $x_1 < t$; and in addition, the point $(t, 0)$ is marked, corresponding to $r = 0$. If we divide our rectangle into $s+1$ rectangles of width $t/(s+1)$ (closed except at the top) by means of vertical lines, then at least one of them contains more than t points, all corresponding to different values of r . The corresponding values of x_2 are $t+1$ or more numbers, some two of which differ mod 1 by not more than $1/(t+1)$. Thus we find two points $(r'\xi_1, r'\xi_2)$ and $(r''\xi_1, r''\xi_2)$, whose horizontal distance mod 1 does not exceed $t/(s+1)$ and whose vertical distance mod 1 does not exceed $1/(t+1)$. Putting $b = |r' - r''|$ gives the required result.

UNIVERSITY OF CALIFORNIA