

and

$$(5) \quad v_i(x) = \sum_{k=1}^i B_{ik} w_k(x).$$

Since $B_{ik} \leq \left\{ \sum_{j=1}^{\infty} B_{jk}^2 \right\}^{1/2} = l_{i-1, k}$, in all the cases of the ratio function $r(x)$ considered, the right-hand members of (4) and (5) are absolutely convergent and bounded, wherever, respectively, the v 's and w 's are bounded. Hence, if conditions are such that the right-hand member of (4) converges to the value of the left-hand member and if a set of points is known for which the v 's are bounded, then the w 's are bounded on the same set except where $r(x) = 0$. Similarly, boundedness of the w 's leads through (5) to results on the boundedness of the v 's.

UNIVERSITY OF MINNESOTA

A MAPPING CHARACTERIZATION OF PEANO SPACES

O. G. HARROLD, JR.

The Hahn-Mazurkiewicz theorem states that any Peano space (compact, connected, locally connected, metric space) is a continuous image of the interval $0 \leq t \leq 1$, and conversely. Clearly, the mapping function is not uniquely determined. If the Peano space \mathcal{M} has special topological properties, the mapping may be selected in a simpler fashion than might be expected generally. On the other hand, special properties of \mathcal{M} may impose certain necessary restrictions on the mapping. For example, if \mathcal{M} is a regular continuum in the sense of Menger, then, by a theorem due to Nöbeling,¹ there is a continuous mapping f of the circle² onto \mathcal{M} such that each point of finite order is covered by the mapping a number of times which does not exceed the order of the point. That is, if $o(x)$ is the order of the point x and $m(x)$ is the number of points in $f^{-1}(x)$, then $m(x) \leq o(x)$ for each point for which $o(x)$ is finite. On the other hand, if \mathcal{M} is of dimension n , then *any* continuous mapping of a 1-dimensional compact set onto \mathcal{M} ,

Presented to the Society, September 5, 1941; received by the editors October 21, 1941.

¹ G. Nöbeling, *Reguläre Kurven als Bilder der Kreislinie*, Fundamenta Mathematicae, vol. 20 (1933), pp. 30–46.

² The interval may be used instead of the circle if we make $f(0) = f(1)$ and count inverses on $0 \leq t < 1$.

in particular, of an interval or circle, is such that there is a dense set of points in the n -dimensional part of \mathcal{M} each of which has at least n inverse points in the original set.³

Denote the set of local separating points of the Peano space \mathcal{M} by \mathcal{L} .⁴ If $\overline{\mathcal{M} - \mathcal{L}}$, that is, if \mathcal{M} contains no free arcs,⁵ there is a strongly irreducible mapping of the interval \mathfrak{J} or circle \mathcal{C} onto \mathcal{M} .⁶ That is, for such spaces there exist continuous mappings of \mathfrak{J} or \mathcal{C} onto \mathcal{M} such that no proper closed subset maps onto the whole space. Thus if \mathcal{M} is an n -dimensional sphere and f is a strongly irreducible mapping of \mathfrak{J} onto \mathcal{M} , there is a dense set of points each covered at least $n+1$ times and also a dense set of points each covered just once.

In addition to the symbols f , \mathcal{M} , \mathcal{L} , $m(x)$ and $o(x)$ used above, the following notations will be observed. Let ψ denote the aggregate of points $x \in \mathcal{M}$ lying in an open free arc of $\mathcal{M} - a - b$. If for a (continuous) mapping of \mathfrak{J} into a subset of \mathcal{M} , $y \in \psi$ implies $m(y) \leq 2$, the mapping will be said to be of type \mathfrak{M} .

THEOREM 1. *Let a and b be points of the Peano space \mathcal{X} . There is a continuous mapping of the interval $0 \leq t \leq 1$ onto \mathcal{X} of type \mathfrak{M} such that $f(0) = a$, $f(1) = b$.*

The theorem asserts, essentially, that there is a mapping of \mathfrak{J} onto \mathcal{X} such that every free arc is swept through at most twice.

The following lemmas will be useful in the proof of Theorem 1.

LEMMA 1.1. *If D is a subcontinuum of the dendrite D^0 , to $\epsilon > 0$ there is a finite collection D^1, D^2, \dots, D^n of dendrites in D^0 such that $D = D^1 \subset D^2 \subset \dots \subset D^n = D^0$ and each component of $D^{i+1} - D^i$ has a diameter less than or equal to ϵ .*

Let ρ be a convex metric⁷ on D^0 . Let $d = \text{glb}$ of numbers r such that

³ If the original set is locally euclidean, the phrase *at least n* may be replaced by *at least $n+1$* . See W. Hurewicz, *Über dimensionserhöhende stetige Abbildungen*, Journal für die reine und angewandte Mathematik, vol. 169 (1933), pp. 71-78.

⁴ The point p is called a local separating point of M provided that to every neighborhood U of p there is some pair of points of the component of U containing p which is separated in $U - p$.

⁵ The set A is called a free arc of M provided A is an arc and the interior of A is open in M . An *open* free arc is an open subset of M which is homeomorphic to $0 < x < 1$. A point is said to lie in an open free arc provided there is a neighborhood of the point in M which is an open free arc. It is to be noted that if M is an arc, neither end point lies in an open free arc.

⁶ O. G. Harrold, Jr., *A note on strongly irreducible maps of an interval*, Duke Mathematical Journal, vol. 6 (1940), pp. 750-752.

⁷ The metric ρ is called *convex* after Menger provided that to each pair of distinct points x and y in M there is a point of $M - x - y$ such that $\rho(x, z) + \rho(z, y) = \rho(x, y)$.

$D^0 \subset S(D, r)$. Each of the sets $D^i = S[D, (i-1)e]$, $i = 1, 2, \dots, n$ is a subcontinuum of D^0 . The sets D^i satisfy our requirements, where n is the smallest integer such that $(n-1) \geq d$.

LEMMA 1.2. *Let X be a Peano space. There is a sequence (T_i) of dendritic graphs in X such that (a) $\lim T_i = X$, (b) $T_{i+1} \supset T_i$, and (c) each component of $T_{i+1} - T_i$ has a diameter less than or equal to $1/2^{i+1}$.*

That a sequence of dendrites exists in X satisfying (a) and (b) is well known. Application of Lemma 1.1 to the successive terms of this sequence gives the desired result.

Proof of Theorem 1. The theorem is true for a connected dendritic graph with n end points, as a simple induction shows. Suppose, temporarily, that neither a nor b lies in an open free arc. Let T_1 be a dendritic graph in X containing a and b .⁸ Let f_1 denote a continuous mapping of type \mathfrak{M} of \mathfrak{J} onto T_1 with $f_1(0) = a$, $f_1(1) = b$. Let (T_i) be a sequence of dendritic graphs satisfying Lemma 1.2. Since T_2 is a graph, $T_2 - T_1$ has but a finite number of components which may be denoted by $C_1^1, C_2^1, \dots, C_{n_1}^1$. Each $\bar{C}_i^1 \cdot T_1$ is a point c_i . Let $x_i \in f_1^{-1}(c_i)$. By a rearrangement of notation it may be supposed that $0 \leq x_1 < x_2 < \dots < x_{p_1} \leq 1$, $p_1 \leq n_1$, where each x_i corresponds to a distinct c_i . Set $d_1 = \min |x_i - x_j|$, $i \neq j$, $|x_i|$, $x_i \neq 0$, $|1 - x_i|$, $x_i \neq 1$. To $\epsilon = 1/2$ there is a $d_2 > 0$ such that $|x - y| < d_2$ implies $\rho[f_1(x), f_1(y)] < \epsilon/2 = 1/2^2$, where ρ denotes the metric of X . Put $W = S(x_1 + x_2 + \dots + x_{p_1}, d/3)$, where $d = \min(d_1, d_2)$. Let J_i be the component of W containing x_i . Let $I_1, I_2, \dots, I_{p_1+1}$ be the intervals on \mathfrak{J} complementary to W , where I_1 becomes degenerate if $x_1 = 0$ and I_{p_1+1} degenerate if $x_{p_1} = 1$. The interval \mathfrak{J} is now subdivided into the intervals (in order) $I_1, \bar{J}_1, I_2, \bar{J}_2, \dots, I_{p_1+1}$. Let t denote the piecewise linear map obtained by sending I_1 onto $(0x_1)$ with order preserved, I_2 onto (x_1x_2) , \dots , I_{p_1+1} onto $(x_{p_1}1)$. For $x \in \sum I_i$, put $f_2(x) = f_1[t(x)]$. On \bar{J}_i define a map g_i of the desired type so that $g_i(\bar{J}_i) = D_i$, where D_i is the enclosure of all components C_i^1 having c_i as a limit point. The set D_i is a dendrite of diameter less than or equal to $1/2^2$. The map g_i may be so selected that for the end points of J_i , $g_i = f_2$.

See K. Menger, *Untersuchungen über allgemeine Metrik*, *Mathematische Annalen*, vol. 100 (1928), pp. 81 ff. For the existence of the metric assumed here, see C. Kuratowski and Whyburn, *Sur les elements cycliques et leurs applications*, *Fundamenta Mathematicae*, vol. 16 (1930), pp. 305-331.

⁸ T_1 could be taken to be an arc but in order that the discussion to follow be general it is assumed only to be a connected linear graph containing no simple closed curve.

The definition of f_2 is now completed: if $x \in \bar{J}_i, f_2(x) = g_i(x)$. Clearly, $f_2(\mathfrak{J}) = T_2$ and f_2 is continuous. For $x \in \sum I_i, |x - t(x)| < d$, hence $\rho[f_1(x), f_2(x)] < 1/2^2$. If $x \in \bar{J}_i, \rho[f_1(x), f_1(x_i)] < 1/2^2$, and since the diameter of D_i is less than or equal to $1/2^2, \rho[f_2(x), f_2(t^{-1}(x_i))] < 1/2^2$. But $f_2(t^{-1}(x_i)) = f_1(x_i)$, hence by the triangle inequality $\rho[f_1(x), f_2(x)] < 1/2$. Thus if σ denotes the usual metric of the function space $\mathcal{X}^{\mathfrak{J}}$, $\sigma(f_1, f_2) \leq 1/2$.⁹

To show that f_2 is of type \mathfrak{M} consider a point $y \in \psi \cdot T_2$. It is to be shown that $f_2^{-1}(y)$ contains at most two points, that is, $m(y, f_2) \leq 2$. If $y \in T_2 - T_1, y$ lies in a unique D_i , hence $m(y, f_2) \leq 2$. If $y \in T_1 - \sum D_i, f_2^{-1}(y) = t^{-1}f_1^{-1}(y)$. Since f_1 has the desired property and t is 1-1 on the set composed of the interiors of the intervals $I_i, m(y, f_2) \leq 2$. Consider the remaining case $y = c_i$. Here $m(y, f_1) = 1$, for suppose, on the contrary, $f_1^{-1}(y) \supset q_1 + q_2$. Since $y \in \psi, y \neq a, b$, hence q_1 and q_2 divide \mathfrak{J} into three subintervals A, B and C . But each of these subintervals has as an image under f_1 a nondegenerate continuum containing y . Hence points of T_1 near y have three inverses on \mathfrak{J} , which denies the property of f_1 . Since $y \in \psi$, only one component $C_i' = D_i$ of $T_2 - T_1$ can have y as a limit point. The mapping g_i has the desired property, thus $f_2^{-1}(y) = g_i^{-1}(y) + f_1^{-1}[t^{-1}(y)]$ is precisely a pair of points. Hence for $y \in \psi \cdot T_2, m(y, f_2) \leq 2$, and f_2 is of type \mathfrak{M} .

The general inductive hypothesis is now clear.

To the dendrite T_n there is a continuous mapping $f_n, f_n(\mathfrak{J}) = T_n$, of type \mathfrak{M} and such that $f_n(0) = a, f_n(1) = b$. Further, $\sigma(f_{i-1}, f_i) \leq 1/2^{i-1}, i = 2, 3, \dots, n$. The construction of f_{n+1} from f_n is accomplished precisely as above.

There is thus determined a sequence of points (f_n) of the space $\mathcal{X}^{\mathfrak{J}}$ such that to $\epsilon > 0$ there is an index N such that for $i, j > N, \sigma(f_i, f_j) < \epsilon$. The space $\mathcal{X}^{\mathfrak{J}}$ being complete, let $\lim f_n = f$. Clearly, $f(\mathfrak{J}) = \mathcal{X}$. To complete the proof of Theorem 1 it will suffice to show that $y \in \psi$ implies $f^{-1}(y)$ has at most two components. For, if we grant this, the factorization $f = k[h(x)]$, where h is the monotone transformation obtained by shrinking the components of $f^{-1}(y)$ into points and k is the corresponding light transformation, yields $k(\mathfrak{J}) = \mathcal{X}, m(y \in \psi, k) \leq 2$, hence k is of type \mathfrak{M} .¹⁰

Suppose, on the contrary, $y \in \psi$ and X_1, X_2 and X_3 are three components of $f^{-1}(y)$. Since $y \in \psi, y \neq a, b$, hence $\mathfrak{J} - \sum X_i$ has precisely 4 components $R_i, i = 1, 2, 3$ and 4. Suppose w_i and t_i are the left and right end points of X_i , respectively. (If X_i is a point, $w_i = t_i$.) Let the

⁹ If $f, g \in \mathcal{X}^{\mathfrak{J}}, \sigma(f, g) = \text{lub } \rho[f(x), g(x)], x \in \mathfrak{J}$.

¹⁰ This is an application of a factor theorem for continuous transformations due to Eilenberg and Whyburn.

notation be arranged so that $w_i = \bar{R}_i \cdot X_i$, $i = 1, 2$ and 3 . The points w_i and w_j , $i \neq j$, are, of course, distinct. Let A_i , $i = 1, 2$ and 3 be a sub-interval of \bar{R}_i containing w_i such that $f(A_i) \subset U$, where U is any fixed open free arc of X containing y . Let B_{i+1} $i = 1, 2$ and 3 be a proper sub-interval of $R_{i+1} - A_{i+1}$ containing t_i such that $f(B_i) \subset U$, where $A_4 = 0$ by definition. Since X_i is a component of $f^{-1}(y)$, the sets of $f(A_i), f(B_i)$ are nondegenerate subarcs of U with at least the point y common. Some point y^0 of $U - y$ must be covered by at least three of these six sets. Denote three of the corresponding sets $A_i (B_i)$, $i = 1, 2$ and 3 by C_1, C_2 and C_3 . One end point of C_i , say a_i , maps into y . As C_i is traversed from a_i let b_i be the first point in $f^{-1}(y^0)$. Let G denote any subarc of $U - y - y^0$ which lies between y and y^0 . Set $d = \rho(G, y + y^0) > 0$. Then for n large enough $\sigma(f_n, f) < d/3$. But $f_n(a_i b_i)$ is a connected subset of U which contains a point from each component of $U - G$, hence $f_n(a_i b_i) \supset G$. This denies that f_n is of type \mathfrak{M} . The proof of Theorem 1 under the special restriction that neither a nor b lies in a free arc has been completed.

To remove the restriction suppose first that only a lies in a free arc of X . Imagine that X is situated in the Hilbert cube and let \mathcal{A} be an arc which is joined onto X at a and has no other point in X . Let a^1 be the other end point of \mathcal{A} . Construct a mapping as above with $f(0) = a^1, f(1) = b, f(\mathfrak{J}) = X + \mathcal{A}$. Since neither a^1 nor b lies in an open free arc of $X + \mathcal{A}$, such a mapping will exist. Let x^1 be the least x for which $f(x) = a$. Then the mapping f on the interval $x^1 \leq t \leq 1$ satisfies our requirements. A similar modification suffices to treat the case in which b is an open free arc and also the case in which both a and b have this property.

Set $\mathcal{W} = X - \bar{\psi}$. The set \mathcal{W} is open. Put $\mathcal{Q} = \psi + (\mathcal{W} \cdot \mathcal{N})$, where \mathcal{N} is the set of nonlocal separating points of X . We come now to the principal result.

THEOREM 2. *Let a and b be points of the metric space X . In order that X be a Peano space it is necessary and sufficient that for any countable subset \mathcal{P} of $\mathcal{Q} - a - b$ there be a continuous mapping f of $0 \leq t \leq 1$ onto X such that $f(0) = a, f(1) = b$ and $y \in \mathcal{P}$ implies $m(y) \leq 2$.*

PROOF. The sufficiency is clear. If $\psi = 0$, the result is known, in fact, in this case a mapping of the described type exists such that for $y \in \mathcal{P}$, $m(y) = 1$.⁶ It is supposed, then, that $\psi \neq 0$. By application of Theorem 1, there is a mapping of type \mathfrak{M} of \mathfrak{J} onto X with $f(0) = a, f(1) = b$. The desired map will be obtained by a modification of f .

To facilitate the discussion it will be supposed that X has an \mathcal{S} -metric, that is, a metric ρ such that for each $r > 0$ and $x \in X$, $\bar{\mathfrak{S}}(x, r)$ is a lo-

cally connected continuum.¹¹ Let $A = a_1 + a_2 + \dots = \mathcal{P} \cdot \mathcal{W} = \mathcal{P} \cdot \mathcal{W} \cdot \mathcal{N}$. Set $d_1 = \rho(a_1, \mathcal{X} - \mathcal{W})$. Choose a number e_1 such that $d_1/2 < e_1 < d_1$ and $A \cdot \{\overline{S}(a_1, e_1) - S(a_1, e_1)\} = 0$. This is possible since A is countable. Let a_{k_2} be the first point of A in $\mathcal{W} - \overline{S}_1$, where $S_1 = S(a_1, e_1)$. Let $d_2 = \rho[a_{k_2}, (\mathcal{X} - \mathcal{W}) + S_1]$. The number e_2 is chosen so that $d_2/2 < e_2 < d_2$ and $A \cdot \{\overline{S}(a_{k_2}, e_2) - S(a_{k_2}, e_2)\} = 0$. Continuing in this way a sequence of spheres (S_i) is determined such that (a) \overline{S}_i is a Peano space, (b) $\overline{S}_i \cdot \overline{S}_j = 0, i \neq j$, (c) $\delta(S_i) \rightarrow 0$, (d) $\sum \overline{S}_i = \overline{\mathcal{W}}$ and (e) $A \cdot (\overline{S}_i - S_i) = 0$.

Set $V_i = f^{-1}(S_i)$. Let $V_{ij}, j = 1, 2, \dots$ be the components of V_i . Let V_{i1} be a component of V_i such that $V_{i1} \cdot f^{-1}(a_{k_i}) \neq 0$. Every point of \overline{S}_i is either a nonlocal separating point of \overline{S}_i or a limit point of such points. This is clear if $x \in S_i$, for $S_i \subset \mathcal{W}$. If $x \in \overline{S}_i - S_i$, x is a limit point of points of S_i and hence a limit point of nonlocal separating points of \overline{S}_i . Thus, having shown that \overline{S}_i is a Peano space with no free arcs, there is a strongly irreducible mapping, $f_{i1}(V_{i1}) = \overline{S}_i$, such that $f = f_{i1}$ on $\overline{V}_{i1} - V_{i1}$ and $y \in \mathcal{P} \cdot \overline{S}_i$ implies $f_{i1}^{-1}(y)$ is a single point.

On $V_{ij}, j > 1$, two cases are distinguished according as f maps the end points of V_{ij} into the same point or not. If f carries the end points of V_{ij} into x , define $f_{ij} \equiv x$ on \overline{V}_{ij} . If f carries the end points of V_{ij} into distinct points x and y , proceed as follows. The set $\overline{S}_i - A$ is a connected and locally connected G_δ set¹² in a complete space, hence there is an arc $R_{ij} \subset \overline{S}_i - A$ which joins x and y .¹³ On \overline{V}_{ij} define f_{ij} to be a homeomorphism into R_{ij} such that f_{ij} agrees with f on $\overline{V}_{ij} - V_{ij}$.

The new mapping g will now be defined. On $\mathfrak{J} - \sum V_{ij}$, set $g(x) \equiv f(x)$. On V_{ij} , set $g(x) = f_{ij}(x)$. Since f agrees with g on the end points of V_{ij} and each f_{ij} is continuous, g is continuous (we use here the condition (c) on the spheres (S_i)). Clearly, $g(\mathfrak{J}) = \mathcal{X}$. If $y \in \mathcal{P} \cdot \psi$, $m(y, f) = m(y, g) \leq 2$, by virtue of the fact that f is of type \mathfrak{M} . If $y \in \mathcal{P} \cdot \mathcal{W}$, y lies in a unique S_i and $g^{-1}(y) = f_{i1}^{-1}(y)$, hence $m(y, g) = 1$.

NORTHWESTERN UNIVERSITY

¹¹ J. L. Kelley, *A metric connected with property S*, American Journal of Mathematics, vol. 61 (1939), pp. 764-768.

¹² The complement of a countable set of nonlocal separating points in a Peano space is connected and locally connected, see G. T. Whyburn, *Semi-closed sets and collections*, Duke Mathematical Journal, vol. 2 (1936), pp. 685-690.

¹³ This is the well known Moore-Menger generalization of the arcwise connectivity theorem for regions in a Peano space.