

ON THE EXTENSION OF DIFFERENTIABLE FUNCTIONS

HASSLER WHITNEY

The author has shown previously how to extend the definition of a function of class C^m defined in a closed set A so it will be of class C^m throughout space (see [1]).¹ Here we shall prove a uniformity property: If the function and its derivatives are sufficiently small in A , then they may be made small throughout space. Besides being bounded, we assume that A has the following property:

(P) There is a number ω such that any two points x and y of A are joined by an arc in A of length less than or equal to ωr_{xy} (r_{xy} being the distance between x and y).

This property was made use of in [2]; its necessity in the theorem is shown by two examples below.

A second theorem removes the boundedness condition in the first theorem, and weakens the hypothesis (P); its proof makes use of the proof of the first theorem. We remark that in each theorem, as in [1], the extended function is a linear functional of its values in A .

The proof of Theorem I is obtained by examining the proof in [1]; hence we assume that the reader has this paper before him, and we shall follow its notations closely.

THEOREM I. *Let A be a bounded closed set in n -space E with the property (P), and let m be a positive integer. Then there is a number α with the following property. Let $f(x)$ be any function of class C^m in A , with derivatives $f_k(x)$ ($\sigma_k = k_1 + \dots + k_n \leq m$). Suppose*

$$|f_k(x)| < \eta \quad (x \in A, \sigma_k \leq m).$$

Then $f(x)$ may be extended throughout E so that

$$|f_k(x)| < \alpha \eta \quad (x \in E, \sigma_k \leq m).$$

Let d be the diameter of A , or 1 if this is larger, and let R be a spherical region of radius $2d$ with its center at a point of A . Set $f(x) = 0$ in $E - R$. Then the extension of f in $R - A$ given in [1] will be shown to have the property, using

$$\alpha = 2n(m!)^n(m+1)^{8n}(433n^{1/2}d\omega)^{mc}N,$$

where N and c are as given in [1, §§11, 12]. Note that $433 = 4 \cdot 108 + 1$.

Presented to the Society, September 13, 1943; received by the editors November 27, 1943.

¹ Numbers in brackets refer to the references cited at the end of this paper.

Set $B = A \cup (\bar{R} - R)$. We show first that for any points x', x'' of B ,

$$|R_k(x'; x'')| < \beta r_{x'x''}^{m-\sigma_k}, \quad \beta = 2n(m+1)^n \omega^m.$$

Suppose first that x' and x'' are in A . Let C be a curve in A joining them, of length less than or equal to $\omega r_{x'x''}$. The inequality is then a consequence of [2, Lemma 3]. Suppose next that one of the points is in A , and the other is in $\bar{R} - R$ (the case that both are in $\bar{R} - R$ is trivial). By [1, (3.1)], since $r_{x'x''} \geq d \geq 1$,

$$\begin{aligned} |R_k(x'; x'')| &\leq \eta + \sum_{\sigma_i \leq m - \sigma_k} \eta r_{x'x''}^{\sigma_i} \leq r_{x'x''}^{m-\sigma_k} \left[1 + \sum_{\sigma_i \leq m - \sigma_k} 1 \right] \\ &\leq (m+1)^n r_{x'x''}^{m-\sigma_k} \eta. \end{aligned}$$

Now take any x in $R - B$. Let $\delta^*/4$ be the distance from x to B , and let x^* be a point of B distant $\delta^*/4$ from x . Say x is in the cube C of the set of cubes K_s ; let $I_{\lambda_1}, \dots, I_{\lambda_l}$ be those I_λ with points in C (see [1, §11]). Now y^v is the center of I_v , and x^v is a nearest point of B to y^v . As noted in [1, (9.1)], $r_{y^v x^*}$ and $r_{y^v x^v}$ each lie between $\delta^*/8$ and $\delta^*/2$. Since $r_{x^v y^v} < \delta^*/2$, we have

$$r_{x^v x^*} < \delta^*, \quad r_{x^v x^v} < \delta^*.$$

The definition of ζ in [1, §11] together with [1, (6.3)] gives

$$\zeta_{v;k}(x) = \psi_k(x; x^v) - \psi_k(x; x^*) = \sum_{\sigma_i \leq m - \sigma_k} \frac{R_{k+i}(x^v; x^*)}{i!} (x - x^v)^i.$$

Hence

$$|\zeta_{v;k}(x)| < (m+1)^n \beta r_{x^v x^*}^{m-\sigma_k-\sigma_i} r_{x^v x^v}^{\sigma_i} < (m+1)^n \beta \delta^{*m-\sigma_k} \eta.$$

Following [1, §11] still, we find

$$|D_k g(x) - \psi_k(x; x^*)| < c \sum_{\sigma_i \leq m - \sigma_k} (m!)^n 2^{\sigma_i} N(m+1)^n \beta \delta^{*m-\sigma_k+\sigma_i} \eta.$$

As in [1], $2^{\sigma_i} < 108n^{1/2}/\delta^*$; hence

$$|D_k g(x) - \psi_k(x; x^*)| < c(m!)^n N(m+1)^{2n} (108n^{1/2})^m \beta \delta^{*m-\sigma_k} \eta.$$

Moreover, since $r_{xx^*} < 3d$, [1, (6.1)] gives

$$|\psi_k(x; x^*)| < 3^m (m+1)^n d^m \eta.$$

Since $\delta^* \leq 4d$ and $f(x) = g(x)$ in $R - B$, the theorem follows.

We turn now to the second theorem. We shall say A satisfies (P) locally if for each $x \in A$ there is a neighborhood U of x and a number

ω such that any two points y and z of $A \cap U$ are joined by an arc in A of length not greater than ωr_{xy} .

THEOREM II. *Let A be a closed subset of an open set R in E , satisfying (P) locally, and let m be a positive integer. Then for any continuous function $\epsilon(x)$ defined and greater than 0 in R there is a continuous function $\delta(x)$ defined and greater than 0 in A with the following property. Let $f(x)$ be any function of class C^m in A , such that*

$$|f_k(x)| < \delta(x) \quad (x \in A, \sigma_k \leq m).$$

Then $f(x)$ may be extended throughout R so that

$$|f_k(x)| < \epsilon(x) \quad (x \in R, \sigma_k \leq m).$$

REMARKS. The preceding theorem is easily seen to be a consequence of this one. The present theorem holds if E is replaced by a differentiable manifold M , in which a fixed set of coordinate systems (each one intersecting but a finite number of others) is used to measure the size of derivatives. To show this, we imbed M in a Euclidean space E' (see [3, Theorem 1]), giving $A \subset R \subset R' \subset E'$ (R' open in E' ; we let R' contain no points of the limit set of M), extend f throughout R' (see the proof of [3, Lemma 4]), and consider its values in R .

To prove the theorem, we begin by choosing spherical regions U_1, U_2, \dots , each $\bar{U}_i \subset R$, with the following properties:

- (a) Each U_i is in a neighborhood U as described above.
- (b) Each \bar{U}_i intersects but a finite number of other \bar{U}_j .
- (c) If U_i is of radius ρ_i , and $U_i^!$ is the concentric region of radius $\rho_i/2$, then $R = \sum U_i^!$.

Let $\psi^i(x)$ be a function of class C^m in E such that

$$\psi^i(x) > 0 \quad (x \in U_i^!),$$

$$\psi^i(x) = 0 \quad (x \in E - U_i^!).$$

Set

$$\phi^i(x) = \psi^i(x) / \sum \psi^i(x) \quad (x \in R);$$

then $\phi^i(x)$ is of class C^m in R , and

$$\phi^i(x) = 0 \quad (x \in R - U_i^!),$$

$$\sum \phi^i(x) = 1 \quad (x \in R).$$

The extension of $f(x)$ is defined as follows. Set

$$f^i(x) = \phi^i(x)f(x) \quad (x \in A),$$

$$f^i(x) = 0 \quad (x \in R - U_i).$$

Then f^i is of class C^m in $A \cup (R - U_i)$. Extend it as in [1] (using a fixed

subdivision of $U_i - A$; we could set $f^i(x) = 0$ in $E - R$ to be of class C^m in R (or E). (Note that if $A \cap U_i' = 0$, then $f^i(x) = 0, x \in R$.) Set

$$f(x) = \sum f^i(x) \quad (x \in R).$$

Then f is an extension of class C^m of its values in A . We must show that it satisfies the condition of smallness.

Choose $a_i \geq 1$ so that

$$|\phi_k^i(x)| \leq a_i \quad (x \in R, \sigma_k \leq m),$$

then if $|f_k(x)| < \eta$ ($x \in A \cap U_i'$),

$$|f_k^i(x)| = \left| \sum_{\sigma_i \leq \sigma_k} \phi_i^i(x) f_{k-i}(x) \right| \leq (m + 1)^n a_i \eta \quad (x \in A).$$

By the choice of U_i , there is an ω_i such that any x' and x'' in $A \cap U_i$ are joined by an arc in A of length not greater than $\omega_i r_{x'x''}$. Set $\sigma_i = \max(1, 2/\rho_i)$. If R_k^i is the remainder for f_k^i , we shall show that for any x' and x'' in $A \cup (R - U_i)$,

$$|R_k^i(x'; x'')| < 2n(m + 1)^{2n} \omega_i^m a_i \sigma_i^m r_{x'x''}^{m-\sigma_k} \eta.$$

If x' and x'' are both in U_i , we apply [2, Lemma 3]. If $x' \in R - U_i$ and $x'' \in U_i'$, or vice versa, then $r_{x'x''} \geq \rho_i/2$, and the proof in the preceding theorem applies; we consider separately the cases $\rho_i/2 \geq 1$, $\rho_i/2 < 1$, using $r_{x'x''} \geq 1$ and $\sigma_i r_{x'x''} \geq 1$ respectively. If $x' \in R - U_i$ and $x'' \in R - U_i'$, or vice versa, $R_k^i = 0$, since $\phi_i^i(x') = \phi_i^i(x'') = 0$. The proof of the preceding theorem now shows that for some α_i , if

$$|f_k(x')| < \eta \quad (x' \in A \cap U_i', \sigma_k \leq m),$$

then

$$|f_k^i(x)| < \alpha_i \eta \quad (x \in R, \sigma_k \leq m).$$

(We may set $\alpha_i = 1$ if $A \cap U_i' = 0$.)

Given $\epsilon(x)$, we determine $\delta(x)$ as follows. For each $x \in R$ there is a set of numbers $\lambda_1, \dots, \lambda_s, s = s(x)$, such that

$$x \in \text{each } U_{\lambda_i}', \quad x \in \text{no other } U_i'.$$

Because of (b), s is finite. Set $\alpha(x) = \alpha_{\lambda_1} + \dots + \alpha_{\lambda_s}$.

We can clearly choose a continuous function $\beta(x)$ in R such that

$$\alpha(x) < \beta(x) \quad (x \in R).$$

We may now choose a continuous function $\delta(x') > 0$ in A such that

for any $x' \in A$, if $U'_{\mu_1}, \dots, U'_{\mu_t}$ are the U'_j containing x' , then

$$\delta(x') \leq \epsilon(x)/\beta(x) \quad (x \in U'_{\mu_1} \cup \dots \cup U'_{\mu_t}).$$

Now take any f of class C^m in A , with $|f_k(x)| < \delta(x)$ ($x \in A, \sigma_k \leq m$); the extension of f through R has been defined. Take any $x \in R$; define $\lambda_1, \dots, \lambda_s$ as above. Then

$$|f_k(x')| < \delta(x') \leq \epsilon(x)/\beta(x) \quad (x' \in A \cap U'_{\lambda_j}, \sigma_k \leq m),$$

and hence

$$|f_k^{\lambda_j}(x)| < \alpha_{\lambda_j} \epsilon(x)/\beta(x) \quad (\sigma_k \leq m).$$

Since $f_k(x) = f_k^{\lambda_1}(x) + \dots + f_k^{\lambda_s}(x)$,

$$|f_k(x)| < \alpha(x) \epsilon(x)/\beta(x) < \epsilon(x)$$

for $\sigma_k \leq m$, which completes the proof.

EXAMPLES. (1) Let A consist of a point, together with a sequence of points approaching it. Letting $f(x) = 1$ at a finite number of points of the sequence, and $f(x) = 0$ in the rest of A shows (with $m = 1$) that the theorem fails here.

(2) Let A be the closed region of the plane defined by (a) $x^2 + y^2 \leq 1$, and (b) either $x \leq 0$ or $|y| \geq x^{3/2}$. Let $f(x, y) = 0$ if $x \leq 0$, and set

$$f(x, y) = \begin{cases} \gamma x^2/(1 + \gamma^2 x^2) & \text{if } x \geq 0, y > 0, \\ -\gamma x^2/(1 + \gamma^2 x^2) & \text{if } x \geq 0, y < 0. \end{cases}$$

We see easily that f is of class C^1 in A . (It would not be if, in (b), we used $|y| \geq x^2$.) The maximum $\partial f/\partial x$ occurs at $x = 1/(3^{1/2}\gamma)$, and has the value $9/(8 \cdot 3^{1/2})$. Set

$$p = (1/3^{1/2}\gamma, 1/3^{3/4}\gamma^{3/2}), \quad q = (1/3^{1/2}\gamma, -1/3^{3/4}\gamma^{3/2}).$$

Then

$$\frac{f(p) - f(q)}{r_{pq}} = \frac{2\gamma/3\gamma^2}{1 + \gamma^2/3\gamma^2} \div \frac{2}{3^{3/4}\gamma^{3/2}} = \frac{3^{3/4}}{4} \gamma^{1/2}.$$

Hence, in any extension of f through the plane, we must have $|\partial f/\partial y| \geq 3^{3/4}\gamma^{1/2}/4$ at some point (between p and q); yet $|f|$, $|\partial f/\partial x|$ and $|\partial f/\partial y|$ are uniformly bounded for all $\gamma > 1$. Taking γ arbitrarily large shows that the conclusion of the theorem does not hold.

REFERENCES

1. H. Whitney, *Analytic extensions of differentiable functions defined in closed sets*, Trans. Amer. Math. Soc. vol. 36 (1934) pp. 63-89.

2. ———, *Functions differentiable on the boundaries of regions*, Ann. of Math. vol. 35 (1934) pp. 482–485.
3. ———, *Differentiable manifolds*, Ann. of Math. vol. 37 (1936) pp. 645–680.
4. ———, *Differentiable functions defined in arbitrary subsets of Euclidean space*, Trans. Amer. Math. Soc. vol. 40 (1936) pp. 309–317. Further references are given here.
5. H. O. Hirschfeld, *Continuation of differentiable functions through the plane*, Quart. J. Math. Oxford Ser. vol. 7 (1936) pp. 1–15.
6. M. R. Hestenes, *Extension of the range of a differentiable function*, Duke Math. J. vol. 8 (1941) pp. 183–192.

HARVARD UNIVERSITY

THE SYMMETRIC JOIN OF A COMPLEX

C. E. CLARK

1. **The definition of J .** Let K be a polyhedron. With each pair of distinct points p, q of K we associate a closed line segment pq . No distinction is made between p and q and the corresponding end points of pq . The length of pq is a continuous function of p and q , and the length approaches zero if p and q approach a common limit. Distinct segments do not intersect except at a common end point. The points of these segments with their obvious natural topology make up J , the symmetric join of K . This space arises in [4]¹ in connection with the problem of finding the chords of a manifold that are orthogonal to the manifold.

2. **The subdivision of J .** Let the mid-point of pq be denoted by $\Delta p \times q = \Delta q \times p$, and let $p = \Delta p \times p$. These points $\Delta p \times q$ make up the symmetric product S of K . Let the mid-point of the segment from p to $\Delta p \times q$ be denoted by $p \times q$, and let $p = p \times p$. These points $p \times q$ make up the topological product $P = K \times K$. Consider the closed segment of pq from $p \times q$ to $q \times p$, it being understood that this segment is the point p when $p = q$. All such segments form the "neighborhood" N_S . Clearly N_S can be homotopically deformed in N_S along the segments pq upon S with S remaining pointwise invariant. Finally consider the closed segment of pq from p to $p \times q$, it being understood that this segment is the point p when $p = q$. All such segments form the "neighborhood" N_K . Clearly N_K can be homotopically deformed in N_K along the segments pq upon K with K remaining pointwise invariant.

Received by the editors July 23, 1943.

¹ Numbers in brackets refer to the References at the end of the paper.