

## A DIFFERENTIAL INEQUALITY

R. P. BOAS, JR.

The following theorem was discovered by S. B. Jackson in connection with a problem in differential geometry.

**THEOREM 1.** *If  $f(x)$  is of class  $C^2$  in  $(0, a)$ ,  $f(0) = f'(0) = 0$ , and  $f''(x) \leq K|f'(x)| + L|f(x)|$  in  $(0, a)$ , where  $K$  and  $L$  are constants, then  $f(x) \leq 0$  in some interval  $(0, b)$ .*

If  $f(x)$  is in addition analytic at 0, Theorem 1 becomes trivial, since if  $a_n$  is the first nonvanishing coefficient of the power series of  $f(x)$ ,  $x^{2-n}f''(x)$  approaches a nonzero limit, while  $x^{2-n}f'(x)$  and  $x^{2-n}f(x)$  approach zero, as  $x \rightarrow 0$ , and consequently  $a_n < 0$ .

I shall prove the following more general theorem.

**THEOREM 2.** *If  $f'(x)$  is absolutely continuous in  $(0, a)$ ,  $f(0) = f'(0) = 0$ , and*

$$(1) \quad f''(x) \leq K(x)|f'(x)| + x^{-1}L(x)|f(x)|$$

*almost everywhere in  $(0, a)$ , where  $K(x)$  and  $L(x)$  are non-negative and integrable in  $(0, a)$ , then either  $f(x) \equiv 0$  in some interval  $(0, b)$ , or  $f'(x) < 0$  in  $0 < x < \min(a, c)$ , where  $c$  is such that*

$$(2) \quad \int_0^x \{K(t) + L(t)\} dt < 1, \quad 0 < x < c.$$

Since  $f(0) = 0$ ,  $f(x)$  is negative in  $(0, c)$  when  $f'(x) < 0$  in  $(0, c)$ . Theorem 1 is contained in the special case  $K(t) = K$ ,  $L(t) = Lt$ .

Theorem 2 is the best possible result of its kind; for, if  $\int_0^x K(t) dt$  diverges and  $K(t)$  is positive and continuous in  $t > 0$ , the function  $f(x)$  defined by  $f'(x) = 1/\int_x^1 K(t) dt$ ,  $f'(0) = f(0) = 0$ , is positive in  $x > 0$  and satisfies (1) for all  $x$  such that  $\int_x^1 K(t) dt > 1$ .

Assume that  $f(x)$  is not identically zero in any interval  $(0, b)$ , and write  $M(x) = \max_{0 \leq t \leq x} |f'(t)|$ , so that  $M(x) > 0$  for  $x > 0$  and  $M(x)$  is nondecreasing.

We observe that for  $0 \leq x \leq a$ ,

$$(3) \quad |f(x)| = \left| \int_0^x f'(t) dt \right| \leq xM(x).$$

There are points  $x_n$  such that  $x_n \downarrow 0$  and  $M(x_n) = |f'(x_n)|$ . Suppose that  $f'(x_n) > 0$  for some  $n$ . Let  $(a_n, b_n)$  be the largest interval, contain-

Received by the editors June 1, 1944.

ing  $x_n$ , in which  $f'(x) > 0$ ; since  $f'(x)$  is continuous, there is such an interval, and  $f'(a_n) = 0$ . Consequently we have, using (1) and (3),

$$\begin{aligned} 0 < M(x_n) &= f'(x_n) = \int_{a_n}^{x_n} f''(t) dt \\ &\leq \int_{a_n}^{x_n} \{ K(t)f'(t) + t^{-1}L(t) | f(t) | \} dt \\ &\leq M(x_n) \int_0^{x_n} \{ K(t) + L(t) \} dt. \end{aligned}$$

This leads to a contradiction if  $x_n < c$ , where  $c$  is defined by (2). Hence we must have  $f'(x_n) < 0$  for  $x_n < c$ .

There is a largest interval  $(a_n, b_n)$ , containing  $x_n$ , in which  $f'(x) < 0$ . Suppose first that  $a_n > 0$  for every  $n$ ; then the intervals  $(a_n, b_n)$  are separated by other intervals in which  $f'(x) \geq 0$ , and consequently  $f'(b_n) = 0, b_n \rightarrow 0$ . Then we have for  $a_n < x < b_n$

$$\begin{aligned} (4) \quad 0 < -f'(x) &= \int_x^{b_n} f''(t) dt \\ &\leq \int_x^{b_n} \{ -K(t)f'(t) + t^{-1}L(t) | f(t) | \} dt \\ &\leq M(b_n) \int_0^{b_n} \{ K(t) + L(t) \} dt. \end{aligned}$$

Since  $f'(b_n) = 0$ , there is a point  $x'_n$  in  $(x_n, b_n)$  such that  $-f'(x'_n) = M(b_n)$ . Taking  $x = x'_n$  in (4), we have

$$(5) \quad 0 < M(b_n) \leq M(b_n) \int_0^{b_n} \{ K(t) + L(t) \} dt,$$

and again there is a contradiction when  $b_n < c$ .

Hence  $a_n = 0$  for some  $n$ . If  $b_n \geq a$ , we have  $f'(x) < 0$  in  $0 < x < a$ ; if  $b_n < a$ , we have  $f'(b_n) = 0$ , and then (4) and (5) hold. But (5) is contradictory unless  $b_n \geq c$ . Hence we have  $f'(x) < 0$  in  $0 < x < \min(a, c)$ .

We have incidentally established the following result about a function and its first derivative.

**THEOREM 3.** *If  $f(x)$  is absolutely continuous in  $(0, a)$ ,  $f(0) = 0$ , and  $f'(x) \leq K(x)|f(x)|$ , where  $K(x)$  is non-negative and integrable in  $(0, a)$ , then either  $f(x) \equiv 0$  in some interval  $(0, b)$ , or  $f(x) < 0$  in  $0 < x < \min(a, c)$ , where  $\int_0^x K(t) dt < 1$  if  $x < c$ .*

HARVARD UNIVERSITY