

A REMARK ON A RESULT DUE TO BLICHFELDT

LOO-KENG HUA

Let $\sigma \geq 1$ and ξ_1, \dots, ξ_n be $n \geq 3$ linear forms of the real variables x_1, \dots, x_n of nonvanishing determinant Δ . For simplicity's sake we assume $|\Delta| = 1$. Let $2s$ of the forms be pairwise conjugate complex and the remaining $n - 2s$ be real. Then

$$|\xi_1|^\sigma + \dots + |\xi_n|^\sigma \leq 1$$

defines a symmetric convex body in the x -space, the volume $V(\sigma)$ of which equals

$$2^n \cdot \frac{\{\Gamma(1 + \alpha)\}^{n-2s} \{\pi\Gamma(1 + 2\alpha)/2^{1+2\alpha}\}^s}{\Gamma(1 + n\alpha)} \quad (\alpha = 1/\sigma).$$

Minkowski's principle states that there is a lattice point $(x_1, \dots, x_n) \neq (0, \dots, 0)$ satisfying the inequality

$$|\xi_1|^\sigma + \dots + |\xi_n|^\sigma \leq r^\sigma$$

provided

$$(1) \quad r^n \geq 2^n V^{-1}(\sigma).$$

By means of Blichfeldt's method, van der Corput and Schaake¹ obtained a sharpening of this result for $\sigma \geq 2$. Decisive in this procedure is an inequality of the following form

$$(2) \quad \sum_{p,q=1}^k |z_p - z_q|^\sigma \leq \epsilon(\sigma) k \cdot \sum_{p=1}^k |z_p|^\sigma,$$

where the factor $\epsilon(\sigma)$ depends neither on the arbitrary complex numbers z_p nor on k . Once such an inequality is known, (1) may be replaced by

$$(3) \quad r^n \geq (\epsilon(\sigma))^{n/\sigma} \cdot \frac{n + \sigma}{\sigma} \cdot V^{-1}(\sigma).$$

The elementary relation

$$|u - v|^\sigma \leq 2^{\sigma-1} (|u|^\sigma + |v|^\sigma)$$

(following from the fact that x^σ is a convex function of $x > 0$) implies (2) with $\epsilon(\sigma) = 2^\sigma$. Substituted in (3) this does not improve, but on

Received by the editors February 24, 1945.

¹ Acta Arithmetica vol. 2 (1936) pp. 152-160.

the contrary worsens, Minkowski's inequality. However, van der Corput and Schaake obtained the better value $2^{\sigma-1}$ for $\sigma \geq 2$. I shall show here that $\epsilon(\sigma) = 2$ is a legitimate choice for $1 \leq \sigma \leq 2$ and that both facts follow almost immediately from Marcel Riesz's convexity theorem.

Indeed, specialize this theorem (Theorem 296 on p. 219 of Hardy, Littlewood and Pólya's *Inequalities*) by taking $\gamma = \alpha$ and the X as the linear forms $X_{pq} = z_p - z_q$. It then turns out that the logarithm of the maximum $M_k(\alpha)$ of

$$\left\{ \sum_{p,q=1}^k |z_p - z_q|^{1/\alpha} / k \sum_{p=1}^k |z_p|^{1/\alpha} \right\}^\alpha$$

for fixed k and variable z_1, \dots, z_k is a convex function of α in the interval $0 \leq \alpha \leq 1$. One readily verifies that

$$M_k(0) = 2, \quad M_k(1/2) = 2^{1/2}, \quad M_k(1) = 2(1 - 1/k) \leq 2.$$

As

$$\left\{ \sum_p |z_p|^{1/\alpha} \right\}^\alpha \rightarrow \max |z_p| \quad \text{for } \alpha \rightarrow 0,$$

the first equation follows from $\max |z_p - z_q| \leq 2 \cdot \max |z_p|$ together with the observation that the upper bound 2 is attained for $z_1 = 1, z_2 = -1, z_3 = \dots = z_k = 0$. Similarly the two other equations are immediate consequences of the elementary inequalities

$$\begin{aligned} \sum_{p,q} |z_p - z_q|^2 &= 2k \sum_p |z_p|^2 - 2 \left| \sum_p z_p \right|^2 \leq 2k \sum_p |z_p|^2, \\ \sum_{p \neq q} |z_p - z_q| &\leq \sum_{p \neq q} (|z_p| + |z_q|) = 2(k-1) \sum_p |z_p|, \end{aligned}$$

and the corresponding obvious observations about the z_p for which the upper bound is reached.

Let us use 2 as the basis of our logarithms. Then the values of $\log_2 M_k(\alpha)$ are 1, 1/2 and less than or equal to 1 for $\alpha = 0, 1/2, 1$ respectively, and hence the broken line consisting of $1 - \alpha$ for $0 \leq \alpha \leq 1/2$ and α for $1/2 \leq \alpha \leq 1$ gives an upper bound for the convex function $\log_2 M_k(\alpha)$. We thus obtain the promised result that (2) holds with

$$(4) \quad \epsilon(\sigma) = 2^{\sigma-1} \quad \text{for } \sigma \geq 2 \quad \text{and} \quad \epsilon(\sigma) = 2 \quad \text{for } 1 \leq \sigma \leq 2.$$

Both choices are the best possible of their kinds, as, for $0 \leq \alpha \leq 1/2$, is shown by the example $k = 2, z_1 = -z_2 = 1$, and, for $1/2 \leq \alpha \leq 1$, by the example $z_1 = -z_2 = 1, z_3 = \dots = z_k = 0$, with large k .

Consider the case $1 \leq \sigma \leq 2$. If we substitute the value $\epsilon(\sigma) = 2$ in (3), we shall find that it does not always improve Blichfeldt's known inequality, in particular not for the most interesting case $\sigma = 1$. We observe that

$$\left(\frac{|\xi_1|^\sigma + \cdots + |\xi_n|^\sigma}{n} \right)^{1/\sigma}$$

is an increasing function of the exponent σ , while the upper bound for its lattice minimum as derived from (3), namely,

$$(5) \quad \left(\frac{2}{n} \right)^{1/\sigma} \left(\frac{n + \sigma}{\sigma} \right)^{1/n} (V(\sigma))^{-1/n},$$

is not. For $s=0$ the expression (5) tends to a limit with $n \rightarrow \infty$, namely

$$\frac{1}{2} \left(\frac{2}{\sigma e} \right)^{1/\sigma} \Big/ \Gamma \left(1 + \frac{1}{\sigma} \right) = 2^{\alpha-1} \left(\frac{\alpha}{e} \right)^\alpha \Big/ \Gamma(1 + \alpha).$$

The logarithmic derivative of this function with respect to α is negative for $\alpha = 1/2$ and positive for $\alpha = 2/3$, and hence this function has a minimum between $\sigma = 2$ and $\sigma = 1.5$; numerical computation gives as its location $\sigma = \sigma_0 = 1.8653 \cdots$.² At this point the value of the function is

$$\leq 1/(3.146e)^{1/2}$$

which is slightly better than the constant

$$1/(\pi e)^{1/2}$$

due to Blichfeldt.³

In conclusion, for $2 \geq \sigma \geq \sigma_0$, (1) may be replaced by

$$r^n \geq 2^{n/\sigma} \left(\frac{n + \sigma}{\sigma} \right) V^{-1}(\sigma),$$

and, for $1 \leq \sigma \leq \sigma_0$, (1) may be replaced by

$$r^n \geq 2^{n/\sigma_0} \left(\frac{n + \sigma_0}{\sigma_0} \right) V^{-1}(\sigma_0).$$

This would be true however σ_0 were chosen within the limits $1 \leq \sigma_0 \leq 2$; our special choice approaches the best possible for $n \rightarrow \infty$ (and $s=0$) and is sharp enough to beat Blichfeldt's record by a slight margin, even for small n .

NATIONAL TSING HUA UNIVERSITY

² The author is indebted to Mr. Sze for this numerical value.

³ Trans. Amer. Math. Soc. vol. 15 (1914) pp. 227-235.