

has a topology much like before. Thus, if S were originally metric, it would remain metric. This concept permits use of known properties of non-compact continuous curves in certain problems concerning non-locally connected spaces. Several examples of this technique are given, one of which is the following: Let M be a set which contains no arc, let D be a dendrite, and let G be a minimal collection of arcs such that each point of M is joined to D by an arc of G . Then if $M+D+G^*$ contains no simple closed curve, it has the fixed-point property. In order to obtain this last, a study is made of the properties of a type of generalized dendrite. (Received July 30, 1945.)

NEW PUBLICATIONS

- BIEBERBACH, L. Lehrbuch der Funktionentheorie. Vol. 1. Reprint of 4th ed. 14+322 pp. Vol. 2. Reprint of 2d ed. 6+370 pp. New York, Chelsea, 1945. \$3.25 each.
- CARTER, G. W. The simple calculation of electrical transients. New York, Macmillan; Cambridge University Press, 1945. 8+120 pp. \$1.75.
- DONNAY, J. D. H. Spherical trigonometry after the Cesàro method. New York, Interscience, 1945. 12+84 pp. \$1.75.
- KAGAN, V. F. Lobachevsky. Academy of Sciences of the U.S.S.R., 1944. 348 pp.
- SMITH, C. E. Applied mathematics for radio and communication engineers. New York and London, McGraw-Hill, 1945. 10+336 pp. \$3.50.
- TCHEBYCHEV, P. L. Collected works. Vol. 1. Academy of Sciences of the U.S.S.R., 1944. 342 pp.