

336. H. E. Salzer: *Further remarks on the approximation of numbers as sums of reciprocals.*

The present work consists of three main parts. (1) Comparison of R -expansions with simple continued fractions for rational numbers a/b leads to the analogue of the Euclidean algorithm, with a multiple of the g.c.d. of a and b in place of the g.c.d. One practical result is that for a/b , as a rule, fewer partial fractions are required in the R -expansion than in the s.c.f. (2) Proof that if p/q is an approximation to x obtained by the R -expansion, then the remainder $(x - p/q) < 1/q$. (3) Proof that if p/q is an approximation to x obtained by the \bar{R} -expansion, then $|x - p/q| < 1/2q$ except when $x = 3/4$, when the \leq relation may hold. Both theorems are best possible ones. It is shown that when in an R - or \bar{R} -expansion the denominator of the $(n+1)$ th partial fraction is about k times the minimum value that could arise in an R - or \bar{R} -expansion, then not only the closeness of the n th, but of all ensuing approximations p/q will be about $1/kq$ or $1/2kq$ respectively. A critical bibliography is provided, which reviews all work by other authors on R - or \bar{R} -expansions. (Received June 18, 1947.)

337. H. E. Salzer: *Polynomials of best approximation in an infinite interval.* Preliminary report.

Chebyshev polynomials $C_n(x)$ are useful for approximating polynomials of high degree in a finite interval $[a, b]$, by polynomials of much lower degree, because of the property that of all polynomials with leading coefficient 1, $C_n(x)$ has the least value of the greatest deviation from 0 in the interval $[-1, 1]$. To approximate functions of the form $e^{-x}p(x)$ and $e^{-x^2}q(x)$, where $p(x)$, $q(x)$ are polynomials, over $[0, \infty]$ and $[-\infty, \infty]$ respectively, it is useful to know: (I) polynomials $P_n(x)$, degree n , leading coefficient 1, such that the greatest absolute value of $e^{-x}P_n(x)$ differs least from 0 in $[0, \infty]$; (II) polynomials $Q_n(x)$, degree n , leading coefficient 1, such that the greatest absolute value of $e^{-x^2}Q_n(x)$ differs least from 0 in $[-\infty, \infty]$. $P_n(x) \equiv x^n + a_{n-1}x^{n-1} + \dots + a_0$ satisfies $2n$ transcendental equations $P_n'(x_i) = P_n(x_i)$, $e^{-x_i}P_n(x_i) = (-1)^i a_0$, $i = 1, \dots, n$, where x_i are the abscissae of the extrema of $e^{-x}P_n(x)$. $Q_n(x) \equiv x^n + b_{n-1}x^{n-1} + \dots + b_0$ satisfies $2n+1$ equations $Q_n'(x_i) = 2x_i Q_n(x_i)$, $e^{-x_i^2}Q_n(x_i) = (-1)^{i-1} e^{-x_i^2} Q_n(x_1)$, $i = 1, \dots, n, n+1$. When suitably normalized, $P_n(x)$ and $Q_n(x)$ are characterized by being tangent alternately to $\pm e^x$ and $\pm e^{x^2}$ respectively. $Q_n(x) \equiv P_{n/2}(x^2)$ for n even, and is an odd function for n odd. (Received July 11, 1947.)

GEOMETRY

338. W. R. Utz: *The properties of geodesics on certain n -dimensional manifolds.* Preliminary report.

Let S denote the interior of an $(n-1)$ -dimensional unit sphere in Euclidean n -space and let G denote a properly discontinuous group of homeomorphisms of S onto itself that preserve the hyperbolic metric $\int(dx_i dx_i)^{1/2}/(1-x_i x_i)$. The action of this group is discussed in a manner similar to that of Poincaré (*Théorie des groupes Fuchsien*s, Acta Math. vol. 1 (1882) pp. 1-62) for the case $n=2$. By the identification of congruent points of S under G an n -dimensional manifold, Σ , is secured. An investigation of the geodesics on Σ leads to results concerning geodesics and hyperbolic lines of the same type, asymptotic geodesics and limit geodesics analogous to certain results of Morse (*A fundamental class of geodesics on any closed surface of genus greater than one*, Trans. Amer. Math. Soc. vol. 26 (1924) pp. 25-60) in the case of certain two-dimensional manifolds. (Received May 24, 1947.)