

## RESEARCH PROBLEMS

### 1. Paul Slepian: *Problems on polynomials.*

(1) Let  $0 < A < 1$ . Let  $B$  be the set of all positive integers  $n$  such that there exist  $n$  positive numbers  $a_1, a_2, \dots, a_n$  such that the polynomial

$$(x^2 - 2Ax + 1) \prod_{i=1}^n (x + a_i)$$

has all non-negative coefficients. It is known that  $B$  is nonempty. (See P. M. Lewis, *The concept of the one in voltage transfer synthesis*, IRE Trans. Vol. CT-2, pp. 316-319, December, 1955.) Find the least element of  $B$ .

(2) Let  $0 < A < 1$  and let  $N$  be the smallest integer in  $B$ , as described in (1) above. Does there exist  $b > 0$  such that

$$(x^2 - 2Ax + 1)(x + b)^N$$

has all non-negative coefficients?

(3) Generalize the questions raised in (1) and (2) above to the case where the factor  $x^2 - 2Ax + 1$  is replaced by an arbitrary real polynomial, say  $\sum_{i=0}^m C_i x^i$ , having no positive real roots. (Received September 13, 1957.)

### 2. Louis Weinberg: *Decomposition of Hurwitz polynomials.*

Let  $q(s) = \sum_{k=0}^n a_k s^k$  represent a Hurwitz polynomial with real coefficients, i.e., all of its zeros have negative real parts. Can  $q(s)$  be divided into the arithmetic sum of two polynomials,

$$q(s) = q_1(s) + q_2(s)$$

each of which has positive coefficients and only nonpositive real roots? This can easily be done in particular cases; for example, if  $q(s) = (s^2 + 2s + 5)(s + 4) = s^3 + 6s^2 + 13s + 20$ , then  $q_1(s) = s^3 + 6s^2 + 11s + 6 = (s + 1)(s + 2)(s + 3)$  and  $q_2(s) = 2s + 14$ . If this can be shown to be impossible in the general case, can the decomposition always be carried out with three polynomials,

$$q(s) = q_1(s) + q_2(s) + q_3(s),$$

each of which again has positive coefficients and only nonpositive real roots? (Received September 19, 1957.)

### 3. R. E. Bellman: *Number theory. I.*

The problem of generating the integer solutions of the equation  $x^2 + y^2 = 1 \pmod{p}$  by means of the formula  $x_n = \cos n\theta$ ,  $y_n = \sin n\theta$ , where  $(x_1, y_1)$  is a fundamental solution which we can write symbolically in the form  $x_1 = \cos \theta_1$ ,  $y_1 = \sin \theta_1$ , has been extensively studied. What are the corresponding results for the equations  $x_1^2 + x_2^2 + \dots + x_n^2 = 1 \pmod{p}$ ?

In particular, for the equation  $x_1^2 + x_2^2 + x_3^2 = 1 \pmod{p}$ , what subset of solutions do we obtain by means of the formulas

$$\begin{aligned} x_1 &= \cos k\theta_1 \cos l\theta_2, \\ x_2 &= \cos k\theta_1 \sin l\theta_2, \\ x_3 &= \sin k\theta_1, \end{aligned}$$

where  $k, l = 0, 1, \dots$ , and  $\theta_1, \theta_2$  correspond to certain primitive solutions? (Received May 22, 1957.)

4. R. E. Bellman: *Number theory. II.*

Consider the same type of problem for the multiplicative form

$$x^3 + y^3 + z^3 - 3xyz$$

and for the circulant functions of higher order. (Received May 22, 1957.)

5. R. E. Bellman: *Number theory. III.*

Consider the equation  $y^2 = 4x^3 - g_2x - g_3$  which may be uniformized by means of the Weierstrassian elliptic functions  $x = p(u)$ ,  $y = p'(u)$ . What subset of solutions of the congruence  $y^2 = 4x^3 - g_2x - g_3 \pmod{p}$  can be obtained by means of the formulas  $x = p(mu + nv)$ ,  $y = p'(mu + nv)$ ,  $m, n = 0, 1, 2, \dots$ , (not both zero simultaneously), where  $u$  and  $v$  correspond to certain primitive solutions?

Consider the similar problem for  $y^2 = (1-x^2)(1-k^2x^2)$  which can be uniformized by means of Jacobian elliptic functions. (Received May 22, 1957.)

6. R. E. Bellman: *Number theory. 1.*

Let  $x$  be an irrational number in  $[0, 1]$  and let  $g(y; a, b)$ ,  $0 \leq a < b \leq 1$  be a periodic function of  $y$  with period 1 defined by the conditions  $g(y; a, b) = 1$ ,  $a \leq y \leq b$ ,  $g(y; a, b) = 0$  elsewhere for  $y$  in  $[0, 1]$ . Define the function

$$f_N(z, x) = g(z; a, b) + g(z+x; a, b) + \dots + g(z+Nx; a, b)$$

for  $1 \geq z \geq 0$ , equal to the number of elements of the finite sequence  $\{nx+z\}$ ,  $n = 0, 1, \dots, N$ , falling inside  $[a, b]$ , modulo one.

The Weyl equidistribution theorem asserts that  $f_N(z, x)/(N+1) \sim b-a$  as  $N \rightarrow \infty$ . It is easy to show via Fourier series that

$$\int_0^1 \int_0^1 [f_N(z, x) - (N+1)(b-a)]^2 dz dx \sim (N+1)c_1(a, b)$$

as  $N \rightarrow \infty$ .

This suggests that the quantity

$$u_N(z, x) = \frac{f_N(z, x) - (N+1)(b-a)}{(N+1)^{1/2}}$$

possesses asymptotic moments of all orders.

Does

$$\lim_{N \rightarrow \infty} \int_0^1 \int_0^1 \left( \frac{f_N(z, x) - (N+1)(b-a)}{(N+1)^{1/2}} \right)^{2k} dz dx,$$

$k=2, 3, \dots$  exist, and if so what is the limiting distribution? (Received July 15, 1957.)

7. R. E. Bellman: *Number theory. 2.*

Consider the same problem for the function

$$f_N(z, y, x) = g(z; a, b) + g(z+2y+x; a, b) + \dots + g(z+2Ny+N^2x; a, b)$$

with  $x$  irrational,  $y$  and  $z$  in  $[0, 1]$ . As above, it is easy to show via Fourier series that

$$\int_0^1 \int_0^1 \int_0^1 [f_N(z, y, x) - (N+1)(b-a)]^2 dx dy dz \sim (N+1)c_1(a, b)$$

as  $N \rightarrow \infty$ .

Does

$$\lim_{N \rightarrow \infty} \int_0^1 \int_0^1 \int_0^1 \left( \frac{f_N(z, y, x) - (N+1)(b-a)}{(N+1)^{1/2}} \right)^{2k} dx dy dz$$

exist for  $k=1, 2, \dots$ , and if so, what is its value?

There are corresponding versions of this problem for polynomials of all orders.

(Received July 15, 1957.)

#### 8. R. E. Bellman: *Differential equations.*

Consider the second order linear differential equation  $u'' + (1 + \lambda g(x))u = 0$ , where  $\lambda$  is a real constant and  $\int_0^\infty |g(x)| dx < \infty$ . Let  $u_1(x)$  be the solution specified by  $u_1(0) = 0$ ,  $u_1'(0) = 1$ . It is known that  $u \sim r(\lambda) \sin(x + \theta(\lambda))$  as  $x \rightarrow \infty$ .

Taking  $\lambda$  to be complex variable, what are the analytic properties of the functions  $r(\lambda)$  and  $\theta(\lambda)$ ? In particular, where are the singularities nearest the origin?

If  $g(x) > 0$  for  $x \geq 0$ , is the singularity nearest the origin on the negative axis?  
(Received August 16, 1957.)