

APPROXIMATION OF SMOOTH FUNCTIONS

BY G. G. LORENTZ¹

Communicated by R. P. Boas, January 13, 1960

Theorems of Jackson and S. Bernstein about the approximation of smooth functions are usually interpreted in the way that all functions with a prescribed degree of smoothness have a definite degree of approximation. They can be viewed in another way, which reveals their susceptibility to generalization.

Let $\omega(h)$ be an increasing continuous subadditive function defined for $h \geq 0$ with $\omega(0) = 0$, A a compact metric space with infinitely many points. By C_1^ω we denote the set of all real valued functions f on A with $|f(x)| \leq 1$, $|f(x) - f(x')| \leq \omega(h)$, $h = \rho(x, x')$. If A is a q -dimensional cube, p a natural number and $0 < \alpha \leq 1$, we denote by $C_1^{p+\alpha}$ the set of all functions on A with continuous partial derivatives of orders not exceeding p and bounded by 1, and with the derivatives of order p satisfying a Lipschitz condition of order α and with coefficient 1. Let $G = \{g_n\}$ be a sequence of continuous functions on A . Then, with some norm, for example the uniform norm on A ,

$$E_n(f) = E_n^G(f) = \inf \left\| f - \sum_{i=1}^n a_i g_i \right\|$$

is the degree of approximation of f by linear combinations of g_1, \dots, g_n ; and

$$\varepsilon_n(W) = \sup_{f \in W} E_n(f)$$

is the degree of approximation of a class W .

The theorems of Jackson and Bernstein state that for periodic $f \in C_1^{p+\alpha}$, and the trigonometric approximation, $E_n(f)$ has the exact order $n^{-(p+\alpha)/q}$; exceptions occur only if f has a higher degree of smoothness. We regard this as a statement about a certain massivity of $C_1^{p+\alpha}$, which prevents better approximation by linear combinations of only n functions. One can hope that an estimate of $\varepsilon_n(C_1^{p+\alpha})$ from below can be given for an arbitrary system G , and that the trigonometric system is close to the best possible. That this is true, is shown by the following results:

THEOREM 1. *Let A be a compact metric space, and $\delta = \delta(n)$ the largest*

¹ This research was in part supported by the United States Air Force under Contract No. AF49(638)-619 monitored by the AFOSR of the Air Research and Development Command.

number such that there exist n points of A with mutual distances $\geq \delta$. Then for each G ,

$$\varepsilon_n(C_1^\omega) \geq \frac{1}{2} \omega(\delta(n+1)).$$

This cannot be essentially improved, for there exists a G with $\varepsilon_n(C_1^\omega) \leq \omega(\delta_1)$, if A can be covered by n open balls of radius $\delta_1 = \delta_1(n)$.

THEOREM 2. *If A is a q -dimensional cube, then for some constant B , and each G , in the uniform and the L^1 norm*

$$(1) \quad \varepsilon_n(C_1^{p+\alpha}) \geq Bn^{-(p+\alpha)/q}, \quad p = 0, 1, \dots; 0 < \alpha < 1.$$

From these and similar theorems one can obtain by a method of condensation of singularities:

THEOREM 3. *If A is as in Theorem 2, $p=0, 1, \dots$ and $0 < \alpha < 1$, then there exists a constant B such that for each system G one can find a function $f_0 \in C_1^{p+\alpha}$ such that, in the uniform and the L^1 norm,*

$$E_n(f_0) \geq Bn^{-(p+\alpha)/q}$$

for an infinite number of values of n .

THEOREM 4. *Let A_ρ be the ellipse with foci $-1, +1$ and the sum of the half-axes 2ρ . For each G and each sequence $\epsilon_n \rightarrow 0$ there exists a function $f_0(z)$, analytic inside A_ρ , with $|f_0(z)| \leq 1$ such that the degree of approximation of f_0 on $(-1, +1)$ satisfies, in the L^1 norm,*

$$(2) \quad E_n(f_0) \geq \epsilon_n \rho^{-n}$$

for infinitely many n .

Similar and more general results were recently obtained by A. G. Vituškin [1]. However, his lower bounds (for $s = n+1, m = 0$ in [1]) are of orders $(n \log n)^{-(p+\alpha)/2q}$ and $n^{-a_n}, a > 0$, for the problems of types (1), and (2), respectively.

REFERENCE

1. A. G. Vituškin, *Best approximation of differentiable and analytic functions*, Dokl. Akad. Nauk SSSR vol. 119 (1958) pp. 418-420.

SYRACUSE UNIVERSITY