

## AREA OF DISCONTINUOUS SURFACES

BY CASPER GOFFMAN<sup>1</sup>

Communicated by J. W. T. Youngs, January 19, 1961

1. A general theory of surface area, [1; 2], exists for the non-parametric case. Thus, area is defined for all measurable  $f$  on the unit square  $Q = I \times J$ . The area functional is lower semi-continuous with respect to almost everywhere convergence and agrees with the Lebesgue area for continuous  $f$ . On the other hand, for continuous parametric mappings  $T$  of the closed unit square  $Q$  into euclidean 3-space  $E_3$ , Lebesgue area is not lower semi-continuous with respect to almost everywhere convergence nor even, as C. J. Neugebauer has shown, [3], with respect to pointwise convergence.

It thus appears that a theory of parametric surface area must be restricted to surfaces which cannot deviate too far from the ones given by continuous mappings. In this paper, we develop the beginnings of a theory for a class of surfaces which we call linearly continuous.

2. Let  $f$  be a real function defined on  $Q$  and, for every  $u$ , let  $f_u$  be defined by  $f_u(v) = f(u, v)$  and let  $f_v$  be defined similarly. Then  $f$  is linearly continuous if  $f_u$  is continuous for almost all  $u$  and  $f_v$  is continuous for almost all  $v$ . A mapping  $T: x = x(u, v), y = y(u, v), z = z(u, v)$  of  $Q$  into  $E_3$  is linearly continuous if  $x, y, z$  are linearly continuous.

A sequence  $\{f_n\}$  of functions converges linearly to a function  $f$  if  $(f_n)_u$  converges uniformly to  $f_u$  for almost all  $u$ , and  $(f_n)_v$  converges uniformly to  $f_v$  for almost all  $v$ . A sequence  $T_n: x = x_n(u, v), y = y_n(u, v), z = z_n(u, v)$  converges linearly to a mapping  $T: x = x(u, v), y = y(u, v), z = z(u, v)$  if  $\{x_n\}, \{y_n\}, \{z_n\}$  converge linearly to  $x, y, z$ , respectively.

Let  $P$  be the set of quasi linear mappings from  $Q$  into  $E_3$ . For  $p, q \in Q$  let

$$d(p, q) = \inf[k: \text{there are sets } A_k \subset I, B_k \subset J, \\ m(A_k) > 1 - k, m(B_k) > 1 - k, \text{ and } |p(u, v) - q(u, v)| < k \\ \text{on } (A_k \times J) \cup (I \times B_k)].$$

It is easy to verify that  $P$  is a metric space and that  $\{p_n\}$  converges to  $p$  in this space if and only if it converges linearly. Let  $E$  be the elementary area functional on  $P$ . It is not hard to prove

---

<sup>1</sup> Research supported by National Science Foundation Grant No. NSF G-5867.

THEOREM 1.  $E$  is lower semi-continuous on  $P$ . In other words, if  $\{p_n\}$  converges linearly to  $p$  then  $\liminf E(p_n) \geq E(p)$ .

By the Fréchet extension theorem,  $E$  is extended to a lower semi-continuous functional  $\Phi$  on the completion  $\mathcal{L}$  of  $P$ .

THEOREM 2. The completion  $\mathcal{L}$  of  $P$  is the space of linearly continuous mappings with the metric corresponding (as above) to linear convergence.

3. It is obvious that for every continuous mapping  $T$ ,  $A(T) \geq \Phi(T)$  where  $A(T)$  is the Lebesgue area. The inverse inequality holds so that the functional  $\Phi$  constitutes a legitimate extension of Lebesgue area to substantially wider class of mappings than the continuous ones. We outline the proof.

For a continuous  $T: x = x(u, v)$ ,  $y = y(u, v)$ ,  $z = z(u, v)$ , the lower area  $V(T)$  is defined as follows:

Let  $T_1: y = y(u, v)$ ,  $z = z(u, v)$ ,  $T_2: x = x(u, v)$ ,  $z = z(u, v)$ , and  $T_3: x = x(u, v)$ ,  $y = y(u, v)$  be the associated flat mappings. For every simple polygonal region  $P$  in  $Q^0$ , let

$$v_1(P) = \int |O(\xi, T_1 P^*)|,$$

where the integration is over the  $yz$  plane, and  $O(\xi, T_1 P^*)$  is the topological index of  $T_1 P^*$  at  $\xi$  ( $A^0$  and  $A^*$  are the interior and boundary, respectively, of a set  $A$ ). Define  $v_2(P)$  and  $v_3(P)$ , similarly, and let

$$v(P) = [v_1(P)^2 + v_2(P)^2 + v_3(P)^2]^{1/2}.$$

Let  $\pi = (P_1, \dots, P_n)$  be a finite set of pair-wise disjoint simple polygonal regions in  $Q^0$  and

$$v(\pi) = \sum_{i=1}^n v(P_i).$$

Finally, let

$$V(T) = \sup[v(\pi): \pi].$$

Cesari has shown (e.g. [4]) that  $A(T) = V(T)$  for every continuous  $T$ .

The distance between 2 sets  $A$  and  $B$  is defined by

$$d(A, B) = \sup[d(x, B): x \in A] + \sup[d(y, A): y \in B].$$

With this metric, the set  $\alpha$  of simple polygonal regions is a separable metric space. Let  $\beta \subset \alpha$  be dense in  $\alpha$  and

$$V_\beta = \sup[v(\pi): \pi \subset \beta].$$

LEMMA 1.  $V_\beta(T) = V(T)$ .

Now, let  $\{T_n\}$  be a sequence of continuous mappings which converges linearly to a continuous mapping  $T$ . Let  $\gamma$  be the set of simple polygonal regions whose boundaries consist of line segments parallel to the coordinate axes for which  $T$  and  $T_n, n = 1, 2, \dots$  are continuous and on each of which  $\{T_n\}$  converges uniformly to  $T$ . For each  $\pi \subset \gamma$ ,  $\liminf v(\pi, T_n) \geq v(\pi, T)$ . Since  $\gamma$  is dense in  $\alpha$ , it follows that  $\liminf V(T_n) \geq V(T)$ . This proves

THEOREM 3.  $A(T)$  is lower semi-continuous with respect to linear convergence on the set of continuous mappings.

COROLLARY 1.  $A(T) = \Phi(T)$  for every continuous  $T$ .

PROOF. For every sequence  $\{p_n\}$  of quasi-linear mappings converging linearly to  $T$ ,  $\liminf E(P_n) \geq A(T)$ . Choose  $\{p_n\}$  so that  $\lim E(p_n) = \Phi(T)$ . Then  $A(T) \leq \Phi(T)$ ,

4. A set  $S$  will be called negligible if  $S \subset Z_1 \times Z_2$  where  $Z_1$  and  $Z_2$  have linear measure zero. Kolmogoroff's principle holds in the following form.

THEOREM 4. If  $T_1$  and  $T_2$  are linearly continuous mappings from  $Q$  into  $E_3$  and if for every pair of points  $\xi, \eta$  not belonging to a negligible set

$$|T_1\xi - T_1\eta| \leq |T_2\xi - T_2\eta|,$$

then  $\Phi(T_1) \leq \Phi(T_2)$ .

5. A real function  $f$  on  $Q$  is BVC if for almost all  $u$  and almost all  $v, f_u$  and  $f_v$  are equivalent to functions of bounded variation and the corresponding variation functions are summable.  $f$  is ACE if for almost all  $u$  and almost all  $v, f_u$  and  $f_v$  are equivalent to absolutely continuous functions.

For functions which are BVT and ACT it is a simple known fact that the integral means commute with the partial derivatives. This also holds almost everywhere for functions which are BVC and ACE. Using this fact and the fact, [5], that if  $f$  is BVC and linearly continuous then the integral means of  $f$  converge linearly to  $f$ , the proof of the following generalization of a theorem of Morrey, [4], may be obtained in somewhat standard fashion. The generalization is in two directions. Instead of holding only for conjugate Lebesgue spaces, the theorem holds for conjugate Köthe spaces, [6; 7], and the theorem

holds for linearly continuous mappings rather than just for continuous ones.

**THEOREM 5.** *If the functions  $x, y, z$  of a linearly continuous  $T$  are BVC and ACE and if the pairs of partial derivatives  $(x_u, y_v), (x_v, y_u), (x_u, z_v), (x_v, z_u), (y_u, z_v), (y_v, z_u)$  belong to conjugate Köthe spaces, the area  $\Phi(T)$  is given by the formula*

$$\Phi(T) = \int J \, dudv$$

where  $J = [J_1^2 + J_2^2 + J_3^2]^{1/2}$  and  $J_1, J_2, J_3$  are the jacobians of  $T_1, T_2, T_3$ , respectively.

6. We define an equivalence relation for linearly continuous mappings.  $T$  is equivalent to  $T'$  ( $T \approx T'$ ) if there are sequences  $\{p_n\}$  and  $\{q_n\}$  of quasi linear mappings such that, for every  $n$ ,  $p_n \approx q_n$  in the Lebesgue sense and  $\{p_n\}$  converges linearly to  $T$ ,  $\{q_n\}$  converges linearly to  $T'$ .

The following simple facts hold:

- (a) The relation " $\approx$ " has the properties of an equivalence relation.
- (b) If  $T$  and  $T'$  are continuous and Fréchet equivalent then  $T \approx T'$ .
- (c) If  $T \approx T'$  then  $\Phi(T) = \Phi(T')$ .

We refer to an equivalence class as a surface and to its elements as representations.

$D$  mappings, the Dirichlet integral, and almost conformal mappings are defined as for the continuous case, [4], with BVT and ACT replaced by BVC and ACE.

We say that a mapping  $T$  is simple if there is a negligible set  $S$  such that  $\xi \in Q - S, \eta \in Q - S, \xi \neq \eta$  implies  $T(\xi) \neq T(\eta)$ .

The following holds:

**THEOREM 6.** *If  $T'$  is a linearly continuous simple mapping and  $\Phi(T') < \infty$ , the surface given by  $T'$  has a representation  $T$ , with jacobian  $J$ , such that*

$$\Phi(T') = \Phi(T) = \int J \, dudv.$$

**COROLLARY.** *Every linearly continuous nonparametric surface of finite area has a parametric representation  $T$ , with jacobian  $J$ , such that*

$$\Phi(T) = \int J \, dudv$$

## BIBLIOGRAPHY

1. L. Cesari, *Sulle funzioni a variazione limitata*, Ann. Scuola Norm. Sup. Pisa vol. 5 (1936) pp. 299–313.
2. C. Goffman, *Lower semi-continuity and area functionals. I, The non-parametric case*, Rend. Circ. Mat. Palermo vol. 2 (1954) pp. 203–235.
3. C. J. Neugebauer, *Lebesgue area and pointwise convergence*, Abstract 554-13, Notices Amer. Math. Soc. vol. 6 (1959) p. 80.
4. L. Cesari, *Surface area*, Princeton, University Press, 1956.
5. C. Goffman, *Area of linearly continuous functions*, Acta Math. vol. 103 (1960) pp. 269–290.
6. J. Dieudonné, *Sur les espaces de Köthe*, J. Anal. Math. vol. 1 (1951) pp. 81–115.
7. C. Goffman, *Completeness in topological vector lattices*, Amer. Math. Monthly vol. 66 (1959) pp. 87–92.

PURDUE UNIVERSITY