

A DUALITY THEORY FOR CONVEX PROGRAMS WITH CONVEX CONSTRAINTS

BY A. CHARNES, W. W. COOPER AND K. KORTANEK¹

Communicated by Edwin Hewitt, April 15, 1962

The existence of a solution to the problem of minimizing a convex function subject to restriction of the variables to a closed convex set in n -space ("convex programming") has been characterized (for suitable differentiability conditions) by the Kuhn-Tucker theorem [5]. In general, no dual programming problem (not involving the variables of the direct problem) has been associated with this situation except in the linear programming case, and very recently by E. Eisenberg in [3], for homogeneity of order one in the function and linear inequality constraints, and by R. J. Duffin [2] in an inverse manner for a highly specialized problem.

Starting with a little known paper of A. Haar [4] in the light of current linear programming constructs (e.g., "regularization" [1]), we effect a generalization of these ideas (with maximal finite algebra and minimal topology) so that a dual theory practically as straightforward as linear programming theory is obtained, and which includes a dual theorem covering the most general convex programming situation (e.g. no differentiability conditions qualifying the convex function or constraints, or homogeneity, etc.).

This general theorem is made possible by associating a suitably restricted, usually infinite-dimensional space problem with the minimization problem in n -space instead of the usual association of another finite m -space problem. The space we use is a "generalized finite sequence space" (g.f.s.s.), defined with respect to an index set I of arbitrary cardinality as the vector space, S , of all vectors $\lambda = [\lambda_i; i \in I]$ over an ordered field F with only finitely many nonzero entries.

Such spaces possess the following key characteristics for linear programming of ordinary n -spaces. Let V be a vector space over F and consider a collection of vectors: $P_0, P_i; i \in I$ in V . Let R be the subspace spanned by these vectors, and let

¹ The research of A. Charnes and K. Kortanek at Northwestern University has been supported by O.N.R. contract Nonr-1228(10); that of W. W. Cooper at Carnegie Institute of Technology has been supported by O.N.R. contract Nonr-760(01). Reproduction in whole or in part is permitted for any purpose of the United States Government.

$$\Lambda = \left\{ \lambda \in S: \sum_{i \in I} \lambda_i P_i = P_0, \lambda \geq 0 \right\}.$$

Clearly Λ is convex in S and we have (assuming V finite-dimensional):

THEOREM 1. $\lambda \neq 0$ is an extreme point of Λ in S if and only if the nonzero coordinates of λ correspond to coefficients of linearly independent vectors in R .

THEOREM 2. Λ is generated by its extreme points if and only if for any $\alpha \in S$, $\alpha \neq 0$, $\sum_{i \in I} \alpha_i P_i = 0$ implies some α_r and some α_s are of opposite signs.

REMARK. Λ need not be bounded as in n -space. ("Bounded" means there exists $M \in F$ such that $\sum_i |\lambda_i| \leq M$ for all λ in the set.)

These theorems can be proved in similar fashion to their finite space forms due respectively to Charnes and to Charnes-Cooper (see [1]).

By "dual semi-infinite programs" we mean the following pair of problems formed from the same data:

<p style="text-align: center;">I</p> $\min u^T P_0$ <p style="text-align: center;">subject to $u^T P_i \geq c_i \quad i \in I$</p>	<p style="text-align: center;">II</p> $\max \sum_{i \in I} c_i \lambda_i$ <p style="text-align: center;">subject to $\sum_{i \in I} P_i \lambda_i = P_0$</p> $\lambda \in S, \lambda \geq 0.$
---	--

We restrict ourselves now to the real field and to semi-infinite programs whose $\{P_i, c_i\}$ are "canonically closed" in the sense that in an equivalent inequality system in which the new $\{P_i, c_i\}$ form a bounded set, e.g. by dividing each inequality by some $d_i > 0$, the set is also closed. We call such programs "dual Haar programs."

We require next the inhomogeneous (inequality system) theorem of Haar [4].

THEOREM 3. Let $u^T P_i \geq c_i$, $i \in I$ be a canonically closed system. If $u^T P \geq c$ holds whenever $u^T P_i \geq c_i$ for all $i \in I$, then there exist $\lambda_k \geq 0$, $\lambda_0 \geq 0$, with at most $n+1$ nonzero such that

$$u^T P - c = \sum_k \lambda_k (u^T P_k - c_k) + \lambda_0.$$

Haar does not specifically use the notion of canonical closure, but as counter-examples show he must have intended something of this sort. By use of Theorem 3 we obtain the following lemma.

LEMMA 1. *For Haar programs if both I and II are consistent, then*

$$\inf u^T P_0 = \sup \sum_{i \in I} u^T P_i \lambda_i = \sum_{i \in I} c_i \lambda_i^* \quad \text{for some } \lambda^* \in \Lambda.$$

Hence we conclude

THEOREM 4 (EXTENDED DUAL THEOREM). *For any pair of dual Haar programs precisely one of the following occurs.*

- (i) $\sup \sum_{i \in I} c_i \lambda_i = \infty$ and I is inconsistent.
- (ii) $\inf u^T P_0 = -\infty$ and II is inconsistent.
- (iii) I and II are both inconsistent.
- (iv) $\inf u^T P_0 = \sup \sum_{i \in I} c_i \lambda_i^*$ for some $\lambda^* \in \Lambda$.

REMARK. Only the Farkas-Minkowski property of Theorem 3 is employed to obtain Theorem 4. Canonical closure is a sufficient but not a necessary condition for this.

To obtain the general convex programming dual theorem, we move the functional into the constraints and replace it with a linear function as follows. Suppose the direct problem is: $\min C(u)$ subject to $G(u) \geq 0$, where $G^T = (\dots, G_i(u), \dots)$ is a finite vector of concave functions which defines the convex set W of the u 's. Let $u^T P_i \geq c_i$, $i \in I$ be a system of supports for W , and $z - u^T Q_\alpha \geq d_\alpha$, $\alpha \in A$ be a system of supports for $z - C(u) \geq 0$. Then the direct problem may be rewritten as:

$$\min z, \quad \text{subject to } z - u^T Q_\alpha \geq d_\alpha, \quad u^T P_i \geq c_i, \quad \alpha \in A, i \in I.$$

Thus we have

THEOREM 5. *Assuming the Farkas-Minkowski property for this system, the extended dual theorem applies to the following dual programs:*

$$\begin{array}{ll} \min z & \max \sum_{\alpha} d_{\alpha} \eta_{\alpha} + \sum_i c_i \lambda_i \\ \text{subject to } z - u^T Q_{\alpha} \geq d_{\alpha} & \text{subject to } \sum_{\alpha} \eta_{\alpha} = 1 \\ & - \sum_{\alpha} Q_{\alpha} \eta_{\alpha} + \sum_i P_i \lambda_i = 0 \\ & \eta_{\alpha}, \lambda_i \geq 0. \end{array}$$

Complete generality may now be obtained since an arbitrary semi-infinite program may be replaced by a Haar program according to the following observation:

THEOREM 6. *The canonical closure $u^T P_i \geq c_i$, $i \in \hat{I}$ of the system $u^T P_i \geq c_i$, $i \in I$, has precisely the same set of solutions $\{u\}$, where $\hat{I} \supseteq I$*

denotes the increased index set to index limit points of the (P_i, c_i) not indexed by I .

REFERENCES

1. A. Charnes and W. W. Cooper, *Management models and industrial applications of linear programming*, Vols. I and II, Wiley, New York, 1961.
2. R. J. Duffin, *Dual programs and minimum cost*, J. Soc. Indust. Appl. Math. **10** (1962), 119–123.
3. E. Eisenberg, *Duality in homogeneous programming*, Proc. Amer. Math. Soc. **12** (1961), 783–787.
4. A. Haar, *Über lineare Ungleichungen*, Acta. Math. Szeged **2** (1924), 1–14.
5. H. W. Kuhn and A. W. Tucker, *Non-linear programming*, Proc. 2nd Berkeley Sympos. Math. Statist. Prob., 1950, pp. 481–492, Univ. of California Press, Berkeley, Calif., 1951.

NORTHWESTERN UNIVERSITY AND
CARNEGIE INSTITUTE OF TECHNOLOGY