

## APPROXIMATION OF BOUNDED FUNCTIONS BY CONTINUOUS FUNCTIONS

BY B. R. KRIPKE AND R. B. HOLMES

Communicated by V. Klee, June 15, 1965

We shall show that every bounded function on a paracompact space has a best approximation by continuous functions, and characterize the functions whose best approximators are unique. This is a special case of a measure-theoretic problem, whose setting is as follows. Let  $X$  be a topological space and  $\mu$  a Borel measure on  $X$  which assigns positive mass to each nonempty open set, and has the property that  $\mu(Y) = 0$  if  $Y$  intersects a neighborhood of each point in a  $\mu$ -null set. The latter condition is automatically fulfilled if each open cover of  $X$  has a countable subcover. Let  $L^\infty$  be the space of essentially bounded real-valued  $\mu$ -measurable functions on  $X$ , and give it the semi-norm  $\|f\| = \text{essential sup } |f|$ . The bounded continuous functions on  $X$  form a closed subspace  $C$  of  $L^\infty$ . We say that  $g \in C$  is a *best approximator* to  $f \in L^\infty$  if  $\|f - g\| = \text{dist}(f, C) = \inf \{\|f - h\| : h \in C\}$ .

If  $f \in L^\infty$  and  $x \in X$ ,  $f^*(x) = \limsup_{y \rightarrow x} f(y) = \inf \{\text{ess sup of } f \text{ over } U : U \text{ is a neighborhood of } x\}$ ;  $f_* = \liminf_{y \rightarrow x} f(y)$  has a similar definition. It is easy to verify that the functions  $f^*$  and  $f_*$  are defined everywhere, and are upper semi-continuous (usc) and lower semi-continuous (lsc) respectively.

**PROPOSITION.** *If  $X$  is any topological space and  $f \in L^\infty$ , then  $2 \text{ dist}(f, C) \geq d(f) \equiv \sup \{f^*(y) - f_*(y) : y \in X\}$ .*

**PROOF.** If  $f^*(x) - f_*(x) > d(f) - \epsilon$  and  $g \in C$  then one or the other of  $\limsup_{y \rightarrow x} (f(y) - g(y))$  and  $\limsup_{y \rightarrow x} (g(y) - f(y))$  is greater than  $\frac{1}{2}(d(f) - \epsilon)$ .

**THEOREM 1.** *If  $X$  is paracompact, then  $g \in C$  is a best approximator to  $f \in L^\infty$  if, and only if,  $f^* - \frac{1}{2}d(f) \leq g \leq f_* + \frac{1}{2}d(f)$ ; every  $f \in L^\infty$  has such a best approximator; and  $\text{dist}(f, C) = 1/2d(f)$ .*

**PROOF.** Since  $f_* + \frac{1}{2}d(f) \geq f^* - \frac{1}{2}d(f)$ , the first pair of inequalities is equivalent to the condition that for every  $\epsilon > 0$  and every  $x \in X$ , there be a neighborhood  $U$  of  $x$  such that  $(\text{ess sup } |f(y) - g(y)| : y \in U) \leq \frac{1}{2}d(f) + \epsilon$ . This in turn is equivalent to the assertion that for every  $\epsilon > 0$ ,  $|f(y) - g(y)| > \frac{1}{2}d(f) + \epsilon$  only on a  $\mu$ -null set, which says that  $\|f - g\| \leq \frac{1}{2}d(f)$ . It remains only to show that there is a continuous function which satisfies these inequalities. Since  $f^* - \frac{1}{2}d(f)$

is usc and  $f_* + \frac{1}{2}d(f)$  is lsc, this follows from the Interposition Theorem of Dieudonné [1, p. 75].

**THEOREM 2.** *If  $X$  is a normal Hausdorff space then an element  $f \in L^\infty$  has exactly one best approximator in  $C$  if, and only if,  $f^* - f_*$  is a constant function.*

**PROOF.** If  $f^* - f_*$  is constant, then the function  $g = f^* - \frac{1}{2}d(f) = f_* + \frac{1}{2}d(f)$  is both lsc and usc, and hence is continuous. As in Theorem 1,  $\|f - g\| = \text{dist}(f, C)$ , and no other element of  $C$  has this property. Conversely, we must show that if  $f^* - f_*$  is not constant and  $f$  has a best approximator  $g$  in  $C$ , then it has more than one. If  $f^* - f_*$  is not constant, we can choose an  $\epsilon > 0$  and an  $x \in X$  such that  $f_*(x) + \frac{1}{2}d(f) - (f^*(x) - \frac{1}{2}d(f)) = \epsilon$ . Since  $g$  is continuous and  $f^*$  and  $f_*$  are semi-continuous, there is a neighborhood  $U$  of  $x$  on which  $|g(y) - g(x)| < \epsilon/6$ ,  $f_*(y) > f_*(x) - \epsilon/6$  and  $f^*(y) < f^*(x) + \epsilon/6$ . Since  $\{x\}$  is closed and  $X$  is normal, Urysohn's Lemma asserts the existence of a non-negative function  $p \in C$  such that  $\|p\| = \epsilon/6$  and that  $p$  vanishes outside  $U$ . One or the other of the inequalities  $f_*(x) + \frac{1}{2}d(f) - \epsilon/2 \geq g(x)$  and  $g(x) \geq f^*(x) - \frac{1}{2}d(f) + \epsilon/2$  must hold, so that either  $f_*(y) + \frac{1}{2}d(f) - \epsilon/6 > g(y)$  or  $g(y) > f^*(y) - \frac{1}{2}d(f) + \epsilon/6$  on  $U$ . According to which is the case, put  $h = g + p$  so that  $g \leq h \leq f_* + \frac{1}{2}d(f)$  on  $U$ , or  $h = g - p$  so that  $f^* - \frac{1}{2}d(f) \leq h \leq g$  on  $U$ ;  $h = g$  on the complement of  $U$ . Then  $h$  is also a best approximator to  $f$  out of  $C$ .

If  $\mu$  is the measure which assigns mass 1 to every point in  $X$ , then it certainly assigns positive mass to each nonempty open set, and mass 0 to each set which intersects a neighborhood of every point in a set of measure 0. In this case,  $L^\infty$  is the Banach space of all bounded functions on  $X$ , and  $\|\cdot\|$  is the supremum norm. Theorems 1 and 2 thus solve, as a special case, the problem of approximating bounded functions by continuous functions in the uniform norm.

#### REFERENCE

1. J. Dieudonné, *Une généralisation des espaces compacts*, J. Math. Pures Appl. **23** (1944), 65-76.

UNIVERSITY OF CALIFORNIA, BERKELEY AND  
WRIGHT-PATTERSON AIR FORCE BASE, OHIO