

2. Jean Geffroy, *Contribution à la théorie des valeurs extrêmes*, Publ. Inst. Statist. Univ. Paris 7 (1958), 36–123; 8 (1959) 3–52.
3. ———, *Localisation asymptotique du polyèdre d'appui d'un échantillon Laplacien à  $k$  dimensions*, Publ. Inst. Statist. Univ. Paris 10 (1961), 213–228.
4. B. V. Gnedenko, *Sur la distribution limite du terme maximum d'une série aléatoire*, Ann. of Math. 44 (1943), 423–453.
5. Ulf Grenander, *Probabilities on algebraic structures*, Wiley, New York, 1963.
6. A. Rényi and R. Sulanke, *Über die konvexe Hülle von  $n$  zufällig gewählten Punkten I and II*, Z. Wahrscheinlichkeitstheorie und Verw. Gebiete 2 (1963), 75–84; 3 (1964), 138–148.
7. A. V. Skorohod, *Limit theorems for stochastic processes*, Theor. Probability Appl. 1 (1956), 261–290. (Engl. transl.)
8. V. Strassen, *An invariance principle for the law of the iterated logarithm*, Z. Wahrscheinlichkeitstheorie und Verw. Gebiete 3 (1964), 211–226.

DARTMOUTH COLLEGE

---

## A RELATION BETWEEN A THEOREM OF BOHR AND SIDON SETS

BY DANIEL RIDER<sup>1</sup>

Communicated by W. Rudin, December 17, 1965

1. **Introduction.** In 1913, Bohr [1] proved the following theorem for Dirichlet series: if

$$(1) \quad f(\sigma + it) = \sum_{n=1}^{\infty} c(n)n^{-\sigma-it}$$

and if  $|f(\sigma + it)| \leq 1$  for all  $\sigma > 0$ , then

$$(2) \quad \sum_p |c(p)| \leq 1,$$

the sum in (2) extending over all primes.

A set of positive integers  $E$  will be called a *Bohr set* if there is a finite constant  $B$  such that for every function  $f$  as in (1)

$$(3) \quad \sum_{n \in E} |c(n)| \leq B.$$

---

<sup>1</sup> This research was supported in part by Air Force Office of Scientific Research Grant AF-AOSR 335-63.

It is easily seen that  $E$  is a Bohr set if and only if for every finite sum  $f(t) = \sum c(n)n^{-it}$

$$(4) \quad \sum_{n \in E} |c(n)| \leq B \sup_{-\infty < t < \infty} |f(t)|.$$

Let  $G$  be a compact Abelian group and  $E$  a subset of its dual group  $\Gamma$ . An  $E$ -polynomial is a trigonometric polynomial,  $F$ , such that  $\hat{F}(\gamma) = 0$  for  $\gamma \notin E$  where

$$\hat{F}(\gamma) = \int_G F(x)\gamma(-x) dx, \quad \gamma \in \Gamma.$$

Here  $dx$  is normalized Haar measure on  $G$ .  $E$  is called a *Sidon set* if there is a finite constant  $B$  such that

$$(5) \quad \sum_{\gamma \in E} |\hat{F}(\gamma)| \leq B \sup_{x \in G} |F(x)| = B \|F\|^\infty$$

for every  $E$ -polynomial  $F$ .

Let  $T^\omega$  be the direct product of a countably infinite collection of circles.  $T^\omega$  is a compact Abelian group with dual group  $Z^\omega$ . Each  $\gamma \in Z^\omega$  is given by a sequence of integers  $\{\alpha_k\}$  where only a finite number of the  $\alpha_k$  are not zero.  $M(T^\omega)$  is the space of regular Borel measures,  $\mu$ , on  $T^\omega$  with finite total variation  $\|\mu\|$ .  $\hat{\mu}$  is the Fourier-Stieltjes transform of  $\mu$ .

In this note we give a characterization of Bohr sets in terms of Sidon sets in  $Z^\omega$  and certain measures on  $T^\omega$ . It is then possible to obtain a sufficient arithmetic condition for Bohr sets.

**2. The relation between Bohr sets and Sidon sets.**  $P$  will denote the positive cone of  $Z^\omega$ . Let  $p_1, p_2, \dots$  be the primes. If  $n$  is an integer and  $n = \prod p_j^{\alpha_j}$ , then we associate  $n$  with the element  $\gamma_n = (\alpha_1, \alpha_2, \dots)$  of  $P$ . For a set of positive integers  $E$ ,  $\hat{E} = \{\gamma_n : n \in E\}$ .

To a function  $f(t) = \sum c(n)n^{-it}$  we associate the function  $F(x) = \sum c(n)\gamma_n(x)$  on  $T^\omega$ . Bohr noticed the following: if  $\phi: (-\infty, \infty) \rightarrow T^\omega$  by

$$(6) \quad \phi(t) = (\exp(-it \log p_1), \exp(-it \log p_2), \dots)$$

then  $\gamma_n(\phi(t)) = n^{-it}$  so that  $F(\phi(t)) = f(t)$ . Now since  $\{\log p_j\}$  is linearly independent over the integers,  $\phi(-\infty, \infty)$  is dense in  $T^\omega$ . Thus

$$(7) \quad \|F\|_\infty = \sup_{-\infty < t < \infty} |f(t)|.$$

THEOREM. *A set of positive integers  $E$  is a Bohr set if and only if*

- (a)  $\hat{E}$  is a Sidon set in  $T^\omega$ , and
- (b) there is a measure  $\mu \in M(T^\omega)$  such that

$$(8) \quad \begin{aligned} \hat{\mu}(\gamma) &= 1 && \text{if } \gamma \in \hat{E}, \\ &= 0 && \text{if } \gamma \in P - \hat{E}. \end{aligned}$$

PROOF. Let  $F(x) = \sum \hat{F}(\gamma_n)\gamma_n(x)$  be a  $P$ -polynomial and let  $f(t) = \sum \hat{F}(\gamma_n)n^{-it}$ .

If  $E$  is a Bohr set then by (7)

$$(9) \quad \sum_{\gamma \in \hat{E}} |\hat{F}(\gamma)| = \sum_{n \in E} |\hat{F}(\gamma_n)| \leq B \sup_{-\infty < t < \infty} |f(t)| = B\|F\|_\infty.$$

Thus if  $b$  is a function on  $\hat{E}$  and  $|b(\gamma)| \leq 1$  then  $L(F) = \sum_{\gamma \in \hat{E}} b(\gamma)\hat{F}(\gamma)$  is a bounded linear functional on the  $P$ -polynomials with norm at most  $B$ . By the Hahn-Banach and Riesz representation theorems there is a measure  $\mu \in M(T^\omega)$  with

$$\begin{aligned} \hat{\mu}(\gamma) &= b(\gamma), && \gamma \in \hat{E}, \\ &= 0, && \gamma \in P - \hat{E}. \end{aligned}$$

By [3, Theorem 5.7.3],  $\hat{E}$  is a Sidon set; by taking  $b \equiv 1$  we obtain the measure for (b).

Conversely suppose (a) and (b) are true for  $E$  and let  $f(t) = \sum c(n)n^{-it}$  be a finite sum. By (a) and the proof of [3, Theorem 5.7.3] there is  $\nu \in M(T^\omega)$  with  $\|\nu\| \leq B$  ( $B$  depends only on  $E$ ) and  $c(n)\hat{\nu}(\gamma_n) = |c(n)|$  for  $n \in E$ . Let  $\mu$  be as in (b) and  $*$  denote ordinary convolution. Then

$$\begin{aligned} \sum_{n \in E} |c(n)| &= \mu * \nu * \sum c(n)\gamma_n(0) \\ &\leq B\|\mu\|\|\nu\| \|F\|_\infty \\ &\leq B' \sup_{-\infty < t < \infty} |f(t)|. \end{aligned}$$

COROLLARY. *Let  $E = \{n_1, n_2, \dots\}$  be a set of positive integers satisfying*

- (c)  $\{\log n_j\}$  are linearly independent over the integers, and
- (d) if  $n$  is a positive integer,  $\{\beta_j\}$  is a collection of integers,  $\sum \beta_j = 1$ , and  $n = \prod n_j^{\beta_j}$  then  $n \in E$ .

Then  $E$  is a Bohr set.

PROOF. It follows from (c) that if  $k_1 < k_2 < \dots < k_s$  then  $0 \neq \pm \gamma_{n_{k_1}} \pm \gamma_{n_{k_2}} \pm \dots \pm \gamma_{n_{k_s}}$ . Thus by [2, Theorem 1.5],  $\hat{E}$  is a Sidon set.

Let  $H = \{\gamma \in Z^\omega : \gamma = \sum \beta_j \gamma_{n_j}, \beta_j \text{ integers, } \sum \beta_j = 1\}$ .  $H$  is a coset of a subgroup of  $Z^\omega$  and by (d)  $\hat{E} = H' \cap P$ . By [3, p. 60] there is  $\mu \in M(T^\omega)$  such that  $\hat{\mu}$  is the characteristic function of  $H'$ .  $\mu$  satisfies condition (b) of the theorem.

**3. Examples.** The corollary shows that there are Bohr sets which are not the finite union of sets with pairwise relatively prime elements. For example,  $p_1 p_2, p_1 p_3, p_4 p_5, p_4 p_6, p_4 p_7, p_8 p_9, \dots$ . It is known [3, p. 126] that every infinite subset of a discrete group contains an infinite Sidon subset. However this is not true of Bohr sets.

EXAMPLE. Let  $F = \{n_j = (p_1 p_2 \cdots p_j)^i\}$ . Then  $F$  contains no infinite Bohr subset.

In fact  $F$  contains no infinite subset for which there is a measure satisfying (8). For suppose  $E = \{n_{j_1}, n_{j_2}, \dots\}$  and  $\hat{\mu}$  satisfies (8). Let  $\mu_k$  be the translation of  $\mu$  such that

$$(10) \quad \hat{\mu}_k(\gamma) = \hat{\mu}(\gamma + \gamma_{n_{j_k}}).$$

$\{\mu_k\}$  has a weak star convergent subsequence to a measure  $\nu \in M(T^\omega)$  which by a lemma of Helson [3, p. 66] must be singular with respect to Haar measure.

But this is impossible since it is easily seen that

$$\begin{aligned} \hat{\nu}(\gamma) &= 1 && \text{if } \gamma = 0, \\ &= 0 && \text{if } \gamma \neq 0 \end{aligned}$$

so that  $\nu$  must be the Haar measure.

This example also shows that the corollary is false without (d).

#### BIBLIOGRAPHY

1. H. Bohr, *Über die Bedeutung der Potenzreihen unendlich vieler Variablen in der Theorie der Dirichletschen Reihen*, Nachr. Akad. Wiss. Göttingen Math.-Phys. Kl. (1913), 441-448.
2. D. Rider, *Gap series on groups and spheres*, Canad. J. Math. (to appear).
3. W. Rudin, *Fourier analysis on groups*, Interscience, New York, 1962.

UNIVERSITY OF WISCONSIN AND  
MASSACHUSETTS INSTITUTE OF TECHNOLOGY