

GLOBAL CONTINUOUS SOLUTIONS OF HYPERBOLIC SYSTEMS OF QUASI-LINEAR EQUATIONS¹

BY J. L. JOHNSON

Communicated by Jürgen Moser, May 16, 1967

Recently there have appeared a number of results on global solutions of the Cauchy problem for hyperbolic systems of quasi-linear equations [2], [3], [4], [5]. These solutions are in general discontinuous. In certain cases, however, such as the interaction of two rarefaction waves in gas dynamics, it is known that the Cauchy problem has a global continuous solution [1, pp. 191–197]. In this announcement we outline a proof that a global continuous solution exists and is unique for a two-dimensional system provided the Riemann invariants associated with the initial data satisfy certain monotonicity and continuity conditions.

Let $\lambda^+(r, s)$, $\lambda^-(r, s)$ be C^1 real-valued functions on a domain $D \subset R_2$, with

$$(1) \quad \lambda^+(r, s) > \lambda^-(r, s), \quad \partial\lambda^+(r, s)/\partial r > 0, \quad \partial\lambda^-(r, s)/\partial s > 0$$

for $(r, s) \in D$. Consider the two-dimensional system of quasi-linear equations in Riemann invariant form

$$(2) \quad r_t + \lambda^+(r, s)r_x = 0, \quad s_t + \lambda^-(r, s)s_x = 0$$

where $r(t, x)$ and $s(t, x)$ are real-valued functions of two scalar variables. We seek a solution of the Cauchy problem in the halfplane $\{(t, x) \in R_2: t \geq 0\}$ with initial conditions

$$(3) \quad r(0, x) = r^0(x), \quad s(0, x) = s^0(x), \quad -\infty < x < +\infty.$$

Let $G_T = \{(t, x) \in R_2: 0 \leq t < T\}$ for $0 < T \leq +\infty$. A pair of Lipschitz continuous functions $(r(t, x), s(t, x))$, $(t, x) \in G_T$, is called a Lipschitz continuous solution of the Cauchy problem (2), (3) if $r(t, x)$ is constant on the integral curves

$$(4) \quad x'(t) = \lambda^+(r(t, x), s(t, x)),$$

$s(t, x)$ is constant on the integral curves

$$(5) \quad x'(t) = \lambda^-(r(t, x), s(t, x)),$$

¹ These results are part of the author's Ph.D. thesis written under Joel A. Smoller at the University of Michigan.

and $r(0, x)$, $s(0, x)$ satisfy the initial conditions (3). The pair $(r(t, x), s(t, x))$ is called a global Lipschitz continuous solution of (2), (3) if the functions are defined and Lipschitz continuous on G_∞ .

THEOREM 1. *If $r^0(x)$, $s^0(x)$, $-\infty < x < +\infty$, are bounded, Lipschitz continuous, and nondecreasing, satisfying*

$$(6) \quad [r^0(-\infty), r^0(+\infty)] \times [s^0(-\infty), s^0(+\infty)] \subset D,$$

then the Cauchy problem (2), (3) with initial functions $r^0(x)$, $s^0(x)$, $-\infty < x < +\infty$, has a global Lipschitz continuous solution which takes its values in the rectangle (6).

OUTLINE OF PROOF FOR THEOREM 1. For each finite subset A of R_1 we construct an approximate solution $(r(t, x; A), s(t, x; A))$, $(t, x) \in G_\infty$, with the property that $r(t, x; A)$ is constant on curves of the form (4) and $s(t, x; A)$ is constant on a finite number of curves of the form (5). Using condition (1) and the assumed properties of the initial functions, we show that $r(t, x; A)$ is Lipschitz continuous in G_∞ with Lipschitz constant independent of t, x , and A . If $\{B_n: n=1, 2, \dots\}$ is an increasing sequence of finite sets whose union is dense in R_1 , then by the Ascoli theorem the sequence of functions $r(t, x; B_n)$ contains a subsequence converging to a Lipschitz continuous function $r(t, x)$. Having this function, we construct $s(t, x)$, $(t, x) \in G_\infty$, so that the pair $(r(t, x), s(t, x))$, $(t, x) \in G_\infty$, is a global Lipschitz continuous solution of (2), (3).

We have also obtained the following result regarding the dependence of Lipschitz continuous solutions on initial data.

THEOREM 2. *Let $r_i^0(x)$, $s_i^0(x)$, $-\infty < x < +\infty$, $i=1, 2$, be bounded real-valued functions with*

$$a_i \leq r_i^0(x) \leq b_i, \quad c_i \leq s_i^0(x) \leq d_i, \quad -\infty < x < +\infty,$$

and suppose that $[a_i, b_i] \times [c_i, d_i] \subset D$, $i=1, 2$. Let

$$m = \sup_{-\infty < x < +\infty} (|r_1^0(x) - r_2^0(x)| + |s_1^0(x) - s_2^0(x)|).$$

If $(r_i(t, x), s_i(t, x))$, $(t, x) \in G_T$, is a Lipschitz continuous solution of the Cauchy problem for the equations (2) with initial vector $(r_i^0(x), s_i^0(x))$, $-\infty < x < +\infty$, $i=1, 2$, then there is a constant $L(T)$ such that

$$\sup_{(t, x) \in G_T} (|r_1(t, x) - r_2(t, x)| + |s_1(t, x) - s_2(t, x)|) \leq mL(T).$$

It follows easily from this that Lipschitz continuous solutions are unique.

Theorems 1 and 2, together with results of Lax [3], can be applied to the conservation law

$$(7) \quad u_t + (p(v))_x = 0, \quad v_t - u_x = 0$$

to yield the following corollaries.

COROLLARY 1. *Let $p(v) \in C^2$ on the halfline ($v > 0$) with $p'(v) < 0$, $p''(v) > 0$ and $\int_1^\infty [-p'(v)]^{1/2} dv = \infty$. If the functions $u^0(x)$, $v^0(x)$, $-\infty < x < +\infty$, are bounded and Lipschitz continuous, with $v^0(x)$ positive and bounded away from 0, and satisfy*

$$(8) \quad u^0(x_2) - u^0(x_1) \cong \left| \int_{v^0(x_1)}^{v^0(x_2)} [-p'(v)]^{1/2} dv \right| \quad \text{for } x_2 > x_1,$$

then the Cauchy problem for the equations (7) with initial vector $(u^0(x), v^0(x))$ has a unique Lipschitz continuous weak solution.

COROLLARY 2. *Let $p(v)$ be as in Corollary 1, and let $u^0(x)$, $v^0(x)$, $-\infty < x < +\infty$, be bounded and piecewise constant real-valued functions, with $v^0(x)$ positive and bounded away from 0, which satisfy (8). If the set A of discontinuities of the vector function $(u^0(x), v^0(x))$ has the property*

$$\inf\{|a - b| : a, b \in A, a \neq b\} > 0,$$

then the Cauchy problem for the system (7) with initial vector $(u^0(x), v^0(x))$ has a solution which is Lipschitz continuous in each of the sets $\{(t, x) \in G_\infty : t > t_0\}$, for $t_0 > 0$.

Corollary 2 provides a solution for the interaction of simple waves centered on the line ($t = 0$).

REFERENCES

1. R. Courant and K. O. Friedrichs, *Supersonic flow and shock waves*, Interscience, New York, 1948.
2. James Glimm, *Solutions in the large for nonlinear hyperbolic systems of equations*, Comm. Pure Appl. Math. **18** (1965), 697-715.
3. P. D. Lax, *Hyperbolic systems of conservation laws*. II, Comm. Pure Appl. Math. **10** (1957), 537-566.
4. B. L. Roždestvenskiĭ, *Construction of discontinuous solutions to systems of quasi-linear equations*. I, II, Ž. Vyčisl. Mat. i Mat. Fiz. **2** (1962), 1019-1043; **3** (1963), 79-98.
5. Zhang Tong and Guo Yu-fa, *A class of initial value problems for systems of aerodynamic equations*, Acta Math. Sinica **15** (1965), no. 3, 386-396; English transl., Chinese Math. **7** (1965), 90-101.