

THE OBSTRUCTION TO FIBERING A MANIFOLD OVER A CIRCLE

BY F. T. FARRELL¹

Communicated by S. Smale, April 6, 1967

1. Introduction. In [1], Stallings considers the following question. When does a 3-manifold fiber a circle? Browder and Levine generalized Stallings' result to differentiable and piecewise linear manifolds M of dimension greater than five under the restriction that $\pi_1(M) \cong \mathbb{Z}$. Their theorem is purely *homotopic* in nature. That is if $h: M' \rightarrow M$ is a homotopy equivalence and $f: M \rightarrow S^1$ is a smooth fiber map then there always exists a smooth fiber map $f': M' \rightarrow S^1$ such that f' is homotopic to $f \circ h$.

This result is *false* if we drop their restriction on the fundamental group. In particular let N be the cartesian product of a 3-dimensional lens space L with fundamental group Z_p^2 and the torus T^{n-3} where $n \geq 5$. Let $M = N \times S^1$ and $f: M \rightarrow S^1$ denote projection onto the second factor. Then there exists a manifold M' and a homotopy equivalence $h: M' \rightarrow M$ (in fact we may take M' to be h -cobordant to M) such that a smooth fiber map $f': M' \rightarrow S^1$ homotopic to $f \circ h$ cannot exist. This example is based on recent deep results of Bass and Murthy [3] concerning the structure of the projective class group. In a joint paper with W. C. Hsiang [4] we use this example to construct an *h-cobordism* (W, M, M') which is not homeomorphic to $M \times [0, 1]$.

In this paper we will state necessary and sufficient conditions, in terms of a new obstruction theory, for a manifold M^n ($n \geq 6$) to fiber a circle. No restrictions will be placed on the fundamental group of M . We will always work in the differential category, but the corresponding theorem is also true in the piecewise-linear category.

2. Description of obstructions. Let M^n be a closed connected smooth manifold with $n \geq 6$. Let $f: M \rightarrow S^1$ be a continuous map. (Recall that the homotopy class of f is an element of $H^1(M, \mathbb{Z})$.) We will state three properties about f which are necessary and sufficient to guarantee the existence of a smooth fiber map $\tilde{f}: M \rightarrow S^1$ homotopic to f . For convenience we restrict our attention to maps f such that $f_{\#}: \pi_1(M) \rightarrow \pi_1(S^1)$ is onto. (This is equivalent to considering only indi-

¹ The author holds a National Science Foundation Graduate Fellowship. This note is a summary of a doctoral dissertation written under the direction of Professor Wu-Chung Hsiang and submitted to Yale University in partial fulfillment of the requirements for the Ph.D. degree.

visible elements in $H^1(M, Z)$.) This corresponds geometrically to considering fibrations with connected fiber. Let $G = \ker f_{\#}$ and X denote the covering space of M corresponding to G . If \tilde{f} exists then it is clear that the fiber of \tilde{f} is homotopically equivalent to X . But the fiber of \tilde{f} would be a closed smooth manifold. In particular it would be a finite C.W. complex. Hence we obtain

CONDITION 1. X is dominated by a finite C.W. complex.

Let (N^{n-1}, ν) be a framed submanifold of M which represents f under the Pontrjagin-Thom construction. Let M_N denote the manifold obtained by "cutting" M along N . Then ∂M_N consists of two copies of N which we label N' and N'' . (See Figure 1.)



FIGURE 1

M_N is a cobordism from N' to N'' . The pair (N, ν) is called a splitting of M . Let $1 < s < n - 2$, s an integer. When Condition 1 holds it is always possible to find a splitting (N, ν) such that (M_N, N') has a handlebody decomposition with handles of only two dimension s and $s + 1$. The proof of this uses essentially the same arguments as in [2]. Note that the existence of a smooth fiber map is equivalent to the existence of a splitting (N, ν) such that M_N is diffeomorphic to $N \times [0, 1]$. Conditions 2 and 3 will guarantee the existence of such a splitting. Condition 2 will hold if and only if there exists a splitting (N, ν) such that (M_N, N', N'') is an h -cobordism. Condition 3 will hold if and only if some such h -cobordism is a product.

We proceed to formulate Condition 2. From the exact sequence $0 \rightarrow G \rightarrow \pi_1(M) \rightarrow Z \rightarrow 0$ we see that $\pi_1(M)$ is a semidirect product of G and Z with respect to an automorphism α of G . (α is only well defined up to an inner automorphism but this is all right for our purposes.) If Condition 1 is satisfied then we can define an element $c(f)$ in an abelian group $C(Z(G), \alpha)$. $c(f)$ has the following property: $c(f) = 0$ if and only if there exists a splitting (N, ν) such that (M_N, N', N'') is an

h -cobordism. The proof of this fact is quite long and relies heavily on handle body theory.

CONDITION 2. $c(f) = 0$.

If Condition 2 is satisfied then $\tau(M_N, N') \in \text{Wh}(G)$ is defined. But it may be possible to have a second splitting (N_1, v_1) such that (M_{N_1}, N'_1, N''_1) is an h -cobordism and $\tau(M_{N_1}, N'_1) \neq \tau(M_N, N')$. Let α_* denote the automorphism of $\text{Wh}(G)$ induced by α (see [5]). Let $\tau(f)$ be the image of $\tau(M_N, N')$ in the group $\text{Wh}(G) / \{x - \alpha_*(x) \mid x \in \text{Wh}(G)\}$ under the quotient homomorphism. We can show that $\tau(f)$ is well defined. (i.e. $\tau(f)$ is independent of the splitting (N, v)). Also $\tau(f) = 0$ if and only if there exists a splitting (N, v) such that $\tau(M_N, N') = 0$. The proof of this fact makes use of Stallings' realizability theorem for h -cobordisms (see [6]). But the s -cobordism theorem of Barden-Mazur-Smale states that M_N is diffeomorphic to $N \times [0, 1]$ if and only if $\tau(M_N, N') = 0$. Therefore

CONDITION 3. $\tau(f) = 0$.

Summarizing we have the following theorem.

THEOREM. *There exists a smooth fiber map $\tilde{f}: M \rightarrow S^1$ homotopic to f if and only if*

1. X is dominated by a finite C.W. complex,
2. $c(f) = 0$,
3. $\tau(f) = 0$.

NOTE. There exists a version of this theorem for manifolds with boundary where the boundary already fibers a circle.

3. Properties of $C(R, \alpha)$. If R is a ring with identity and α is an automorphism of R then by a Grothendieck construction we can define an abelian group $C(R, \alpha)$. $\tilde{K}_0(R)$ is a direct summand of $C(R, \alpha)$. Denote by $\tilde{C}(R, \alpha)$ the complementary summand. Write $c(f) = \sigma(f) + \tilde{c}(f)$ where $\sigma(f) \in \tilde{K}_0(R)$ and $\tilde{c}(f) \in \tilde{C}(R, \alpha)$. Then $\sigma(f)$ is the Novikov-Siebenmann-Wall obstruction to X splitting differentiably as a cartesian product $N \times R$.

R is called regular if it is Noetherian and every finitely generated R module has a resolution of finite length by projective R modules. If R is regular then $\tilde{C}(R, \alpha) = 0$. But this is not the general situation since Bass and Murthy have shown that $\tilde{C}(Z(G), \text{id}) \neq 0$ for certain finitely generated abelian groups G . A particular example is $G = Z \oplus Z \oplus Z_4$.

As an example of the fibering theorem consider the case where $G = Z^n$. Then $Z(G)$ is regular and hence $\tilde{C}(Z(G), \alpha) = 0$. Also it is

known that $\tilde{K}_0(Z(G))=0$ and $\text{Wh}(G)=0$ (see [5]). Therefore Conditions 2 and 3 become vacuous. Also observe that Condition 1 is only a homotopy theoretic condition. In particular if M and M' are homotopically equivalent manifolds such that $\pi_1(M)$ is free abelian then M fibers a circle if and only if M' fibers a circle.

BIBLIOGRAPHY

1. J. Stallings, *On fibering certain 3-manifolds*, Topology of 3-manifolds, Prentice-Hall, Englewood Cliffs, N. J., 1962, pp. 95–100.
2. W. Browder and J. Levine, *Fibering manifolds over a circle*, Comment. Math. Helv. **40** (1966), 153–160.
3. H. Bass and M. P. Murthy, *Grothendieck groups and Picard groups of abelian group rings*, (to appear).
4. F. T. Farrell and W. C. Hsiang, *H-cobordant manifolds are not necessarily homeomorphic*, Bull. Amer. Math. Soc. **73** (1967), 741–756.
5. J. Milnor, *Whitehead torsion*, Bull. Amer. Math. Soc. **72** (1966), 358–426.
6. J. Stallings, *On infinite processes leading to differentiability in the complement of a point*, Differential and Combinatorial Topology, Princeton Univ. Press, Princeton, N. J.
7. F. T. Farrell, Ph.D. Thesis, Yale University, New Haven, Conn.,
8. L. Siebenmann, *The obstruction to finding a boundary for an open manifold of dimension greater than five*, Ph.D. Thesis, Princeton University, Princeton, N. J.
9. C. T. C. Wall, *Finiteness conditions for C. W. complexes*, Ann. of Math. **81** (1965), 56–69.

YALE UNIVERSITY