

# COHOMOLOGY OF CERTAIN STEINBERG GROUPS

BY DAVID HERTZIG

Communicated by D. Buchsbaum, March 25, 1968

In [3] Steinberg considers the relations satisfied by generators of the Chevalley groups and defines certain abstract groups  $\Delta$  and  $\Gamma$  via presentation. Let  $\Sigma$  be a root system of a simple complex Lie algebra  $\mathfrak{G}_{\mathbb{C}}$  and let  $K$  be a field of characteristic  $p \geq 0$ . We consider a set of generators  $x_r(t)$  ( $r \in \Sigma, t \in K$ ) and the relations

$$(A) \quad x_r(t)x_r(u) = x_r(t+u) \quad (r \in \Sigma; t, u \in K),$$

$$(B) \quad x_r(t)x_s(u)x_r(t)^{-1} = x_s(u) \prod x_{ir+js}(C_{ij;rs}t^i u^j) \\ (r, s \in \Sigma, r+s \neq 0; t, u \in K).$$

The product in (B) is over all integers  $i, j \geq 1$  for which  $ir+js \in \Sigma$ , taken in lexicographic order. The  $C_{ij;rs}$  are certain integers depending only on the structure of  $\mathfrak{G}_{\mathbb{C}}$  (cf. [1]). Steinberg defines  $w_r(t) = x_r(t)x_{-r}(-t^{-1})x_r(t)$  and  $h_r(t) = w_r(t)w_r(-1)$  ( $r \in \Sigma; t \in K^*$ ) and considers also the relations

$$(B') \quad w_r(t)x_r(u)w_r(t^{-1}) = x_{-r}(-t^{-2}u) \quad (r \in \Sigma; t \in K^*, u \in K),$$

$$(C) \quad h_r(t)h_r(u) = h_r(tu) \quad (r \in \Sigma; t, u \in K^*).$$

The *Steinberg group*  $\Delta$  is the abstract group generated by the symbols  $x_r(t)$  ( $r \in \Sigma; t \in K$ ) subject to the relations (A) and (B) if the rank of  $\Sigma$  is  $> 1$ , to the relations (A) and (B') if the rank of  $\Sigma = 1$ . The *Steinberg group*  $\Gamma$  is the abstract group with the same generators as  $\Delta$  subject to the relations of  $\Delta$  and in addition subject to the relations (C).

In [1] Chevalley constructs a corresponding Lie algebra  $\mathfrak{G}$  over the field  $K$  and there is a natural action of the Steinberg groups on  $\mathfrak{G}$ . One is then led to consideration of the cohomology  $H^1(\Delta, \mathfrak{G})$  and  $H^1(\Gamma, \mathfrak{G})$ .

The author has developed a technique for computation of such cohomology (cf. [2]). This is applied successfully to obtain the following results. Proofs will appear elsewhere.

We denote  $\mathfrak{D}(K)$  the module of derivations of  $K$ . In the case of characteristic  $p=2$  we denote  $\mathfrak{L}(K)$  the  $K^2$ -linear transformations  $L$  of  $K$  such that  $L(1)=0$ . Since  $p=2$ ,  $\mathfrak{D}(K) \subset \mathfrak{L}(K)$ , but in general  $\mathfrak{D}(K) \neq \mathfrak{L}(K)$ .

**THEOREM 1.**  $H^1(\Delta, \mathfrak{G}) = H^1(\Gamma, \mathfrak{G}) \cong \mathfrak{D}(K)$  in the following cases:

- (i) type  $A_1$ ,  $p \neq 2$  and  $K \neq F_5$ ;

- (ii) type  $A_n$  ( $n \geq 2$ ),  $p \nmid n+1$ ;
- (iii) type  $D_n$  ( $n \geq 4$ ),  $p \neq 2$ ;
- (iv) type  $E_6$ ,  $p \neq 3$ ;
- (v) type  $E_7$ ,  $p \neq 2$ ;
- (vi) type  $E_8$ ;
- (vii) type  $F_4$ .

THEOREM 2.  $H^1(\Delta, \mathcal{G}) = H^1(\Gamma, \mathcal{G}) \cong K \oplus \mathfrak{D}(K)$  in the following cases:

- (i) type  $A_n$  ( $n \geq 2$ ),  $p \mid n+1$ ;
- (ii) type  $D_n$  ( $n \geq 4$ ,  $n$  odd),  $p = 2$ ;
- (iii) type  $E_6$ ,  $p = 3$ ;
- (iv) type  $E_7$ ,  $p = 2$ .

THEOREM 3.  $H^1(\Delta, \mathcal{G}) = H^1(\Gamma, \mathcal{G}) \cong K \oplus K \oplus \mathfrak{D}(K)$  in the case: type  $D_n$  ( $n \geq 4$ ,  $n$  even),  $p = 2$ .

THEOREM 4.  $H^1(\Delta, \mathcal{G}) = H^1(\Gamma, \mathcal{G}) \cong K$  in the case: type  $A_1$ ,  $K = F_6$ .

THEOREM 5.  $H^1(\Delta, \mathcal{G}) \cong K \oplus \mathfrak{L}(K)$ ,  $H^1(\Gamma, \mathcal{G}) \cong K \oplus \mathfrak{D}(K)$  in the case: type  $A_1$ ,  $p = 2$ .

#### REFERENCES

1. C. Chevalley, *Sur certains groupes simples*, Tôhoku Math. J. (2) 7 (1956), 14–66.
2. D. Hertz, *Cohomology of algebraic groups*, J. Algebra 6 (1967), 317–334.
3. R. Steinberg, *Générateurs, relations et revêtements de groupes algébriques*, Colloque Théorie des Groupes Algébriques, Brussels Libraire Universitaire Louvain and Gauthier-Villars, Paris 1962, pp. 113–127.

PURDUE UNIVERSITY, LAFAYETTE, INDIANA 47907