

A NORMAL SPACE X FOR WHICH $X \times I$ IS NOT NORMAL

BY MARY ELLEN RUDIN

Communicated by E. A. Michael, September 28, 1970

In [1] C. Dowker gave a number of interesting characterizations of normal Hausdorff spaces whose cartesian product with the closed unit interval is not normal. Thus, such a space is often called a Dowker space; a Dowker space X will be described below. It was previously known only that the existence of a Dowker space is consistent with the usual axioms of set theory [2], [3]. The proof that X is a Dowker space uses no set theoretic assumptions beyond the axiom of choice.

We use the convention that an ordinal λ is the set of all ordinals less than λ . An ordinal γ is said to be *cofinal* with λ if there is a subset Γ of λ order isomorphic with γ such that $\alpha \in \lambda$ implies $\alpha \leq \beta$ for some $\beta \in \Gamma$; let $\text{cf}(\lambda)$ be the smallest ordinal cofinal with λ .

Define $F = \{f: \omega_0 \rightarrow \omega_\omega \mid \forall n \in \omega_0, f(n) \leq \omega_n + 1\}$.

Define $X = \{f \in F \mid \exists k \in \omega_0 \text{ such that } \forall n \in \omega_0, \omega_0 < \text{cf}(f(n)) < \omega_k\}$.

For f and g in F , define $U_{fg} = \{h \in X \mid \forall n \in \omega_0, f(n) < h(n) \leq g(n)\}$. Then topologize X by using the set of all U_{fg} for f and g in F as a basis for the topology. The resulting space is a collectionwise normal Dowker space.

REFERENCES

1. C. H. Dowker, *On countably paracompact spaces*, Canad. J. Math. **3** (1951), 219–224. MR **13**, 264.
2. M. E. Rudin, *Countable paracompactness and Souslin's problem*, Canad. J. Math. **7** (1953), 543–547. MR **17**, 391.
3. S. Tennenbaum, *Souslin's problem*, Proc. Nat. Acad. Sci. U. S. A. **59** (1968), 60–63. MR **37** #55.

UNIVERSITY OF WISCONSIN, MADISON, WISCONSIN 53706

AMS 1969 subject classifications. Primary 5420, 0415, 0430.

Key words and phrases. Dowker space, cartesian product, normal, binormal, collectionwise normal, countably paracompact.