

## WEAKLY CONTINUOUS ACCRETIVE OPERATORS

BY W. E. FITZGIBBON

Communicated by Fred Brauer, September 28, 1972

We shall be concerned with the autonomous differential equation

$$(1.1) \quad u'(t) + Au(t) = 0, \quad u(0) = x,$$

where  $A$  is a weakly continuous possibly nonlinear operator mapping a reflexive Banach space  $X$  to itself. Recently S. Chow and J. D. Schuur [2] have considered existence theory for ordinary differential equations involving weakly continuous operators on separable, reflexive Banach spaces.

We now make clear our notion of strong solutions to (1.1).

**DEFINITION 1.2.** A function  $u: [0, T) \rightarrow X$  is said to be a *strong solution* to the Cauchy problem

$$u'(t) + Au(t) = 0, \quad u(0) = x,$$

provided that  $u$  is Lipschitz continuous on each compact subset of  $[0, T)$ ,  $u(0) = x$ ,  $u$  is strongly differentiable almost everywhere and  $u'(t) + Au(t) = 0$  for a.e.  $t \in [0, T)$ .

By employing a variant of the Peano method we provide local solution to (1.1).

**LEMMA 1.3.** *Let  $X$  be a reflexive Banach space and suppose that  $A$  is a weakly continuous operator with  $D(A) = X$ . Then there is a finite interval  $[0, T)$  such that the Cauchy problem (1.1) has a strong solution on  $[0, T)$ .*

**DEFINITION 1.4.** An operator  $A$  is said to be *accretive* provided that  $\|x + \lambda Ax - (y + \lambda Ay)\| \geq \|x - y\|$  for all  $\lambda \geq 0$  and  $x, y \in D(A)$ . T. Kato [5] has shown that this definition is equivalent to the statement that  $\operatorname{Re}(Ax - Ay, f) \geq 0$  for some  $f \in F(x - y)$  where  $F$  is the duality map from  $X$  to  $X^*$ .

If we require that the operator  $A$  be accretive we are able to extend the local solution of Lemma 1.3 to a global solution.

**THEOREM 1.5.** *Let  $X$  be a reflexive Banach space and suppose that  $A$  is a weakly continuous accretive operator with  $D(A) = X$ . Then the Cauchy problem (1.1) has a unique strong global solution on  $[0, \infty)$ .*

---

AMS (MOS) subject classifications (1970). Primary 47H15, 34H05; Secondary 47B44, 47D05.

Key words and phrases. Accretive, weakly continuous, semigroup of nonexpansive non-linear operators.

If we set  $u(t) = T(t)x$  we obtain a semigroup of nonlinear nonexpansive operators  $\{T(t): t \geq 0\}$  which map  $X$  to  $X$ . We can say that  $\{T(t): t \geq 0\}$  is the semigroup associated with  $A$ . The next theorem provides an exponential representation for  $\{T(t): t \geq 0\}$ .

**THEOREM 1.6.** *Let  $A$  and  $X$  satisfy the conditions of Theorem 1.5. Then the operator  $A$  is  $m$ -accretive, i.e.,  $R(I + \lambda A) = X$  for all  $\lambda \geq 0$ . If  $\{T(t): t \geq 0\}$  is a semigroup associated with  $A$  then  $T(t)$  may be represented as the pointwise limit*

$$T(t)x = \lim_{n \rightarrow \infty} (I + t/nA)^{-n}x.$$

Moreover, for each fixed  $t_0 > 0$ , the operator  $T(t_0)$  is weakly continuous.

The  $m$ -accretiveness of  $A$  is obtained by considering the equation  $u'(t) + A'u(t) = 0$  where  $A' = A + I$ . Once the  $m$ -accretiveness of  $A$  has been established the exponential representation of  $\{T(t): t \geq 0\}$  follows immediately from a theorem of M. Crandall and T. Liggett [1]. The fact that  $T(t_0)$  is weakly continuous is obtained by showing that  $(I + \lambda A)^{-1}$  is weakly continuous for all  $\lambda \geq 0$  and employing estimates of Crandall and Liggett. The foregoing results may be applied to the rest point theory developed by C. Yen [10].

#### REFERENCES

1. M. Crandall and T. Liggett, *Generation of semigroups of nonlinear transformations on general Banach spaces*, Amer. J. Math. **93** (1971), 265–298.
2. S. Chow and J. D. Schuur, *An existence theorem for ordinary differential equations in Banach spaces*, Bull. Amer. Math. Soc. **77** (1971), 1018–1020.
3. E. Hille and R. S. Phillips, *Functional analysis and semi-groups*, rev. ed., Amer. Math. Soc. Colloq. Publ., vol. 31, Amer. Math. Soc., Providence, R.I., 1957. MR **19**, 664.
4. T. Kato, *Accretive operators and nonlinear evolution equations in Banach spaces*, Proc. Sympos. Pure Math., vol. 18, part 1, Amer. Math. Soc., Providence, R.I., 1970, pp. 138–161. MR **42** #6663.
5. ———, *Nonlinear semigroups and evolution equations*, J. Math. Soc. Japan **19** (1967), 508–520. MR **37** #1820.
6. R. H. Martin, Jr., *A global existence theorem for autonomous differential equations in a Banach space*, Proc. Amer. Math. Soc. **26** (1970), 307–314. MR **41** #8791.
7. J. W. Neuberger, *An exponential formula for one-parameter semi-groups of nonlinear transformations*, J. Math. Soc. Japan **18** (1966), 154–157. MR **34** #622.
8. S. Ôharu, *Note on the representation of semi-groups of non-linear operators*, Proc. Japan Acad. **42** (1966), 1149–1154. MR **36** #3167.
9. G. F. Webb, *Continuous perturbations of linear accretive operators in Banach spaces*, J. Functional Analysis (to appear).
10. C. Yen, *The convergence, periodicity and rest point behaviour of orbits in nonlinear semigroups of contractions*, Thesis, Vanderbilt University, Nashville, Tenn., 1971.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HOUSTON, HOUSTON, TEXAS 77004